

# UC Davis

## UC Davis Previously Published Works

### Title

Wolbachia Acquisition by Drosophila yakuba-clade Hosts and Transfer of Incompatibility Loci Between Distantly Related Wolbachia

### Permalink

<https://escholarship.org/uc/item/5dc7k6jj>

### Journal

Genetics, 212(4)

### ISSN

0016-6731

### Authors

Cooper, Brandon S  
Vanderpool, Dan  
Conner, William R  
et al.

### Publication Date

2019-08-01

### DOI

10.1534/genetics.119.302349

Peer reviewed

# *Wolbachia* Acquisition by *Drosophila yakuba*-Clade Hosts and Transfer of Incompatibility Loci Between Distantly Related *Wolbachia*

Brandon S. Cooper,<sup>\*1</sup> Dan Vanderpool,<sup>†</sup> William R. Conner,<sup>\*</sup> Daniel R. Matute,<sup>‡</sup> and Michael Turelli<sup>§</sup>

<sup>\*</sup>Division of Biological Sciences, University of Montana, Missoula, Montana 59812, <sup>†</sup>Department of Biology, Indiana University, Bloomington, Indiana 47405, <sup>‡</sup>Biology Department, University of North Carolina at Chapel Hill, North Carolina 27510, and <sup>§</sup>Department of Evolution and Ecology, University of California, Davis, California 95616

ORCID IDs: 0000-0002-8269-7731 (B.S.C.); 0000-0002-6856-5636 (D.V.); 0000-0002-7597-602X (D.R.M.); 0000-0003-1188-9856 (M.T.)

**ABSTRACT** Maternally transmitted *Wolbachia* infect about half of insect species, yet the predominant mode(s) of *Wolbachia* acquisition remains uncertain. Species-specific associations could be old, with *Wolbachia* and hosts codiversifying (*i.e.*, cladogenic acquisition), or relatively young and acquired by horizontal transfer or introgression. The three *Drosophila yakuba*-clade hosts [*D. santomea*, *D. yakuba*, *D. teissieri*] diverged ~3 MYA and currently hybridize on the West African islands Bioko and São Tomé. Each species is polymorphic for nearly identical *Wolbachia* that cause weak cytoplasmic incompatibility (CI)—reduced egg hatch when uninfected females mate with infected males. *D. yakuba*-clade *Wolbachia* are closely related to *wMel*, globally polymorphic in *D. melanogaster*. We use draft *Wolbachia* and mitochondrial genomes to demonstrate that *D. yakuba*-clade phylogenies for *Wolbachia* and mitochondria tend to follow host nuclear phylogenies. However, roughly half of *D. santomea* individuals, sampled both inside and outside of the São Tomé hybrid zone, have introgressed *D. yakuba* mitochondria. Both mitochondria and *Wolbachia* possess far more recent common ancestors than the bulk of the host nuclear genomes, precluding cladogenic *Wolbachia* acquisition. General concordance of *Wolbachia* and mitochondrial phylogenies suggests that horizontal transmission is rare, but varying relative rates of molecular divergence complicate chronogram-based statistical tests. Loci that cause CI in *wMel* are disrupted in *D. yakuba*-clade *Wolbachia*; but a second set of loci predicted to cause CI are located in the same WO prophage region. These alternative CI loci seem to have been acquired horizontally from distantly related *Wolbachia*, with transfer mediated by flanking *Wolbachia*-specific ISWpi1 transposons.

**KEYWORDS** cytoplasmic incompatibility; horizontal gene transfer; introgression; transposable elements; WO phage

**E**NDOSYMBIOTIC *Wolbachia* bacteria infect many arthropods (Bouchon *et al.* 1998; Hilgenboecker *et al.* 2008), including about half of all insect species (Werren and Windsor 2000; Weinert *et al.* 2015). *Wolbachia* often manipulate host reproduction, facilitating spread to high frequencies within host species (Laven 1951; Yen and Barr 1971; Turelli and Hoffmann 1991; Rousset *et al.* 1992; O'Neill *et al.* 1998; Weeks *et al.* 2007; Kriesner *et al.* 2016; Turelli *et al.* 2018). In *Drosophila*, reproductive manipulations include cytoplasmic incompatibility (CI) and male killing

(Hoffmann *et al.* 1986; Hoffmann and Turelli 1997; Hurst and Jiggins 2000). CI reduces the egg hatch of uninfected females mated with *Wolbachia*-infected males, and recent work has demonstrated that WO prophage-associated loci cause CI (Beckmann and Fallon 2013; Beckmann *et al.* 2017, 2019; LePage *et al.* 2017). Although reproductive manipulations are common, some *Wolbachia* show little or no reproductive manipulation (*e.g.*, *wMel* in *Drosophila melanogaster*, Hoffmann 1988; Hoffmann *et al.* 1994; Kriesner *et al.* 2016; *wMau* in *D. mauritiana*, Giordano *et al.* 1995; Meany *et al.* 2019; *wAu* in *D. simulans*, Hoffmann *et al.* 1996; *wSuz* in *D. sukukii* and *wSp* in *D. subpulchrella*, Hamm *et al.* 2014; Cattel *et al.* 2018). These *Wolbachia* presumably spread by enhancing host fitness in various ways, with some support for viral protection, fecundity enhancement, and supplementation of host nutrition (Weeks *et al.* 2007; Hedges *et al.* 2008; Teixeira *et al.* 2008; Brownlie

Copyright © 2019 by the Genetics Society of America

doi: <https://doi.org/10.1534/genetics.119.302349>

Manuscript received May 22, 2019; accepted for publication June 4, 2019; published Early Online June 21, 2019.

Supplemental material available at FigShare: <https://doi.org/10.25386/genetics.8301035>.

<sup>1</sup>Corresponding author: Division of Biological Sciences, University of Montana, 32 Campus Drive, Missoula, MT 59812. E-mail: [brandon.cooper@umontana.edu](mailto:brandon.cooper@umontana.edu)

*et al.* 2009; Gill *et al.* 2014; Martinez *et al.* 2014; Moriyama *et al.* 2015; Kriesner and Hoffmann 2018). Better understanding of *Wolbachia* effects, transmission and evolution should facilitate using *Wolbachia* for biocontrol of human diseases by either transforming vector populations with virus-blocking *Wolbachia* (e.g., McMeniman *et al.* 2009; Hoffmann *et al.* 2011; Schmidt *et al.* 2017; Ritchie 2018) or using male-only releases of CI-causing *Wolbachia* to suppress vector populations (Laven 1967; O'Connor *et al.* 2012).

There is a burgeoning literature on *Wolbachia* frequencies and dynamics in natural populations (e.g., Kriesner *et al.* 2013; Kriesner *et al.* 2016; Cooper *et al.* 2017; Bakovic *et al.* 2018; Meany *et al.* 2019), but fewer studies elucidate the modes and timescales of *Wolbachia* acquisition by host species (O'Neill *et al.* 1992; Rousset and Solignac 1995; Huigens *et al.* 2004; Baldo *et al.* 2008; Raychoudhury *et al.* 2009; Ahmed *et al.* 2015; Schuler *et al.* 2016; Turelli *et al.* 2018). Sister hosts could acquire *Wolbachia* from their most recent ancestors. Such cladogenic acquisition seems to be the rule for the obligate *Wolbachia* found in filarial nematodes (Bandi *et al.* 1998); and there are also examples in at least two insect clades, *Nasonia* wasps (Raychoudhury *et al.* 2009) and *Nomada* bees (Gerth and Bleidorn 2016). In contrast, infections can be relatively young and acquired through introgressive or horizontal transfer (Raychoudhury *et al.* 2009; Schuler *et al.* 2016; Conner *et al.* 2017; Turelli *et al.* 2018). Comparisons of host nuclear and mitochondrial genomes with the associated *Wolbachia* genomes enable discrimination among cladogenic, introgressive, and horizontal acquisition (Raychoudhury *et al.* 2009; Turelli *et al.* 2018; see Supplemental Material, Figure S1). Concordant nuclear, mitochondrial, and *Wolbachia* cladograms—including consistent divergence-time estimates for all three genomes—support cladogenic acquisition and codivergence of host and *Wolbachia* lineages. Concordant *Wolbachia* and mitochondrial phylogenies and consistent *Wolbachia* and mitochondrial divergence-time estimates that are more recent than nuclear divergence support introgressive acquisition. In this case, mitochondrial and *Wolbachia* relationships may or may not recapitulate the host phylogeny. Finally, if *Wolbachia* diverged more recently than either nuclear or mitochondrial genomes, horizontal transfer (or paternal transmission; Hoffmann and Turelli 1988; Turelli and Hoffmann 1995) is indicated. This is often associated with discordance between host and *Wolbachia* phylogenies (O'Neill *et al.* 1992; Rousset and Solignac 1995; Turelli *et al.* 2018).

Introgressive *Wolbachia* acquisition may be common in *Drosophila*. About half of all closely related *Drosophila* species have overlapping geographical ranges and show pervasive evidence of reinforcement (Coyne and Orr 1989, 1997; Yukilevich 2012; Nosil 2013), indicating that hybridization must be common (Turelli *et al.* 2014). Several instances of sporadic contemporary hybridization and interspecific gene flow have been documented in the genus (Carson *et al.* 1989; Shoemaker *et al.* 1999; Jaenike *et al.* 2006; Kulathinal *et al.* 2009; Garrigan *et al.* 2012; Brand *et al.* 2013; Matute and

Ayroles 2014; Lohse *et al.* 2015), but only two stable hybrid zones (HZs) have been well described, and both involve *D. yakuba*-clade species. In West Africa, *D. yakuba* hybridizes with endemic *D. santomea* on the island of São Tomé, and with *D. teissieri* on the island of Bioko (Lachaise *et al.* 2000; Comeault *et al.* 2016; Cooper *et al.* 2018). The ranges of *D. yakuba* and *D. teissieri* overlap throughout continental Africa, but contemporary hybridization has not been observed outside of Bioko (Cooper *et al.* 2018). Genomic analyses support both mitochondrial and nuclear introgression in the *D. yakuba* clade (Lachaise *et al.* 2000; Bachtrog *et al.* 2006; Llopart *et al.* 2014; Turissini and Matute 2017; Cooper *et al.* 2018), and the *Wolbachia* infecting all three species (*wSan*, *wTei*, and *wYak*) are at intermediate frequencies and identical with respect to commonly used typing loci (Lachaise *et al.* 2000; Charlat *et al.* 2004; Cooper *et al.* 2017).

Horizontal *Wolbachia* transmission has been repeatedly demonstrated since its initial discovery by O'Neill *et al.* (1992) (e.g., Baldo *et al.* 2008; Schuler *et al.* 2016), and it may be common in some systems (e.g., Huigens *et al.* 2000; Huigens *et al.* 2004; Ahmed *et al.* 2015; Li *et al.* 2017). Phylogenomic analyses indicate recent horizontal *Wolbachia* transmission, on the order of 5000–27,000 years, among relatively distantly related *Drosophila* species of the *D. melanogaster* species group (Turelli *et al.* 2018), but there is little evidence for nonmaternal transmission within *Drosophila* species (Richardson *et al.* 2012; Turelli *et al.* 2018). Horizontal *Wolbachia* transfer could occur via a vector (Vavre *et al.* 2009; Ahmed *et al.* 2015) and/or shared food sources during development (Huigens *et al.* 2000; Li *et al.* 2017). Rare paternal *Wolbachia* transmission has been documented in *D. simulans* (Hoffmann and Turelli 1988; Turelli and Hoffmann 1995). However, the most common mode of *Wolbachia* acquisition by *Drosophila* species (and most other hosts) remains unknown, and all three modes seem plausible in the *D. yakuba* clade (Lachaise *et al.* 2000; Bachtrog *et al.* 2006; Cooper *et al.* 2017).

Distinguishing acquisition via introgression vs. horizontal or paternal transmission requires estimating phylogenies and relative divergence times of mitochondrial DNA (mtDNA) and *Wolbachia* (Raychoudhury *et al.* 2009; Conner *et al.* 2017; Turelli *et al.* 2018). To convert sequence divergence to divergence-time estimates requires understanding relative rates of divergence for mtDNA, nuclear genomes, and *Wolbachia*. If each genome followed a constant-rate molecular clock, taxa with cladogenic *Wolbachia* transmission, such as *Nasonia* wasps (Raychoudhury *et al.* 2009) and *Nomada* bees (Gerth and Bleidorn 2016), would provide reference rates for calibration. However, like Langley and Fitch (1974), Turelli *et al.* (2018) found significantly varying relative rates of *Wolbachia* and mtDNA divergence. This variation, which we document across *Drosophila*, confounds attempts to understand *Wolbachia* acquisition, as discussed below.

The discovery of weak CI in the *D. yakuba* clade (Cooper *et al.* 2017) motivates detailed comparative analysis of loci

associated with CI (CI factors or *cifs*) in WO prophage regions of *Wolbachia* genomes. Beckmann and Fallon (2013) first associated wPip\_0282 and wPip\_0283 proteins in wPip-infected *Culex pipiens* with *Wolbachia*-modified sperm. Later work confirmed that these proteins induce toxicity and produce rescue when expressed/co-expressed in *Saccharomyces cerevisiae* (Beckmann *et al.* 2017), lending support to a toxin-antidote model of CI (Beckmann *et al.* 2019; but see Shropshire *et al.* 2019). Homologs of these genes in wMel (WD0631 and WD0632) recapitulate CI when transgenically expressed in *D. melanogaster* (LePage *et al.* 2017), and transgenic expression of WD0631 in *D. melanogaster* rescues CI (Shropshire *et al.* 2018). A distantly related WO prophage-associated pair, present in wPip, wPip\_0294 and wPip\_0295, causes similar toxicity/rescue in *S. cerevisiae* (Beckmann *et al.* 2017), and causes CI when placed transgenically into a *D. melanogaster* background (M. Hochstrasser, unpublished data). We adopt Beckmann *et al.* (2019)'s nomenclature, which assigns names based on enzymatic activity of the predicted toxin [deubiquitylase (DUB) and nuclease (Nuc)], with superscripts denoting focal *Wolbachia* strains when needed. Specifically, we refer to wPip\_0282-wPip\_0283 and the wPip\_0294-wPip\_0295 pairs as *cidA-cidB*<sup>wPip</sup> and *cinA-cinB*<sup>wPip</sup>, respectively; and we refer to WD0631-WD0632 as *cidA-cidB*<sup>wMel</sup>. This distinguishes *cid* (CI-inducing DUB) from *cin* (CI-inducing Nuc) pairs, with the predicted antidote and toxin denoted “A” and “B,” respectively (Beckmann *et al.* 2019). We acknowledge ongoing disagreement in the literature and direct readers to Beckmann *et al.* (2019) and Shropshire *et al.* (2019) for details. However, none of these debates on terminology or mechanism affect our findings.

Here, we use host and *Wolbachia* genomes from the *D. yakuba* clade to demonstrate introgressive and horizontal *Wolbachia* acquisition. General concordance of mitochondrial and *Wolbachia* phylogenies indicates that horizontal acquisition is rare within this clade. However, tests involving divergence-time estimates are complicated by varying relative rates and patterns of *Wolbachia*, mtDNA, and nuclear sequence divergence, as illustrated by data from more distantly related *Drosophila* (*Drosophila* 12 Genomes Consortium *et al.* 2007). Finally, we demonstrate that *cid* loci underlying CI in closely related wMel (LePage *et al.* 2017) are disrupted in all *D. yakuba*-clade *Wolbachia*. However, these *Wolbachia* also contain a set of *cin* loci, absent in wMel, but very similar to those found in wPip (Beckmann *et al.* 2017), a B-group *Wolbachia* strain that diverged 6–46 MYA from A-group wYak and wMel (Werren *et al.* 1995; Meany *et al.* 2019). This is the first discovery of two sets of loci implicated in CI co-occurring within the same prophage region. Several analyses implicate *Wolbachia*-specific insertion sequence (IS) transposable elements, specifically ISWpi1, in the horizontal transfer of these loci (and surrounding regions) between distantly related *Wolbachia*. Horizontal movement of incompatibility factors between prophage regions of *Wolbachia* variants adds another layer to what is already known about horizontal movement of prophages within and between

*Wolbachia* variants that themselves move horizontally between host species.

## Materials and Methods

### Genomic data

The *D. yakuba*-clade isofemale lines included in our study were sampled over several years in West Africa (Comeault *et al.* 2016; Cooper *et al.* 2017; Turissini and Matute 2017). Each line within each species used in our analyses exhibits little nuclear introgression (<1%) (Turissini and Matute 2017), and no hybrids were included. Reads from *D. yakuba* ( $N = 56$ ), *D. santomea* ( $N = 11$ ), and *D. teissieri* ( $N = 13$ ) isofemale lines were obtained from the data archives of Turissini and Matute (2017) and aligned to the *D. yakuba* nuclear and mitochondrial reference genomes (*Drosophila* 12 Genomes Consortium *et al.* 2007) with bwa 0.7.12 (Li and Durbin 2009), requiring alignment quality scores of at least 50. Because many of the read archives were single end, all alignments were completed using single-end mode for consistency.

**mtDNA:** Consensus mtDNA sequences for each of the 80 lines were extracted with samtools v.1.3.1 and bcftools v.1.3.1 (Li 2011). Coding sequences for the 13 protein-coding genes were extracted, based on their positions in the *D. yakuba* reference. We also extracted the 13 protein-coding genes from additional unique *D. yakuba* ( $N = 28$ ) and *D. santomea* ( $N = 15$ ) mitochondrial genomes (Llopart *et al.* 2014), and from the *D. melanogaster* reference (Hoskins *et al.* 2015). Genes were aligned using MAFFT v.7 and concatenated (Katoh and Standley 2013). Lines identical across all 13 genes were represented by one sequence for the phylogenetic analyses.

**Wolbachia:** Reads from each of the 80 Turissini and Matute (2017) lines were aligned using bwa 0.7.12 (Li and Durbin 2009) to the *D. yakuba* reference genome (*Drosophila* 12 Genomes Consortium *et al.* 2007) combined with the wMel reference genome (Wu *et al.* 2004). We calculated the average depth of coverage across the wMel genome. We considered lines with <1× coverage uninfected and lines with >10× coverage infected. No lines had between 1× and 10× coverage. To test our genomic analyses of infection status, we used a polymerase chain reaction (PCR) assay on a subset of lines. We extracted DNA using a “squish” buffer protocol (Gloor *et al.* 1993), and infection status was determined using primers for the *Wolbachia*-specific *wsp* gene (Braig *et al.* 1998; Baldo *et al.* 2006). We amplified a *Drosophila*-specific region of chromosome 2L as a positive control (Kern *et al.* 2015) (primers are listed in Table S1). For each run, we also included a known *Wolbachia*-positive line and a water blank as controls.

To produce a draft *Wolbachia* “pseudoreference” genome for each species, we first determined the isofemale line from

each species that produced the greatest average coverage depth over *wMel*. We trimmed the reads with Sickle v.1.33 (Joshi and Fass 2011) and assembled with ABySS v.2.0.2 (Jackman *et al.* 2017). *K* values of 51, 61...91 were tried. Scaffolds with best nucleotide BLAST matches to known *Wolbachia* sequences with *E*-values  $<10^{-10}$  were extracted as the draft *Wolbachia* assembly. For each species, the assembly with the highest N50 and fewest scaffolds was kept as our *Wolbachia* pseudoreference genome for that host species, denoted *wYak*, *wSan*, and *wTei* (Table S2). To assess the quality of these three draft assemblies, we used BUSCO v.3.0.0 to search for homologs of the near-universal, single-copy genes in the BUSCO proteobacteria database (Simão *et al.* 2015). For comparison, we followed Conner *et al.* (2017) and performed the same search on the reference genomes for *wMel* (Wu *et al.* 2004), *wRi* (Klasson *et al.* 2009), *wAu* (Sutton *et al.* 2014), and *wHa* and *wNo* (Ellegaard *et al.* 2013) (Table S3).

Using these draft *Wolbachia* pseudoreference genomes, reads from all other genotypes were aligned to the *D. yakuba* reference (nuclear and mitochondrial) plus the species-specific *Wolbachia* draft assembly with bwa 0.7.12 (Li and Durbin 2009). Consensus sequences were extracted with samtools v.1.3.1 and bcftools v.1.3.1 (Li 2011).

To test whether the choice of *Wolbachia* pseudoreference influenced the *Wolbachia* draft sequence obtained for each infected line, we arbitrarily selected three infected lines of each host species and aligned reads from each of those lines independently to our *wYak*, *wSan*, and *wTei* pseudoreferences. This resulted in 27 alignments (nine per host species). Among the three *Wolbachia* consensus sequences generated for each line, we assessed single-nucleotide variants between different mappings of the same line within loci used for downstream analyses. Irrespective of the pseudoreference used, we obtained identical *Wolbachia* sequences. This indicates the robustness of our approach to generating population samples of draft *Wolbachia* genomes, without having a high-quality *Wolbachia* reference genome from any of the host species. Because Llopart *et al.* (2014) released only assembled mitochondrial genomes, we could not assess the *Wolbachia* infection status of their lines.

#### **Loci for phylogenetic and comparative-rate analyses**

***Wolbachia* genes:** For each phylogenetic analysis of *Wolbachia* data, the draft genomes were annotated with Prokka v.1.11 (Seemann 2014), which identifies homologs to known bacterial genes. To avoid pseudogenes and paralogs, we used only genes present in a single copy and with identical lengths in all of the sequences analyzed. Genes were identified as single copy if they uniquely matched a bacterial reference gene identified by Prokka v.1.11. By requiring all homologs to have identical length in all of our draft *Wolbachia* genomes, we removed all loci with indels across any of our sequences.

Given that many loci accumulate indels over time, the number of loci included in our phylogenetic analyses

depended on the number of strains included. We first estimated phylograms for the A-group *Wolbachia* from the *D. yakuba*-clade, *wMel* from *D. melanogaster* (Wu *et al.* 2004), *wInc* from *D. incompta* (Wallau *et al.* 2016), *wSuz* from *D. suzukii* (Siozios *et al.* 2013), *wAna* from *D. anannasae* (Choi *et al.* 2015), *Wolbachia* that infect *Nomada* bees (*wNFe*, *wNPa*, *wNLeu*, and *wNFa*; Gerth and Bleidorn 2016), and *Wolbachia* that infect *D. simulans* (*wRi*, *wAu*, and *wHa*; Klasson *et al.* 2009; Ellegaard *et al.* 2013; Sutton *et al.* 2014). We also analyzed distantly related B-group *Wolbachia*: *wNo* from *D. simulans* (Ellegaard *et al.* 2013), *wPip\_Pel* from *C. pipiens* (Klasson *et al.* 2008), and *wAlbB* from *Aedes albopictus* (Mavingui *et al.* 2012). To increase our phylogenetic resolution within the *D. yakuba* *Wolbachia* clade (by increasing the number of loci), we estimated a phylogram that included only *D. yakuba*-clade *Wolbachia* and *wMel*. Our phylogram with both A- and B-group *Wolbachia* included 146 genes (containing 115,686 bp), and our phylogram that included only *D. yakuba*-clade *Wolbachia* and *wMel* included 643 genes (containing 644,586 bp).

We also constructed an absolute chronogram to estimate divergence between *D. yakuba*-clade *Wolbachia* strains. To illustrate how divergence-time estimates vary with changing patterns of molecular divergence, we estimated two additional *Wolbachia* chronograms: one that considered only the *wMel* reference plus the two most-diverged *wMel* lines included in Richardson *et al.* (2012), and one that included these three *wMel* variants and our *D. yakuba*-clade *Wolbachia*. Our chronogram that included only *D. yakuba*-clade *Wolbachia* included 678 genes (containing 695,118 bp), our chronogram that included only *wMel* *Wolbachia* included 692 genes (containing 709,599 bp), and our chronograms with *wMel* plus *D. yakuba*-clade *Wolbachia* included 621 genes (containing 624,438 bp). Independent estimates were obtained using relaxed-clock analyses with  $\Gamma(2,2)$  and  $\Gamma(7,7)$  branch-rate priors. We also estimated divergence between *D. yakuba*-clade *Wolbachia* and *wMel* using a strict-clock analysis, corresponding to  $\Gamma(n,n)$  as  $n \rightarrow \infty$ .

***mtDNA and nuclear genes from diverse Drosophila:*** To assess variation in the relative rates of divergence of mitochondrial and nuclear genes, we analyzed the canonical 12 - *Drosophila* genomes (*Drosophila* 12 Genomes Consortium *et al.* 2007) plus *D. suzukii* (Chiu *et al.* 2013). We excluded *D. sechellia* and *D. persimilis* which show evidence of introgression with *D. simulans* and *D. pseudoobscura*, respectively (Kulathinal *et al.* 2009; Brand *et al.* 2013; Schrider *et al.* 2018). Coding sequences for 20 nuclear genes used in the analyses of Turelli *et al.* (2018) (*aconitase*, *aldolase*, *bicoid*, *ebony*, *Enolase*, *esc*, *g6pdh*, *GlyP*, *GlyS*, *ninaE*, *pepck*, *Pgi*, *Pgm1*, *pic*, *ptc*, *Tpi*, *Transaldolase*, *white*, *wingless*, and *yellow*) were obtained from FlyBase for each species. The genes were aligned with MAFFT v.7 (Katoh and Standley 2013). Coding sequences for the 13 protein-coding mitochondrial genes in the inbred reference strains were also obtained from FlyBase for each species and were aligned with MAFFT v.7 and concatenated.

**Phylogenetic analyses:** All of our analyses used RevBayes v.1.0.9 (Höhna *et al.* 2016), following the procedures of Turelli *et al.* (2018). For completeness, we summarize those methods below. For additional details on the priors and their justifications, consult Turelli *et al.* (2018). Four independent runs were performed for each phylogenetic tree we estimated, and in all cases, the runs converged to the same topologies. Nodes with posterior probability <0.95 were collapsed into polytomies.

**Wolbachia phylograms:** We estimated a phylogram for A- and B-group *Wolbachia* and for only *D. yakuba*-clade and *wMel* *Wolbachia* using the same methodology as Turelli *et al.* (2018). We used a GTR +  $\Gamma$  model with four rate categories, partitioning by codon position. Each partition had an independent rate multiplier with prior  $\Gamma(1,1)$  [*i.e.*,  $\text{Exp}(1)$ ], as well as stationary frequencies and exchangeability rates drawn from flat, symmetrical Dirichlet distributions [*i.e.*,  $\text{Dirichlet}(1,1,1\dots)$ ]. The model used a uniform prior over all possible topologies. Branch lengths were drawn from a flat, symmetrical Dirichlet distribution, thus they summed to 1. Since the expected number of substitutions along a branch equals the branch length times the rate multiplier, the expected number of substitutions across the entire tree for a partition is equal to the partition's rate multiplier.

**Wolbachia chronograms:** We first created a relaxed-clock relative chronogram with the root age fixed to 1 using the GTR +  $\Gamma$  model, partitioned by codon position, using the same birth-death prior as Turelli *et al.* (2018). Each partition had an independent rate multiplier with prior  $\Gamma(1,1)$ , as well as stationary frequencies and exchangeability rates drawn from flat, symmetrical Dirichlet distributions. The branch-rate prior for each branch was  $\Gamma(2,2)$ , normalized to a mean of 1 across all branches (Table S4). We also tried a strict-clock tree and a relaxed-clock tree with branch-rate prior  $\Gamma(7,7)$ , which produced no significant differences. We used the scaled distribution  $\Gamma(7,7) \times 6.87 \times 10^{-9}$  to model substitutions per third-position site per year. This transforms the relative chronogram into an absolute chronogram. This scaled distribution was chosen to replicate the upper and lower credible intervals of the posterior distribution estimated by Richardson *et al.* (2012), assuming 10 *Drosophila* generations per year, normalized by their median substitution-rate estimate. Branch lengths in absolute time were calculated as the relative-branch length times the third-position rate multiplier divided by the substitutions per third-position site per year estimate above.

We illustrate how divergence-time estimates depend on changing patterns of *Wolbachia* molecular evolution observed over different timescales, specifically the relative rates of third-site substitutions *vs.* first- and second-site substitutions. We compare divergence times estimated for variants within *D. melanogaster*-group host species, with a timescale of only hundreds or thousands of years, with the same divergence times estimated when more distantly related *Wolbachia*, with

divergence times over tens of thousands of years, are included in the analyses. We present three separate analyses: one using only *D. yakuba*-clade *Wolbachia* variants, a second using only *wMel* variants analyzed by Richardson *et al.* (2012), and a third simultaneously analyzing both sets of data.

***Drosophila* phylogeny for 11 reference species:** We estimated a phylogram from our 20 nuclear loci using the GTR +  $\Gamma$  model, with partitioning by gene and codon position. The model was identical to the *Wolbachia* phylogram model above except for the partitioning (there are too few *Wolbachia* substitutions to justify partitioning by gene).

**mtDNA phylogeny for the 11 reference species:** We estimated a phylogram from the 13 protein-coding mitochondrial loci using the GTR +  $\Gamma$  model, with partitioning only by codon position. The model was identical to the *Wolbachia* phylogram model above.

**mtDNA chronogram for the *D. yakuba* clade:** We estimated a relative chronogram for the *D. yakuba*-clade mitochondria from the 13 protein-coding loci using the GTR +  $\Gamma$  model, partitioning only by codon position. To test the sensitivity of our results to priors, we ran a strict clock, a relaxed clock with a  $\Gamma(7,7)$  branch-rate prior, and a relaxed clock with a  $\Gamma(2,2)$  branch-rate prior, corresponding to increasing levels of substitution-rate variation across branches. The model was identical to the *Wolbachia* chronogram model above, except that we did not transform it into an absolute chronogram.

**Ratios of divergence rates for mtDNA *vs.* nuclear loci:** To quantify ratios of mtDNA to nuclear substitution rates, we estimated relative substitution rates for host nuclear genes *vs.* mtDNA using the GTR +  $\Gamma$  model. The (unrooted) topology was fixed to the consensus topology for the nuclear and mitochondrial data. The data from 20 nuclear loci were partitioned by locus and codon position, the mtDNA data were partitioned only by codon position. All partitions shared the same topology, but the nuclear partitions were allowed to have branch lengths different from the mtDNA partitions. The sum of the branch lengths for each partition was scaled to 1. Assuming concurrent nuclear-mtDNA divergence (because we used only species showing no evidence of introgression), we imposed the same absolute ages for all nodes of the nuclear and mtDNA chronograms. For each nuclear locus, the third-position rate ratio for the mtDNA *vs.* nuclear genomes was calculated as follows: (mitochondrial branch length  $\times$  mitochondrial third-position rate multiplier)/(nuclear branch length  $\times$  nuclear third-position rate multiplier). We summarized the relative rates of mtDNA *vs.* nuclear substitutions along each branch using the arithmetic average of the 20 ratios obtained from the individual nuclear loci.

**Introgressive *vs.* horizontal *Wolbachia* transfer-concordance of phylograms:** Horizontal transfer of *Wolbachia* was initially detected by discordance between *Wolbachia* and host

phylogenies (O'Neill *et al.* 1992). When considering samples within species and closely related species, a comparison of mitochondrial and *Wolbachia* phylogenies provides a natural test for horizontal transmission. We first look for significant discrepancies between strongly supported nodes in the mitochondrial vs. *Wolbachia* phylogenies. To test for less obvious differences, we follow Richardson *et al.* (2012) and compute Bayes factors to assess the support for models that assume that mitochondrial vs. *Wolbachia* follow the same topology vs. distinct phylogenies. For these calculations, we omit *D. san-tomea* line Quija 37 due to its clear discordance.

We calculated the marginal likelihood in RevBayes for two models: one with a shared mitochondrial and *Wolbachia* phylogeny, and another with independent topologies. The mtDNA and *Wolbachia* were each partitioned by codon position, for a total of six partitions. In the shared phylogeny model, all six share the same topology; in the independent model, mitochondria and *Wolbachia* have separate topologies. All priors were the same as the *Wolbachia* phylogram model. Lines that were identical across the mtDNA and *Wolbachia* were collapsed into one sample. We ran two independent replicates of each model with 50 stepping stones per run. The Bayes factor is computed as the difference between the marginal likelihoods of each model.

**Introgressive vs. horizontal *Wolbachia* transfer–relative rates:** Within species, the phylogenies of mitochondria and *Wolbachia* are fully resolved. Thus we consider an alternative approach for distinguishing between introgressive vs. horizontal transfer that depends on estimating divergence times for *Wolbachia* genomes and host mtDNA. This is complicated by variation in the relative rates of mtDNA vs. *Wolbachia* divergence and by systematic changes over time in the relative rates of substitutions at the three codon positions for *Wolbachia*. Following the procedures in Turelli *et al.* (2018), we estimated the mtDNA:*Wolbachia* third-position substitution-rate ratio for each branch in the *D. yakuba* clade. For each analysis, we computed the marginal likelihood of the model where all branches shared the same ratio and the same model, except allowing different ratios on each branch. We then calculated Bayes factors (*i.e.*, differences in the log of the marginal likelihoods) to determine which model was favored. We repeated these analyses after including three *wMel*-infected *D. melanogaster* lines with the *D. yakuba*-clade data set. Because *D. yakuba*-clade species and *D. melanogaster* do not produce fertile hybrids (Sánchez and Santamaria 1997; Turissini *et al.* 2017), introgressive transfer of *Wolbachia* is not possible between these species. Hence, including *wMel* provides a control for our proposed test of introgressive transfer. For these controls, we calculated the mitochondrial vs. *Wolbachia* substitution-rate ratio for the interspecific branches separately, regardless of whether the shared rate ratio model was favored.

### ***Wolbachia* loci associated with CI**

*Wolbachia* that infect all three *D. yakuba*-clade hosts cause weak intra- and interspecific CI (Cooper *et al.* 2017), but the

genetic basis of CI in this clade remains unknown. We used tBLASTN to search for *cif* homologs in each of our *D. yakuba*-clade *Wolbachia* assemblies, querying *cidA-cidB* variants and the *cinA-cinB<sup>wRi</sup>* pair (*wRi\_006720* and *wRi\_006710*) found in *wRi* and some *wRi*-like *Wolbachia* (Beckmann and Fallon 2013; Turelli *et al.* 2018), and the *cinA-cinB<sup>wPip</sup>* pair (Beckmann and Fallon 2013; Beckmann *et al.* 2017; LePage *et al.* 2017; Lindsey *et al.* 2018). We identified both *cid* and *cin* homologs, but we found no close matches to the *cinA-cinB<sup>wRi</sup>* pair (denoted Type II *cif* loci by LePage *et al.* 2017) in any of our genomes. For all of the samples, we extracted consensus sequences from our assemblies and alignments for the *cidA-cidB<sup>wYak-clade</sup>* and *cinA-cinB<sup>wYak-clade</sup>* gene pairs. The genes were aligned with MAFFT v.7 (Katoh and Standley 2013). We examined variation in these regions relative to *cidA-cidB<sup>wMel</sup>* and *cinA-cinB<sup>wPip</sup>*, respectively.

Because we unexpectedly identified homologs of *cinA-cinB* in all *wYak*-clade *Wolbachia*, we took additional measures to understand the origin and placement of these loci in the genomes. The *cinA-cinB<sup>wYak-clade</sup>* open reading frames (ORFs) are located on a ~11,500 bp scaffold in the fragmented *wYak* assembly (*wYak* scaffold “702380” from ABySS output; we use quotes to indicate names assigned by ABySS). The first ~4000 bp of this scaffold contain *cinA* and *cinB* genes that have ~97% identity with those in *wPip*. We performed a BLAST search using all contigs in the *wYak* assembly as queries against the *wMel* genome (Camacho *et al.* 2009). The *cinA-cinB<sup>wYak-clade</sup>* scaffold was placed on the *wMel* genome (~617,000–623,000) adjacent to and downstream of the *wYak* contig containing *cidA-cidB<sup>wYak-clade</sup>* loci (*wYak* contig “187383”). However, only ~7000 bp of the 11,500 bp of the scaffold containing *cinA-cinB<sup>wYak-clade</sup>* align to this region of *wMel*. The ~4000 bp sequence that contains the ORFs for *cinA-cinB<sup>wYak-clade</sup>* has no significant hit against the *wMel* genome.

To verify the placement of the scaffold containing *cinA-cinB<sup>wYak-clade</sup>*, we first performed targeted assembly using the *cinA-cinB<sup>wYak-clade</sup>* contig and the adjacent contigs as references (Hunter *et al.* 2015). Iterative targeted assembly often extends assembled scaffolds (Langmead and Salzberg 2012). Reads are mapped to the target of interest, and the mapped reads are then assembled using SPADES (Bankevich *et al.* 2012; Nurk *et al.* 2013). The newly assembled scaffolds serve as the target in a subsequent round of mapping. The procedure is repeated until no new reads are recruited to the assembled scaffold from the previous round. The extended scaffolds enabled us to merge flanking *wYak* contigs downstream of the *cinA-cinB<sup>wYak-clade</sup>* scaffold but failed to connect the fragment with the *cidA-cidB<sup>wYak-clade</sup>* contig. We used PCR to amplify the intervening region (Table S1), and Sanger sequencing to evaluate this product.

Subsequent mapping of paired-end reads to the merged scaffolds confirmed the correct order and orientation of the contigs containing the *cidA-cidB<sup>wYak-clade</sup>* and *cinA-cinB<sup>wYak-clade</sup>* genes. Observed, uncorrected, pairwise distances between focal regions in *wYak*, *wMel*, and *wPip* and between focal regions in



wYak, wPip, wAlbB, and wNleu were calculated using a sliding window (window size = 200 bp, step size = 25 bp).

### Data availability

The *Wolbachia* assemblies (BioProject PRJNA543889, accession numbers GCA\_005862115.1, GCA\_005862095.1, and GCA\_005862135.1) and Sanger sequences (MK950151 and MK950152) are archived in GenBank. All three assemblies were manually corrected using Sanger sequence to accurately represent variation in the *cin* region. Relevant genetic data and scripts have been uploaded to DRYAD (10.5061/dryad.8n1n677). Supplemental material available at FigShare: <https://doi.org/10.25386/genetics.8301035>.

## Results

### Unidirectional introgression of *D. yakuba* mitochondria into *D. santomea*

Previous analyses suggest that *D. yakuba* and *D. santomea* carry very similar mtDNA due to ongoing hybridization and introgression (Lachaise *et al.* 2000; Bachtrog *et al.* 2006; Llopart *et al.* 2014). However, our mtDNA relative chronogram (Figure 1), based on mitochondrial whole-proteome data sampled from throughout the ranges of all three species, supports three mtDNA clades that largely agree with the nuclear topology; *D. teissieri* mtDNA sequences are outgroup to sister *D. yakuba* and *D. santomea* mtDNA (Figure 1). Consistent with introgression, 12 out of 26 *D. santomea* isofemale lines have *D. yakuba*-like mtDNA (indicated by blue branches and letters in Figure 1). Yet, our samples showed no evidence of *D. santomea* mtDNA introgressed into *D. yakuba* and no mtDNA introgression involving *D. teissieri*. The 12 *D. santomea* isofemales with introgressed *D. yakuba*-like mtDNA were sampled from both within ( $N = 8$ ) and outside ( $N = 4$ ) the well-described Pico de São Tomé HZ (Llopart *et al.* 2014; Turissini and Matute 2017). The *D. santomea* HZ samples are indicated by “HZ” in Figure 1 in blue, and they are found throughout the clade that includes all *D. yakuba* mtDNA. This suggests that hybridization occurs in other areas of the island or that introgressed *D. santomea* genotypes migrate outside of the HZ.

Previous analysis of nuclear genomes (Turissini and Matute 2017) also found more introgression from *D. yakuba* into *D. santomea* when populations of *D. yakuba* from near the Gulf of Guinea, Cameroon, and Kenya were included; however, excluding Cameroon and Kenya indicated similar amounts of introgression in each direction. Matings between *D. santomea* females and *D. yakuba* males are rare relative to the reciprocal cross (Coyne *et al.* 2002; Matute 2010). When these matings do occur, F1 females produce fewer progeny than do F1 females produced by *D. yakuba* females and *D. santomea* males (Matute and Coyne 2010). F1 hybrids produced by *D. santomea* females also have a shortened life span (Matute and Coyne 2010) due to copulatory wounds inflicted by *D. yakuba* males during mating (Kamimura and

Mitsumoto 2012). These observations are consistent with our finding of preferential introgression of *D. yakuba* mitochondria into *D. santomea* backgrounds.

### *Wolbachia* frequencies and draft genomes

As expected, we found *Wolbachia* in all three *D. yakuba*-clade species sampled by Turissini and Matute (2017). For *D. yakuba*, 21 out of 56 lines were infected, yielding an infection frequency of  $P = 0.36$ , with 95% binomial confidence interval (0.25–0.51). For *D. santomea*, 10 out of 11 were infected, yielding  $P = 0.91$  (0.59–1.0); and for *D. teissieri*, 11 out of 13 were infected, yielding  $P = 0.85$  (0.55–0.98). Additional frequency estimates are reported in Cooper *et al.* (2017), which found that *Wolbachia* frequencies vary through time and space in West Africa. All PCR tests for *Wolbachia* matched our coverage-based genomic analyses of infection status.


The average coverage across the genome, calculated as the total number of bases aligned to the wMel reference divided by its length, was 1940 for our wYak pseudoreference genome (yak-CY17C), 40 for wSan (san-Quija630.39), and 489 for wTei (teis-cascade\_4\_2). These pseudoreference genomes were included in the phylogram that includes A-group and B-group *Wolbachia* (Figure 2), and they cluster with the *Wolbachia* from their respective hosts (Figure 3), as expected if introgression is rare. In general, the wSan data are lower quality due to the relatively small size of the *D. santomea* libraries (Turissini and Matute 2017). The scaffold count, N50, and total assembly sizes are reported in Table S2.

### Introgressive and horizontal *Wolbachia* transfer among hosts

Our phylogram that includes A-group and B-group *Wolbachia* places the three weak-CI-causing *D. yakuba*-clade *Wolbachia* together and sister to wMel in the A group (Figure 2A). In contrast, *D. simulans* carries diverse *Wolbachia* that do (wRi, wHa, wNo) and do not (wAu) cause CI, spanning *Wolbachia* groups A (wRi, wHa, wAu) and B (wNo). All wSan, wYak, and wTei *Wolbachia* variants included in our analysis are identical at the MLST loci often used to type *Wolbachia* variants (Baldo *et al.* 2006; Cooper *et al.* 2017). Because of the relatively small number of genes that meet our inclusion criteria (146 genes across 115,686 bp), the resulting phylogram does not resolve relationships among wSan, wYak, and wTei (Figure 2A). Our phylogram that includes only *D. yakuba*-clade and wMel *Wolbachia* (based on 643 genes across 644,586 bp) resolves these relationships, placing wTei variants outgroup to sister wSan and wYak in distinct clades (Figure 2B). We observe 0.0039% third-position pairwise differences between wTei and wYak and between wTei and wSan, and 0.0017% between wSan and wYak. The third-position pairwise differences between any of the *D. yakuba*-clade *Wolbachia* and wMel is 0.11%. For reference, the third-position pairwise difference between wYak and wRi is 2.9%, whereas wRi and wSuz differ by only 0.014% (Turelli *et al.* 2018).

Our absolute chronogram that includes wMel and *D. yakuba*-clade *Wolbachia* also indicates three monophyletic






**Figure 1** Bayesian relative chronogram for the *D. yakuba* clade mtDNA. The root age was fixed to 1. Nodes with posterior probability <0.95 were collapsed into polytomies, numbers at the tips show the number of genotypes collapsed. Node labels are the point estimates and 95% credible intervals for the relative ages. Tips and branches showing *D. santomea* individuals with *D. yakuba* mtDNA are colored blue, those sampled from the Pico de São Tomé hybrid zone are denoted by HZ. *D. santomea* line Quija 37 showing evidence of horizontal or paternal *Wolbachia* transmission (or paternal mtDNA transmission) is in bold blue. The chronogram was estimated from the 13 protein-coding mitochondrial genes. A representation of host nuclear relationships with photos of host females at the tips is inset, with the divergence times estimated by Turissini and Matute (2017) superimposed.

groups (Figure 3), whose topology—[*wTei*, (*wYak*, *wSan*)]—agrees with the nuclear topology of the hosts (Turissini and Matute 2017). However, one *D. santomea* line, *D. santomea* Quija 37, sampled from the south side of São Tomé (Matute 2010), carries *Wolbachia* identical to the other eight *wSan* samples across 695,118 bp, yet its mtDNA is nested in the *D.*

*yakuba* mtDNA cluster (bold blue in Figure 1). This single example of *Wolbachia*-mtDNA phylogenetic discordance can be explained by either horizontal *Wolbachia* transfer or paternal transfer of *Wolbachia* or mtDNA. While rare, paternal *Wolbachia* transmission has been observed in other *Drosophila* (Hoffmann and Turelli 1988; Turelli and




**Figure 2** (A) Bayesian phylogram placing the *D. yakuba*-clade *Wolbachia* in the context of other A-group, and also B-group, *Wolbachia*. *wPip*, *wNo*, and *wAlbB* belong to *Wolbachia* group B, while *D. yakuba* clade-*Wolbachia*, *wMel*, *wAu*, *wInc*, *wRi*, *wAna*, *wSuz*, *wHa*, and *Nomada Wolbachia* belong to group A. The phylogram was estimated with 146 single-copy genes of identical length in each genome, spanning 115,686 bp. The limited number of genes across both A-group and B-group *Wolbachia* meeting our criteria precluded resolving relationships of *D. yakuba*-clade *Wolbachia*. (B) Bayesian phylogram of *wYak*, *wSan*, *wTei*, and outgroup *wMel*. The phylogram was estimated with 643 single-copy genes of identical length in each genome, spanning 644,586 bp. These more extensive data provide increased phylogenetic resolution of the *D. yakuba*-clade *Wolbachia*. For both phylograms, nodes with posterior probability <0.95 were collapsed into polytomies.

Hoffmann 1995), as has paternal transmission of mtDNA (Kondo *et al.* 1990). The only other *Wolbachia*-screened *D. santomea* isofemale line with introgressed mtDNA was *Wolbachia* uninfected.

In Figure 3, we present alternative estimates of the divergence times for *Wolbachia* within and between *D. yakuba*-clade species, within *D. melanogaster*, and between the *D. yakuba* clade and *D. melanogaster*. The estimates from the chronogram that includes all of the indicated *Wolbachia* variants are in bold. To demonstrate how these time estimates depend on the sequences included in the analyses (through estimates of relative divergence for each codon position), estimates that included only *D. yakuba*-clade *Wolbachia* or only *wMel* variants are superimposed and not in bold. These estimates were all obtained using a relaxed-clock analysis with a  $\Gamma(2,2)$  branch-rate prior [results using  $\Gamma(7,7)$  and a strict clock are mentioned below]. When considering only the *D. yakuba*-clade *Wolbachia* (Figure 3), we estimate the root at 2510 years (95% credible interval 773–5867 years) and the *wYak*-*wSan* split at 1556 years (95% credible interval 411–3662 years). The *wMel* chronogram estimates the most-recent common ancestor (MRCA) of *wMel* at 4890 years ago (95% credible interval 1258–11,392 years). Our *wMel* result is consistent with the Richardson *et al.* (2012) estimate of 8008 years (95% credible interval 3263–13,998). The alternative branch-rate  $\Gamma(7,7)$  prior has little effect on these divergence-times estimates, with credible intervals that overlap with those produced using a  $\Gamma(2,2)$  branch-rate prior (Figure 3 and Table S5). In contrast, Figure 3 shows that including more divergent variants influences our time estimates, although the credible intervals are overlapping in each case. Table S4 shows how estimates of the relative

amounts of the divergence for the three codon positions vary across the data sets. The underlying model for divergence times assumes constant relative rates at the three positions. Hence, changes in relative rates affect divergence-time estimates.

As noted above, our estimates of the time of the MRCA for *Wolbachia* within the *D. yakuba* clade or within *D. melanogaster* are relatively insensitive to whether the branch-rate prior is  $\Gamma(2,2)$  or  $\Gamma(7,7)$ . The choice of genes used to estimate divergence also has little effect—maximal distance between *D. yakuba*-clade strains was identical for analyses that included only the *D. yakuba*-clade ( $N = 678$  genes) or *wYak*-clade plus *wMel* ( $N = 621$  genes) gene sets (*wYak*-*wTei* distance = 0.0035%). The estimated divergence time between *wMel* and the *D. yakuba*-clade *Wolbachia* is less robust, although again the credible intervals generated using alternative branch-rate priors are overlapping (Figure 3 and Table S5). A strict-clock analysis produces an estimate of 72,612, with 95% credible interval (23,412–132,276), again overlapping with our relaxed-clock estimates. An alternative point estimate can be obtained from pairwise differences. Averaging over the *D. yakuba*-clade sequences, we observe an average third-position difference of 0.107% between the *D. yakuba*-clade *Wolbachia* and *wMel*, or 0.0535% from tip to root. The “short-term evolutionary rate” of divergence within *wMel*, estimated by Richardson *et al.* (2012), produces a point estimate of 78,000 years, which overlaps with the credible interval of our relaxed-tree estimate with a  $\Gamma(7,7)$  branch-rate prior (Table S5). This molecular-clock estimate closely agrees with our strict-clock chronogram analysis (Figure S2). Given that these estimates are much shorter than the divergence time between their reproductively isolated hosts,



**Figure 3** Bayesian chronogram for the *D. yakuba*-clade *Wolbachia* plus three *wMel* genomes with absolute age estimates based on the calibration in Richardson *et al.* (2012) in bold. The chronogram was estimated with 621 single-copy genes of identical length in all of the genomes, spanning 624,438 bp. Nodes with posterior probability <0.95 were collapsed into polytomies. Node labels are the point estimates and 95% credible intervals for the ages. We also constructed two other chronograms: one that included only *D. yakuba*-clade *Wolbachia* and one that included only *wMel* *Wolbachia*. The divergence estimates for the chronogram including only *D. yakuba*-clade *Wolbachia* (\*) and only *wMel* *Wolbachia* (§) are superimposed to illustrate that the inclusion of both groups in the chronogram alters the absolute divergence estimates within each group. Our chronogram with only *D. yakuba*-clade *Wolbachia* included 678 genes (containing 695,118 bp), and our chronogram including only *wMel* *Wolbachia* included 692 genes (709,599 bp).


horizontal *Wolbachia* transmission must have occurred. As discussed below, the horizontal transmission probably involved intermediate hosts.

**mtDNA and nuclear concordance across *Drosophila*:** The topologies of the nuclear and mtDNA trees for the 11 reference *Drosophila* species are completely concordant (Figure 4) and agree with the neighbor-joining result presented in *Drosophila* 12 Genomes Consortium *et al.* (2007). We find the average ratio of mtDNA to nuclear substitution rates across branches vary from  $\sim 1\times$  to  $10\times$ , consistent with the results of Havird and Sloan (2016) across Diptera.

**mtDNA and *Wolbachia* concordance in the *D. yakuba* clade:** *D. santomea* line Quija 37 unexpectedly carries

mitochondria belonging to the clade associated with *D. yakuba* but has *wSan* *Wolbachia*. This discordance requires horizontal or paternal transmission. Apart from this, we find no phylogenetic discordance between strongly supported nodes (*i.e.*, posterior probability at least 0.95) in the mitochondrial and *Wolbachia* trees (Figure S3). For a more refined test of concordance, we removed the Quija 37 line from our mitochondrial and *Wolbachia* data and computed Bayes factors. The shared topology model was favored by a Bayes factor of  $e^{55}$ , indicating strong support for concordance of the mitochondrial and *Wolbachia* phylogenies.

**Variation in the ratio of mtDNA:*Wolbachia* substitution rates:** Following the methods of Turelli *et al.* (2018) to look for different mitochondrial vs. *Wolbachia* branch lengths, we



**Figure 4** A Bayesian phylogram, with branch lengths fixed to 1, displaying relative rate variation for mtDNA vs. nuclear loci. Branches are labeled with quartiles of the estimated third-position mtDNA to third-position nuclear substitution rate. The nuclear substitution rate was taken as the arithmetic mean third-position rate over the 20 loci used. The mtDNA and nuclear phylograms are topologically identical when run separately.

also found no evidence of variation in the ratio of mtDNA:*Wolbachia* substitution rates within the *D. yakuba* clade (median third-position ratio = 223, quartiles = 186 and 268); the model with all branches sharing the same ratio is favored over the model where each branch has its own ratio by a Bayes factor of  $e^7$  (Figure S4). For comparison, Turelli *et al.* (2018) found a median ratio of 566 within *D. suzukii* and 406 within *w*Ri-infected *D. ananassae*-subgroup species. These differences in relative rates are broadly compatible with the 10-fold variance in relative rates noted above for nuclear vs. mtDNA divergence across *Drosophila* species. Hence, our data seem compatible with *Wolbachia* transmission via introgression within the *D. yakuba* clade.

Although this result seems to strongly support purely maternal transmission of *Wolbachia* within species and introgressive transfer between species (with more recent introgression between the sister species *D. yakuba* and *D. santomea*), this interpretation is severely weakened by our “control” analysis that includes *w*Mel-infected *D. melanogaster* (for which introgression is impossible). Including the *D. melanogaster* data, the constant-ratio model was still favored over variable ratios by a Bayes factor of  $e^{54}$  (median third-position ratio = 297, quartiles = 276 and 323). We discuss the implications of this anomalous result below. The key observation is that significantly declining rates of substitution for *Wolbachia* [and mtDNA, see Ho *et al.* (2005)]

over time, together with heterogeneity of relative rates (as illustrated in Figure 4), limit the power of relative substitution ratios to differentiate between introgression and horizontal transmission.

#### **Transposon-mediated transfer of CI factors independent of WO phage**

In contrast to *w*Mel, which contains only the *cidA-cidB* gene pair (LePage *et al.* 2017), the *D. yakuba*-clade *Wolbachia*, which also belong to *Wolbachia* group A, have both *cidA-cidB* and a *cinA-cinB* pair homologous to CI loci originally identified in group-B *w*Pip (Beckmann *et al.* 2017). The A and B *Wolbachia* groups diverged ~ 6–46 MYA (Werren *et al.* 1995; Meany *et al.* 2019). Among our *w*Yak-clade sequences, there are no single-nucleotide variants within any of the *cif* loci. The *cidB<sup>w</sup>Yak-clade* locus has an inversion from amino acids 37–103 relative to the same region in *cidB<sup>w</sup>Mel* in every *w*Yak-clade *Wolbachia* variant we analyzed. This introduces several stop codons, which might render this gene nonfunctional. On the other hand, RNA polymerase should still transcribe a complete polycistronic transcript. Therefore translation of an N-terminally truncated *CidB<sup>w</sup>Yak-clade* protein cannot be ruled out. With the exception of a 236 bp tandem duplication in *cinB<sup>w</sup>Yak-clade* (Figure 5 and Figure 6), the sequence differences between *cinA<sup>w</sup>Yak-clade* and *cinB<sup>w</sup>Yak-clade* regions compared to *cinA<sup>w</sup>Pip* and *cinB<sup>w</sup>Pip* homologs are 1.35 and 0.59%,

respectively. In contrast, the average difference between *wYak* and *wPip* genomes across all 194 genes (161,655 bp) present in single copy and with identical lengths in both genomes is 11.60%. Conversely, outside of the prophage regions, *wMel* and *wYak* differ by ~1%. This is consistent with data that indicate WO phage regions have a different evolutionary history than the bulk of the *Wolbachia* genome (LePage *et al.* 2017; Lindsey *et al.* 2018). The 236 bp tandem duplication in the *cinB*<sup>*wYak-clade*</sup> introduces a frameshift in the transcript at position 588. It is unclear whether the *cinB*<sup>*wYak-clade*</sup> protein retains functionality.

The absence of *cin* loci in *wMel*, combined with the similarity of this region between *wYak* and distantly related *wPip*, led us to assess how these *Wolbachia* acquired *cin* loci. Targeted assembly extended the scaffold *wYak* “702380” containing the *cin* loci (Figure 5) but did not definitively place it relative to the *wMel* genome. BLAST searches indicated that contig *wYak* “187383” was likely flanked by *wYak* “702380.” PCR primers designed in both contigs amplified the intervening region (labeled “A” in Figure 5), confirming we have discovered the first WO prophage with two sets of *cif* loci. Subsequent Sanger sequencing revealed that this region contains an ISWpi1 element, found in many *Wolbachia* genomes, but not present at this location in the *wMel* reference sequence (Cordaux 2008).

IS elements encode a transposase gene that mediates their movement (Chandler and Mahillon 2002). ISWpi1 elements are related to the IS5 family, which seems to be restricted to *Wolbachia* (Duron *et al.* 2005; Cordaux 2008). ISWpi1 elements occur in more than half of the *Wolbachia* strains that have been evaluated, including *wYak* and *wMel* (Cordaux 2008); and these elements occur at variable copy number, potentially facilitating horizontal transmission of ISWpi1 elements and their intervening sequence between *Wolbachia* variants/strains (Cordaux 2008). Following placement of contig *wYak* “702380” relative to *wMel*, we aligned these regions and calculated pairwise differences along the chromosome using a sliding window.

In addition to containing the *cinA-cinB* loci, absent in *wMel*, contig *wYak* “702380” is on average ~10% different from *wMel*. In contrast, most of *wYak* is <1% different from *wMel* (for instance, across the 650,559 bp used in our phylogenetic analyses, *wMel* and *wYak* differ by only 0.09%). Downstream of the *cinA-cinB*<sup>*wYak-clade*</sup> region, targeted assembly enabled us to join several more contigs. The junctions were corroborated by mapping paired-end reads (Langmead and Salzberg 2012) and visually inspecting the resulting bam files around joined contigs for reads spanning the new junctions and for concordant read-pair mappings. However, our attempts to fully bridge the gap downstream of the *cinA-cinB*<sup>*wYak-clade*</sup> genes via targeted assembly, scaffolding, and PCR were unsuccessful (see “*wYak* unassembled gap” in Figure 5). The unassembled region in *wYak* contains an ISWpi1 element in *wMel*, Mel #9 (labeled “B” in Figure 5; Cordaux 2008). Although not part of our *wYak* assembly, homologs of this ISWpi1 element appear in assemblies of this region of the


WO prophage in several other A-group *Wolbachia* (Cordaux 2008; Bordenstein and Bordenstein 2016), specifically *wInc* from *D. incompta* (Wallau *et al.* 2016) and *wRi* from *D. simulans* (see Figure 2). In both, we find orthologs to Mel#9 ISWpi1 with corresponding flanking sequence assembled, indicating that this IS element probably occurs in this unassembled region of *wYak*, *wSan*, and *wTei*. The *wYak* sequence between these two ISWpi1 elements is highly diverged from *wMel* in comparison to the rest of the *wYak* genome (~10% difference vs. 1%). It therefore seems plausible that this region experienced a horizontal transfer event in the ancestor of *wYak*, *wTei*, and *wSan*, mediated by the flanking ISWpi1 elements.

We conjectured that horizontal transfer occurred via the excision of the two ISWpi1 elements and the intervening DNA from the donor followed by homologous recombination within the IS elements. To assess the plausibility of this scenario, we used BLAST to compare *cinA-cinB*<sup>*wPip*</sup> genes against all published *Wolbachia* genomes in the NCBI assembly database (<https://www.ncbi.nlm.nih.gov/assembly/?term=Wolbachia>). We found homologs of the *cinA-cinB*<sup>*wPip*</sup> in the group-A *Wolbachia* associated with *Nomada* bees (Gerth and Bleidorn 2016). An unrooted tree of the ORFs for *cinA* and *cinB* (Figure 6) indicates that these genes in *wYak*, *wSan*, and *wTei* are more similar to *cinA-cinB* from group-B *wPip* and *wAlbB* than to *cinA-cinB* found in the fellow group-A *Wolbachia* associated with *Nomada* bees (*wNFe*, *wNPa*, *wNLeu*, and *wNFa*), which each harbor two different *cinA-cinB* copies. This indicates that *cinA-cinB*<sup>*wYak-clade*</sup> were acquired via a horizontal transfer event across *Wolbachia* groups B and A that is independent of the event(s) that placed *cinA-cinB* in the *Wolbachia* associated with *Nomada* bees, suggesting repeated transfers of *cin* loci.

The two *cinA-cinB* copies (denoted *wNLeu1* and *wNLeu2* in Figure 6) in *Nomada Wolbachia* are nearly as distinct from each other as they are from the homologs in *wPip*, *wAlbB*, and *wYak* (~7% diverged from these strains, Figure 6). However, among the four *Wolbachia*-infected *Nomada* species, the orthologs are very similar, with *cinA*<sup>*wNLeu1*</sup> having only 0–0.15% pairwise differences among the four strains and *cinA*<sup>*wNLeu2*</sup> having 0–0.56% pairwise differences (reconstructions for *cinB* gene copies were more complicated as the *cinB*<sup>*wNLeu1*</sup> copy fails to assemble into a single contig on the three-prime end). This pattern suggests that *wNLeu1* and *wNLeu2* *cin* copies were acquired by the common ancestor of the four *Nomada Wolbachia* strains analyzed, followed by cladogenic transfer across *Nomada* species (Gerth and Bleidorn 2016). The highly fragmented assemblies of the four *Nomada Wolbachia* strains, with duplicate copies confounding assembly, make it difficult to determine the relative positions of the *cinA* and *cinB* copies and if they are likewise flanked by ISWpi1 elements.

To determine whether these genes were potentially moved by ISWpi1 elements into the *Wolbachia* of the *D. yakuba* clade and *Nomada*, we searched each genome using both the *cinA-cinB*<sup>*wYak-clade*</sup> contig and the flanking ISWpi1 elements. Long repeated elements like ISWpi1 (916 bp) break most short-read assemblies. Despite this, there is often a small





**Figure 5** The gene structure and observed, uncorrected, pairwise distances among wYak, wMel, and wPip *Wolbachia* spanning the region of the WO prophage island where the hypothesized transfer occurred between the ISWpi1 element “A” present in wYak and the ISWpi1 sequence labeled “B” in wMel. Over the region shown, wYak, wSan, and wTei are identical. The difference between wYak and wMel is <1% between the contigs (wYak “187383” and wYak “694583”) flanking the region containing the *cinA-cinB* homologs. These homologs assembled on the first ~4000 bp of the contig wYak “702380,” but they do not appear to exist in wMel. A BLAST search against the NCBI nucleotide collection using contig wYak “702380” as a query indicated the *cinA-cinB* homologs share high identity with group-B wPip *Wolbachia* (~97%). The remainder of the wYak contig is approximately equally divergent from both wMel and wPip, consistent with the entire region being transferred from an unknown *Wolbachia* variant that had acquired the *cinA-cinB* homologs. Under this scenario, ISWpi1 elements mediate the transfer to wYak via excision of elements “A” and “B” along with the intervening DNA from the donor and subsequent homologous recombination between IS sequences “A” and “B” and WO phage island genes in wYak. The ISWpi1 element “A” present in wYak, but not in either wMel or wPip, was spanned using Sanger sequencing with primers anchored on unique sequence in the flanking contigs (Sanger sequences are denoted by yellow lines with primer locations denoted by red flanking regions, Table S1). Additionally, the presence of a 236 bp tandem duplication in *cinB* (“C”) was confirmed using Sanger sequencing. The ISWpi1 sequence labeled “B” is assembled in wMel and corresponds to wMel#9 (Cordaux 2008) but could not be assembled or spanned with Sanger sequencing in the wYak genome. White gene models (arrows) indicate a partial transposase gene present in the WO prophage island, truncated in wYak. Pairwise differences were calculated using a sliding window (window size = 200 bp, step size = 25 bp).

fragment of the element, the length of the short read, on either end of the broken contig, indicative of the repeat element being responsible for terminating contig extension. We looked for the footprint of these elements at the edges of the contigs on which the *cin* genes were found. We found ISWpi1 elements in the region flanking both *cinA*<sup>wNLeu1</sup> and *cinA*<sup>wNLeu2</sup> copies, consistent with our wYak assembly in which we verified the ISWpi1 element with Sanger sequencing. These data support a role for ISWpi1 in the acquisition of the *cinA-cinB* genes by the *Wolbachia* in the *D. yakuba* clade and the *Nomada* bees. We conjecture that future work will fully confirm ISWpi1 in the horizontal movement of incompatibility loci between *Wolbachia*.

## Discussion

Our results indicate introgressive and horizontal *Wolbachia* acquisition in the *D. yakuba* clade. Evidence for horizontal *Wolbachia* transfer here and elsewhere (Turelli *et al.* 2018) suggests that double infections must be common, even if ephemeral. Such double infections provide an opportunity for ISWpi1 transposable elements to mediate horizontal transfer of incompatibility loci among divergent *Wolbachia*. Importantly, our results highlight that incompatibility factors may move independently of prophage, as evidenced by our discovery of the first prophage documented to have two sets of *cif* loci. We discuss these conclusions below.


### mtDNA introgression

Our relative mitochondrial chronogram provides strong support for three mitochondrial clades, including a monophyletic

*D. teissieri* clade that is outgroup to two sister clades: one clade consisting of mitochondria from 14 *D. santomea* individuals, and the other contains all *D. yakuba* mitochondria plus mitochondria from 12 *D. santomea* individuals. Our results suggest less mitochondrial introgression in the *D. yakuba* clade than a past report that used sequence data from only two mitochondrial loci (*COII* and *ND5*, 1777 bp) and reported a clade with mitochondria sampled from each species represented (Figure 3 in Bachtrog *et al.* 2006) (using sequence data from only *COII* and *ND5*, we can replicate this result, indicating data from additional loci are needed to add resolution). Our results also agree with recent work that demonstrated little nuclear introgression in the *D. yakuba* clade (Turissini and Matute 2017).

Our results broadly agree with the results produced by Llopart *et al.* (2014), who assessed mitochondrial introgression between *D. santomea* and *D. yakuba* using whole mitochondrial genomes. They generated a neighbor-joining tree that produced a clade consisting of all *D. yakuba* individuals and 10 *D. santomea* individuals, and this clade is sister to a clade with 6 *D. santomea* individuals; one *D. santomea* haplotype is outgroup to all other haplotypes included in their analysis (Figure 3 in Llopart *et al.* 2014). Nested within the mixed *D. yakuba* clade, Llopart *et al.* (2014) identified a “HZ clade” that includes *D. yakuba* individuals sampled from São Tomé and from continental Africa and *D. santomea* individuals sampled from both within and outside the Pico de São Tomé HZ. The sister *D. santomea* clade also contains both HZ and non-HZ individuals. Thus, their analysis and ours provide support for hybridization within and outside of the HZ, leading us to question both the existence of a HZ clade and





**Figure 6** The observed pairwise differences between *cinA* and *cinB* copies in *wYak*-clade *Wolbachia* and those in *wAlbB* and *wPip*, not including flanking regions, are shown. We also plot the pairwise differences between the two copies of these loci found in *Nomada Wolbachia* (denoted *wNleu1* and *wNleu2*); we chose *wNleu* to represent the *Nomada*-clade *Wolbachia* because this strain contains the longest assembled contigs for both *cinA-cinB* copies. Homologous *cinA-cinB* copies have relatively low divergence (inset unrooted tree), with the highest divergence in the 5' end of region. The most highly diverged region among any of the copies is the first ~750 bp of *wNleu1*, with pairwise differences between *wNleu1* and *wYak* reaching ~25% in some windows, with similar divergence between *wNleu1* and *wNleu2* in this same region. Pairwise

differences abruptly change 750 bp from the start of the *cinA* gene with the remaining ~1150 bp of the assembled region having a single difference from the *wYak* sequence (only ~1900 bp assembled into a single contig in the *wNleu1* copy). This pattern suggests a recent transfer/recombination event from the same unknown donor of the *wYak cinA-cinB* copy. The gap at position 1705–1940 represents a tandem duplication present in *wYak*. The unrooted tree was generated using RAxML 8.2.9 (Stamatakis 2014), and the representative *cinA-cinB* gene copies were obtained using BLAST to search all *Wolbachia* genomes in the NCBI assemblies database (<https://www.ncbi.nlm.nih.gov/assembly/?term=Wolbachia>).

the claim that these species “share the same mitochondrial genome” (Llopart *et al.* 2014). Instead, both our results and theirs suggest unidirectional introgression of *D. yakuba* mitochondria into *D. santomea*; we find 59% of *D. santomea* individuals having *D. yakuba*-like mitochondria and they found 46%.

Llopart *et al.* (2014) used a strict molecular clock to estimate the mitochondrial MRCA of *D. santomea* and *D. yakuba* at 10,792–17,888 years by calibrating their tree with the *D. yakuba-D. erecta* split, estimated at 10.4 MYA (Tamura *et al.* 2004). A high level of mitochondrial saturation over time, with an expected value of 1.44 substitutions per synonymous site for the *D. yakuba-D. erecta* split, could influence this estimate (Llopart *et al.* 2014). Moreover, Ho *et al.* (2005) demonstrated that the mtDNA substitution rate resembles an exponential curve, with high short-term substitution rates that approach the mutation rate, then slowing to a long-term rate after ~1–2 MY of divergence, far younger than the inferred *D. yakuba-D. erecta* nuclear and mtDNA divergence. Hence, using the slow long-term *D. yakuba-D. erecta* calibration is likely to underestimate the more rapid rates of divergence experienced by *D. yakuba*-clade mtDNA, inflating divergence-time estimates (Ho *et al.* 2005). If we assume that *Wolbachia* and mitochondria were transferred by introgression, which our analyses support, our estimates in Figure 3 suggest that *D. santomea* and *D. yakuba* mtDNA diverged more recently, with our point estimates ranging from ~1500 to 2800 years.

#### **Wolbachia placement, divergence, and acquisition**

**Wolbachia placement:** Despite their similarity, *wSan*, *wTei*, and *wYak* form monophyletic clades with *wTei* outgroup to sisters *wSan* and *wYak*, recapitulating host relationships (Figure 1; Turissini and Matute 2017). *wMel* is sister to the

*D. yakuba*-clade strains in the A group (Figure 2), which also includes *D. simulans* strains (*wRi*, *wAu*, and *wHa*), *wAna*, *wSuz*, and the *Nomada Wolbachia* (*wNFa*, *wNleu*, *wNPa*, and *wNFe*). Our phylogram (Figure 2A) places *wAlbB* outgroup to *wNo* and *wPip* strains that diverged from A-group *Wolbachia* ~6–46 MYA (Meany *et al.* 2019).

**Wolbachia divergence:** Our chronogram analyses (Figure 3) estimate that *D. yakuba*-clade *Wolbachia* and three *wMel* variants diverged ~29,000 years ago, and that *wTei* split from sisters *wSan* and *wYak* ~2500–4500 years ago, with *wSan* and *wYak* diverging ~1600–2800 years ago. We estimate that the two most divergent *wMel* variants from Richardson *et al.* (2012) and the reference *wMel* genome split ~4900–7200 years ago, indicating more divergence among *wMel* variants than among *D. yakuba*-clade *Wolbachia* strains. All of these results depend on the calibration provided by Richardson *et al.* (2012) and the relative accuracy of the underlying models of molecular evolution, which assume constant relative rates of change across data partitions. For the deepest divergence in Figure 3, between *D. yakuba*-clade *Wolbachia* and *wMel*, we find that the estimated time depends on the variance in our prior distribution for substitution rates across branches, with a strict clock putting the divergence at ~73,000 years rather than 29,000 years obtained with the most variable prior. Despite this uncertainty, the quantitative differences of our *Wolbachia* divergence-time estimates do not alter the qualitative conclusion that these *Wolbachia* did not codiverge with these hosts, which split several million years ago.

Our findings here and in Turelli *et al.* (2018) suggest that for several *Drosophila* species, their current *Wolbachia* infections have been in residence for only hundreds to tens of thousands of years. Bailly-Bechet *et al.* (2017) estimated

*Wolbachia* residence times using data from >10,000 arthropod specimens spanning over 1000 species. However, they analyzed DNA sequences from only part of the fast-evolving *fbpA* *Wolbachia* locus and the host *CO1* mtDNA locus. From an initial model-based meta-analysis, they concluded that “... most infections are very recent ...,” consistent with our results. However, they also fit a more complex model with “short” and “long” timescales for acquisition and loss, conjecturing that short-term rates were associated with imperfect maternal transmission. Focusing on long-time rates, they concluded that *Wolbachia* infections persisted in lineages for ~7 MY on average, whereas lineages remained uninfected for ~9 MY. For *Drosophila*, such long infection durations would imply that *Wolbachia*-host associations often persist through speciation [see Coyne and Orr (1997) and Turelli *et al.* (2014) for estimates of speciation times in *Drosophila*, generally  $10^5$ – $10^6$  years]. Extrapolating from limited *Wolbachia* sequence data, Hamm *et al.* (2014) conjectured that cladogenic *Wolbachia* transmission might be common among *Drosophila*, but this extrapolation is refuted by our genomic analyses. The long *Wolbachia* durations proposed Bailly-Bechet *et al.* (2017) depend on their conjecture that their long-term rate estimates accurately reflect acquisition and loss of *Wolbachia* infections across species. This is worth testing with additional analyses of *Wolbachia* and mitochondrial and nuclear genomes from a broad range of arthropods.

***Wolbachia* acquisition—introgression vs. horizontal:** Our divergence-time estimates for the *D. yakuba*-clade *Wolbachia* vs. their hosts preclude cladogenic acquisition. Unlike introgression, horizontal (or paternal) *Wolbachia* acquisition should produce discordance between phylogenies inferred for *Wolbachia* and the associated mitochondria. With the notable exception of *D. santomea* line Quija 37, which has mitochondria belonging to the clade associated with *D. yakuba*, but has *w*San *Wolbachia*, we find no evidence of discordance between the estimated mitochondrial and *Wolbachia* phylogenies. Hence our data indicate that acquisition by introgression is far more common than horizontal transmission between closely related species, consistent with data on acquisition of *w*Ri-like *Wolbachia* in the *D. melanogaster* species group (Turelli *et al.* 2018). Similarly, consistent with extensive data from *D. melanogaster* (Richardson *et al.* 2012) and *D. simulans* (Turelli *et al.* 2018) and smaller samples from *D. sukuzii* and *D. ananassae* (Turelli *et al.* 2018), we find only one possible example of paternal transmission or horizontal transmission within *D. yakuba*-clade species.

We also investigated an alternative approach to distinguish between introgressive and horizontal *Wolbachia* acquisition by estimating substitution ratios for mtDNA vs. *Wolbachia* (Turelli *et al.* 2018). Because we could estimate this ratio on each branch, we conjectured that this approach might have greater resolving power than our incompletely resolved mitochondrial and *Wolbachia* phylogenies. We expect higher ratios with horizontal transmission because mtDNA would

have been diverging longer than recently transferred *Wolbachia*. However, this approach assumes that mtDNA and *Wolbachia* substitution rates remain relatively constant. This is contradicted by the finding of Ho *et al.* (2005) that mtDNA substitution rates decline substantially with increasing divergence time, reaching an asymptote after ~1–2 MY. To calibrate their *Wolbachia* substitution rate estimates, Richardson *et al.* (2012) used an experimentally observed mitochondrial mutation rate in *D. melanogaster* ( $6.2 \times 10^{-8}$  mutations per third-position site per generation) that extrapolates to 62% third-position divergence per million years. This extrapolation is nonsensical as a long-term substitution rate. As summarized by Ho *et al.* (2005), typical mtDNA substitution rates are 0.5–1.5% per coding site per million years. Nevertheless, using the ratio of short-term rates for mtDNA and *Wolbachia*, Richardson *et al.* (2012) produced an estimate of the long-term *Wolbachia* substitution rate that agrees with independent estimates from *Nasonia* wasps (Raychoudhury *et al.* 2009) and *Nomada* bees (Gerth and Bleidorn 2016) derived from much longer divergence times (assuming cladogenic *Wolbachia* acquisition) (Conner *et al.* 2017). This paradox is resolved if *Wolbachia* molecular evolution is not subject to the dramatic slowdown in rates seen for mtDNA.

The apparent difference between mtDNA molecular evolution (dramatic slowdown over longer timescales; Ho *et al.* 2005) and *Wolbachia* molecular evolution (relative constancy, as inferred from similar rates of differentiation over very different timescales) suggests why our relative-rate test does not reject introgressive transmission of *Wolbachia* between *D. melanogaster* and the *D. yakuba* clade, even though it is clearly impossible. A roughly 50-fold slowdown in mtDNA substitution rates over the timescale of the divergence of *D. melanogaster* from the *D. yakuba* clade, relative to the rate of differentiation within the *D. yakuba* clade, produces comparable mtDNA-*Wolbachia* substitution ratios for comparisons within the *D. yakuba* clade and between *D. melanogaster* and the *D. yakuba* clade. Because of this complication and our conjecture that relative rates of mtDNA vs. *Wolbachia* substitutions over longer periods are likely to mirror the ten-fold differences we see for mtDNA vs. nuclear genes, phylogenetic discordance between mitochondria and *Wolbachia* is clearly a much more robust indicator of horizontal (or paternal) *Wolbachia* transmission. Nevertheless, additional examples of cladogenic *Wolbachia* acquisition are needed to better understand relative rates and patterns of *Wolbachia*, mtDNA and nuclear differentiation over different timescales.

Our divergence-times estimate of the *D. yakuba*-clade *Wolbachia* vs. their hosts precludes cladogenic transmission; and our phylogenetic analyses suggest that these species share very similar *Wolbachia* because of introgression, as originally argued by Lachaise *et al.* (2000). However, under either introgressive or horizontal transfer of *Wolbachia*, we expect the donor species *Wolbachia* sequences would appear paraphyletic when analyzed jointly with the *Wolbachia* from

the recipient. Paraphyly allowed Turelli *et al.* (2018) to infer that *D. simulans* likely obtained its *Wolbachia* from *D. ananassae* and *D. subpulchrella* likely obtained its *Wolbachia* from *D. sukuzii*. Paraphyly is generally expected soon after gene flow stops between populations. As noted by Hudson and Coyne (2002; Figure 1), the timescale expected to produce reciprocal monophyly for mitochondria (and *Wolbachia*) under a neutral model of molecular evolution is on the order of the effective size of the species. Our results in Figure 3 indicate that, at least for our small samples, reciprocal monophyly for the *Wolbachia* in these three species has been achieved within a few thousand years. This suggests that reciprocal monophyly has been accelerated by species-specific selective sweeps within the *Wolbachia* or mitochondria of these species. This conjecture may be testable from estimates of host fitness using transinfected *vs.* native *Wolbachia*.

### **IS transposable elements mediate horizontal transfer of incompatibility loci between divergent *Wolbachia***

*Wolbachia* in all three *D. yakuba*-clade hosts cause both intra- and interspecific CI (Cooper *et al.* 2017), despite originally being characterized as non-CI causing (Charlat *et al.* 2004; Zabalou *et al.* 2004). CI is relatively weak, and its strength can vary among *wTei* variants and *D. teissieri* backgrounds (Table 3 in Cooper *et al.* 2017). Differences in CI among *Wolbachia* variants has also been demonstrated in interspecific backgrounds where *wTei* caused stronger CI in a *D. simulans* background ( $97.2 \pm 1.3$  SE, percent embryo mortality) than either *wYak* ( $26.5 \pm 4.2$  SE, percent embryo mortality) or *wSan* ( $24.0 \pm 4.1$  SE, percent embryo mortality) (Zabalou *et al.* 2008). Both *wYak* and *wSan* induced CI in *D. simulans* comparable in intensity to that found by Cooper *et al.* (2017; Figure 3) in their original hosts. Surprisingly, Zabalou *et al.* (2008) found that the strength of CI induced by *wTei* in *D. simulans* even eclipsed that of *wRi* ( $89.8 \pm 4.5$  SE, percent embryo mortality). These results must be reconciled with the fact that loci known to underlie CI do not vary within or among *D. yakuba*-clade *Wolbachia* variants we examined. The nearly complete CI induced by *wTei* in *D. simulans* may depend on CI-causing factors yet to be identified or differences in gene expression.

In each *D. yakuba*-clade *Wolbachia* variant included in our analyses, we find a disruption of *cidB*<sup>*wYak*-clade</sup> with an inversion from amino acids 37–103 relative to the same region in sister *wMel*. The inversion introduces multiple stop codons that could render this gene nonfunctional. Fixation of loss-of-function mutations in CI-causing loci is consistent with theoretical analyses showing that selection on *Wolbachia* within host lineages does not act to increase or maintain CI (Prout 1994; Turelli 1994; Haygood and Turelli 2009); indeed, we have also recently observed a single mutation that disrupts *cidB* in non-CI causing *wMau* *Wolbachia* that infect *D. mauritiana* on the island of Mauritius (Meany *et al.* 2019). In both *wMau* and the *D. yakuba*-clade *Wolbachia*, we find fixation of defects in the putative toxin gene. We expect that future genomic analyses will produce additional examples.

All *D. yakuba*-clade *Wolbachia* genomes included in our analysis harbor *cinA-cinB* loci originally discovered in the *wPip* strain that diverged from A-group *Wolbachia*, including the *D. yakuba*-clade variants, ~6–46 MYA (Meany *et al.* 2019). *cin* loci are also present in B-group *wAlbB* that infects *Ae. albopictus* and in A-group *wNFe*, *wNPa*, *wNLeu*, and *wNFa* *Wolbachia* that infect *Nomada* bees. *cin* loci are absent from *wMel*, but the *wYak* contig containing these loci is ~10% diverged from *wMel*, while observed divergence between *wYak* and *wMel* across the rest of the genome is <1%. The *wYak*-clade *cin* loci share ~97% similarity with the divergent B-group *wPip* strain. *wYak*-clade *cin* loci are more similar to *cinA-cinB* from the B-group *Wolbachia* *wPip* and *wAlbB* than to those in A-group *Nomada* *Wolbachia* strains, which have two sets of *cin* loci that are as diverged from each other as they are from these regions in *wYak* and in B-group *Wolbachia*. These observations suggest independent horizontal transfer of *cin* loci into *wYak* and *Nomada* *Wolbachia*.

Our results indicate that independent of prophage movement, ISWpi1 element paralogs can move incompatibility loci via the excision of flanking ISWpi1 elements, followed by homologous recombination within the elements. Horizontal *Wolbachia* acquisition is common in *Drosophila* (Turelli *et al.* 2018) and other species (O'Neill *et al.* 1992), suggesting that double infections, which could provide the opportunity for ISWpi1-mediated transfer of incompatibility loci, may be common, even if transient (a second infection need not become stably transmitted for horizontal gene transfer via ISWpi1 elements to occur). In contrast, phage particles or virions could be introduced by a vector and provide the opportunity for ISWpi1-mediated transfer (Ahmed *et al.* 2015; Brown and Lloyd 2015), without the presence of a double *Wolbachia* infection. Determining whether the insertion of these loci was derived from a prophage region of the *Wolbachia* genome, or from a phage genome encapsulated in a phage particle, remains an open question. While the ISWpi1 element in *wMel* (Mel #9 labeled “B” in Figure 5; Cordaux 2008) is not part of our *wYak* assembly, homologs of this element are present in assemblies of several other A-group *Wolbachia* including *wInc* and *wRi* (Cordaux 2008; Bordenstein and Bordenstein 2016). We predict this element occurs in the unassembled region of our *wYak* assembly. Footprints of ISWpi1 elements in the region flanking the *cinA* genes for both copies of the gene in the *Nomada* *Wolbachia* provide further support for our hypothesis. Long-read-based *Wolbachia* assemblies from many infected host systems will elucidate the role of ISWpi1 elements in horizontal transfer of CI loci. Overall, the ecology of horizontal *Wolbachia* transmission is crucial to understanding *Wolbachia* acquisition, and the transfer and dynamics of CI loci are crucial to understanding *Wolbachia* evolution.

### **Acknowledgments**

We thank Tim Wheeler for help in the laboratory and for taking pictures of host species. John Beckmann, Amelia

Lindsey, Emily Delaney, Sylvain Charlat and three anonymous reviewers provided helpful comments. Research reported in this publication was supported by the National Institute Of General Medical Sciences of the National Institutes of Health (NIH) under award numbers R35GM124701 to B.S.C., R01GM121750 to D.R.M., and R01GM104325 to M.T. The content is solely the responsibility of the authors and does not necessarily represent the official views of the NIH.

## Literature Cited

- Ahmed, M. Z., S.-J. Li, X. Xue, X.-J. Yin, S.-X. Ren *et al.*, 2015 The intracellular bacterium *Wolbachia* uses parasitoid wasps as phoretic vectors for efficient horizontal transmission. *PLoS Pathog.* 11: e1004672. <https://doi.org/10.1371/journal.ppat.1004672>
- Bachtrog, D., K. Thornton, A. Clark, and P. Andolfatto, 2006 Extensive introgression of mitochondrial DNA relative to nuclear genes in the *Drosophila yakuba* species group. *Evolution* 60: 292–302. <https://doi.org/10.1111/j.0014-3820.2006.tb01107.x>
- Bailly-Bechet, M., P. Martins-Simoes, G. J. Szollosi, G. Mialdea, M.-F. Sagot *et al.*, 2017 How long does *Wolbachia* remain on board. *Mol. Biol. Evol.* 34: 1183–1193. <https://doi.org/10.1093/molbev/msx073>
- Bakovic, V., M. Schebeck, A. Telschow, C. Stauffer, and H. Schuler, 2018 Spatial spread of *Wolbachia* in *Rhagoletis cerasi* populations. *Biol. Lett.* 14: 20180161 (erratum: *Biol. Lett.* 14: 20180683). <https://doi.org/10.1098/rsbl.2018.0161>
- Baldo, L., J. C. D. Hotopp, K. A. Jolley, S. R. Bordenstein, S. A. Biber *et al.*, 2006 Multilocus sequence typing system for the endosymbiont *Wolbachia pipientis*. *Appl. Environ. Microbiol.* 72: 7098–7110. <https://doi.org/10.1128/AEM.00731-06>
- Baldo, L., N. A. Ayoub, C. Y. Hayashi, J. A. Russell, J. K. Stahlhut *et al.*, 2008 Insight into the routes of *Wolbachia* invasion: high levels of horizontal transfer in the spider genus *Agelenopsis* revealed by *Wolbachia* strain and mitochondrial DNA diversity. *Mol. Ecol.* 17: 557–569. <https://doi.org/10.1111/j.1365-294X.2007.03608.x>
- Bandi, C., T. J. Anderson, C. Genchi, and M. L. Blaxter, 1998 Phylogeny of *Wolbachia* in filarial nematodes. *Proc. Biol. Sci.* 265: 2407–2413. <https://doi.org/10.1098/rspb.1998.0591>
- Bankevich, A., S. Nurk, D. Antipov, A. A. Gurevich, M. Dvorkin *et al.*, 2012 SPAdes: a new genome assembly algorithm and its applications to single-cell sequencing. *J. Comput. Biol.* 19: 455–477. <https://doi.org/10.1089/cmb.2012.0021>
- Beckmann, J. F., and A. M. Fallon, 2013 Detection of the *Wolbachia* protein WPIP0282 in mosquito spermathecae: implications for cytoplasmic incompatibility. *Insect Biochem. Mol. Biol.* 43: 867–878. <https://doi.org/10.1016/j.ibmb.2013.07.002>
- Beckmann, J. F., J. A. Ronau, and M. Hochstrasser, 2017 A *Wolbachia* deubiquitylating enzyme induces cytoplasmic incompatibility. *Nat. Microbiol.* 2: 17007. <https://doi.org/10.1038/nmicrobiol.2017.7>
- Beckmann, J. F., M. Bonneau, H. Chen, M. Hochstrasser, D. Poinot *et al.*, 2019 The toxin-antidote model of cytoplasmic incompatibility: genetics and evolutionary implications. *Trends Genet.* 35: 175–185. <https://doi.org/10.1016/j.tig.2018.12.004>
- Bordenstein, S. R., and S. R. Bordenstein, 2016 Eukaryotic association module in phage WO genomes from *Wolbachia*. *Nat. Commun.* 7: 13155. <https://doi.org/10.1038/ncomms13155>
- Bouchon, D., T. Rigaud, and P. Juchault, 1998 Evidence for widespread *Wolbachia* infection in isopod crustaceans: molecular identification and host feminization. *Proc. Biol. Sci.* 265: 1081–1090. <https://doi.org/10.1098/rspb.1998.0402>
- Braig, H. R., W. Zhou, S. L. Dobson, and S. L. O'Neill, 1998 Cloning and characterization of a gene encoding the major surface protein of the bacterial endosymbiont *Wolbachia pipientis*. *J. Bacteriol.* 180: 2373–2378.
- Brand, C. L., S. B. Kingan, L. Wu, and D. Garrigan, 2013 A selective sweep across species boundaries in *Drosophila*. *Mol. Biol. Evol.* 30: 2177–2186. <https://doi.org/10.1093/molbev/mst123>
- Brown, A. N., and V. K. Lloyd, 2015 Evidence for horizontal transfer of *Wolbachia* by a *Drosophila* mite. *Exp. Appl. Acarol.* 66: 301–311. <https://doi.org/10.1007/s10493-015-9918-z>
- Brownlie, J. C., B. N. Cass, M. Riegler, J. J. Witsenburg, I. Iturbe-Ormaetxe *et al.*, 2009 Evidence for metabolic provisioning by a common invertebrate endosymbiont, *Wolbachia pipientis*, during periods of nutritional stress. *PLoS Pathog.* 5: e1000368. <https://doi.org/10.1371/journal.ppat.1000368>
- Camacho, C., G. Coulouris, V. Avagyan, N. Ma, J. Papadopoulos *et al.*, 2009 BLAST+: architecture and applications. *BMC Bioinformatics* 10: 421. <https://doi.org/10.1186/1471-2105-10-421>
- Carson, H. L., K. Y. Kaneshiro, and F. C. Val, 1989 Natural hybridization between the sympatric Hawaiian species *Drosophila silvestris* and *Drosophila heteroneura*. *Evolution* 43: 190–203. <https://doi.org/10.1111/j.1558-5646.1989.tb04217.x>
- Cattel, J., K. Nikolouli, T. Andrieux, J. Martinez, F. Jiggins *et al.*, 2018 Back and forth *Wolbachia* transfers reveal efficient strains to control spotted wing *Drosophila* populations. *J. Appl. Ecol.* 55: 2408–2418. <https://doi.org/10.1111/1365-2664.13101>
- Chandler, M., and J. Mahillon, 2002 Insertion sequences revisited, pp. 305–366 in *Mobile DNA II*, edited by N. Craig, R. Craigie, M. Gellert, and A. Lambowitz. ASM Press, Washington, DC. <https://doi.org/10.1128/9781555817954.ch15>
- Charlat, S., J. W. O. Ballard, and H. Merçot, 2004 What maintains noncytoplasmic incompatibility inducing *Wolbachia* in their hosts: a case study from a natural *Drosophila yakuba* population. *J. Evol. Biol.* 17: 322–330. <https://doi.org/10.1046/j.1420-9101.2003.00676.x>
- Chiu, J. C., X. Jiang, L. Zhao, C. A. Hamm, J. M. Cridland *et al.*, 2013 Genome of *Drosophila suzukii*, the spotted wing *Drosophila*. *G3 (Bethesda)* 3: 2257–2271. <https://doi.org/10.1534/g3.113.008185>
- Choi, J. Y., J. E. Bubnell, and C. F. Aquadro, 2015 Population genomics of infectious and integrated *Wolbachia pipientis* genomes in *Drosophila ananassae*. *Genome Biol. Evol.* 7: 2362–2382. <https://doi.org/10.1093/gbe/evv158>
- Comeault, A. A., A. Venkat, and D. R. Matute, 2016 Correlated evolution of male and female reproductive traits drive a cascading effect of reinforcement in *Drosophila yakuba*. *Proc. Biol. Sci.* 283: 20160730. <https://doi.org/10.1098/rspb.2016.0730>
- Conner, W. R., M. L. Blaxter, G. Anfora, L. Ometto, O. Rota-Stabelli *et al.*, 2017 Genome comparisons indicate recent transfer of wRi-like *Wolbachia* between sister species *Drosophila suzukii* and *D. subpulchrella*. *Ecol. Evol.* 7: 9391–9404. <https://doi.org/10.1002/ece3.3449>
- Cooper, B. S., P. S. Ginsberg, M. Turelli, and D. R. Matute, 2017 *Wolbachia* in the *Drosophila yakuba* complex: pervasive frequency variation and weak cytoplasmic incompatibility, but no apparent effect on reproductive isolation. *Genetics* 205: 333–351. <https://doi.org/10.1534/genetics.116.196238>
- Cooper, B. S., A. Sedghifar, W. T. Nash, A. A. Comeault, and D. R. Matute, 2018 A maladaptive combination of traits contributes to the maintenance of a *Drosophila* hybrid zone. *Curr. Biol.* 28: 2940–2947.e6. <https://doi.org/10.1016/j.cub.2018.07.005>
- Cordaux, R., 2008 ISWpi1 from *Wolbachia pipientis* defines a novel group of insertion sequences within the IS5 family. *Gene* 409: 20–27. <https://doi.org/10.1016/j.gene.2007.10.035>



- Coyne, J. A., and H. A. Orr, 1989 Patterns of speciation in *Drosophila*. *Evolution* 43: 362–381. <https://doi.org/10.1111/j.1558-5646.1989.tb04233.x>
- Coyne, J. A., and H. A. Orr, 1997 “Patterns of speciation in *Drosophila*” revisited. *Evolution* 51: 295–303.
- Coyne, J. A., S. Y. Kim, A. S. Chang, D. Lachaise, and S. Elwyn, 2002 Sexual isolation between two sibling species with overlapping ranges: *Drosophila santomea* and *Drosophila yakuba*. *Evolution* 56: 2424–2434. <https://doi.org/10.1111/j.0014-3820.2002.tb00168.x>
- Drosophila* 12 Genomes Consortium, A. G. Clark, M. B. Eisen, D. R. Smith, C. M. Bergman, B. Oliver *et al.*, 2007 Evolution of genes and genomes on the *Drosophila* phylogeny. *Nature* 450: 203–218. <https://doi.org/10.1038/nature06341>
- Duron, O., J. Lagnel, M. Raymond, K. Bourtzis, P. Fort *et al.*, 2005 Transposable element polymorphism of *Wolbachia* in the mosquito *Culex pipiens*: evidence of genetic diversity, superinfection and recombination. *Mol. Ecol.* 14: 1561–1573. <https://doi.org/10.1111/j.1365-294X.2005.02495.x>
- Ellegaard, K. M., L. Klasson, K. Näslund, K. Bourtzis, and S. G. E. Andersson, 2013 Comparative genomics of *Wolbachia* and the bacterial species concept. *PLoS Genet.* 9: e1003381. <https://doi.org/10.1371/journal.pgen.1003381>
- Garrigan, D., S. B. Kingan, A. J. Geneva, P. Andolfatto, A. G. Clark *et al.*, 2012 Genome sequencing reveals complex speciation in the *Drosophila simulans* clade. *Genome Res.* 22: 1499–1511. <https://doi.org/10.1101/gr.130922.111>
- Gerth, M., and C. Bleidorn, 2016 Comparative genomics provides a timeframe for *Wolbachia* evolution and exposes a recent biotin synthesis operon transfer. *Nat. Microbiol.* 2: 16241. <https://doi.org/10.1038/nmicrobiol.2016.241>
- Gill, A. C., A. C. Darby, and B. L. Makepeace, 2014 Iron necessity: the secret of *Wolbachia*'s success? *PLoS Negl. Trop. Dis.* 8: e3224. <https://doi.org/10.1371/journal.pntd.0003224>
- Giordano, R., S. L. O'Neill, and H. M. Robertson, 1995 *Wolbachia* infections and the expression of cytoplasmic incompatibility in *Drosophila sechellia* and *D. mauritiana*. *Genetics* 140: 1307–1317.
- Gloor, G. B., C. R. Preston, D. M. Johnsonschlitz, N. A. Nassif, R. W. Phillis *et al.*, 1993 Type-1 repressors of P-element mobility. *Genetics* 135: 81–95.
- Hamm, C. A., D. J. Begun, A. Vo, C. C. R. Smith, P. Saelao *et al.*, 2014 *Wolbachia* do not live by reproductive manipulation alone: infection polymorphism in *Drosophila suzukii* and *D. subpulchrella*. *Mol. Ecol.* 23: 4871–4885. <https://doi.org/10.1111/mec.12901>
- Havird, J. C., and D. B. Sloan, 2016 The roles of mutation, selection, and expression in determining relative rates of evolution in mitochondrial vs. nuclear genomes. *Mol. Biol. Evol.* 33: 3042–3053. <https://doi.org/10.1093/molbev/msw185>
- Haygood, R., and M. Turelli, 2009 Evolution of incompatibility-inducing microbes in subdivided host populations. *Evolution* 63: 432–447. <https://doi.org/10.1111/j.1558-5646.2008.00550.x>
- Hedges, L. M., J. C. Brownlie, S. L. O'Neill, and K. N. Johnson, 2008 *Wolbachia* and virus protection in insects. *Science* 322: 702. <https://doi.org/10.1126/science.1162418>
- Hilgenboecker, K., P. Hammerstein, P. Schlattmann, A. Telschow, and J. H. Werren, 2008 How many species are infected with *Wolbachia*? A statistical analysis of current data. *FEMS Microbiol. Lett.* 281: 215–220. <https://doi.org/10.1111/j.1574-6968.2008.01110.x>
- Ho, S. Y., M. J. Phillips, A. Cooper, and A. J. Drummond, 2005 Time dependency of molecular rate estimates and systematic overestimation of recent divergence times. *Mol. Biol. Evol.* 22: 1561–1568. <https://doi.org/10.1093/molbev/msi145>
- Hoffmann, A. A., 1988 Partial cytoplasmic incompatibility between two Australian populations of *Drosophila melanogaster*. *Entomol. Exp. Appl.* 48: 61–67. <https://doi.org/10.1111/j.1570-7458.1988.tb02299.x>
- Hoffmann, A. A., and M. Turelli, 1988 Unidirectional incompatibility in *Drosophila simulans*: inheritance, geographic variation and fitness effects. *Genetics* 119: 435–444.
- Hoffmann, A. A., and M. Turelli, 1997 Cytoplasmic incompatibility in insects, pp. 42–80 in *Influential Passengers: Inherited Microorganisms and Arthropod Reproduction*, edited by S. L. O'Neill, A. A. Hoffmann, and J. H. Werren. Oxford University Press, New York.
- Hoffmann, A. A., M. Turelli, and G. M. Simmons, 1986 Unidirectional incompatibility between populations of *Drosophila simulans*. *Evolution* 40: 692–701. <https://doi.org/10.1111/j.1558-5646.1986.tb00531.x>
- Hoffmann, A. A., D. J. Clancy, and E. Merton, 1994 Cytoplasmic incompatibility in Australian populations of *Drosophila melanogaster*. *Genetics* 136: 993–999.
- Hoffmann, A. A., D. Clancy, and J. Duncan, 1996 Naturally-occurring *Wolbachia* infection in *Drosophila simulans* that does not cause cytoplasmic incompatibility. *Heredity* 76: 1–8. <https://doi.org/10.1038/hdy.1996.1>
- Hoffmann, A. A., B. L. Montgomery, J. Popovici, I. Iturbe-Ormaetxe, P. H. Johnson *et al.*, 2011 Successful establishment of *Wolbachia* in *Aedes* populations to suppress dengue transmission. *Nature* 476: 454–457. <https://doi.org/10.1038/nature10356>
- Höhna, S., M. J. Landis, T. A. Heath, B. Boussau, N. Lartillot *et al.*, 2016 RevBayes: bayesian phylogenetic inference using graphical models and an interactive model-specification language. *Syst. Biol.* 65: 726–736. <https://doi.org/10.1093/sysbio/syw021>
- Hoskins, R. A., J. W. Carlson, K. H. Wan, S. Park, I. Mendez *et al.*, 2015 The Release 6 reference sequence of the *Drosophila melanogaster* genome. *Genome Res.* 25: 445–458. <https://doi.org/10.1101/gr.185579.114>
- Hudson, R. R., and J. A. Coyne, 2002 Mathematical consequences of the genealogical species concept. *Evolution* 56: 1557–1565. <https://doi.org/10.1111/j.0014-3820.2002.tb01467.x>
- Huigens, M. E., R. F. Luck, R. H. Klaassen, M. F. Maas, M. J. Timmermans *et al.*, 2000 Infectious parthenogenesis. *Nature* 405: 178–179. <https://doi.org/10.1038/35012066>
- Huigens, M. E., R. P. de Almeida, P. A. Boons, R. F. Luck, and R. Stouthamer, 2004 Natural interspecific and intraspecific horizontal transfer of parthenogenesis-inducing *Wolbachia* in *Trichogramma* wasps. *Proc. Biol. Sci.* 271: 509–515. <https://doi.org/10.1098/rspb.2003.2640>
- Hunter, S. S., R. T. Lyon, B. A. J. Sarver, K. Hardwick, L. J. Forney *et al.*, 2015 Assembly by reduced complexity (ARC): a hybrid approach for targeted assembly of homologous sequences. *bioRxiv*. doi:10.1101/014662 (Preprint posted January 31, 2015).
- Hurst, G. D. D., and F. M. Jiggins, 2000 Male-killing bacteria in insects: mechanisms, incidence, and implications. *Emerg. Infect. Dis.* 6: 329–336. <https://doi.org/10.3201/eid0604.000402>
- Jackman, S. D., B. P. Vandervalk, H. Mohamadi, J. Chu, S. Yeo *et al.*, 2017 ABySS 2.0: resource-efficient assembly of large genomes using a Bloom filter. *Genome Res.* 27: 768–777. <https://doi.org/10.1101/gr.214346.116>
- Jaenike, J., K. A. Dyer, C. Cornish, and M. S. Minhas, 2006 Asymmetrical reinforcement and *Wolbachia* infection in *Drosophila*. *PLoS Biol.* 4: e325 [corrigenda: *PLoS Biol.* 5: e3 (2007)]. <https://doi.org/10.1371/journal.pbio.0040325>
- Joshi, N. A., and J. N. Fass, 2011 Sickle: a sliding-window, adaptive, quality-based trimming tool for FastQ files. Available at: <https://scholar.google.com/scholar?cluster=6706699853004034677&hl=en&oi=scholar>
- Kamimura, Y., and H. Mitsumoto, 2012 Lock-and-key structural isolation between sibling *Drosophila* species. *Entomol. Sci.* 15: 197–201. <https://doi.org/10.1111/j.1479-8298.2011.00490.x>

- Katoh, K., and D. M. Standley, 2013 MAFFT Multiple sequence alignment software version 7: improvements in performance and usability. *Mol. Biol. Evol.* 30: 772–780. <https://doi.org/10.1093/molbev/mst010>
- Kern, A. D., D. A. Barbash, J. Chang Mell, A. Jensen, and D. Hupalo, 2015 Highly constrained intergenic *Drosophila* ultraconserved elements are candidate ncRNAs. *Genome Biol. Evol.* 7: 689–698. <https://doi.org/10.1093/gbe/evv011>
- Klasson, L., T. Walker, M. Sebahia, M. J. Sanders, M. A. Quail *et al.*, 2008 Genome evolution of *Wolbachia* strain wPip from the *Culex pipiens* group. *Mol. Biol. Evol.* 25: 1877–1887. <https://doi.org/10.1093/molbev/msn133>
- Klasson, L., J. Westberg, P. Sapountzis, K. Näslund, Y. Lutnaes *et al.*, 2009 The mosaic genome structure of the *Wolbachia* wRi strain infecting *Drosophila simulans*. *Proc. Natl. Acad. Sci. USA* 106: 5725–5730. <https://doi.org/10.1073/pnas.0810753106>
- Kondo, R., Y. Satta, E. T. Matsuura, H. Ishiwa, N. Takahata *et al.*, 1990 Incomplete maternal transmission of mitochondrial DNA in *Drosophila*. *Genetics* 126: 657–663.
- Kriesner, P., and A. A. Hoffmann, 2018 Rapid spread of a *Wolbachia* infection that does not affect host reproduction in *Drosophila simulans* cage populations. *Evolution* 72: 1475–1487. <https://doi.org/10.1111/evo.13506>
- Kriesner, P., A. A. Hoffmann, S. F. Lee, M. Turelli, and A. R. Weeks, 2013 Rapid sequential spread of two *Wolbachia* variants in *Drosophila simulans*. *PLoS Pathog.* 9: e1003607. <https://doi.org/10.1371/journal.ppat.1003607>
- Kriesner, P., W. R. Conner, A. R. Weeks, M. Turelli, and A. A. Hoffmann, 2016 Persistence of a *Wolbachia* infection frequency cline in *Drosophila melanogaster* and the possible role of reproductive dormancy. *Evolution* 70: 979–997. <https://doi.org/10.1111/evo.12923>
- Kulathinal, R. J., L. S. Stevison, and M. A. F. Noor, 2009 The genomics of speciation in *Drosophila*: diversity, divergence, and introgression estimated using low-coverage genome sequencing. *PLoS Genet.* 5: e1000550. <https://doi.org/10.1371/journal.pgen.1000550>
- Lachaise, D., M. Harry, M. Solognac, F. Lemeunier, V. Benassi *et al.*, 2000 Evolutionary novelties in islands: *Drosophila santomea*, a new *melanogaster* sister species from São Tomé. *Proc. Sci. Biol.* 267: 1487–1495. <https://doi.org/10.1098/rspb.2000.1169>
- Langley, C. H., and W. M. Fitch, 1974 An examination of the constancy of the rate of molecular evolution. *J. Mol. Evol.* 3: 161–177. <https://doi.org/10.1007/BF01797451>
- Langmead, B., and S. L. Salzberg, 2012 Fast gapped-read alignment with Bowtie 2. *Nat. Methods* 9: 357–359. <https://doi.org/10.1038/nmeth.1923>
- Laven, H., 1951 Crossing experiments with *Culex* strains. *Evolution* 5: 370–375. <https://doi.org/10.1111/j.1558-5646.1951.tb02795.x>
- Laven, H., 1967 Eradication of *Culex pipiens fatigans* through cytoplasmic incompatibility. *Nature* 216: 383–384. <https://doi.org/10.1038/216383a0>
- LePage, D. P., J. A. Metcalf, S. R. Bordenstein, J. On, J. I. Perlmutter *et al.*, 2017 Prophage WO genes recapitulate and enhance *Wolbachia*-induced cytoplasmic incompatibility. *Nature* 543: 243–247. <https://doi.org/10.1038/nature21391>
- Li, H., 2011 A statistical framework for SNP calling, mutation discovery, association mapping and population genetical parameter estimation from sequencing data. *Bioinformatics* 27: 2987–2993. <https://doi.org/10.1093/bioinformatics/btr509>
- Li, H., and R. Durbin, 2009 Fast and accurate short read alignment with Burrows-Wheeler transform. *Bioinformatics* 25: 1754–1760. <https://doi.org/10.1093/bioinformatics/btp324>
- Li, S.-J., M. Z. Ahmed, N. Lv, P.-Q. Shi, X.-M. Wang *et al.*, 2017 Plant-mediated horizontal transmission of *Wolbachia* between whiteflies. *ISME J.* 11: 1019–1028. <https://doi.org/10.1038/ismej.2016.164>
- Lindsey, A. R. I., D. W. Rice, S. R. Bordenstein, A. W. Brooks, S. R. Bordenstein *et al.*, 2018 Evolutionary genetics of cytoplasmic incompatibility genes *cifA* and *cifB* in prophage WO of *Wolbachia*. *Genome Biol. Evol.* 10: 434–451 [corrigenda: *Genome Biol. Evol.* 11: 1320 (2019)]. <https://doi.org/10.1093/gbe/evy012>
- Llopert, A., D. Herrig, E. Brud, and Z. Stecklein, 2014 Sequential adaptive introgression of the mitochondrial genome in *Drosophila yakuba* and *Drosophila santomea*. *Mol. Ecol.* 23: 1124–1136. <https://doi.org/10.1111/mec.12678>
- Lohse, K., M. Clarke, M. G. Ritchie, and W. J. Etges, 2015 Genome-wide tests for introgression between cactophilic *Drosophila* implicate a role of inversions during speciation. *Evolution* 69: 1178–1190. <https://doi.org/10.1111/evo.12650>
- Martinez, J., B. Longdon, S. Bauer, Y.-S. Chan, W. J. Miller *et al.*, 2014 Symbionts commonly provide broad spectrum resistance to viruses in insects: a comparative analysis of *Wolbachia* strains. *PLoS Pathog.* 10: e1004369. <https://doi.org/10.1371/journal.ppat.1004369>
- Matute, D. R., 2010 Reinforcement of gametic isolation in *Drosophila*. *PLoS Biol.* 8: e1000341. <https://doi.org/10.1371/journal.pbio.1000341>
- Matute, D. R., and J. F. Ayroles, 2014 Hybridization occurs between *Drosophila simulans* and *D. sechellia* in the Seychelles archipelago. *J. Evol. Biol.* 27: 1057–1068. <https://doi.org/10.1111/jeb.12391>
- Matute, D. R., and J. A. Coyne, 2010 Intrinsic reproductive isolation between two sister species of *Drosophila*. *Evolution* 64: 903–920. <https://doi.org/10.1111/j.1558-5646.2009.00879.x>
- Mavingui, P., C. Valiente Moro, V. Tran-Van, F. Wisniewski-Dye, V. Raquin *et al.*, 2012 Whole-genome sequence of *Wolbachia* strain wAlbB, an endosymbiont of tiger mosquito vector *Aedes albopictus*. *J. Bacteriol.* 194: 1840. <https://doi.org/10.1128/JB.00036-12>
- McMeniman, C. J., R. V. Lane, B. N. Cass, A. W. C. Fong, M. Sidhu *et al.*, 2009 Stable introduction of a life-shortening *Wolbachia* infection into the mosquito *Aedes aegypti*. *Science* 323: 141–144. <https://doi.org/10.1126/science.1165326>
- Meany, M. K., W. R. Conner, S. V. Richter, J. A. Bailey, M. Turelli *et al.*, 2019 Loss of cytoplasmic incompatibility and minimal fecundity effects explain relatively low *Wolbachia* frequencies in *Drosophila mauritiana*. *Evolution* 73: 1278–1295. <https://doi.org/10.1111/evo.13745>
- Moriyama, M., N. Nikoh, T. Hosokawa, and T. Fukatsu, 2015 Riboflavin provisioning underlies *Wolbachia*'s fitness contribution to its insect host. *MBio* 6: e01732–15. <https://doi.org/10.1128/mBio.01732-15>
- Nosil, P., 2013 Degree of sympatry affects reinforcement in *Drosophila*. *Evolution* 67: 868–872. <https://doi.org/10.1111/j.1558-5646.2012.01817.x>
- Nurk, S., A. Bankevich, D. Antipov, A. Gurevich, A. Korobeynikov *et al.*, 2013 Assembling genomes and mini-metagenomes from highly chimeric reads, pp. 158–170 in *Research in Computational Molecular Biology*, edited by M. Deng, R. Jiang, F. Sun, and X. Zhang. Springer, Berlin, Heidelberg.
- O'Connor, L., C. Plichart, A. C. Sang, C. L. Brelsfoard, H. C. Bossin *et al.*, 2012 Open release of male mosquitoes infected with a *Wolbachia* biopesticide: field performance and infection containment. *PLoS Negl. Trop. Dis.* 6: e1797. <https://doi.org/10.1371/journal.pntd.0001797>
- O'Neill, S. L., R. Giordano, A. M. Colbert, T. L. Karr, and H. M. Robertson, 1992 16S rRNA phylogenetic analysis of the bacterial endosymbionts associated with cytoplasmic incompatibility in insects. *Proc. Natl. Acad. Sci. USA* 89: 2699–2702. <https://doi.org/10.1073/pnas.89.7.2699>



- O'Neill, S. L., A. A. Hoffmann, and J. H. Werren, 1998 *Influent Passengers: Inherited Microorganisms and Arthropod Reproduction*. Oxford University Press, Oxford.
- Prout, T., 1994 Some evolutionary possibilities for a microbe that causes incompatibility in its host. *Evolution* 48: 909–911. <https://doi.org/10.1111/j.1558-5646.1994.tb01371.x>
- Raychoudhury, R., L. Baldo, D. C. S. G. Oliveira, and J. H. Werren, 2009 Modes of acquisition of *Wolbachia*: horizontal transfer, hybrid introgression, and codivergence in the *Nasonia* species complex. *Evolution* 63: 165–183. <https://doi.org/10.1111/j.1558-5646.2008.00533.x>
- Richardson, M. F., L. A. Weinert, J. J. Welch, R. S. Linheiro, M. M. Magwire *et al.*, 2012 Population genomics of the *Wolbachia* endosymbiont in *Drosophila melanogaster*. *PLoS Genet.* 8: e1003129. <https://doi.org/10.1371/journal.pgen.1003129>
- Ritchie, S. A., 2018 *Wolbachia* and the near cessation of dengue outbreaks in Northern Australia despite continued dengue importations via travellers. *J. Travel Med.* 25: tay084.
- Rousset, F., and M. Solignac, 1995 Evolution of single and double *Wolbachia* symbioses during speciation in the *Drosophila simulans* complex. *Proc. Natl. Acad. Sci. USA* 92: 6389–6393. <https://doi.org/10.1073/pnas.92.14.6389>
- Rousset, F., D. Bouchon, B. Pintureau, P. Juchault, and M. Solignac, 1992 *Wolbachia* endosymbionts responsible for various alterations of sexuality in arthropods. *Proc. Biol. Sci.* 250: 91–98. <https://doi.org/10.1098/rspb.1992.0135>
- Sánchez, L., and P. Santamaria, 1997 Reproductive isolation and morphogenetic evolution in *Drosophila* analyzed by breakage of ethological barriers. *Genetics* 147: 231–242.
- Schmidt, T. L., N. H. Barton, G. Rašić, A. P. Turley, B. L. Montgomery *et al.*, 2017 Local introduction and heterogeneous spatial spread of dengue-suppressing *Wolbachia* through an urban population of *Aedes aegypti*. *PLoS Biol.* 15: e2001894. <https://doi.org/10.1371/journal.pbio.2001894>
- Schrider, D. R., J. Ayroles, D. R. Matute, and A. D. Kern, 2018 Supervised machine learning reveals introgressed loci in the genomes of *Drosophila simulans* and *D. sechellia*. *PLoS Genet.* 14: e1007341. <https://doi.org/10.1371/journal.pgen.1007341>
- Schuler, H., K. Köppler, S. Daxböck-Horvath, B. Rasool, S. Krumböck *et al.*, 2016 The hitchhiker's guide to Europe: the infection dynamics of an ongoing *Wolbachia* invasion and mitochondrial selective sweep in *Rhagoletis cerasi*. *Mol. Ecol.* 25: 1595–1609. <https://doi.org/10.1111/mec.13571>
- Seemann, T., 2014 Prokka: rapid prokaryotic genome annotation. *Bioinformatics* 30: 2068–2069. <https://doi.org/10.1093/bioinformatics/btu153>
- Shoemaker, D. D., V. Katju, and J. Jaenike, 1999 *Wolbachia* and the evolution of reproductive isolation between *Drosophila recens* and *Drosophila subquinaria*. *Evolution* 53: 1157–1164. <https://doi.org/10.1111/j.1558-5646.1999.tb04529.x>
- Shropshire, J. D., J. On, E. M. Layton, H. Zhou, and S. R. Bordenstein, 2018 One prophage WO gene rescues cytoplasmic incompatibility in *Drosophila melanogaster*. *Proc. Natl. Acad. Sci. USA* 115: 4987–4991. <https://doi.org/10.1073/pnas.1800650115>
- Shropshire, J. D., B. Leigh, S. R. Bordenstein, A. Duploux, M. Riegler *et al.*, 2019 Models and nomenclature for cytoplasmic incompatibility: caution over premature conclusions – a response to Beckmann *et al.* *Trends Genet.* 35: 397–399. <https://doi.org/10.1016/j.tig.2019.03.004>
- Simão, F. A., R. M. Waterhouse, P. Ioannidis, E. V. Kriventseva, and E. M. Zdobnov, 2015 BUSCO: assessing genome assembly and annotation completeness with single-copy orthologs. *Bioinformatics* 31: 3210–3212. <https://doi.org/10.1093/bioinformatics/btv351>
- Siozios, S., A. Cestaro, R. Kaur, I. Pertot, O. Rota-Stabelli *et al.*, 2013 Draft genome sequence of the *Wolbachia* endosymbiont of *Drosophila suzukii*. *Genome Announc.* 1: e00032-13. <https://doi.org/10.1128/genomeA.00032-13>
- Stamatakis, A., 2014 RAXML version 8: a tool for phylogenetic analysis and post-analysis of large phylogenies. *Bioinformatics* 30: 1312–1313. <https://doi.org/10.1093/bioinformatics/btu033>
- Sutton, E. R., S. R. Harris, J. Parkhill, and S. P. Sinkins, 2014 Comparative genome analysis of *Wolbachia* strain wAu. *BMC Genomics* 15: 928. <https://doi.org/10.1186/1471-2164-15-928>
- Tamura, K., S. Subramanian, and S. Kumar, 2004 Temporal patterns of fruit fly (*Drosophila*) evolution revealed by mutation clocks. *Mol. Biol. Evol.* 21: 36–44.
- Teixeira, L., A. Ferreira, and M. Ashburner, 2008 The bacterial symbiont *Wolbachia* induces resistance to RNA viral infections in *Drosophila melanogaster*. *PLoS Biol.* 6: e1000002. <https://doi.org/10.1371/journal.pbio.1000002>
- Turelli, M., 1994 Evolution of incompatibility-inducing microbes and their hosts. *Evolution* 48: 1500–1513. <https://doi.org/10.1111/j.1558-5646.1994.tb02192.x>
- Turelli, M., and A. A. Hoffmann, 1991 Rapid spread of an inherited incompatibility factor in California *Drosophila*. *Nature* 353: 440–442. <https://doi.org/10.1038/353440a0>
- Turelli, M., and A. A. Hoffmann, 1995 Cytoplasmic incompatibility in *Drosophila simulans*: dynamics and parameter estimates from natural populations. *Genetics* 140: 1319–1338.
- Turelli, M., J. R. Lipkowitz, and Y. Brandvain, 2014 On the Coyne and Orr-igin of species: effects of intrinsic postzygotic isolation, ecological differentiation, X chromosome size, and sympatry on *Drosophila* speciation. *Evolution* 68: 1176–1187. <https://doi.org/10.1111/evo.12330>
- Turelli, M., B. S. Cooper, K. M. Richardson, P. S. Ginsberg, B. Peckenpaugh *et al.*, 2018 Rapid global spread of wRi-like *Wolbachia* across multiple *Drosophila*. *Curr. Biol.* 28: 963–971.e8. <https://doi.org/10.1016/j.cub.2018.02.015>
- Turissini, D. A., and D. R. Matute, 2017 Fine scale mapping of genomic introgressions within the *Drosophila yakuba* clade. *PLoS Genet.* 13: e1006971. <https://doi.org/10.1371/journal.pgen.1006971>
- Turissini, D. A., J. A. McGirr, S. S. Patel, J. R. David, and D. R. Matute, 2017 The rate of evolution of postmating-prezygotic reproductive isolation in *Drosophila*. *Mol. Biol. Evol.* 35: 312–334.
- Vavre, F., L. Mouton, and B. A. Pannebakker, 2009 Chapter 12 *Drosophila*–parasitoid communities as model systems for host–*Wolbachia* interactions, pp. 299–331 in *Advances in Parasitology*. Academic Press, San Diego, CA.
- Wallau, G. L., M. T. da Rosa, F. C. De Re, and E. L. Loreto, 2016 *Wolbachia* from *Drosophila incompta*: just a hitchhiker shared by *Drosophila* in the new and old world? *Insect Mol. Biol.* 25: 487–499. <https://doi.org/10.1111/imb.12237>
- Weeks, A. R., M. Turelli, W. R. Harcombe, K. T. Reynolds, and A. A. Hoffmann, 2007 From parasite to mutualist: rapid evolution of *Wolbachia* in natural populations of *Drosophila*. *PLoS Biol.* 5: e114. <https://doi.org/10.1371/journal.pbio.0050114>
- Weinert, L. A., E. V. Araujo-Jnr, M. Z. Ahmed, and J. J. Welch, 2015 The incidence of bacterial endosymbionts in terrestrial arthropods. *Proc. Biol. Sci.* 282: 20150249. <https://doi.org/10.1098/rspb.2015.0249>
- Werren, J. H., and D. M. Windsor, 2000 *Wolbachia* infection frequencies in insects: evidence of a global equilibrium? *Proc. Biol. Sci.* 267: 1277–1285. <https://doi.org/10.1098/rspb.2000.1139>
- Werren, J. H., W. Zhang, and L. R. Guo, 1995 Evolution and phylogeny of *Wolbachia*: reproductive parasites of arthropods. *Proc. Biol. Sci.* 261: 55–63. <https://doi.org/10.1098/rspb.1995.0117>

- Wu, M., L. V. Sun, J. Vamathevan, M. Riegler, R. Deboy *et al.*, 2004 Phylogenomics of the reproductive parasite *Wolbachia pipientis* wMel: a streamlined genome overrun by mobile genetic elements. *PLoS Biol.* 2: e69. <https://doi.org/10.1371/journal.pbio.0020069>
- Yen, J. H., and A. R. Barr, 1971 New hypothesis of the cause of cytoplasmic incompatibility in *Culex pipiens* L. *Nature* 232: 657–658. <https://doi.org/10.1038/232657a0>
- Yukilevich, R., 2012 Asymmetrical patterns of speciation uniquely support reinforcement in *Drosophila*. *Evolution* 66: 1430–1446. <https://doi.org/10.1111/j.1558-5646.2011.01534.x>
- Zabalou, S., S. Charlat, A. Nirgianaki, D. Lachaise, H. Merçot *et al.*, 2004 Natural *Wolbachia* infections in the *Drosophila yakuba* species complex do not induce cytoplasmic incompatibility but fully rescue the wRi modification. *Genetics* 167: 827–834. <https://doi.org/10.1534/genetics.103.015990>
- Zabalou, S., A. Apostolaki, S. Pattas, Z. Veneti, C. Paraskevopoulos *et al.*, 2008 Multiple rescue factors within a *Wolbachia* strain. *Genetics* 178: 2145–2160. <https://doi.org/10.1534/genetics.107.086488>

Communicating editor: M. Johnston