

Lawrence Berkeley National Laboratory

Recent Work

Title

THE DECAYS E- \rightarrow AoE-v

Permalink

<https://escholarship.org/uc/item/5f61c52t>

Authors

Hubbard, J. Richard

Berge, J. Peter

Dauber, Philip M.

Publication Date

1968-01-08

cy. J

University of California Ernest O. Lawrence Radiation Laboratory

THE DECAY $\Xi^- \rightarrow \Lambda^0 e^- \bar{\nu}$

J. Richard Hubbard, J. Peter Berge, and Philip M. Dauber

January 8, 1968

RECEIVED
LAWRENCE
RADIATION

LIBRARY
DOCUMENTS

TWO-WEEK LOAN COPY

*This is a Library Circulating Copy
which may be borrowed for two weeks.
For a personal retention copy, call
Tech. Info. Division, Ext. 5545*

UCRL-18025
cy. J

UCRL

DISCLAIMER

This document was prepared as an account of work sponsored by the United States Government. While this document is believed to contain correct information, neither the United States Government nor any agency thereof, nor the Regents of the University of California, nor any of their employees, makes any warranty, express or implied, or assumes any legal responsibility for the accuracy, completeness, or usefulness of any information, apparatus, product, or process disclosed, or represents that its use would not infringe privately owned rights. Reference herein to any specific commercial product, process, or service by its trade name, trademark, manufacturer, or otherwise, does not necessarily constitute or imply its endorsement, recommendation, or favoring by the United States Government or any agency thereof, or the Regents of the University of California. The views and opinions of authors expressed herein do not necessarily state or reflect those of the United States Government or any agency thereof or the Regents of the University of California.

Submitted to Physical Review Letters

UCRL-18025
Preprint

UNIVERSITY OF CALIFORNIA

Lawrence Radiation Laboratory
Berkeley, California

AEC Contract No. W-7405-eng-48

THE DECAY $\Xi^- \rightarrow \Lambda^0 e^- \bar{\nu}$

J. Richard Hubbard, J. Peter Berge, and Philip M. Dauber

January 8, 1968

THE DECAY $\Xi^- \rightarrow \Lambda^0 e^- \bar{\nu}$ *J. Richard Hubbard,[†] J. Peter Berge,[‡] and Philip M. DauberLawrence Radiation Laboratory
University of California
Berkeley, California

January 8, 1968

In the course of a search for leptonic decays of the Ξ hyperon, we have observed two unambiguous examples of

$$\Xi^- \rightarrow \Lambda^0 e^- \bar{\nu}, \quad (1)$$

and obtain a branching fraction of 1.0×10^{-3} for this mode. The Ξ^- were produced in a 27-event/ μb exposure of the Lawrence Radiation Laboratory 72-inch hydrogen bubble chamber to an incident K^- beam with momentum 1.7, 2.1, and 2.40 to 2.75 GeV/c. We have considered production events of the types

$$K^- p \rightarrow \Xi^- K^+ \quad (2a)$$

$$\Xi^- K^+ \pi^0 \quad (2b)$$

$$\Xi^- K^0 \pi^+ \quad (2c)$$

$$\Xi^- K^+ \pi^+ \pi^- \quad (2d)$$

$$\Xi^- K^0 \pi^+ \pi^0 \quad (2e)$$

where the decay kink of the Ξ^- and the decay Λ^0 were observed. Some 2823 events fitted one of reactions (2a) through (2e) as well as the normal decay sequence

$$\Xi^- \rightarrow \Lambda^0 \pi^-, \quad (3a)$$

$$\Lambda^0 \rightarrow p \pi^-, \quad (3b)$$

with confidence level $\geq 0.5\%$ for each of the three one-vertex fits.

Candidates for the beta decay mode (1) satisfied the following criteria:

- (a) Lambda decay, (3b), fits with confidence level $\geq 0.5\%$;
- (b) normal Ξ^- decay, (3a), does not fit; confidence level $\leq 0.5\%$;
- (c) a two-vertex fit to one of reactions (2a) through (2e) followed by the decay (1) is obtained with confidence level $\geq 0.5\%$.

Only nine events of the topologies giving rise to reactions (2a) through (2e) satisfied the criteria (a), (b), and (c). Of these, three events have negative tracks from the Ξ^- decay which are nearly flat in the chamber, have measured laboratory-system momenta less than 200 MeV/c, but are clearly darker than minimum-ionizing. A pion of 200 MeV/c or less has bubble density > 1.5 times that for the minimum-ionizing beam tracks, a difference distinguishable in our pictures for any but steeply dipping tracks. Another four events have charged Ξ^- decay tracks with momenta greater than 200 MeV/c. We have imposed the additional requirement that this momentum be less than 200 MeV/c for an event to be a candidate for (1). The remaining two events are shown in Fig. 1.

Each of the events in Fig. 1 is an unambiguous example of the decay (1). Event A fits $K^- p \rightarrow \Xi^- K^+$ with subsequent $\Xi^- \rightarrow \Lambda e^- \bar{\nu}$ decay with a four-constraint $\chi^2 = 1.45$. Event B¹ fits the same production-decay sequence with $\chi^2 = 0.34$. Neither event fits muonic decay $\Xi^- \rightarrow \Lambda \mu^- \bar{\nu}$, although four of the seven rejected events do fit this mode. None of these four events has been unambiguously identified as an example of Ξ muonic decay. In each of events A and B the electron tracks are nearly flat in the chamber, with measured momenta

100.3±1.2 and 134.6±1.8 MeV/c respectively. If the negative decay tracks were pions their relative ionizations would be 2.6 and 2.0 respectively (as muons 2.0 and 1.6); therefore their identification is unambiguous by ionization alone. In addition, event B has an 8-MeV "delta ray" on the negative decay track which also requires the track to be an electron. Neither event is a candidate for Ξ^- production and decay via $\Xi^- \rightarrow \Lambda e^- \bar{\nu}$. The measured transverse electron momenta (component of momentum of the decay track perpendicular to the Ξ^- direction) in each event--82.4±1.3 MeV/c for event A and 86.9±1.6 MeV/c for event B--are greater than the maximum of 79 MeV/c for $\Xi^- \rightarrow \Lambda$ decay. An attempt was nevertheless made to fit event A, in particular, to $\Xi^- \rightarrow \Lambda$ decay with production via both $K^- p$ and $\pi^- p$ interactions (the beam at 2.6 GeV/c included 13±3% π^-). No such fits were obtained. The relevant quantities in the identification of the events are summarized in Table I.

In order to calculate a branching fraction for the decay (1) based on our sample of two events we need only know the number of events (2823) fitting reactions (2a) through (2e), and the efficiency for detecting $\Xi^- \rightarrow \Lambda e^- \bar{\nu}$ with our selection criteria. The most important effect is the requirement which we impose that the electron momentum be less than 200 MeV/c. This eliminates about 25% of the events, according to a sample of the decay (1) Monte Carlo-generated by using a realistic Ξ^- momentum distribution and phase space for the electron momentum distribution in the Ξ^- rest frame. Only 6% of the Monte Carlo events fitted the normal decay hypothesis (3a). Neglecting possible small differences in the scanning efficiency for normal and leptonic Ξ^- decays, we arrive at an overall detection efficiency of 0.7±0.1 for the decay (1). Thus we obtain for the branching fraction

$$\frac{\Xi^- \rightarrow \Lambda e^- \bar{\nu}}{\text{all } \Xi^-} = \frac{2}{1976} = (1.0^{+1.3}_{-0.65}) \times 10^{-3}.$$

For such small numbers of events the usual $(N)^{1/2}$ approximation for the error is inappropriate, so we have used the Poisson distribution directly. The quoted errors correspond to fractions for which the observed number of events would represent a one-standard-deviation fluctuation from the true branching fraction.

Previous knowledge of the branching fraction for Ξ^- beta decay was based on one certain event found at UCLA² and one unambiguous plus one ambiguous event found at Brookhaven.³ A compilation⁴ based on the two unambiguous events and including several other experiments in which no leptonic Ξ^- decays were found yielded 790 events as an effective denominator. An additional sample of events at UCLA⁵ with incident K^- momentum 2.0 GeV/c has yielded no leptonic Ξ^- events, with an effective denominator of 717. If the available data are combined, there are four events in an effective sample of 3483 Ξ^- , which results in a fraction of $(1.15_{-0.55}^{+0.90}) \times 10^{-3}$. This value is probably a slight overestimate of the true fraction because experimenters with small samples and no leptonic events tend not to report their results. In addition, one should consider that an event with sufficiently low electron momentum might be detected in some experiments, including this one, despite the fact that it failed to satisfy all the criteria used to calculate the effective denominator. Such events have not been reported; this suggests that the true branching fraction is still lower than that obtained above. A value of 1.0×10^{-3} is probably realistic for the world average.

Our result for $\Xi^- \rightarrow \Lambda e^- \bar{\nu}$ / all Ξ^- may be compared with the prediction of the Cabibbo theory of leptonic decays.⁶ Fits to the theory for baryonic decays are tightly constrained by the data on Λ and Σ^- decay; the predicted

branching fraction for Ξ^- leptonic decay is only slightly affected by an input value for this fraction with a large uncertainty. Willis et al.,⁷ using $(2.4 \pm 1.4) \times 10^{-3}$ as input for the leptonic Ξ^- decay fraction, obtained two solutions in a fit to Cabibbo's theory. Solution A predicted 0.66×10^{-3} for this fraction; solution B gave 1.06×10^{-3} . Since then, solution B has been essentially eliminated by new data⁸ on the decay branching fraction for $\Sigma^- \rightarrow \Lambda e^- \bar{\nu}$. Several fits that closely correspond to solution A have been published; new data on leptonic baryon decay and modified versions of the theory were used. Brene et al.⁹ use a parameterization in terms of two angles θ_V and θ_A for the vector and axial-vector baryon currents, respectively, instead of Cabibbo's one angle θ . With an input fraction for $\Xi^- \rightarrow \Lambda e^- \bar{\nu}$ of $(1.2 \pm 0.8) \times 10^{-10}$ they obtain $(0.43 \pm 0.03) \times 10^{-3}$ for the predicted fraction. Carlson,¹⁰ using a one-angle theory, momentum-dependent form factors, and no input $\Xi^- \rightarrow \Lambda$ fraction, predicts 0.56×10^{-3} . Finally, a fit has been obtained,¹¹ including the $\Sigma^- \rightarrow \Lambda e^- \bar{\nu}$ data of Barash et al.⁸ which were not used in the earlier fits; this fit predicts 0.62×10^{-3} for the $\Xi^- \rightarrow \Lambda$ fraction, again using no input value.

The experimental branching fraction for $\Xi^- \rightarrow \Lambda e^- \bar{\nu}$ is therefore consistent with the Cabibbo theory and with data for other baryon leptonic decays.

Improved upper limits for other unusual decay modes of Ξ^- and Ξ^0 will be presented in a later paper along with a detailed analysis of the normal $\Xi \rightarrow \Lambda \pi$ decay.

The authors wish to thank Miss Jo Canada Cochran for her assistance in carrying out this work. The support and encouragement of Professor Luis W. Alvarez are gratefully acknowledged.

Footnotes and References

*Work done under the auspices of the U. S. Atomic Energy Commission.

† Now at Departement de Physique des Particules Elémentaires, CEN, Saclay, France.

‡ Now at the Nuclear Physics Laboratory, University of Oxford, Oxford, England.

1. Event B was discovered under a rather unusual circumstance. One of the authors (JRH) had just found event A on the scanning table after working halfway through a list of some 150 events of various kinds. Another of us (PMD) jokingly drew an arrow completely at random next to one of the remaining 81 events and declared that to be the next leptonic decay. Six hours later the event with the arrow was reached. It was event B.
2. D. Duane Carmony and Gerald M. Pierrou, Phys. Rev. Letters 10, 381 (1963).
3. G. W. London, R. R. Rau, N. P. Samios, S. S. Yamamoto, M. Goldberg, S. Lichtman, M. Primer, and J. Leitner, Phys. Rev. 143, 1034 (1966).
4. A. H. Rosenfeld, A. Barbaro-Galtieri, W. J. Podolsky, L. R. Price, P. Soding, C. G. Wohl, M. Roos, and W. J. Willis, Rev. Mod. Phys. 39, 1 (1967).
5. Thomas Trippe (UCLA), private communication.
6. N. Cabibbo, Phys. Rev. Letters 10, 531 (1963).
7. W. Willis et al., Phys. Rev. Letters 13, 291 (1964).
8. N. Barash, T. B. Day, R. G. Glasser, B. Kehoe, R. Knop, B. Sechi-Zorn, and G. A. Snow, Phys. Rev. Letters 19, 181 (1967).
9. N. Brene, L. Veje, M. Roos, and C. Cronström, Phys. Rev. 149, 1288 (1966).

10. C. Carlson, Phys. Rev. 152, 1433 (1966).
11. Lawrence K. Gershwin, (Lawrence Radiation Laboratory), private communication. The version of the theory used was that of Willis et al.


Figure Caption

Fig. 1. Two examples of $K^- p \rightarrow \Xi^- K^+$ followed by the decay $\Xi^- \rightarrow \Lambda e^- \bar{\nu}$; on the left, event A and on the right, event B.

Table I. Fitted quantities^a for Ξ^- beta decay events.

Quantity	Event A	Event B
Beam K^- momentum (MeV/c)	2633±24	1718±13
Ξ^- momentum	1209±10	1666±13
K^+ momentum	1758±18	341±5
e^- momentum	100± 1	135±2
Transverse e^- momentum		
(unfitted)	82.4 ±1.3	86.9 ±1.6
e^- dip	-4±1°	11 ±1°
χ^2 (3C) $\Lambda \rightarrow p \pi^-$	1.65	1.89
χ^2 (4C) $K^- p \rightarrow \Xi^- K^+$	1.45	0.34
↳ $\Lambda e^- \bar{\nu}$		
↳ $\Lambda \mu^- \bar{\nu}$	162.40	69.20
χ^2 (7C) 3 vertex fit to		
$K^- p \rightarrow \Xi^- K^+$		
↳ $\Lambda e^- \bar{\nu}$		
↳ $p \pi^-$	3.86	2.54
Overall confidence level	80%	92%
χ^2 (3C) $\Xi^- \rightarrow \Lambda \pi^-$	380.38	314.38

a. The fitted momenta and angles are insignificantly different from the measured, or unfitted, quantities except for the momentum of the short Ξ^- , which is essentially undetermined without fitting.



XBB 6711-6521

Fig. 1

This report was prepared as an account of Government sponsored work. Neither the United States, nor the Commission, nor any person acting on behalf of the Commission:

- A. Makes any warranty or representation, expressed or implied, with respect to the accuracy, completeness, or usefulness of the information contained in this report, or that the use of any information, apparatus, method, or process disclosed in this report may not infringe privately owned rights; or
- B. Assumes any liabilities with respect to the use of, or for damages resulting from the use of any information, apparatus, method, or process disclosed in this report.

As used in the above, "person acting on behalf of the Commission" includes any employee or contractor of the Commission, or employee of such contractor, to the extent that such employee or contractor of the Commission, or employee of such contractor prepares, disseminates, or provides access to, any information pursuant to his employment or contract with the Commission, or his employment with such contractor.

