

# Lawrence Berkeley National Laboratory

## Recent Work

### **Title**

THE PHYSICS OF A REGGE TRAJECTORY WITH A FINITE ASYMPTOTIC PHASE

### **Permalink**

<https://escholarship.org/uc/item/5fd2838d>

### **Author**

Sivers, Dennis.

### **Publication Date**

1970-11-01

Submitted to Physical Review;  
also filed as a Ph. D. Thesis

UCRL-20162  
Preprint

THE PHYSICS OF A REGGE TRAJECTORY WITH A  
FINITE ASYMPTOTIC PHASE

Dennis Sivers

November 17, 1970

AEC Contract No. W-7405-eng-48

TWO-WEEK LOAN COPY

*This is a Library Circulating Copy  
which may be borrowed for two weeks.  
For a personal retention copy, call  
Tech. Info. Division, Ext. 5545*

LAWRENCE RADIATION LABORATORY  
UNIVERSITY of CALIFORNIA BERKELEY

UCRL-20162

## **DISCLAIMER**

This document was prepared as an account of work sponsored by the United States Government. While this document is believed to contain correct information, neither the United States Government nor any agency thereof, nor the Regents of the University of California, nor any of their employees, makes any warranty, express or implied, or assumes any legal responsibility for the accuracy, completeness, or usefulness of any information, apparatus, product, or process disclosed, or represents that its use would not infringe privately owned rights. Reference herein to any specific commercial product, process, or service by its trade name, trademark, manufacturer, or otherwise, does not necessarily constitute or imply its endorsement, recommendation, or favoring by the United States Government or any agency thereof, or the Regents of the University of California. The views and opinions of authors expressed herein do not necessarily state or reflect those of the United States Government or any agency thereof or the Regents of the University of California.

THE PHYSICS OF A REGGE TRAJECTORY WITH A FINITE  
ASYMPTOTIC PHASE

Contents

Abstract . . . . .	v
I. Introduction . . . . .	1
II. Parametrization of a Regge Trajectory in the Phase Representation . . . . .	3
III. The Kinematics of the Decay of a High Spin Resonance . . . . .	10
IV. Implications for Dynamical Models . . . . .	15
V. Acknowledgment . . . . .	21
Appendix A: The Theorem of Brower and Harte . . . . .	22
Footnotes and References . . . . .	25
Figure Captions . . . . .	28

THE PHYSICS OF A REGGE TRAJECTORY WITH A FINITE  
ASYMPTOTIC PHASE\*

Dennis Sivers†

Department of Physics and Lawrence Radiation Laboratory  
University of California, Berkeley, California 94720

November 17, 1970

ABSTRACT

The hypothesis that the width of a resonance on a leading Regge trajectory is proportional to its mass is compared to the alternative that high-spin resonances become very stable. When the width grows with the mass, it is observed that the kinematics allow decay channels to have low orbital angular momentum. The dynamical implications of this observation are discussed.

## I. INTRODUCTION

Taking the limit of zero-width resonances is an extremely delicate process. In spite of the attractive simplicity of the Veneziano model,<sup>1</sup> detailed analyses of its predictions always conclude that a description of hadronic scattering processes involving meromorphic amplitudes is strained and artificial.<sup>2</sup> As an example of the subtle nature of the narrow resonance limit, consider a Regge trajectory with a small but finite asymptotic phase,

$$\delta(\infty) = \lim_{t \rightarrow \infty} \arctan[\text{Im } \alpha(t)/\text{Re } \alpha(t)]. \quad (1.1)$$

A high-spin particle on such a trajectory can communicate with open channels of low angular momentum while, if the asymptotic phase is zero, these channels are closed kinematically.<sup>3</sup>

It is entirely possible that the dynamics of two cases will be completely different. Observations based on coupled-channel unitarity approximations suggest strongly that the dynamics are dominated by low-orbital-angular-momentum channels. In order to maintain itself, an asymptotically real Regge trajectory could require a rich spectrum of nonleading singularities.<sup>4</sup> In contrast, the dynamics of a leading trajectory with finite asymptotic phase can be compatible with a model proposed by Carruthers<sup>5</sup> where a particle of spin  $J$  on the leading trajectory is primarily a bound state in the system formed by a particle of spin  $J-1$  on the same trajectory and an arbitrary, low-mass particle. In this sort of model, there is no need for all nonleading Regge singularities to rise to high values of  $J$ .

Experimental evidence on baryon widths is consistent with these states lying on trajectories with finite asymptotic phase. There is also some indication that this notion is applicable to meson trajectories even though it implies that the very narrow peaks in the CERN missing mass spectrometer<sup>6</sup> should not be identified with recurrances of the  $\rho$ -f trajectory. Instead, these states would represent fine structure in the meson spectrum, perhaps connected with the dip in the  $A_2$  peak.<sup>7</sup>

## II. PARAMETRIZATION OF A PHYSICAL REGGE TRAJECTORY

Consider a Regge trajectory which, at  $t = 0$ , is the leading J-plane singularity in a nonvacuum channel. For simplicity, ignore the inessential complications of signature and assume the channel has baryon number zero. A parametrization which is particularly convenient for a discussion of the asymptotic behavior of the trajectory function is the phase representation.<sup>8,9</sup>

Let

$$\alpha(t \pm i0) = |\alpha(t)| \exp\{\pm i\delta(t)\}, \quad t \geq 4m^2, \quad (2.1)$$

where  $\delta(t)$  is the phase of the Regge trajectory above the physical cut beginning at  $t = 4m^2$ . The phase representation for  $\alpha(t)$  is then given by

$$\alpha(t) = P(t) \exp \left\{ \frac{t}{\pi} \int_{4m^2}^{\infty} \frac{\delta(x) dx}{x(x-t)} \right\} \quad (2.2)$$

where  $P(t)$  is a polynomial. The representation (2.2) is valid if:<sup>10</sup>

(a)  $\alpha(t)$  is analytic in  $t$  except for a cut along the real axis.

(b)  $\alpha(t)$  is real in the sense  $\alpha^*(t) = \alpha(t^*)$ .

(c)  $\alpha(t)$  is bounded as  $|t| \rightarrow \infty$  on the physical sheet by a finite polynomial in  $t$ .

(d) The phase above the physical cut,  $\delta(t)$ , goes to a finite limit as  $t \rightarrow \infty + i0$ .



The conditions (a) and (b) above are quite generally true for meson trajectories<sup>11</sup> if the singularity surface represented by  $\alpha(t)$  does not collide with any other singularity in the partial wave amplitude. For fermion trajectories, the analyticity condition for the trajectory function is usually derived in terms of the variable,  $W = t^{\frac{1}{2}}$ , but the situation is complicated by the cuts which seem to be present to shield the parity partners implied by MacDowell symmetry.<sup>12</sup> It is possible that, even in meson channels, trajectory functions develop extra cuts associated with the collision of different singularity surfaces.<sup>13,14</sup> The representation, (2.2), can be extended to include contributions from other cuts but it is a sensible first step to deal with trajectory functions having only the physical threshold cut and a discussion of possible complications due to collisions will be deferred to the final section.

Condition (c) seems a reasonable restriction on the asymptotic behavior of physically interesting trajectory functions. Although there is no a priori reason to prohibit exponential growth of the trajectory function in certain sections of the  $t$  plane, such behavior is unexpected from experimental grounds. The phase,  $\delta(t)$ , of the trajectory function is well defined along the physical cut, except, possibly at those points where  $|\alpha(t)| = 0$ . The phase can be shown to be piecewise continuous, and it makes physical sense to limit the magnitude of discontinuities in  $\delta(t)$  to be less than  $\pi$ ,

$$\lim_{\eta \rightarrow 0} |\delta(t_0 - \eta) - \delta(t_0 + \eta)| < \pi. \quad (2.3)$$

Consistency with the analytic continuation of the unitarity condition requires

$$\text{Im } \alpha(t + i0) > 0, \quad t > 4m^2 \quad (2.4)$$

unless at the point where  $\text{Im } \alpha = 0$  on the physical cut, the residue function associated with  $\alpha(t)$  also vanishes. In conjunction with the requirement that there can be no resonance poles on the physical sheet, it is frequently conjectured that (2.4) holds for all  $t$  above threshold.<sup>11</sup> Except at a point where  $\text{Re } \alpha = 0$ , this constraint then leads to the bound

$$0 \leq \delta(t) \leq \pi. \quad (2.5)$$

Condition (d) seems a reasonable assumption in view of (2.5). This assumption only eliminates the possibility that the phase oscillates as  $t \rightarrow \infty$  without approaching a definite limit.

Using the conditions, (a) - (d), plus the extra assumption that the phase is consistent with the usual unitarity conditions at the elastic threshold,<sup>15</sup> Childers<sup>8</sup> was able to show that the polynomial,  $P(t)$ , in (2.2) has one and only one zero so that it must be of order one,

$$P(t) = a + bt. \quad (b \neq 0) \quad (2.6)$$

Except for possible logarithmic factors, the asymptotic behavior of the exponential in (2.2) is determined by a single parameter,  $\delta(\infty)$ , the asymptotic limit of the phase.<sup>9</sup> Making use of the result (2.6), the asymptotic behavior of the trajectory function in (2.2) is given by

$$\lim_{|t| \rightarrow \infty} \alpha(t) = b e^{i\delta(\infty)} t^{1-\delta(\infty)/\pi} \quad (2.7)$$

uniformly on the first sheet. This means, for example, that if the trajectory rises indefinitely ( $\text{Re } \alpha \rightarrow +\infty$  as  $t \rightarrow +\infty$ ) faster than a power of a logarithm, then it also falls indefinitely ( $\text{Re } \alpha \rightarrow -\infty$  as  $t \rightarrow -\infty$ ).

Dynamical trajectory functions found in potential scattering<sup>16</sup> and in those approaches such as the N/D model<sup>17</sup> which attempt to saturate the unitarity condition with a small number of internal channels are consistent with the prediction (2.7) in that they have

$$\delta(\infty) = \pi \quad (2.8)$$

and they approach negative integers as  $|t| \rightarrow \infty$ . The strictly real, linear trajectory functions used as input in the Veneziano model are also trivially consistent with (2.7). The convenience of the phase representation for  $\alpha(t)$  is seen in the fact that a wide class of possibilities are covered by the single formula, (2.2). This contrasts to the parametrization of  $\alpha(t)$  in terms of a dispersion relation<sup>18,19</sup> such as

$$\alpha(t) = \alpha(0) + t \left[ d + \frac{1}{\pi} \int_{4m^2}^{\infty} dx \frac{\text{Im } \alpha(x)}{x(x-t)} \right] \quad (2.9)$$

with  $\alpha(0)$ ,  $d$  real and  $d \geq 0$ , where the form changes according to whether or not  $d = 0$ . The conditions under which  $d$  can be nonzero are expressed in terms of  $\text{Im}(\alpha)$  and are quite complicated.<sup>20</sup>

Figure 1 shows a plot of the phase of the  $\Delta$  trajectory against  $t$ . The points are determined from the resonance parameters given in the Particle Data Group<sup>21</sup> compilations. The plot is consistent with a phase which quickly approaches a small constant although, on the basis of the data, it is not possible to eliminate the situation where the phase continues to rise so that the trajectory eventually turns over. The linear extrapolation included on this graph gives an approximate lower bound on the energy where such a turnover could occur.<sup>22</sup> In view of the fact that the analyticity properties of fermion trajectories are expected to be more complicated than those assumed in (2.2), the relation between the phase of the  $\Delta$  trajectory and the asymptotic behavior implied by (2.7) should not be taken too seriously.

Figure 2 shows a plot of the phase of an exchange-degenerate  $\rho$ -f trajectory with the points again determined from the Particle Data Group compilations.<sup>21</sup> If the very narrow peaks, the T(2200) and the U(2375), found by the CERN missing mass spectrometer,<sup>6</sup> are identified as recurrences of this trajectory then the data suggests the phase may go to zero. However, there is some reason to doubt that a missing mass spectrometer could discriminate peaks with widths 190 - 220 MeV from the background in this region so the evidence is not conclusive that there are not wide resonances in the T and U regions.<sup>23</sup> In fact, the  $\rho(2275)$ , the  $\bar{N}\bar{N}_{I=1}(2345)$ , and the  $\bar{N}\bar{N}_{I=0}(2380)$  are candidates. The question then becomes one of understanding what the narrow peaks are. In view of present models for the  $A_2$  fine structure, it seems plausible that there should exist many narrow "doorway" states in the meson

spectrum which coexist with wide resonances.<sup>7,23</sup> Perhaps these are what the CERN spectrometer is seeing. It would certainly be an unattractive theoretical situation if the  $A_2$  were unique, but the question has ultimately to be resolved experimentally. What is important here is that there may be two alternatives for the asymptotic phase of the  $\rho$ -f trajectory. If

$$\delta(\infty)/\pi = \epsilon \ll 1 \quad (2.10)$$

the states on this trajectory would grow to be very wide. If

$$\delta(\infty)/\pi = 0 \quad (2.11)$$

then the large widths of the low-spin resonances such as the  $\rho$ ,  $f$ , and  $g$  are anomalous and the high-spin recurrences will become very stable. The physical consequences of the two alternatives, (2.10) and (2.11), are quite distinct. In particular, a high-spin state on a trajectory with a finite phase can decay into open channels with low orbital angular momentum while in the case (2.11), a theorem due to Brower and Harte<sup>3</sup> shows that such decays are kinematically forbidden.

This brings up one of the anomalies of the Veneziano model<sup>1,2</sup> which may give a hint that the concept of an asymptotic phase has physical relevance. Although the model requires strictly real, non-physical trajectories, it only has Regge asymptotic behavior in a region excluding a wedge of finite axis along the real axis of the  $t$  plane where there is a line of poles.<sup>24</sup> There may be some confusion on this point since Roskies,<sup>24</sup> who bases his argument on the requirement

that the Beta function  $B[-\alpha(t), -\alpha(u)]$  decreases exponentially at fixed  $s$ , has quoted this requirement in the form

$$\text{Im } \alpha(t)/|t|^{1+\mu} \rightarrow \infty, \quad \text{all } \mu > 0 \quad (2.12)$$

which depends on the fact that he used the dispersion relation (2.9), for  $\alpha(t)$  so that  $\text{Re } \alpha \sim ct$ . Since the argument of the Beta function depends on  $t$  only parametrically on  $t$  through  $\alpha(t)$ , it is obvious that the bound

$$\text{Im } \alpha(t)/|\text{Re } \alpha(t)|^{1+\mu} \rightarrow \infty, \quad \text{all } \mu > 0 \quad (2.13)$$

would satisfy Roskies' constraint. Equation (2.12) is essentially the condition that the trajectory phase be finite asymptotically or fall no faster than an inverse power of a logarithm.

In terms of the interpretation of the role of small parameters in physics presented by Chew,<sup>25</sup> it is the smallness of  $\epsilon$  which justifies the use of the Veneziano model. It is not surprising that the physics of the two cases (2.10) and (2.11) should be quite different so that a simple extrapolation procedure should fail.

### III. THE KINEMATICS OF A DECAY

Assume that the behavior of the Regge trajectory is given asymptotically by (2.7) and (2.9). This then gives the expression for the mass of the particle of spin  $J$

$$\lim_{J \rightarrow \infty} M_J = \left[ \frac{J}{b \cos(\pi\epsilon)} \right]^{1/(2-2\epsilon)} \quad (3.1)$$

Because the phase,  $\pi\epsilon = \delta(\infty)$ , of the trajectory is assumed to be small so that

$$(\text{Im } \alpha)^2 / (\text{Re } \alpha)^2 \ll 1, \quad (3.2)$$

the relation between the width of the resonance and the imaginary part of the trajectory is given by

$$\Gamma_J = \frac{\text{Im } \alpha(M_J^2)}{\left[ \frac{d}{dt} \text{Re } \alpha(t) \Big|_{t=M_J^2} \right] \times M_J} \quad (3.3)$$

The reader may wonder about the validity of an expression such as (3.3) which is based on the Breit-Wigner formula when it is used in a situation where the width it gives is large compared to the spacing of singularities. A discussion of overlapping resonances which relies on simple quasi-two body unitarity has been presented by Coleman.<sup>26</sup> The situation for the case of many body unitarity may be more complicated but it remains plausible that (3.3) should give approximately the imaginary part of the pole in the complex energy plane in spite of the spacing of nearby singularities. Of course, as the "lifetime" predicted by

(3.3) becomes shorter, it becomes harder and harder to design an experiment which would measure the quantum numbers of the resonance so that the "particle" connection of the pole is gradually lost and it becomes indistinguishable from "nonresonant background."

Combining (3.3) with (2.7) gives

$$\Gamma_J = \frac{\tan(\pi\epsilon)}{1-\epsilon} \left[ \frac{J}{b \cos(\pi\epsilon)} \right]^{1/(2-2\epsilon)} \equiv \mathcal{M}_J. \quad (3.4)$$

Under the assumption that (3.4) does not hold, but instead the trajectory is asymptotically real in the sense

$$\lim_{J \rightarrow \infty} \Gamma_J / \mathcal{M}_J = 0, \quad (3.5)$$

Brower and Harte<sup>3</sup> showed that kinematic constraints decouple a high spin resonance from all available channels with low orbital angular momentum unless

$$\lim_{t \rightarrow \infty + i0} \operatorname{Re} \alpha(t) = o(t^{\frac{1}{2}}). \quad (3.6)$$

Of course, the behavior implied by (3.5) and (3.6) is inconsistent with the asymptotic behavior implied by the phase representation (2.2) or the dispersion relation (2.9) and so, in the absence of left-hand cuts in the trajectory function, the bound (3.6) will not be considered further.

Consider the decay of a particle of spin  $J$  on the trajectory given by (2.7). Suppose it decays into two particles of spin  $j_1$  and  $j_2$  on the same trajectory where



$$j_1 = \eta J \quad (3.7a)$$

and

$$j_2 = (1 - \eta)J. \quad (3.7b)$$

Conservation of energy permits the decay if

$$M_J + \Gamma_J \geq M_{j_1} - \Gamma_{j_1} + M_{j_2} - \Gamma_{j_2}, \quad (3.8)$$

$$\begin{aligned} (1 + r) \left[ \frac{J}{b \cos(\pi\epsilon)} \right]^{1/(2-2\epsilon)} \\ \geq (1 - r) \left[ \eta^{1/(2-2\epsilon)} + (1 - \eta)^{1/(2-2\epsilon)} \right] \left[ \frac{J}{b \cos(\pi\epsilon)} \right]^{1/(2-2\epsilon)} \end{aligned} \quad (3.9)$$

$$(1 + r)/(1 - r) \geq \left[ \eta^{1/(2-2\epsilon)} + (1 - \eta)^{1/(2-2\epsilon)} \right]. \quad (3.10)$$

The inequality (3.10) is interesting in that it gives an idea of what size  $\epsilon$  must be in order to have a "fission" decay<sup>27</sup> where

$$j_1 \approx j_2, \quad (3.11a)$$

$$\eta \approx \frac{1}{2}, \quad (3.11b)$$

so that (3.10) and (3.4) become

$$\frac{1 - \epsilon + \tan \pi\epsilon}{1 - \epsilon - \tan \pi\epsilon} \geq 2^{(1-2\epsilon)/(2-2\epsilon)}. \quad (3.12)$$

This inequality has the solution

$$\epsilon \geq 0.049, \quad (3.13a)$$

$$r \geq 0.16, \quad (3.13b)$$

so that a "fission" decay appears impossible for a high spin particle on the  $\rho$ -f trajectory given in Fig. 2 which has approximately

$$\epsilon_{\rho} \approx 0.025 \quad (3.14a)$$

$$\gamma_{\rho} \approx 0.081. \quad (3.14b)$$

Going back to (3.10) and assuming  $\epsilon$ ,  $\gamma$  and  $\eta$  are all small, the decay is permitted if

$$\gamma > \frac{1}{2} \eta^{\frac{1}{2}}. \quad (3.15)$$

Assuming that the asymptotic form (2.7) is adequate at low values of  $J$ , the constraint that  $J \rightarrow (J - 1) + 1[\rho(765)]$  is permitted can be found by putting  $\eta = 1/J$  in (3.15) and using (3.14b)

$$0.08 > \frac{1}{2} J^{-\frac{1}{2}}, \quad (3.16a)$$

$$J > 40. \quad (3.16b)$$

Requiring very low orbital angular momentum in the final state therefore implies the decay into two particles on the same trajectory does not begin to take place until quite high values of spin.

At intermediate values of spin a "cascade" decay<sup>27</sup> into a high mass meson of opposite  $G$  parity plus a pion is possible. Assuming the asymptotic form (2.7) holds for the leading meson trajectory of both  $G$ -parities. Conservation of energy permits the "cascade" decay  $J \rightarrow (J - 1) + \pi$  if

$$M_J + \Gamma_J \geq M_{J-1} + \Gamma_{J-1} + m_{\pi}, \quad (3.17)$$

$$(1 + \gamma) \left[ \frac{J}{\cos(\pi\epsilon)} \right]^{1/(2-2\epsilon)} \geq (1 - \gamma) \left[ \frac{J-1}{\cos(\pi\epsilon)} \right]^{1/(2-2\epsilon)} + m_\pi. \quad (3.18)$$

To see how the theorem of Brower and Harte depends on the assumption (3.5), notice that this decay is prohibited if  $\gamma = 0$ ,

$$\left[ \frac{J}{\cos(\pi\epsilon)} \right]^{1/(2-2\epsilon)} - \left[ \frac{J-1}{\cos(\pi\epsilon)} \right]^{1/(2-2\epsilon)} = \mathcal{O}[J^{-(1-2\epsilon)/(2-2\epsilon)}]. \quad (3.19)$$

A more thorough discussion of the theorem of Brower and Harte is given in the appendix.

H. Goldberg<sup>27</sup> has also examined the kinematics of decays under the assumption (3.5) and has found that the "cascade" decay,  $J \rightarrow (J-L) + \pi$  dominates all other modes. It can take place provided the orbital angular momentum,  $L$ , is bounded by

$$L \geq 2m_\pi J^{\frac{1}{2}}. \quad (3.20)$$

The partial decay width of this mode then behaves like

$$\Gamma_c \sim \left[ \frac{2}{J} \ln J \right]^\pi 2m_\pi J^{\frac{1}{2}} \quad (3.21)$$

and resonances become very stable unless there is a vast number of other states into which they can decay. This contrasts with the alternative (3.4) which does not involve the orbital angular momentum of the decay products growing large at any stage.

#### IV. IMPLICATIONS FOR DYNAMICAL MODELS

To this point, the discussion has been remarkably innocent of dynamics. The reduced residue functions associated with  $\alpha(t)$  have nowhere appeared. Crossing has been ignored in order to discuss poles in one channel and unitarity has only been invoked to obtain the analytic structure of  $\alpha(t)$  and to obtain the bound (2.5) on the phase along the physical cut.

Given the perverse, nonlinear character of the unitarity equation, it may be that a good understanding of its implications can only be obtained after the approximate nature of the resonance spectrum is known. However it is instructive to examine a crude dynamic model based on coupled-channel unitarity which might be expected to produce a leading Regge trajectory corresponding to that given asymptotically by (2.7) and (2.10).

First, consider a simple N/D model for a system of two coupled channels each consisting of two natural parity mesons. One channel has spin zero and the other has spin one. Using a simple, one-pole approximation for the left-hand cut, the N/D model gives the output trajectory functions both in the absence and the presence of coupling. In the absence of coupling, the leading singularity in each channel is a pole trajectory which turns over shortly above threshold before reaching a high value of orbital angular momentum. In the example considered, the potentials were adjusted so that a P-wave resonance was formed a short distance above threshold in each channel. When the coupling was turned on, the two output trajectories "exchanged tails"

to form a leading trajectory which rises above  $J = 2$  ( $L = 1$  in the  $1^- - 0^+$  channel) and a nonleading trajectory which turns over before reaching  $J = 1$ . Figure 3 gives the phase of the two trajectory functions in both the coupled and uncoupled cases.

The suggestion that an extension of this simple two-channel model which included an infinite number of channels of increasing spin could account for a leading Regge trajectory which is indefinitely rising has been frequently discussed.<sup>5,28</sup> Since the model at this stage ignores crossing, it is not a bootstrap attempt. The difficulties in bootstrapping the scheme by getting a self-consistent set of trajectories which provide the forces on the left-hand cut as well as containing the particles on the right will not be discussed here.<sup>29</sup> The goals of this discussion are more modest. The idea is that connecting the existence of high-spin resonances with the simultaneous existence of high-spin channels may have some merit whether or not it can lead immediately to a successful bootstrap. The two-channel N/D model illustrated in Fig. 3 gives some indication already that the dynamics of a trajectory is dominated by channels coupled to it with low orbital angular momentum and it seems plausible that this would be true quite generally.

Two points must be made about the possibility that a mechanism involving coupled-channel unitarity supports a leading singularity which rises indefinitely. First, it seems unlikely that such a mechanism could produce high-spin resonances which are very stable since once the imaginary part of the output trajectory function has become

substantial the "cascade" decay mode becomes available so that there is at least one decay channel with no orbital angular momentum barrier as well as an increasing number of other decay channels open as the spin is increased. From the point of view of the N/D model just discussed, the high-spin channels which keep the real part of  $\alpha$  growing also keep the imaginary part growing and there appears to be no mechanism which would turn over the phase plotted in Fig. 4. If this type of dynamics is relevant, it provides an argument against the identification of the  $U(2375)$  as a recurrence of the rho-trajectory.<sup>18</sup> Quite simply, the model says that if there are open decay channels of low orbital angular momentum, then resonances should have large widths.

A second point about this mechanism for producing an indefinitely rising trajectory involves the nature of the nonleading singularities. The mechanism seems to require an infinite number of nonleading singularities. The model implies the collision or near collision of the various singularity surfaces with the leading trajectory. The analytic form of the leading trajectory given by (2.2) is based on the assumption of the absence of such collisions since the collisions can introduce extra branch cuts into the function  $\alpha(t)$ . The existence of left-hand branch cuts destroys the relation between the asymptotic phase and the asymptotic behavior, (2.7), and changes it to (again ignoring possible logarithmic factors)

$$\lim_{t \rightarrow \infty} \alpha(t) = b e^{i\delta(\infty)} t^{1-\delta(\infty)/\pi+\delta(-\infty)/\pi} \quad (4.1)$$

where  $\delta(-\infty)$  is the asymptotic phase on the left-hand cut. The problem is that the phase on the left-hand cut is not directly measurable and there is no way to put a bound on it. Notice that if

$$\delta(\infty) = \delta(-\infty) \quad (4.2)$$

it is possible for both the real and imaginary parts of the trajectory function to be asymptotically linear.<sup>14</sup> The behavior of the phase along the left-hand cuts depends crucially on the dynamic mechanism which generates the pole surfaces,  $\alpha(t)$ . The important thing is that the existence of open, low angular momentum decay channels depends on the width growing with the mass,

$$\Gamma_J = \gamma M_J \quad (4.3)$$

which holds for (4.1) as well as for (2.7). The credibility of the coupled channel mechanism for producing rising trajectories is not destroyed. Of course, understanding the nature of all of the singularities produced by such a model, cuts as well as poles, is crucial to understanding questions such as the existence of superconvergence relations.<sup>29</sup>

An alternative dynamical model for indefinitely rising trajectories which has been recently discussed is the energy-dependent potential approach.<sup>30,31</sup> Since the output trajectories in such an approach would have the cut structure implied by potential theory, much of what has been said is immediately applicable to them. However, since the mechanism which forced the trajectories to rise is at least

a priori independent of the existence of inelastic channels it seems that such an approach could produce a finite asymptotic phase or a zero asymptotic phase equally well. This approach does have an advantage in that it can be constrained to produce a crossing-symmetric amplitude<sup>30</sup> while crossing can be put into the coupled channel approach only by a generalization of the strip-model.<sup>29</sup>

Finally, consider the problem of interpreting Veneziano model results and extrapolating away from the limit  $\text{Im } \alpha(t) = 0$  (often optimistically referred to as unitarizing the Veneziano model). The only internally consistent approach has been to treat the 0-width model as a Born term and construct a field theory based upon its particle spectrum.<sup>32</sup> If this scheme can be carried out, it will answer a lot of questions concerning the role of quarks, internal symmetries and spin in hadron scattering. It may also provide a counterexample to quash the controversial idea of the bootstrap. Without trying to anticipate the results of such a complicated endeavor involving such a large number of physicists, it might be worthwhile to speculate on the basis of the previous discussion about the two options (2.11) and (2.10) for the phase of the renormalized trajectory. If, after renormalization, the leading trajectory has a finite phase then the output particle spectrum may look considerably different than the input "bare" spectrum. Conversely, if the renormalized trajectory has zero phase then it seems unlikely that the degeneracy of the daughter singularities will be grossly broken.



One argument for the existence of an exponentially growing number of states is that, because of the kinematics of the angular momentum barrier, states on rising trajectories find it increasingly difficult to decay into any particular state.<sup>33</sup> If the output trajectory has a finite phase, this argument fails and the leading trajectory can be largely self-supporting. Instead of requiring nonleading trajectories which are parallel to the leading one, this suggests the possibility that nonleading singularities have a completely different behavior.

The fact that this kind of behavior might have physical significance is suggested by the large width of the  $\epsilon(700,0^+)$  and the absence of  $\rho'(1250,1^-)$  predicted by the Veneziano model. A careful analysis of the data of Crennel et al., Fig. 4, shows no evidence of the  $\rho'$ .<sup>34</sup> This is inconsistent with the predictions of the Veneziano model unless the width of this resonance is greater than 1 GeV. This suggests that the phase of the first daughter trajectory predicted by the Veneziano model behaves something like that shown in Fig. 5 much like the behavior of the nonleading pole in the coupled channel N/D problem in Fig. 3 so that the trajectory turns over. If daughter trajectories do turn over shortly above threshold this would reconcile some of the qualitative success of the Veneziano model with the nonappearance of those daughter states it predicts.

The possibility that daughter trajectories turn over could upset the prediction of a number of resonances which increases exponentially with mass.<sup>4</sup> Since Hagedorn<sup>35</sup> is careful to include nonresonant states in his prediction for the thermodynamic model, this type of model is not necessarily inconsistent with his predictions.

V. ACKNOWLEDGMENT

I am extremely grateful for the guidance and suggestions of G. Chew and S. Mandelstam. I would also like to thank F. von Hippel and C. Sorensen for some informative discussions.

APPENDIX A: THE THEOREM OF BROWER AND HARTE

Consider the kinematics for the decay of a high-spin resonance on the leading Regge trajectory in a particular channel. The mass of the initial resonance is  $M$ , its width  $\Gamma$  and its spin is  $\text{Re } \alpha_0(M^2)$ . It can decay into a multiparticle channel consisting of  $n_f$  particles on each of  $N$  trajectories,  $\alpha_f$ . The  $v$ th particle on the  $f$ th trajectory then has mass  $m_v^f$ , width  $\Gamma_v^f$  and spin  $\text{Re } \alpha_f[(m_v^f)^2]$ . Conservation of energy for the decay process can then be written

$$M + \Gamma \geq \sum_{f=1}^N \sum_{v=1}^{n_f} (m_v^f - \Gamma_v^f). \quad (\text{A.1})$$

If  $L$  is the total orbital angular momentum of the decay channel, conservation of angular momentum gives

$$\begin{aligned} L + \sum_{f=1}^N \sum_{v=1}^{n_f} \text{Re } \alpha_f[(m_v^f)^2] &\geq \text{Re } \alpha_0(M^2) \\ &\geq \min |L + \sum_{f=1}^N \sum_{v=1}^{n_f} (\pm) \text{Re } \alpha_f[(m_v^f)^2]|. \end{aligned} \quad (\text{A.2})$$

Assuming that the asymptotic behavior of the real parts of the trajectory functions are given by

$$\text{Re } \alpha_0(s) \sim c_0 s^{a_0} \quad (\text{A.3})$$

and

$$\text{Re } \alpha_f(s) \sim c_f s^{a_f} \quad (\text{A.4})$$

with

$$a_f \leq a_0 \quad f = (1, \dots, N) \quad (\text{A.5})$$

then the theorem of Brower and Harte<sup>3</sup> generalized to include the effects of resonance widths can be stated in the following form.

Theorem: If  $a_0 > \frac{1}{2}$  then the decay process defined by (A.1) and (A.2) above must fail to satisfy one of the following conditions as  $M \rightarrow \infty$ :

$$(i) \quad \frac{L}{\text{Re } \alpha_0(M^2)} \rightarrow 0$$

$$(ii) \quad \left( \Gamma + \sum_{f=1}^N \sum_{v=1}^{n_f} \Gamma_v^f \right) / M \rightarrow 0$$

$$(iii) \quad m_v^f / M < 1.$$

It is apparent that if the asymptotic phase of the trajectory is not zero, condition (ii) does not hold for simple decay channels. Also, it is possible for condition (iii) to be violated so that one of the "decay products" has mass equal to or greater than the original resonance without violating (A.1). On the other hand, if the asymptotic phase of the trajectories is zero, so that

$$a_0 = 1 \quad (\text{A.6})$$

and

$$\frac{\Gamma}{M} \rightarrow 0 \quad (\text{A.7})$$

condition (ii) is forced to hold. If condition (i) fails and the orbital angular momentum grows with  $\alpha_0(M^2) \propto M^2$  then the result of Jones and Teplitz<sup>36</sup> forces the partial width of the resonance to this particular decay channel to decrease exponentially with  $M^2$ . If the total width is not to decrease exponentially, then there must be an exponentially growing number of available decay channels.

FOOTNOTES AND REFERENCES

- \* This work was supported in part by the U.S. Atomic Energy Commission.
- † NSF Predoctoral Fellow. Present address: Argonne National Laboratory.
1. G. Veneziano, Nuovo Cimento 57A, 190 (1968).
  2. D. Sivers and J. Yellin, Review of Recent Work on Narrow Resonance Models, to be published in Rev. Mod. Phys., April, 1971; M. L. Paciello, L. Sertorio, and B. Taglienti, Nuovo Cimento 65A, 167 (1969); C. Boldighini and L. Sertorio, CERN preprint TH.1173 (1970), to be published.
  3. R. C. Brower and J. Harte, Phys. Rev. 164, 1841 (1967).
  4. S. Fubini and G. Veneziano, Nuovo Cimento 64A, 811 (1969); K. Bardakci and S. Mandelstam, Phys. Rev. 184, 1640 (1969).
  5. P. A. Carruthers, Phys. Rev. 154, 1399 (1967).
  6. G. Chikovani, et al., Phys. Letters 22, 233 (1966).
  7. C. F. Chan, A Model of the Interaction Between Two Overlapping Resonances, Lawrence Radiation Laboratory Report UCRL-20051, to be published.
  8. R. W. Childers, Phys. Rev. D2, 1178 (1970).
  9. G. Frye and R. L. Warnock, Phys. Rev. 130, 478 (1963).
  10. M. Sugawara and A. Tubis, Phys. Rev. 130, 2127 (1963).
  11. P. D. B. Collins and E. J. Squires, Regge Poles in Particle Physics (Springer-Verlag, New York, 1968) p. 70 ff.
  12. S. MacDowell, Phys. Rev. 116, 774 (1959); R. Carlitz and M. Kislinger, Phys. Rev. Letters 24, 186 (1970).
  13. R. Oehme, Phys. Rev. D2, 801 (1970).

14. J. Ball and F. Zachariasen, Phys. Rev. Letters 23, 346 (1969).
15. J. B. Bronzan, Phys. Rev. 187, 2253 (1969).
16. A. Ahmadzadeh, P. G. Burke, and C. Tate, Phys. Rev. 131, 1315 (1963).
17. G. Chew and S. Mandelstam, Phys. Rev. 119, 476 (1960).
18. P. D. B. Collins, R. C. Johnson, and E. J. Squires, Phys. Letters 26B, 223 (1968).
19. D. V. Shirkov, Phys. Letters 32B, 635 (1970); R. M. Spector, Phys. Rev. 173, 1761 (1968).
20. The constant  $d$  in (2.9) can be nonzero only if:

- (a)  $\text{Im } \alpha \rightarrow \infty$  in upper half  $t$  plane;
- (b)  $\lim |\alpha(t)| = \infty$  for directions not parallel to real axis;
- (c)  $\int_1^{\infty} dy y^{-x} \text{Im } \alpha(iy) = \infty$  for all  $x \in (0, 2)$ .

H. Melder, Nuovo Cimento 51A, 882 (1967) footnote 10. See also N. Aronszhan and W. F. Donoghue, J. Anal. Math. (Israel) 5, 321 (1956-57).

21. Particle Data Group, Phys. Letters 33B, 1 (1970).
22. The condition for a maximum in  $\text{Re } \alpha$  is

$$\tan \delta(t) \frac{\delta(t)}{dt} = \frac{d}{dt} \ln |\alpha(t)|$$

and may take place with a phase,  $\delta(t)$ , anywhere between 0 and  $\pi/2$ . For  $\delta(t) \geq \pi/4$  so that  $\text{Im } \alpha \geq \text{Re } \alpha$  the identification of the point  $\text{Re } \alpha(t) = J$  with a spin- $J$  resonance is uncertain.

23. J. Rosner, Wisconsin preprint, to be published; R. C. Arnold and J. L. Uretsky, Phys. Rev. Letters 23, 444 (1969).

24. R. Z. Roskies, Phys. Rev. Letters 21, 1851 (1968).
25. G. F. Chew, Phys. Rev. Letters 22, 364 (1969).
26. S. Coleman, in Theory and Phenomenology in Particle Physics, II, A. Zichichi ed. (Academic Press, New York, 1969).
27. C. Quigg and F. von Hippel, Centrifugal Barrier Effects in Meson Decays, Lawrence Radiation Laboratory Report UCRL-19826, May, 1970, submitted to International Conference of Meson Spectroscopy, Philadelphia, May 1-2, 1970; H. Goldberg, Phys. Rev. Letters 21, 778 (1968).
28. S. Mandelstam, in 1966 Tokyo Summer Lectures in Theoretical Physics, II, G. Takeda and A. Fujii, eds. (W. A. Benjamin, Inc., New York, 1966); See also S. Mandelstam, Phys. Rev. 166, 1539 (1968).
29. P. D. B. Collins and R. C. Johnson, Phys. Rev. 182, 1755 (1969).
30. L. A. P. Balázs, Phys. Rev. D2, 999 (1970).
31. H. H. Aly and H. J. W. Müller, Southern Illinois University preprint (1970).
32. A. R. Swift and R. W. Tucker, Massachusetts University preprint, to be published.
33. A. Kryzwicki, Phys. Rev. 187, 1964 (1969).
34. D. Crennell, et al., Phys. Letters 28B, 136 (1968).
35. R. Hagedorn, Nuovo Cimento 56A, 1027 (1968).
36. C. E. Jones and V. Teplitz, Phys. Rev. Letters 19, 135 (1967).




FIGURE CAPTIONS

- Fig. 1. The phase of the  $\Delta$  trajectory calculated from the masses and widths of the  $I = \frac{3}{2}$  baryons listed in the Particle Data Group<sup>21</sup> compilations. A linear extrapolation of the phase gives an approximate lower limit of the energy at which this trajectory might turn over to be  $S = 160 \text{ GeV}^2$  but the data are perfectly consistent with an indefinitely rising trajectory.
- Fig. 2. The phase of an exchange-degenerate trajectory containing the  $\rho(765)$ ,  $I^G J^P 1^+ 1^-$ , and the  $f(1260)$ ,  $0^+ 2^+$ . The  $g(1660)$  and the  $\rho(1710)$  are listed separately by dashed lines and as a single  $g(1690)$  state by a solid bar. The  $I = 0$  enhancement degenerate with the  $I = 1$ ,  $\rho(1900)$  is chosen to be the recurrence of the  $f$ . The solid, straight line extrapolation predicts wide resonances in the T and U regions with  $\Gamma = 190\text{-}220 \text{ MeV}$  while the dashed extrapolation goes through the CERN missing mass spectrometer peaks.
- Fig. 3. The phases of output trajectories in the two-channel N/D model. In the absence of coupling, trajectory 1 is associated with the  $0^+ 0^+$  channel and trajectory 2 with the  $1^- 0^+$  channel. When the coupling is turned on trajectory 3 is the leading singularity while trajectory 4 turns over quickly.
- Fig. 4. The data of Crennel et al. on pion mass distributions for  $\pi^- p \rightarrow \pi^+ \pi^- n$ ,  $\pi^- p \rightarrow \pi^- \pi^0 p$ , and  $\pi^+ p \rightarrow \pi^+ \pi^+ n$ . This data gives a lower bound on the elasticity of the  $\rho'$  meson predicted by the Veneziano model.


Fig. 5. The phase of the  $\epsilon$ - $\rho'$  trajectory compared to the phase of the  $\rho$ - $f$  indicating the possibility that nonleading singularities are very wide and turn over quickly. For convenience in plotting, the real part of the  $\epsilon$ - $\rho'$  trajectory was moved by one unit to coincide with the  $\rho$ - $f$ .

Phase of  $\Delta(1236)$  Regge trajectory



XBL7010-4075


Fig. 1



XBL7010-4076


Fig. 2

Phase of Regge trajectories from coupled channel N/D




XBL7010-4073

Fig. 3



XBL 696-596

Fig. 4



XBL7010-4072

Fig. 5

LEGAL NOTICE

*This report was prepared as an account of Government sponsored work. Neither the United States, nor the Commission, nor any person acting on behalf of the Commission:*

- A. Makes any warranty or representation, expressed or implied, with respect to the accuracy, completeness, or usefulness of the information contained in this report, or that the use of any information, apparatus, method, or process disclosed in this report may not infringe privately owned rights; or*
- B. Assumes any liabilities with respect to the use of, or for damages resulting from the use of any information, apparatus, method, or process disclosed in this report.*

*As used in the above, "person acting on behalf of the Commission" includes any employee or contractor of the Commission, or employee of such contractor, to the extent that such employee or contractor of the Commission, or employee of such contractor prepares, disseminates, or provides access to, any information pursuant to his employment or contract with the Commission, or his employment with such contractor.*



TECHNICAL INFORMATION DIVISION  
LAWRENCE RADIATION LABORATORY  
UNIVERSITY OF CALIFORNIA  
BERKELEY, CALIFORNIA 94720