

1 This is the pre-peer reviewed version of the following article: Owen JJ, Silver WS. Greenhouse  
2 gas emissions from dairy manure management: a review of field-based studies. Global Change  
3 Biology., which will be published in final form at doi: 10.1111/gcb.12687.

4

5 **Greenhouse gas emissions from dairy manure management: a review of field-based studies**

6

7 Running head: GHG emissions from dairy manure management

8

9 Justine J. Owen\* and Whendee L. Silver

10

11 Department of Environmental Science, Policy & Management, 130 Mulford Hall #3114,

12 University of California, Berkeley, CA 94720

13

14 \*telephone: 510-642-3874, fax: 510-643-5438, email: justine\_owen@berkeley.edu

15

16 **Keywords:** dairy, manure, greenhouse gas, emissions, IPCC, emissions modeling, livestock

17

18 Research Review

19 **Abstract**

20 Livestock manure management accounts for almost 10% of greenhouse gas emissions from  
21 agriculture globally, and contributes an equal proportion to the US methane emission inventory.  
22 Current emissions inventories use emissions factors determined from small-scale laboratory  
23 experiments that have not been compared to field-scale measurements. We compiled published  
24 data on field-scale measurements of greenhouse gas emissions from working and research dairies  
25 and compared these to rates predicted by the IPCC Tier 2 modeling approach. Anaerobic  
26 lagoons were the largest source of methane ( $368 \pm 193 \text{ kg CH}_4 \text{ hd}^{-1} \text{ y}^{-1}$ ), more than three times  
27 that from enteric fermentation ( $\sim 100 \text{ kg CH}_4 \text{ hd}^{-1} \text{ y}^{-1}$ ). Corrals and solid manure piles were large  
28 sources of nitrous oxide ( $1.5 \pm 0.8$  and  $1.1 \pm 0.7 \text{ kg N}_2\text{O hd}^{-1} \text{ y}^{-1}$ , respectively). Nitrous oxide  
29 emissions from anaerobic lagoons ( $0.9 \pm 0.5 \text{ kg N}_2\text{O hd}^{-1} \text{ y}^{-1}$ ) and barns ( $10 \pm 6 \text{ kg N}_2\text{O hd}^{-1} \text{ y}^{-1}$ )  
30 were unexpectedly large. Modeled methane emissions underestimated field-measurement means  
31 for most manure management practices. Modeled nitrous oxide emissions underestimated field-  
32 measurement means for anaerobic lagoons and manure piles, but overestimated emissions from  
33 slurry storage. Revised emissions factors nearly doubled slurry  $\text{CH}_4$  emissions for Europe and  
34 increased  $\text{N}_2\text{O}$  emissions from solid piles and lagoons in the US by an order of magnitude. Our  
35 results suggest that current greenhouse gas emission factors generally underestimate emissions  
36 from dairy manure and highlight liquid manure systems as promising target areas for greenhouse  
37 gas mitigation.

38

39 **Introduction**

40 Animal agricultural currently accounts for 20% of non- $\text{CO}_2$  greenhouse gas (GHG) emissions  
41 globally (EPA, 2012). The majority of these emissions are derived from enteric fermentation by

42 ruminants, especially beef and dairy cattle; however, as livestock agriculture is industrialized,  
43 manure management contributes an increasingly large proportion of GHG emissions. This is  
44 particularly the case for dairy production which, unlike beef production, occurs predominantly  
45 on feedlots in most industrialized countries. In the US, approximately 43% of CH<sub>4</sub> emissions  
46 from dairies were from manure management (USDA, 2011), whereas in California, the state with  
47 the greatest dairy production, 54% of dairy CH<sub>4</sub> was estimated to come from manure  
48 management (CARB, 2011a). Manure management can also be an important source of nitrous  
49 oxide (N<sub>2</sub>O) emissions, accounting for an estimated 5% of global (EPA, 2012) and US (EPA,  
50 2013a) N<sub>2</sub>O emissions. Modeling estimates suggested that N<sub>2</sub>O emissions from manure  
51 management globally played a dominant role in the atmospheric increase in N<sub>2</sub>O over the last  
52 140 y (Davidson, 2009).

53  
54 Emissions from dairy manure management are challenging to measure and model due to the  
55 variability in management systems. Greenhouse gas sources associated with manure  
56 management include solid and liquid manure storage systems and dairy surfaces in corrals and  
57 barns (Figure 1). To facilitate estimates of GHG emissions from dairies, the Intergovernmental  
58 Panel on Climate Change (IPCC) developed a Tier 2 model based on emission factors based on  
59 manure composition, manure production rates, biogeochemical reaction rates, temperature, pH,  
60 and moisture content (IPCC, 2006). Emission factors developed by the IPCC for dairies were  
61 largely based on a few lab or pilot-scale studies (IPCC, 2006; Sedorovich *et al.*, 2007; Chadwick  
62 *et al.*, 2011). However, the relationship between small-scale studies and actual field emissions is  
63 poorly constrained, with only one study making a qualitative comparison (Jungbluth *et al.*,

64 2001). Comparing emission rates calculated using the Tier 2 model with field measurements  
65 provides a valuable test of current emission factors.

66  
67 Previous reviews of animal agriculture emissions have pooled a variety of livestock systems and  
68 scales of studies, i.e. laboratory, pilot, and field scales (Jungbluth *et al.*, 2001; Monteny *et al.*,  
69 2001; Sedorovich *et al.*, 2007; Chadwick *et al.*, 2011; Borhan *et al.*, 2012). However, GHG  
70 emissions from dairies likely differ from other livestock industries due to differences in animal  
71 and manure management. For example, California dairy feedlots had 1.84 million milk cows  
72 and 0.78 million dairy heifers in 2009 which produced 16.4 million kg of volatile solids  $d^{-1}$  and  
73 0.92 million kg N  $d^{-1}$ , over 55% of which was managed in anaerobic lagoons (CARB, 2011b). In  
74 contrast, California beef feedlots had 0.46 million heifers and steers which produced just 0.85  
75 million kg volatile solids  $d^{-1}$  and 0.07 million kg N  $d^{-1}$ , 1% of which was managed as liquid  
76 slurry (CARB, 2011b). Accurately estimating the GHG production from this large stock of dairy  
77 manure is critical for designing successful climate change mitigation programs.

78  
79 The goals of this study were to synthesize a global dataset on GHG emissions from manure  
80 management on dairies, compare the data with modeled values, and identify the greatest  
81 mitigation opportunities. Published field measurements of CH<sub>4</sub> and N<sub>2</sub>O emissions from on-  
82 dairy manure management on working and research dairies globally were reviewed. Carbon  
83 dioxide emission rates were also compiled but because they are not considered to contribute to  
84 climate change (IPCC, 2006; with some contention, e.g., Goodland, 2013) they are not discussed  
85 further. We compared mean emission rates from the field data with values calculated using the  
86 IPCC Tier 2 model to identify discrepancies between measured and modeled values. We used

87 the field data to derive revised emission factors and used these to calculate new emissions  
88 estimates for dairy GHG emissions for the US and Europe.

89

### 90 **Sources of GHG on dairies**

91 Many areas on dairies are potential sources of GHGs, in addition to the direct emissions from  
92 cows (Figure 1). Manure is stored in solid or liquid form. Solid manure piles are composed of  
93 the solids scraped from dairy surfaces (manure and bedding) and/or the solids separated from  
94 slurry. They are heterogeneous in composition and can have both aerobic and anaerobic zones  
95 within the piles, depending on moisture content and management practices. Liquid manure  
96 systems were split into two groups: 1) anaerobic lagoons and 2) slurry tanks and settling ponds,  
97 following the approach of the IPCC (2006). Lagoons are earthen and hold the liquid fraction  
98 after mechanical or gravity-driven separation of the manure plus wash water. They are not  
99 stirred and anaerobic conditions develop rapidly. Slurry tanks and settling ponds are filled with  
100 unseparated, minimally-diluted manure. Slurry manure has more solids than anaerobic lagoon  
101 contents, some of which typically floats on the surface and forms a surface crust. The crust is  
102 important because it provides a substrate that spans anaerobic and aerobic environments where  
103 N<sub>2</sub>O production and CH<sub>4</sub> oxidation can both occur (Petersen *et al.*, 2005; Petersen & Sommer,  
104 2011). Anaerobic digesters are another liquid manure management system, but no studies have  
105 attempted to measure greenhouse gas emissions from functioning anaerobic digesters. Biogas  
106 production from anaerobic digesters has been widely studied, but the literature has focused on  
107 the potential reduction compared to other manure storage, rather than quantifying greenhouse gas  
108 emissions from the anaerobic digester systems themselves. Emissions from digesters are likely  
109 to be dependent upon the type of system and operation practices (ex. retention times,

110 effectiveness of seals in preventing gas escape, and composition of material entering the  
111 digester) (Massé *et al.*, 2011; Tauseef *et al.*, 2013).

112  
113 Corrals included dry lots, loafing pens, and hardstandings. Dry lot corrals are dirt-floored pens  
114 in which manure is deposited and occasionally scraped into piles and/or removed. Loafing pens  
115 are commonly dirt-floored and spread with some sort of bedding material, often dried manure  
116 solids. Milk cows are in loafing pens only when they are not in the milking parlor, dry lots, or  
117 freestalls, thus, loafing pens do not accumulate much manure. Some pasture-based dairies use  
118 standoff pads to hold cows during wet periods when the cows can't be on the pastures. These are  
119 small corrals in which a thick (60-100 cm) layer of sawdust and bark chips is laid over plastic  
120 sheeting (Luo & Saggar, 2008). The sheeting allows the leachate from the pad to be collected  
121 and treated in liquid storage systems. Hardstandings are areas with solid surfaces, such as  
122 concrete, which may be used as corrals or as temporary holding pens, depending on their size  
123 and location on the dairy.

124  
125 Barns were measured as entire barns or only barn floors, depending on measurement approach.  
126 Measurements of whole barns include pens and/or freestalls, manure removal and feeding alleys,  
127 and often the cows themselves. Barn floors are heterogeneous and typically have paved or  
128 slatted-floor areas for livestock movement, farmer access, and manure management, as well as  
129 stalls or pens with some sort of soft bedding where the cows can rest. Emissions for entire  
130 dairies were reported by two studies and were also included.

131

132 **Field measurement data compilation**

133 Thirty-eight studies met our criteria (Table 1), most of which were located in North America and  
134 Europe (Figure 2, Table S1). Emission rates were measured using flux chambers or  
135 micrometeorological techniques. Measurement techniques varied by dairy source area, which  
136 was expected given the different spatial scales (piles versus whole barns) or materials (liquid  
137 versus solid) involved. Measurements were typically carried out every 1 to 2 months over 1 to 5  
138 days for up to a year. Data compiled from the studies included farm characteristics such as the  
139 surface area of the pens and lagoons, and number of cows, as available; measurement and gas  
140 analysis technique; sampling duration and frequency; and climate data as mean annual  
141 temperature [MAT], mean annual precipitation [MAP], and temperature during sampling (Tables  
142 S2-S8). A difficulty in comparing literature data was the difference in, or lack of, information  
143 reported. When possible we remedied this by contacting the authors or providing reasonable  
144 estimates of missing information. Missing MAP and MAT data were estimated using data from  
145 the nearest city on [www.worldclimate.com](http://www.worldclimate.com). Air temperature during sampling periods (“sampling  
146 temperature”) was estimated using either the monthly averages from [www.worldclimate.com](http://www.worldclimate.com) or  
147 the almanac feature on [www.wunderground.com](http://www.wunderground.com). Methane fluxes that included enteric  
148 fermentation-derived emissions from barns, corrals, or whole dairies were corrected for enteric  
149 emissions by subtracting the IPCC regional estimate for enteric fermentation (IPCC, 2006).  
150 Specifically, we used  $128 \text{ kg hd}^{-1} \text{ d}^{-1}$  for North American studies and  $117 \text{ kg hd}^{-1} \text{ d}^{-1}$  for  
151 European and modern, high-producing Chinese dairies (rather than the default of  $68 \text{ kg hd}^{-1} \text{ d}^{-1}$   
152 for Asia, which assumes low-producing cows on small farms).

153

154 Most studies included measurements of the same area at different times (e.g., seasonally) and/or  
155 measurements from areas in which some management component was different (e.g., mixed vs.

156 static manure piles, barns with different flooring and scraping mechanisms). Each permutation  
157 was included in the compilation. The mean emission rate for a given dairy area was calculated  
158 by first averaging the emission rates compiled from each paper, then averaging those values,  
159 such that  $n$  is the number of studies rather than the number of measurements. This method  
160 avoided weighting the mean towards studies, management practices, and measurement  
161 techniques with more measurements. Some studies used climate data to extrapolate between  
162 measurements to calculate an annual emission rate. We included these annual estimates in the  
163 appendices, but they were excluded from the calculation of mean emission rates and the  
164 statistical analyses. Statistical analyses were performed using JMP 10.0.2 (SAS Institute, 2012).  
165 Correlations between GHG emission rates, climate variables, cow populations, manure volume,  
166 and other variables in Tables S2-S8 were explored using multiple linear regressions, with  
167 statistical significance determined as  $P < 0.10$ .

168  
169 Measurement technique may have affected emissions measured from all manure management  
170 systems; large footprint techniques generally measured higher CH<sub>4</sub> and N<sub>2</sub>O emissions than  
171 studies using dynamic or static chambers, with the exceptions of CH<sub>4</sub> from anaerobic lagoons  
172 and corrals. The varied composition and oxygen availability of manure stores creates CH<sub>4</sub> and  
173 N<sub>2</sub>O emission hotspots in space and time which can be missed by smaller footprint techniques  
174 (Parkin & Kaspar, 2004; Sommer *et al.*, 2004). Concurrent measurements using different  
175 techniques have not been made (with one exception, (Sommer *et al.*, 2004)), but are needed to  
176 resolve their impact on reported emissions. Because of the uncertainty in the extent of the  
177 impact of measurement technique, in the analysis below we calculated mean emissions from  
178 each area using all available data. In most cases, this likely produced a conservative estimate.



179

180 Emission rates are presented as the mass of trace gas emitted per head per time (kg trace gas  $\text{hd}^{-1}$   
181  $\text{y}^{-1}$ ) and per unit area per time (kg trace gas  $\text{m}^{-2} \text{y}^{-1}$ ). The discussion focuses on per head numbers  
182 for several reasons. The goal of this study was to evaluate our ability to estimate dairy emissions  
183 from manure management at regional to global scales; therefore, emissions factors needed to use  
184 units that were widely known. Most countries have fairly good estimates of the number of  
185 animals present, but estimates of the area of the various manure handling systems have not been  
186 attempted with few exceptions (Chung *et al.*, 2013). Others (Place & Mitloehner, 2010; O'Brien  
187 *et al.*, 2012) have argued that reporting emissions in terms of the mass of milk produced gives a  
188 better sense of the GHG-efficiency of production. This is a useful approach for comparing  
189 different production systems. However, milk production varies significantly by breed, feed, cow  
190 age and stage in lactation cycle (ex. McCandlish, 1920; Zimmerman *et al.*, 1991); furthermore, it  
191 is not relevant to the emissions from different manure management approaches in which we are  
192 interested here. Few studies (only 11 of the 38 studies included here, with 6 of the studies  
193 measuring whole barns) reported milk production.

194

195 To compare the global warming potential (GWP) of the measured areas,  $\text{N}_2\text{O}$  and  $\text{CH}_4$  emission  
196 rates were converted to 100-year  $\text{CO}_2\text{e}$  emission rates by multiplying by 298 and 34, respectively  
197 (Myhre *et al.*, 2013), and summing the two. When other inventories used the older GWPs for  
198  $\text{N}_2\text{O}$  and  $\text{CH}_4$  of 310 and 21 (IPCC, 1996), respectively, those emission rates were recalculated  
199 using the revised values to be comparable to ours.

200

201 **Summary of field measurements**

202 Anaerobic lagoons and slurry systems had the highest per head GWP on dairies, averaging 12.8  
203  $\pm 7$  Mg CO<sub>2</sub>e hd<sup>-1</sup> y<sup>-1</sup> and  $3.5 \pm 1.7$  Mg CO<sub>2</sub>e hd<sup>-1</sup> y<sup>-1</sup>, respectively (Table 2a). Mean lagoon  
204 GWP was about 20 times higher than mean solid manure storage GWP. When expressed on an  
205 area basis, lagoons and slurry systems were similar, averaging  $703 \pm 195$  kg CO<sub>2</sub>e m<sup>-2</sup> y<sup>-1</sup> and  
206  $827 \pm 320$  kg CO<sub>2</sub>e m<sup>-2</sup> y<sup>-1</sup>, respectively (Table 2b). These rates were high; for comparison, the  
207 highest landfill CH<sub>4</sub> emissions rates reported in Bogner et al. (1995) were  $248$  kg CO<sub>2</sub>e m<sup>-2</sup> y<sup>-1</sup>,  
208 less than half those from liquid manure systems. Barn floors had the lowest GWP ( $38 \pm 7$  kg  
209 CO<sub>2</sub>e hd<sup>-1</sup> y<sup>-1</sup>) of all the dairy environments studied. Methane emissions were the largest  
210 component of total GWP for all sources except for barns and corrals.

211

212 Liquid manure storage systems were the greatest source of CH<sub>4</sub>, with anaerobic lagoons and  
213 slurry stores emitting  $368 \pm 193$  kg CH<sub>4</sub> hd<sup>-1</sup> y<sup>-1</sup> and  $101 \pm 47$  kg CH<sub>4</sub> hd<sup>-1</sup> y<sup>-1</sup>, respectively  
214 (Table 2a). Barns were the next largest source with  $33 \pm 19$  kg CH<sub>4</sub> hd<sup>-1</sup> y<sup>-1</sup>. This was  
215 unexpected given that only one study reported subfloor (deep pit) storage and that most others  
216 reported relatively frequent scraping and/or flushing that removed substrate for GHG production.

217

218 Barns had the greatest N<sub>2</sub>O emissions by nearly an order of magnitude, with  $10.3 \pm 6.2$  kg N<sub>2</sub>O  
219 hd<sup>-1</sup> y<sup>-1</sup> (Table 2a), although field data were highly variable (Table S6). Corrals and solid  
220 manure piles were the next largest N<sub>2</sub>O source with  $1.5 \pm 0.8$  kg N<sub>2</sub>O hd<sup>-1</sup> y<sup>-1</sup> and  $1.1 \pm 0.7$  kg  
221 N<sub>2</sub>O hd<sup>-1</sup> y<sup>-1</sup>, respectively (Table 2a). Nitrous oxide emissions from anaerobic lagoons and slurry  
222 stores were also substantial, with  $0.9 \pm 0.5$  kg N<sub>2</sub>O hd<sup>-1</sup> y<sup>-1</sup> and  $0.3 \pm 0.3$  kg N<sub>2</sub>O hd<sup>-1</sup> y<sup>-1</sup>,  
223 respectively. The relatively large net N<sub>2</sub>O flux from liquid manure storage was surprising given  
224 the predominantly anaerobic conditions typical of unaerated systems. Nitrogen in liquid manure

225 is mostly in the form of ammonium ( $\text{NH}_4^+$ ) and organic N (Harter *et al.*, 2002), and though  
226 anaerobic lagoons are generally anaerobic, aerobic conditions which could promote  
227 denitrification exist at inlets. Other  $\text{N}_2\text{O}$  formation reactions are also feasible, such as  
228 denitrification of nitrate ( $\text{NO}_3^-$ ) produced through annamox (anaerobic  $\text{NH}_4^+$  oxidation, (Mulder  
229 *et al.*, 1995; Maeda *et al.*, 2010)), Feammox (anaerobic  $\text{NH}_4^+$  oxidation coupled to Fe reduction,  
230 (Yang *et al.*, 2012)), or Mnammox (anaerobic  $\text{NH}_4^+$  oxidation coupled to Mn reduction,  
231 (Engström *et al.*, 2005)). Hardstandings and barn floors, surfaces which were scraped or flushed  
232 frequently, had  $\text{CH}_4$  and  $\text{N}_2\text{O}$  emissions generally one to three orders of magnitude lower than  
233 the other sources. These trends were consistent between the per-head and per-area data (Table  
234 2b) and showed that the type of storage or surface measured was the greatest factor controlling  
235 emission rates.

236  
237 Methane emissions from soils are known to be temperature dependent (Conrad, 2007) and  
238 models often assume that manure  $\text{CH}_4$  emissions are positively correlated with MAT (Mangino  
239 *et al.*, 2002; IPCC, 2006). Individual field studies observed greater  $\text{CH}_4$  emissions in summer  
240 and/or with warmer sampling temperatures for manure piles, barns, and whole dairies; however,  
241 there was no significant correlation between  $\text{CH}_4$  emissions and temperature when all the studies  
242 for a given source area were considered. The lack of correlation for liquid systems may be due  
243 to the limited range of MAT represented by the field studies; all studies but one (Todd *et al.*,  
244 2008) sampled liquid systems that were in regions where MAT was 6–15°C. Air temperature  
245 during sampling had a larger range (-10.6 to 34.4°C, Tables S2-S3), but overall liquid systems in  
246 warm climates were under-represented. Differences in volatile solid content, the other key factor  
247 determining  $\text{CH}_4$  production (Mangino *et al.*, 2002), may have also confounded any temperature

248 effect. Insufficient data were available to test the effect of volatile solid content on CH<sub>4</sub>  
249 emissions within or across studies.  
250  
251 Methane and N<sub>2</sub>O emissions were strongly correlated to each other for solid manure piles ( $r^2 =$   
252  $0.73, P < 0.001$ ) and weakly correlated for corrals ( $r^2 = 0.26, P < 0.08$ ). This suggests that in  
253 solid manure management systems, at least a portion of the N<sub>2</sub>O fluxes were derived from  
254 denitrification, which requires the same general environmental conditions as methanogenesis  
255 (warm temperatures, abundant labile C, anaerobic conditions). Corral CH<sub>4</sub> emissions were  
256 negative, indicating soil uptake, in 5 out of 18 cases (Table S4); negative fluxes occurred in late  
257 summer when the soils were dry, or in winter when the soils were cold or frozen. The highest  
258 corral N<sub>2</sub>O emissions were measured in late spring when a combination of warmer temperatures  
259 and moist soils likely promoted nitrification and denitrification (Table S4). The lowest values  
260 were < 50% of the highest emissions and occurred in late fall and winter (Table S4). Despite  
261 these seasonal patterns, neither CH<sub>4</sub> nor N<sub>2</sub>O emissions from corrals were correlated with  
262 temperature and/or precipitation. Leytem et al. (2011) measured higher N<sub>2</sub>O emission rates from  
263 manure piles in warmer months (May and June) than colder ones (September and March, Table  
264 S5), but no correlations were found when all manure pile data were pooled.  
265  
266 Specific management practices could have made it difficult to detect a temperature effect if one  
267 existed. Mixing solid manure piles resulted in increased CH<sub>4</sub> and N<sub>2</sub>O emissions (Yamulki,  
268 2006; Maeda *et al.*, 2010; Ahn *et al.*, 2011; Leytem *et al.*, 2011), contrary to expectations that  
269 mixing would aerate the pile and decrease CH<sub>4</sub> production. The addition or accumulation of  
270 fresh manure was another source of emissions. Addition of fresh material increased pile

271 emissions (Leytem *et al.*, 2011) and the accumulation of fresh material in corrals was likely one  
272 of the most important factors driving positive CH<sub>4</sub> fluxes. Borhan *et al.* (2011a) measured  
273 greater CH<sub>4</sub> and N<sub>2</sub>O emissions from a dry lot corral than from loafing pens (Table S4), probably  
274 due to the corrals having a greater influx of fresh manure and localized, high-moisture urine  
275 patches. Methane emissions from the brick hardstanding were relatively high in the summer  
276 (Table S7) (Gao *et al.*, 2011), likely because scraping was less frequent compared to other dairies  
277 (every 1–4 weeks vs. daily for most other hardstandings, Table S7). Accordingly, Adviento-  
278 Borbe *et al.* (2010) observed a significant, positive correlation between CH<sub>4</sub> emissions and  
279 manure depth on the barn floor. However, Gao *et al.* (2011) was the only study of hardstandings  
280 to use an open path laser rather than flux chambers.

281

### 282 **Emission rate modeling**

283 The field measurements provide a test of emission rate models. The IPCC Tier 2 approach  
284 models CH<sub>4</sub> emissions ( $EF_{CH_4}$ , g CH<sub>4</sub> hd<sup>-1</sup> y<sup>-1</sup>) based on the volatile solid production by the cows  
285 (VS, kg VS hd<sup>-1</sup> y<sup>-1</sup>), a CH<sub>4</sub> conversion factor ( $MCF$ , %) for the manure management practice,  
286 and the maximum possible CH<sub>4</sub> production rate from the volatile solids in the manure ( $B_o$ , m<sup>3</sup>  
287 CH<sub>4</sub> kg VS<sup>-1</sup>):

288

$$289 \quad EF_{CH_4} = VS \times MCF / 100 \times B_o \times 662 \text{ g CH}_4 \text{ m}^{-3} \text{CH}_4 \quad (1).$$

290

291 Volatile solid production by cows can be determined from manure analysis (where volatile solids  
292 are the combustible components of solid manure) or estimated based on feed intake rate,  
293 digestibility, and dry matter content. No studies reviewed here included all the information

294 necessary to calculate dairy specific *VS* so we used the IPCC regional values for international  
295 data (IPCC, 2006) and averages of state values for the US (EPA, 2013b).

296

297 The IPCC Tier 2 approach models direct N<sub>2</sub>O emissions based on annual N excretion rates,  
298 which themselves are a function of energy intake by the cows, crude protein content of feed,  
299 milk production rate, milk protein content, cow growth, typical animal mass, and an emission  
300 factor ( $EF_{N_2O}$ , kg N<sub>2</sub>O-N kg N excreted<sup>-1</sup>) (equations 10.31, 10.32, and 10.33 in IPCC, 2006).

301  $EF_{N_2O}$  can be converted into N<sub>2</sub>O emission rates equivalent to those measured here ( $N_2O_D$ , g N<sub>2</sub>O  
302 hd<sup>-1</sup> d<sup>-1</sup>, where the subscript *D* refers to direct emissions) using the typical animal mass (*TAM*,  
303 kg) and country- or region-specific N excretion rates ( $N_{ex}$ , kg N 1000 kg *TAM*<sup>-1</sup> d<sup>-1</sup>):

304

$$305 \quad N_2O_D = EF_{N_2O} \times 44/28 \times TAM \times N_{ex} \quad (2).$$

306

307 In our calculations, we used a *TAM* of 600 kg, the default for Western Europe (but similar to the  
308 North American default value of 604 kg) (IPCC, 2006). Indirect N<sub>2</sub>O emissions, derived from  
309 the oxidation of gaseous emissions such as ammonia (NH<sub>3</sub>) and nitrous oxides (NO<sub>x</sub>), are  
310 important for calculating the amount of N remaining in manure for its use as an organic fertilizer  
311 (IPCC, 2006). We did not include indirect N<sub>2</sub>O emissions in our calculations, and thus what is  
312 reported here should be considered minimum estimates.

313

314 We used the field measurement means and equations 1 and 2 to derive revised *MCFs* and  $EF_{N_2O}$ s  
315 for the source areas. This is the first time broadly applicable, field measurement-derived *MCFs*

316 and  $EF_{N_2O_S}$  have been calculated. Some revised  $MCF$ s and  $EF_{N_2O_S}$  were very different from  
317 current values.

318

### 319 **Comparisons with modeled emissions**

#### 320 *Measured vs. modeled CH<sub>4</sub> emissions*

321 The means of the field-measured CH<sub>4</sub> emissions from slurry tanks and barns (deep pit storage)  
322 were three times larger than modeled emissions, while the measured CH<sub>4</sub> emissions from solid  
323 manure piles and corrals were lower than modeled values, although there was considerable  
324 variability in measured values (Table 3, Figure 3). The modeled CH<sub>4</sub> emissions from the  
325 remaining sources (anaerobic lagoons and hardstandings) were within the standard error of the  
326 field means or were negligible. Modeled whole dairy CH<sub>4</sub> emissions (calculated using  
327 parameters for Western Europe) were slightly lower than the field measurement mean. The  
328 default  $MCF$ s were within the standard error of the field measurement-derived means except for  
329 slurry tanks and whole barns which had larger revised  $MCF$ s (Table 3).

330

331 The impact of the revised barn/deep pit and slurry store  $MCF$ s was evaluated using data on slurry  
332 storage in Europe because six of 13 barn studies were conducted in Europe, while slurry studies  
333 were distributed in temperate regions globally. We used 1990 and 2011 emissions inventory  
334 data for 12 European countries compiled by the United Nations Framework Convention on  
335 Climate Change (UNFCCC, 2014). Three of the 15 countries in the dataset were excluded due  
336 to lack of data or falling outside the cool MAT temperature zone. The European data did not  
337 distinguish between slurry stored in deep pits and tanks or ponds (i.e., one  $MCF$  was used for all  
338 slurry, that in deep pits and in ponds), whereas we calculated revised  $MCF$ s for each system.

339 Thus, revised European slurry CH<sub>4</sub> emissions were calculated using each revised *MCF* to  
340 provide a range. However, deep pit storage is often a temporary holding for slurry that is  
341 eventually transferred to slurry tanks or ponds, so the *MCF* for slurry stores is likely more  
342 applicable.

343  
344 Calculations using the revised deep pit *MCF* gave total CH<sub>4</sub> emissions from European slurry  
345 storage that were less than those using the country-specific slurry *MCF*s ( $8.4 \pm 4.6$  Tg CO<sub>2</sub>e y<sup>-1</sup>  
346 vs.  $15.2$  Tg CO<sub>2</sub>e y<sup>-1</sup>, respectively, Figure 4). However, the revised slurry pond *MCF* increased  
347 CH<sub>4</sub> emissions from slurry for most countries, with total emissions of  $25.9 \pm 12.2$  Tg CO<sub>2</sub>e y<sup>-1</sup>, a  
348 gain of  $10.7$  Tg CO<sub>2</sub>e y<sup>-1</sup> (Figure 4). Increases were greatest for the countries with the most  
349 manure in liquid systems (Denmark, Germany, The Netherlands, Switzerland, and Sweden). We  
350 found a similar trend using detailed data for the Netherlands (RIVM *et al.*, 2013), with modeled  
351 slurry CH<sub>4</sub> emissions two times larger than those estimated in the current inventory (data and  
352 calculations not shown).

353  
354 The uncertainty in slurry *MCF* has consequences for the evaluation of the European dairy  
355 industry's progress in mitigating its GHG emissions. Between 1990 and 2011, the 13 countries  
356 considered here decreased the total number of cows by nearly 8 million hd leading to a  
357 corresponding decrease in emissions from enteric fermentation by 515 Gg CH<sub>4</sub> (Table 5). The  
358 reduction in cows also decreased VS production by 7.8 Tg so there was less manure to manage  
359 and produce GHG. However, an increase in the proportion of manure in liquid management in  
360 most countries offset some of this decrease in CH<sub>4</sub> production; the current estimates suggest a  
361 total net decrease (combined change in enteric and manure management emissions) of 480 Gg



362 CH<sub>4</sub> (Table 5). Using the revised slurry *MCF* for the 2011 estimates gives a smaller total net  
363 decrease of 166 Gg CH<sub>4</sub>, with some countries (Denmark, Switzerland, and The Netherlands)  
364 having net *increases* of 30–50 Gg CH<sub>4</sub> rather than decreases (Table 5).

365

#### 366 *Measured vs. modeled N<sub>2</sub>O emissions*

367 Modeled N<sub>2</sub>O emissions were less than half of the field measurement means for anaerobic  
368 lagoons, solid manure piles, and barns (Table 6). In contrast, the modeled value for slurry stores  
369 was greater than the field measurement mean. The other sources had modeled emissions that  
370 were within the standard error of the field means or were negligible. The revised *EF<sub>N2O</sub>* values  
371 for anaerobic lagoons, manure piles, and barns were larger than the default values, and the slurry  
372 *EF<sub>N2O</sub>* was the same as the default (Table 6).

373

374 The impact of revised *EF<sub>N2O</sub>* values was evaluated using state-specific data from 2011 for the US  
375 (Table 7) because eight of the nine anaerobic lagoon studies and five of the ten manure pile  
376 studies occurred in the US or North America; therefore, the revised *EF<sub>N2O</sub>* values should be  
377 applicable to this region. The EPA assumed zero N<sub>2</sub>O emissions from anaerobic lagoons,  
378 whereas the revised *EF<sub>N2O</sub>* gave  $1.79 \pm 0.90$  Tg CO<sub>2</sub>e y<sup>-1</sup> (Figure 5). Nitrous oxide emissions  
379 from solid manure piles also increased from 0.51 Tg CO<sub>2</sub>e y<sup>-1</sup> to  $3.36 \pm 2.04$  Tg CO<sub>2</sub>e y<sup>-1</sup> using  
380 the revised *EF<sub>N2O</sub>* (Figure 5). Combined, the revised values increased manure management N<sub>2</sub>O  
381 emissions in the US by more than 4.5 Tg CO<sub>2</sub>e y<sup>-1</sup>, 25% of the 2011 estimate of 17.3 Tg CO<sub>2</sub>e  
382 (EPA, 2013a).

383

384 Whole barn N<sub>2</sub>O emissions varied widely between studies, and the measurements of Leytem et  
385 al. (2013) and Samer et al. (2012) suggested an order of magnitude increase in  $EF_{N_2O}$ . They also  
386 indicated that barns may be significant, largely unaccounted sources of N<sub>2</sub>O from dairies (2 to 3  
387 times more kg N<sub>2</sub>O hd<sup>-1</sup> y<sup>-1</sup> than corrals or solid piles). No standard model has been established  
388 for calculating N<sub>2</sub>O emissions from barns that do not have deep pit manure storage. If we  
389 assume that two-thirds of the cows in the 13 European countries in Table 4 were kept in barns  
390 (11 million hd) for half of the year, and emitted 1 kg N<sub>2</sub>O hd<sup>-1</sup> y<sup>-1</sup> (the approximate mean of the  
391 measurements by Zhang et al. (2005), which was the lowest of the three studies that measured  
392 N<sub>2</sub>O), then barns emitted 1.64 Tg CO<sub>2</sub>e, on the same scale as the revised N<sub>2</sub>O emissions from  
393 anaerobic lagoons in the US (Table 7). Using the field-measurement mean of 10 kg N<sub>2</sub>O hd<sup>-1</sup> y<sup>-1</sup>  
394 for the calculation increased barn emissions by an order of magnitude (to 16.4 Tg CO<sub>2</sub>e),  
395 equivalent to the warming potential of slurry system-derived CH<sub>4</sub> in Europe (Table 4). More data  
396 are needed to assess if barns are actually such large sources of N<sub>2</sub>O.

397

## 398 **Discussion and conclusions**

399 Our results highlight potential issues with the application of IPCC Tier 2 models to estimate  
400 GHG emissions from livestock manure. Emission factors were typically based on few studies,  
401 many of which were not designed for GHG inventory estimation or were small-scale pilot or  
402 laboratory experiments, and spanned various livestock systems (Jungbluth *et al.*, 2001; Chung *et al.*,  
403 2013). These approaches are unlikely to accurately approximate field-scale fluxes from  
404 manure management in a specific livestock system. Our review of field-based research on  
405 dairies suggests that current Tier 2 model parameters are generally underestimating dairy  
406 emissions.

407  
408 Data were conspicuously lacking from India and China, which have the fastest growing dairy  
409 industries in the world (growing by 10.7 and 7.6 million hd, respectively, between 2000 and  
410 2010; FAO, 2014). Though data were not available on manure management practices in the two  
411 countries, estimated CH<sub>4</sub> emissions for each suggest that China is treating more manure in liquid  
412 form; for every million dairy milk cows gained between 2000 and 2010, India's CH<sub>4</sub> emissions  
413 from manure management increased by 5 Gg CH<sub>4</sub>, whereas China's increased by 9 Gg CH<sub>4</sub> per  
414 million hd (FAO, 2014). Accurately modeling these emissions is critical for policy decisions  
415 towards GHG emission reduction.

416  
417 The disagreement between field measurements and modeled values provides mechanistic support  
418 for discrepancies reported by airborne measurements and modeling. In a top-down approach  
419 combining aircraft and tower measurements with an atmospheric transport model, Miller et al.  
420 (2013) calculated total CH<sub>4</sub> emissions for the US that were 1.5 times greater than the EPA  
421 bottom-up approach. Underestimation of emissions from fossil fuel extraction was responsible  
422 for a significant part of this discrepancy, but emissions from livestock enteric fermentation and  
423 manure management were calculated to be twice that of the EPA estimate (Miller *et al.*, 2013).  
424 A smaller scale analysis for the Los Angeles Basin found similar dairy CH<sub>4</sub> fluxes between top-  
425 down and bottom-up approaches (Peischl et al., 2013).

426  
427 Despite the uncertainties in emissions inventories described above, targets for GHG reduction  
428 can be identified. As shown by the European example, decreasing the number of cows can  
429 reduce GHG emissions by decreasing both enteric fermentation and manure production (Ripple

430 *et al.*, 2013). While this is the trend in developed countries, developing nations have growing  
431 livestock populations which must be managed appropriately to be sustainable (Eisler *et al.*,  
432 2014). The most effective GHG mitigation approach for manure management depends on how  
433 manure is handled and stored. Where liquid manure management systems are common,  
434 particularly anaerobic lagoons which were the highest total and per-cow CH<sub>4</sub> source, they  
435 represent the greatest opportunity for GHG emissions reduction. Some estimates suggest that the  
436 total CH<sub>4</sub> and N<sub>2</sub>O emissions per head from anaerobic digesters is about 10% of the emissions  
437 from anaerobic lagoons (CARB, 2011a). The EPA (2011) estimated that adoption of anaerobic  
438 digesters by all US dairies for which this technology is feasible (those with liquid manure  
439 management systems and > 500 hd, or approximately 2,650 farms with 3 million hd) could  
440 reduce US CH<sub>4</sub> emissions by 41.25 Tg CO<sub>2</sub>e y<sup>-1</sup>, or more than 85% of the total CH<sub>4</sub> emissions  
441 from dairy manure management. In addition, these anaerobic digesters would be capable of  
442 producing more than 6.8 million MWh y<sup>-1</sup> (EPA, 2011). According to the GHG equivalency  
443 calculator at [www.epa.gov](http://www.epa.gov), this would offset an additional 17.6 Tg CO<sub>2</sub>e of CO<sub>2</sub> emissions from  
444 energy production.

445

446 Our results show significant disagreement between measured and modeled GHG emissions from  
447 dairies globally. Revised emission factors based on the field data led to greater estimated GHG  
448 emissions from the US and Europe. More field data are needed to refine these models. To  
449 maximize the usefulness of field measurements, better reporting of herd characteristics (number  
450 of milk cows and heifers, average mass, milk production, dry matter and N intake), dairy  
451 characteristics (manure handling practices and storage dimensions, climate parameters, available  
452 land for manure spreading, typical management schedule), and manure characteristics (amount

453 handled by each storage method, volume, volatile solid content, C and N content, manure  
454 temperature) are needed. Emission rates must be reported with the data necessary to convert  
455 between units of per head, per area, and per kg milk production in addition to units specific to  
456 certain types of sources, such as per HPU for barns or per volume for liquid storage. Future  
457 research should focus on GHG emissions from several major dairy industries, particularly China  
458 (the fastest-growing), India (the largest), and California (the largest in the US) (FAO *et al.*,  
459 2006). Each region has unique issues related to climate, development, and legislation that  
460 complicate estimating GHG emissions without direct measurements. Furthermore, longer  
461 monitoring periods are needed to disentangle the effects of management and climate on  
462 emissions and enable more accurate estimates of annual averages.

463

#### 464 **Acknowledgments**

465 This research was partially supported by grant #69-3S75-10-172 from the NRCS-USDA to the  
466 Environmental Defense Fund and W. Silver as part of the Conservation Innovation Program.  
467 Support was also generously provided by grants from the 11th Hour Project, the Lia Fund, and  
468 the Rathmann Family Foundation to W. Silver.

469

#### 470 **References**

- 471 Adviento-Borbe MA, Wheeler EF, Brown NE, Topper PA, Graves RE, Ishler VA, Varga GA  
472 (2010) Ammonia and greenhouse gas flux from manure in freestall barn with dairy cows on  
473 precision fed rations. *Transactions of the ASABE*, **53**, 1251–1266.
- 474 Ahn HK, Mulbry W, White JW, Kondrad SL (2011) Pile mixing increases greenhouse gas  
475 emissions during composting of dairy manure. *Bioresource Technology*, **102**, 2904–2909.
- 476 Amon B, Kryvoruchko V, Amon T, Zechmeister-Boltenstern S (2006) Methane, nitrous oxide  
477 and ammonia emissions during storage and after application of dairy cattle slurry and  
478 influence of slurry treatment. *Agriculture, Ecosystems & Environment*, **112**, 153–162.

- 479 Bjorneberg DL, Leytem AB, Westermann DT, Griffiths PR, Shao L, Pollard MJ (2009)  
480 Measurement of atmospheric ammonia, methane, and nitrous oxide at a concentrated dairy  
481 production facility in southern Idaho using open-path FTIR spectrometry. *Transactions of*  
482 *the ASABE*, **52**, 1749–1756.
- 483 Bogner J, Spokas K, Burton E, Sweeney R, Corona V (1995) Landfills as atmospheric methane  
484 sources and sinks. *Chemosphere*, **31**, 4119–4130.
- 485 Borhan MS, Capareda SC, Mukhtar S, Faulkner WB, McGee R, Parnell CB (2011a) Greenhouse  
486 gas emissions from ground level area sources in dairy and cattle feedyard operations.  
487 *Atmosphere*, **2**, 303–329.
- 488 Borhan MS, Capareda S, Mukhtar S, Faulkner WB, McGee R, Parnell CB (2011b) Determining  
489 seasonal greenhouse gas emissions from ground-level area sources in a dairy operation in  
490 central Texas. *Journal of the Air & Waste Management Association*, **61**, 786–795.
- 491 Borhan S, Mukhtar S, Capareda S, Rahman S (2012) Greenhouse gas emissions from housing  
492 and manure management systems at confined livestock operations. In: *Waste Management -*  
493 *An Integrated Vision* (ed Rebellon LFM). InTech.
- 494 Brown HA, Wagner-Riddle C, Thurtell GW (2002) Nitrous oxide flux from a solid dairy manure  
495 pile measured using a micrometeorological mass balance method. *Nutrient Cycling in*  
496 *Agroecosystems*, **62**, 53–60.
- 497 CARB (2011a) *California Greenhouse Gas Emissions Inventory: 2000-2009*. California Air  
498 Resources Board, Sacramento, CA.
- 499 CARB (2011b) *Annex 3B . Manure Management (IPCC 3A2) to the Technical Support*  
500 *Document for the 2000-2009 California's Greenhouse Gas Emissions Inventory*. California  
501 Air Resources Board.
- 502 Chadwick D, Sommer S, Thorman R, Fanguero D, Cardenas L, Amon B, Misselbrook T (2011)  
503 Manure management: Implications for greenhouse gas emissions. *Animal Feed Science and*  
504 *Technology*, **166-167**, 514–531.
- 505 Chung ML, Shilton AN, Guieysse B, Pratt C (2013) Questioning the accuracy of greenhouse gas  
506 accounting from agricultural waste: a case study. *Journal of Environment Quality*, **42**, 654–  
507 659.
- 508 Conrad R (2007) Microbial ecology of methanogens and methanotrophs. *Advances in Agronomy*,  
509 **96**, 1–63.
- 510 Craggs R, Park J, Heubeck S (2008) Methane emissions from anaerobic ponds on a piggery and  
511 a dairy farm in New Zealand. *Australian Journal of Experimental Agriculture*, **48**, 142.

- 512 Davidson EA (2009) The contribution of manure and fertilizer nitrogen to atmospheric nitrous  
513 oxide since 1860. *Nature Geoscience*, **2**, 659–662.
- 514 Eisler MC, Lee MRF, Tarlton JF et al. (2014) Steps to sustainable livestock. *Nature*, **507**, 32–34.
- 515 Ellis S, Webb J, Misselbrook T, Chadwick D (2001) Emission of ammonia (NH<sub>3</sub>), nitrous oxide  
516 (N<sub>2</sub>O) and methane (CH<sub>4</sub>) from a dairy hardstanding in the UK. *Nutrient Cycling in*  
517 *Agroecosystems*, **60**, 115–122.
- 518 Engström P, Dalsgaard T, Hulth S, Aller RC (2005) Anaerobic ammonium oxidation by nitrite  
519 (anammox): Implications for N<sub>2</sub> production in coastal marine sediments. *Geochimica et*  
520 *Cosmochimica Acta*, **69**, 2057–2065.
- 521 EPA (2011) *Market Opportunities for Biogas Recovery Systems at U.S. Livestock Facilities*. U.S.  
522 Environmental Protection Agency.
- 523 EPA (2012) *Summary Report: Global Anthropogenic Non-CO<sub>2</sub> Greenhouse Gas Emissions:*  
524 *1990 - 2030*. Office of Atmospheric Programs, Climate Change Division, US  
525 Environmental Protection Agency, Washington, DC.
- 526 EPA (2013a) *Inventory of US greenhouse gas emissions and sinks: 1990-2009*. US  
527 Environmental Protection Agency, Washington, DC.
- 528 EPA (2013b) *Inventory of US greenhouse gas emissions and sinks: 1990-2009, Annexes*. US  
529 Environmental Protection Agency, Washington, DC.
- 530 FAO (2014) faostat3.fao.org. *FAOSTAT, Food and Agriculture Organization of the United*  
531 *Nations*.
- 532 FAO, Steinfeld H, Gerber P, Wassenaar T, Castel V, Rosales M, de Haan C (2006) *Livestock's*  
533 *Long Shadow: Environmental Issues and Options*. Food and Agriculture Organization of the  
534 United Nations, Rome.
- 535 Gao Z, Yuan H, Ma W, Liu X, Desjardins RL (2011) Methane emissions from a dairy feedlot  
536 during the fall and winter seasons in Northern China. *Environmental Pollution*, **159**, 1183–  
537 1189.
- 538 Goodland R (2014) A fresh look at livestock greenhouse gas emissions and mitigation potential  
539 in Europe. *Global Change Biology*.
- 540 Gupta PK, Jha AK, Koul S et al. (2007) Methane and nitrous oxide emission from bovine  
541 manure management practices in India. *Environmental pollution (Barking, Essex : 1987)*,  
542 **146**, 219–24.
- 543 Harter T, Davis H, Mathews MC, Meyer RD (2002) Shallow groundwater quality on dairy farms  
544 with irrigated forage crops. *Journal of Contaminant Hydrology*, **55**, 287–315.

- 545 Hensen A, Groot TT, van den Bulk WCM, Vermeulen AT, Olesen JE, Schelde K (2006) Dairy  
546 farm CH<sub>4</sub> and N<sub>2</sub>O emissions, from one square metre to the full farm scale. *Agriculture,*  
547 *Ecosystems & Environment*, **112**, 146–152.
- 548 Husted S (1994) Seasonal Variation in Methane Emission from Stored Slurry and Solid Manures.  
549 *Journal of Environmental Quality*, **23**, 585–592.
- 550 IPCC (1996) *Climate Change 1995: The Science of Climate Change* (eds Joughton JT, Meira  
551 Filho LG, Callander BA, Harris N, Kattenberg A, Maskell K). Cambridge University Press,  
552 Cambridge, United Kingdom.
- 553 IPCC (2006) *IPCC Guidelines for National Greenhouse Gas Inventories, Prepared by the*  
554 *National Greenhouse Gas Inventories Programme* (eds Eggleston S, Buendia L, Miwa K,  
555 Ngara T, Tanabe K). IGES, Japan.
- 556 Jungbluth T, Hartung E, Brose G (2001) Greenhouse gas emissions from animal houses and  
557 manure stores. *Nutrient Cycling in Agroecosystems*, **60**, 133–145.
- 558 Kaharabata SK, Schuepp PH, Desjardins RL (1998) Methane emissions from aboveground open  
559 manure slurry tanks. *Global Biogeochemical Cycles*, **12**, 545–554.
- 560 Kaharabata SK, Schuepp PH, Desjardins RL (2000) Estimating Methane Emissions from Dairy  
561 Cattle Housed in a Barn and Feedlot Using an Atmospheric Tracer. *Environmental Science*  
562 *& Technology*, **34**, 3296–3302.
- 563 Khan RZ, Muller C, Sommer SG (1997) Micrometeorological mass balance technique for  
564 measuring CH<sub>4</sub> emission from stored cattle slurry. *Biology and Fertility of Solis*, **24**, 442–  
565 444.
- 566 Kinsman R, Sauer FD, Jackson HA, Wolynetz MS (1995) Methane and carbon dioxide  
567 emissions from dairy cows in full lactation monitored over a six-month period. *Journal of*  
568 *Dairy Science*, **78**, 2760–6.
- 569 Leytem AB, Dungan RS, Bjorneberg DL, Koehn AC (2011) Emissions of ammonia, methane,  
570 carbon dioxide, and nitrous oxide from dairy cattle housing and manure management  
571 systems. *Journal of Environmental Quality*, **40**, 1383–1394.
- 572 Leytem AB, Dungan RS, Bjorneberg DL, Koehn AC (2013) Greenhouse gas and ammonia  
573 emissions from an open-freestall dairy in southern idaho. *Journal of Environmental Quality*,  
574 **42**, 10–20.
- 575 Luo J, Saggart S (2008) Nitrous oxide and methane emissions from a dairy farm stand-off pad.  
576 *Australian Journal of Experimental Agriculture*, **48**, 179.
- 577 Maeda K, Toyoda S, Shimojima R, Osada T, Hanajima D, Morioka R, Yoshida N (2010) Source  
578 of nitrous oxide emissions during the cow manure composting process as revealed by



- 579 isotopomer analysis of and amoA abundance in betaproteobacterial ammonia-oxidizing  
580 bacteria. *Applied and environmental microbiology*, **76**, 1555–62.
- 581 Mangino J, Bartram D, Brazy A (2002) Development of a methane conversion factor to estimate  
582 emissions from animal waste lagoons. In: *11th International Emission Inventory*  
583 *Conference*. U.S. Environmental Protection Agency.
- 584 Marik T, Levin I (1996) A new tracer experiment to estimate the methane emissions from a dairy  
585 cow shed using sulfur hexafluoride (SF<sub>6</sub>). *Global Biogeochemical Cycles*, **10**, 413–418.
- 586 Massé DI, Talbot G, Gilbert Y (2011) On farm biogas production: A method to reduce GHG  
587 emissions and develop more sustainable livestock operations. *Animal Feed Science and*  
588 *Technology*, **166-167**, 436–445.
- 589 McCandlish AC (1920) Environment and breeding as factors influencing milk production. *The*  
590 *Journal of Heredity*, **11**, 204–214.
- 591 McGinn SM, Beauchemin KA (2012) Dairy farm methane emissions using a dispersion model.  
592 *Journal of Environmental Quality*, **41**, 73–79.
- 593 Miller SM, Wofsy SC, Michalak AM et al. (2013) Anthropogenic emissions of methane in the  
594 United States. *Proceedings of the National Academy of Sciences of the United States of*  
595 *America*, **110**, 20018–20022.
- 596 Misselbrook T, Webb J, Chadwick D, Ellis S, Pain B (2001) Gaseous emissions from outdoor  
597 concrete yards used by livestock. *Atmospheric Environment*, **35**, 5331–5338.
- 598 Monteny GJ, Groenestein CM, Hilhorst MA (2001) Interactions and coupling between emissions  
599 of methane and nitrous oxide from animal husbandry. *Nutrient Cycling in Agroecosystems*,  
600 **60**, 123–132.
- 601 Mulder A, Graaf AA, Robertson LA, Kuenen JG (1995) Anaerobic ammonium oxidation  
602 discovered in a denitrifying fluidized bed reactor. *FEMS Microbiology Ecology*, **16**, 177–  
603 184.
- 604 Myhre G, Shindell D, Breon F-M et al. (2013) Anthropogenic and Natural Radiative Forcing. In:  
605 *Climate Change 2013: The Physical Science Basis. Contribution of Working Group I to the*  
606 *Fifth Assessment Report of the Intergovernmental Panel on Climate Change* (eds Stocker  
607 TF, Qin D, Plattner G-K, Tignor M, Allen SK, Boschung J, Nauels A, Xia Y, Bex V,  
608 Midgley PM). Cambridge University Press, Cambridge, United Kingdom and New York,  
609 NY, USA.
- 610 Ngwabie NM, Jeppsson K-H, Nimmermark S, Swensson C, Gustafsson G (2009) Multi-location  
611 measurements of greenhouse gases and emission rates of methane and ammonia from a  
612 naturally-ventilated barn for dairy cows. *Biosystems Engineering*, **103**, 68–77.

- 613 Ngwabie NM, Jeppsson K-H, Gustafsson G, Nimmermark S (2011) Effects of animal activity  
614 and air temperature on methane and ammonia emissions from a naturally ventilated building  
615 for dairy cows. *Atmospheric Environment*, **45**, 6760–6768.
- 616 O'Brien D, Shalloo L, Patton J, Buckley F, Grainger C, Wallace M (2012) Evaluation of the  
617 effect of accounting method, IPCC v. LCA, on grass-based and confinement dairy systems'  
618 greenhouse gas emissions. *Animal*, **6**, 1512–1527.
- 619 Osada T, Sommer SG, Dahl P, Rom HB (2001) Gaseous emission and changes in nutrient  
620 composition during deep litter composting. *Acta Agriculturae Scandinavica, Section B -  
621 Soil & Plant Science*, **51**, 137–142.
- 622 Parkin TB, Kaspar TC (2004) Temporal variability of soil carbon dioxide flux : Effect of  
623 sampling frequency on cumulative carbon loss estimation. *Soil Science Society of America  
624 Journal*, **68**, 1234–1241.
- 625 Peischl J, Ryerson TB, Brioude J et al. (2013) Quantifying sources of methane using light  
626 alkanes in the Los Angeles basin, California. *Journal of Geophysical Research:  
627 Atmospheres*, **118**, 4974–4990.
- 628 Petersen SO, Sommer SG (2011) Ammonia and nitrous oxide interactions: Roles of manure  
629 organic matter management. *Animal Feed Science and Technology*, **166-167**, 503–513.
- 630 Petersen SO, Amon B, Gattinger A (2005) Methane oxidation in slurry storage surface crusts.  
631 *Journal of environmental quality*, **34**, 455–61.
- 632 Place SE, Mitloehner FM (2010) Invited review: Contemporary environmental issues: a review  
633 of the dairy industry's role in climate change and air quality and the potential of mitigation  
634 through improved production efficiency. *Journal of Dairy Science*, **93**, 3407–16.
- 635 Ripple WJ, Smith P, Haberl H, Montzka SA, McAlpine C, Boucher DH (2013) Ruminants,  
636 climate change and climate policy. *Nature Climate Change*, **4**, 2–5.
- 637 RIVM, Coenen PWHG, van der Maas CWM et al. (2013) *Greenhouse gas emission in The  
638 Netherlands*. National Institute for Public Health and the Environment (RIVM), Bilthoven,  
639 The Netherlands.
- 640 Safley LM, Westerman PW (1988) Biogas production from anaerobic lagoons. *Biological  
641 Wastes*, **23**, 181–193.
- 642 Safley LM, Westerman PW (1992) Performance of a dairy manure anaerobic lagoon.  
643 *Bioresource Technology*, **42**, 43–52.
- 644 Samer M, Ammon C, Loebstin C, Fiedler M, Berg W, Sanftleben P, Brunsch R (2012) Moisture  
645 balance and tracer gas technique for ventilation rates measurement and greenhouse gases

- 646 and ammonia emissions quantification in naturally ventilated buildings. *Building and*  
647 *Environment*, **50**, 10–20.
- 648 Sedorovich DM, Rotz CA, Richard TL (2007) Greenhouse gas emissions from dairy farms. In:  
649 *2007 ASABE Annual International Meeting*, Vol. 0300, p. Paper No. 074096. American  
650 Society of Agricultural and Biological Engineers.
- 651 Sneath RW, Beline F, Hilhorst MA, Peu P (2006) Monitoring GHG from manure stores on  
652 organic and conventional dairy farms. *Agriculture, Ecosystems & Environment*, **112**, 122–  
653 128.
- 654 Snell HGJ, Seipelt F, Van den Weghe HFA (2003) Ventilation rates and gaseous emissions from  
655 naturally ventilated dairy houses. *Biosystems Engineering*, **86**, 67–73.
- 656 Sommer SG, McGinn SM, Hao X, Larney FJ (2004) Techniques for measuring gas emissions  
657 from a composting stockpile of cattle manure. *Atmospheric Environment*, **38**, 4643–4652.
- 658 Tauseef SM, Premalatha M, Abbasi T, Abbasi SA (2013) Methane capture from livestock  
659 manure. *Journal of Environmental Management*, **117**, 187–207.
- 660 Todd RW, Cole NA, Rhoades MB, Parker DB, Casey KD (2008) Daily, monthly, seasonal, and  
661 annual ammonia emissions from Southern High Plains cattle feedyards. *Journal of*  
662 *environmental quality*, **40**, 1090–5.
- 663 Todd RW, Cole N a., Casey KD, Hagevoort R, Auvermann BW (2011) Methane emissions from  
664 southern High Plains dairy wastewater lagoons in the summer. *Animal Feed Science and*  
665 *Technology*, **166-167**, 575–580.
- 666 UNFCCC (2014) 2013 Annex 1 Party GHG National Inventory Submissions. *2013 Annex 1*  
667 *Party GHG National Inventory Submissions*.
- 668 USDA (2011) *U.S. Agriculture and Forestry Greenhouse Gas Inventory: 1990-2008. Technical*  
669 *Bulletin No. 1930*. Climate Change Program Office, Office of the Chief Economist, U.S.  
670 Department of Agriculture.
- 671 VanderZaag AC, Wagner-Riddle C, Park K-H, Gordon RJ (2011) Methane emissions from  
672 stored liquid dairy manure in a cold climate. *Animal Feed Science and Technology*, **166-**  
673 **167**, 581–589.
- 674 Van Vliet LJP, Patni NK, Matin MA (2004) GHG emissions from manure in a naturally  
675 ventilated, freestall dairy barn, comparing sand and sawdust bedding. In: *ASAE/CSAE*  
676 *Meeting Paper No. 044161*, Vol. 0300. ASAE, St. Joseph, MI.
- 677 Wu W, Zhang G, Kai P (2012) Ammonia and methane emissions from two naturally ventilated  
678 dairy cattle buildings and the influence of climatic factors on ammonia emissions.  
679 *Atmospheric Environment*, **61**, 232–243.

- 680 Yamulki S (2006) Effect of straw addition on nitrous oxide and methane emissions from stored  
681 farmyard manures. *Agriculture, Ecosystems & Environment*, **112**, 140–145.
- 682 Yang WH, Weber KA, Silver WL (2012) Nitrogen loss from soil through anaerobic ammonium  
683 oxidation coupled to iron reduction. *Nature Geoscience*, **5**, 538–541.
- 684 Zhang G, Strøm JS, Li B, Rom HB, Morsing S, Dahl P, Wang C (2005) Emission of ammonia  
685 and other contaminant gases from naturally ventilated dairy cattle buildings. *Biosystems  
686 Engineering*, **92**, 355–364.
- 687 Zhu Z, Dong H, Zhou Z (2012) Ammonia and greenhouse gas emissions from a dairy cattle barn  
688 with a daily manure collection system. *Transactions of the ASABE*, **55**, 1959–1964.
- 689 Zimmerman CA, Rakes AH, Jaquette RD, Hopkins BA, Croom WJ (1991) Effects of protein  
690 level and forage source on milk production and composition in early lactation dairy cows.  
691 *Journal of Dairy Science*, **74**, 980–90.
- 692

693 **Supporting Information**

694 Additional Supporting Information may be found in the online version of this article:

695 Table S1. Summary of the measurement techniques, locations, and climate data for each study.

696 Table S2. Anaerobic lagoon study characteristics and GHG emissions.

697 Table S3. Slurry tank and pond study characteristics and GHG emissions.

698 Table S4. Corral study characteristics and GHG emissions.

699 Table S5. Solid manure pile study characteristics and GHG emissions.

700 Table S6. Whole barn and barn floor study characteristics and GHG emissions.

701 Table S7. Hardstanding study characteristics and GHG emissions.

702 Table S8. Whole dairy study characteristics and GHG emissions.

703

704 **Table 1.** Studies included in this review and the sources of greenhouse gases measured by each.

<b>Study</b>	<b>lagoon</b>	<b>slurry tank</b>	<b>manure pile</b>	<b>compost area</b>	<b>corrals</b>	<b>concrete pens</b>	<b>barn</b>	<b>whole dairy</b>
Borhan et al. (2011a)	x			x	x		x	
Borhan et al. (2011b)	x				x		x	
Bjorneberg et al. (2009)	x				x			
Leytem et al. (2011)	x			x	x			
Leytem et al. (2013)	x						x	
Craggs et al. (2008)	x							
Safley and Westerman (1988)	x							
Safley and Westerman (1992)	x							
Todd et al. (2011)	x							
Husted (1994)		x	x					
Sneath et al. (2006) <sup>a</sup>		x	x					
Hensen et al. (2006)		x						x
Kaharabata et al. (1998)		x						
Kahn et al. (1997)		x						
VanderZaag et al. (2011)		x						
Ahn et al. (2011)			x					
Amon et al. (2006)			x					
Brown et al. (2002)			x					
Gupta et al. (2007)			x					
Osada et al. (2001)			x					
Sommer et al. (2004)			x					
Kaharabata et al. (2000)					x			
Luo and Sagar (2008)					x			
Ellis et al. (2001)						x		
Gao et al. (2011)						x		
Misselbrook et al. (2001)						x		
Adviento-Borbe et al. (2010)							x	
Kinsman et al. (1995)							x	
Marik and Levin (1996)							x	
Ngwabie et al. (2009)							x	
Ngwabie et al. (2011)							x	
Samer et al. (2012)							x	
Snell et al. (2003)							x	
van Vliet et al. (2004)							x	
Wu et al. (2012)							x	
Zhang et al. (2005)							x	
Zhu et al. (2012)							x	
McGinn and Beauchemin (2012)								x

<sup>a</sup> excludes slurry tank work which was pilot-scale

705  
706

707 **Table 2a.** Summary of the means and ranges of N<sub>2</sub>O and CH<sub>4</sub> emission rates measured by the studies listed in Table 1, in kg hd<sup>-1</sup> y<sup>-1</sup>.

	Emission rate (kg hd <sup>-1</sup> y <sup>-1</sup> )			Emission global warming potential <sup>b</sup> (kg CO <sub>2</sub> e hd <sup>-1</sup> y <sup>-1</sup> )		
	mean <sup>a</sup> ± standard error (n)			mean <sup>a</sup> ± standard error (n)		
	CH <sub>4</sub>	N <sub>2</sub> O	CO <sub>2</sub>	from CH <sub>4</sub>	from N <sub>2</sub> O	total (CH <sub>4</sub> +N <sub>2</sub> O)
<b>Anaerobic lagoons</b>	368 ± 193 (9) 4 - 2814	0.9 ± 0.5 (4) 0.004 - 3.9	687 ± 266 (6) 4.8 - 2400	12510 ± 7334	264 ± 131	12775 ± 6699
<b>Slurry stores</b>	101 ± 47 (6) 0 - 328	0.3 ± 0.3 (3) 0 - 4.5	nm	3422 ± 1601	81 ± 76	3504 ± 1680
<b>Solid<sup>c</sup></b>	13 ± 11 (4) 0 - 99	1.1 ± 0.7 (4) 0.02 - 7	754 ± 695 (2) 59 - 3546	431 ± 372	315 ± 196	632 ± 470
<b>Corrals<sup>d</sup></b>	-17 ± 24 (6) -128 - 210	1.5 ± 0.8 (4) 0.0 - 12	4242 ± 3040 (3) 134 - 20292	-577 ± 844	454 ± 272	-124 ± 1073
<b>Hard-standings<sup>d</sup></b>	1.2 ± 0.8 (3) -3.8 - 7.1	0.0004 ± 0.0001 (2) 0.0001 - 0.001	nm	40 ± 26	0.13 ± 0.02	40 ± 27
<b>Barn floor</b>	0.9 ± 0.7 (4) 0 - 4.4	0.03 ± 0.01 (4) 0.001 - 0.1	94 ± 39 (4) 25 - 250	30 ± 22	7.5 ± 4.4	38 ± 27
<b>Whole barn<sup>d</sup></b>	33 ± 19 (10) -61 - 289	10 ± 6 (3) 0 - 22	7204 ± 5507 (3) 273 - 35058	1120 ± 931	3076 ± 3154	4197 ± 2496
<b>Whole dairy<sup>d</sup></b>	96 ± 35 (18) -91 - 350	nm	nm	3252 ± 1191	nm	3252 ± 1194

<sup>a</sup> Mean emissions were calculated by first averaging measurements within studies then averaging across studies (n=number of studies). The exception was the whole dairy measurements which were treated individually and not grouped by study (i.e., n=number of dairies measured).

<sup>b</sup> Where 1 g CH<sub>4</sub> = 34 g CO<sub>2</sub>e and 1 g N<sub>2</sub>O = 298 g CO<sub>2</sub>e. CO<sub>2</sub> is not included.

<sup>c</sup> Excludes data from Gupta et al. (2007)

<sup>d</sup> Where necessary, methane emission rates have been corrected for enteric emissions as described in the text. Carbon dioxide emissions were not corrected for respiration

708

709

710 **Table 2b.** Summary of the means and ranges of N<sub>2</sub>O and CH<sub>4</sub> emission rates measured by the studies listed in Table 1, in kg m<sup>-2</sup> y<sup>-1</sup>,  
 711 with same footnotes as Table 2a.

	<b>Emission rate (kg m<sup>-2</sup> y<sup>-1</sup>)</b>			<b>Emission global warming potential<sup>b</sup> (kg CO<sub>2</sub>e m<sup>-2</sup> y<sup>-1</sup>)</b>
	mean <sup>a</sup> ± standard error (n)			mean <sup>a</sup> ± standard error (n)
	<b>CH<sub>4</sub></b>	<b>N<sub>2</sub>O</b>	<b>CO<sub>2</sub></b>	<b>total (CH<sub>4</sub>+N<sub>2</sub>O)</b>
<b>Anaerobic lagoons</b>	20 ± 5 (9) 0.3 - 84	0.09 ± 0.05 (4) 0.001 - 0.4	64 ± 34 (6) 2 - 312	703 ± 195
<b>Slurry stores</b>	24 ± 9 (7) 0 - 74	0.05 ± 0.04 (3) 0 - 0.7	nm	827 ± 320
<b>Solid<sup>c</sup></b>	2.8 ± 0.9 (6) 0 - 13	0.3 ± 0.1 (7) 0.005 - 1.0	268 ± 103 (4) 13 - 461	147 ± 49
<b>Corrals<sup>d</sup></b>	-0.8 ± 0.9 (6) -5.2 - 3.8	0.03 ± 0.014 (4) 0.001 - 0.22	83 ± 52 (3) 12 - 365	-16 ± 35
<b>Hard-standings<sup>d</sup></b>	0.5 ± 0.6 (3) -0.11 - 2.2	0.0003 ± 0.0001 (2) 0 - 0.001	nm	18 ± 21
<b>Barn floor</b>	0.2 ± 0.2 (4) 0 - 1.0	0.01 ± 0.01 (4) 0.00 - 0.04	21 ± 10 (4) 5 - 58	9.4 ± 7.0
<b>Whole barn<sup>d</sup></b>	3.4 ± 2.0 (10) -2.8 - 31	0.9 ± 0.7 (3) 0 - 2.3	774 ± 578 (3) 39 - 3713	381 ± 277
<b>Whole dairy<sup>d</sup></b>	nm	nm	nm	nm

712

713



714 **Table 3.** Methane emissions modeling inputs and results. *VS* = volatile solids, *MCF* = methane conversion factor, *B<sub>o</sub>* = the maximum  
 715 possible CH<sub>4</sub> production rate from the volatile solids in the manure.

	<i>VS</i>		<i>MCF</i>		<i>B<sub>o</sub></i>		<i>CH<sub>4</sub> emission rate</i>		
	( <i>kg-VS</i> <i>hd<sup>-1</sup> y<sup>-1</sup></i> )	<i>VS</i> data source	(%)	<i>MCF</i> data source	( <i>m<sup>3</sup> CH<sub>4</sub></i> <i>kg VS<sup>-1</sup></i> )	<i>B<sub>o</sub></i> data source	Modeled ( <i>kg hd<sup>-1</sup> y<sup>-1</sup></i> )	Field ( <i>kg hd<sup>-1</sup> y<sup>-1</sup></i> )	Field- derived MCF (%)
<b>Anaerobic lagoon</b>	2770	average ID, NM, TX, NC <sup>a</sup>	74	average ID, NM, TX, NC <sup>a</sup>	0.24	default <sup>ab</sup>	326	368 ± 193 (9)	84 ± 44
<b>Slurry stores</b>	1861.5	Western Europe <sup>b</sup>	11	MAT=11, with crust <sup>b</sup>	0.24	Western Europe <sup>b</sup>	33	101 ± 47 (6)	34 ± 16
<b>Manure pile</b>	2750	average TX and ID <sup>a</sup>	5	static pile <sup>a</sup>	0.24	default <sup>a</sup>	22	13 ± 11 (4)	2.9 ± 2.5
<b>Corrals</b>	2800	average TX and ID <sup>a</sup>	1	cool MAT <sup>ab</sup>	0.24	default <sup>ab</sup>	4.4	-17 ± 24 (6)	-3.8 ± 5.5
<b>Paved surfaces and barn floors</b>	1861.5	Western Europe <sup>b</sup>	1	cool MAT <sup>b</sup>	0.24	Western Europe <sup>b</sup>	3.0	0.9 ± 0.5 (6)	0.31 ± 0.16
<b>Brick hardstanding</b>	1022	Asia <sup>b</sup>	1	cool MAT <sup>b</sup>	0.13	Asia <sup>b</sup>	0.9	1.6 ± 7.7 (1)	1.8 ± 8.8
<b>Whole barn</b>	1861.5	Western Europe <sup>b</sup>	3	deep pit, cool MAT <sup>b</sup>	0.24	Western Europe <sup>b</sup>	8.9	33 ± 19 (10)	11 ± 6.4
<b>Whole dairy (= corral + manure pile + slurry tank + whole barn)</b>							56	96 ± 35 (18)	

716 <sup>a</sup> from (EPA, 2013b)

717 <sup>b</sup> from (IPCC, 2006)

718

719 **Table 4.** Comparison of modeled slurry emissions in 13 cool MAT European countries using the 2011 liquid slurry *MCF* s (and other  
720 inputs) for each country (UNFCCC, 2014), the revised deep pit and slurry *MCF*s from Table 3. Countries with 60% liquid manure  
721 management are Denmark, Germany, The Netherlands, Switzerland and Sweden.

	Dairy cows (1000 hd)	VS (kg DM/hd/d)	Bo (m3 CH4/kg VS)	Liquid fraction (%)	Liquid <i>MCF</i> (%)	Revised <i>MCF</i> (deep pit) (%)	Revised <i>MCF</i> (slurry) (%)	Liquid manure emissions (Gg CH <sub>4</sub> y <sup>-1</sup> )			
								Using current <i>MCF</i>	Using revised deep pit <i>MCF</i>	Using revised slurry <i>MCF</i>	
<b>Austria</b>	527.39	4.27	0.24	31.61	8.7	11 ± 6	34 ± 16	3.6	4.5 ± 2.5	14.0 ± 6.6	
<b>Belgium</b>	459.78	4.10	0.24	11.54	19	11 ± 6	34 ± 16	2.4	1.4 ± 0.8	4.3 ± 2.0	
<b>Denmark</b>	565.11	6.09	0.24	88.41	10	11 ± 6	34 ± 16	17.7	19.4 ± 10.6	60.0 ± 28.2	
<b>Finland</b>	285.53	4.94	0.24	46.41	10	11 ± 6	34 ± 16	3.8	4.2 ± 2.3	12.9 ± 6.1	
<b>France</b>	3660.68	4.12	0.24	40.87	39	11 ± 6	34 ± 16	139.3	39.3 ± 21.4	121.4 ± 57.1	
<b>Germany</b>	4190.10	4.01	0.23	73.52	14.4	11 ± 6	34 ± 16	98.6	75.6 ± 41.2	233.6 ± 109.9	
<b>Ireland</b>	1086.11	2.98	0.24	28.60	39	11 ± 6	34 ± 16	20.9	5.9 ± 3.2	18.3 ± 8.6	
<b>Italy</b>	1754.98	6.37	0.14	35.03	13.9	11 ± 6	34 ± 16	19.0	15.0 ± 8.2	46.4 ± 21.9	
<b>Luxembourg</b>	40.45	4.56	0.24	34.20	39	11 ± 6	34 ± 16	1.4	0.4 ± 0.2	1.2 ± 0.6	
<b>Sweden</b>	346.50	5.33	0.24	62.23	3.5	11 ± 6	34 ± 16	2.3	16.0 ± 8.7	22.7 ± 10.7	
<b>Switzerland</b>	589.24	6.24	0.24	68.22	10	11 ± 6	34 ± 16	14.5	40.2 ± 21.9	49.4 ± 23.3	
<b>The Netherlands</b>	1469.72	4.56	0.25	90.38	17	11 ± 6	34 ± 16	62.2	17.1 ± 9.3	124.4 ± 58.5	
<b>United Kingdom</b>	1814.00	3.61	0.24	41.00	39	11 ± 6	34 ± 16	60.8	4.5 ± 2.5	53.0 ± 24.9	
								all	446.5	246.4 ± 134.4	761.7 ± 358.4
								top 5 liquid	195.3	158.6 ± 86.5	490.1 ± 230.6
								all (Tg CO <sub>2</sub> e)	15.2	8.4 ± 4.6	25.9 ± 12.2
								top 5 liquid (Tg CO <sub>2</sub> e)	6.6	5.4 ± 2.9	16.7 ± 7.8

722

723

724 **Table 5.** Comparisons between 1990 and 2011 data for the 13 countries in Table 4 (UNFCCC, 2014). Negative values indicate a  
 725 decrease from 1990 to 2011. The revised slurry *MCF* was used to recalculate 2011 emissions, not the revised deep pit *MCF*, and the  
 726 1990 value were not adjusted.

	Cows (1000 hd)	VS production (Gg)	Liquid fraction (%)	Enteric fermentation emissions (Gg CH <sub>4</sub> )	Manure management emissions (default <i>MCF</i> ) (Gg CH <sub>4</sub> )	Net emissions (default <i>MCF</i> ) (Gg CH <sub>4</sub> )	Manure management emissions (revised <i>MCF</i> ) (Gg CH <sub>4</sub> )	Net emissions (revised <i>MCF</i> ) (Gg CH <sub>4</sub> )
<b>Austria</b>	-377.22	-507.65	-1.04	-26.05	-2.44	-28.49	8.00	-18.05
<b>Belgium</b>	-378.92	-298.63	1.54	-25.54	-0.58	-26.12	1.31	-24.22
<b>Denmark</b>	-188.01	-266.24	18.37	-12.72	0.71	-12.01	43.07	30.35
<b>Finland</b>	-204.37	-162.88	23.87	-11.74	1.37	-10.37	10.47	-1.27
<b>France</b>	-1649.13	-1205.50	14.47	-83.06	29.58	-53.48	11.72	-71.34
<b>Germany</b>	-2164.45	-1937.47	18.61	-205.97	5.90	-200.06	140.89	-65.07
<b>Ireland</b>	-254.84	-168.89	-3.75	-13.47	-6.13	-19.61	-8.82	-22.29
<b>Italy</b>	-886.77	-2063.00	1.39	-40.39	-8.09	-48.48	19.35	-21.04
<b>Luxembourg</b>	-18.39	-12.17	11.20	-0.88	0.29	-0.58	0.11	-0.77
<b>Sweden</b>	-229.51	-400.16	39.63	-23.15	0.98	-22.16	21.33	-1.82
<b>Switzerland</b>	-193.86	-129.64	4.19	-7.01	-0.43	-7.43	34.46	27.45
<b>The Netherlands</b>	-407.96	-145.59	20.82	-18.72	11.48	-7.25	73.67	54.94
<b>United Kingdom</b>	-1034.26	-545.56	8.40	-46.30	1.43	-44.88	-6.37	-52.67
<b>total</b>	-7987.70	-7843.38	-	-514.99	34.06	-480.93	349.20	-165.79
			<b>total Tg CO<sub>2</sub>e</b>	<b>-17.51</b>	<b>1.16</b>	<b>-16.35</b>	<b>11.87</b>	<b>-5.64</b>

727

728 **Table 6.** Nitrous oxide emissions modeling inputs and results.  $EF_{N_2O}$  = emissions factor,  $TAM$  = typical animal mass,  $N_{ex}$  = country-  
 729 or region-specific N excretion rates.  $EF_{N_2O}$  uncertainty range is a factor of 2 for all but anaerobic lagoons and whole barns.

	$EF_{N_2O}$ (kg N <sub>2</sub> O-N kg N excreted <sup>-1</sup> )	$TAM$ (kg hd <sup>-1</sup> )	$N_{ex}$ (kg N 1000 kg $TAM$ <sup>-1</sup> d <sup>-1</sup> )	$EF_{N_2O}$ and $N_{ex}$ sources	N <sub>2</sub> O emission rate		Field-derived $EF_{N_2O}$ (kg N <sub>2</sub> O-N kg N excreted <sup>-1</sup> )
					Modeled (kg hd <sup>-1</sup> y <sup>-1</sup> )	Field (kg hd <sup>-1</sup> y <sup>-1</sup> )	
<b>Anaerobic lagoon</b>	0	600	0.25	average of ID and TX <sup>a</sup>	0	0.9 ± 0.5 (4)	0.010 ± 0.005
<b>Slurry tanks and ponds</b>	0.005	600	0.44	North America, with crust <sup>b</sup>	0.8	0.3 ± 0.3 (3)	0.005 ± 0.005
<b>Manure pile</b>	0.005	600	0.26	ID <sup>a</sup>	0.4	1.1 ± 0.7 (4)	0.033 ± 0.020
<b>Corrals</b>	0.02	600	0.25	average of ID and TX <sup>a</sup>	1.7	1.5 ± 0.8 (4)	0.048 ± 0.026
<b>Barn floors and paved surfaces</b>	0.02	600	0.44	North America <sup>b</sup>	3.0	0.02 ± 0.01 (6)	0.0001 ± 0.0001
<b>Whole barn (deep pit)</b>	0.002	600	0.48	western Europe <sup>b</sup>	0.3	10 ± 6 (3)	0.062 ± 0.038

730 <sup>a</sup> from (EPA, 2013b)

731 <sup>b</sup> from (IPCC, 2006)


732

733 **Table 7.** Comparison of N<sub>2</sub>O emissions from anaerobic lagoons and solid manure piles modeled for the US using the current  $EF_{N_2O}$  (0  
734 and 0.005 kg N<sub>2</sub>O-N kg N excreted<sup>-1</sup>, respectively) (EPA, 2013a) and the revised  $EF_{N_2O}$  (0.01±0.005 and 0.033 ± 0.02 kg N<sub>2</sub>O-N kg  
735 N excreted<sup>-1</sup>, respectively) from Table 5. For all states, dairy cow  $TAM = 680$  kg, heifer  $TAM = 407$ , and heifer  $N_{ex} = 69$  kg N 1000  
736 kg  $TAM^1 d^{-1}$ .

State	Milk cows (hd)	Heifers (hd)	Manure in solid pile (%)	Manure in anaerobic lagoons (%)	$N_{ex}$ dairy cows (kg N hd <sup>-1</sup> y <sup>-1</sup> )	EPA solid pile N <sub>2</sub> O emissions (kg N <sub>2</sub> O y <sup>-1</sup> )	revised solid pile N <sub>2</sub> O emissions (kg N <sub>2</sub> O y <sup>-1</sup> )	EPA	
								lagoons N <sub>2</sub> O emissions (kg N <sub>2</sub> O y <sup>-1</sup> )	revised lagoons N <sub>2</sub> O emissions (kg N <sub>2</sub> O y <sup>-1</sup> )
Alabama	11000	6000	7	16	130	627	4141 ± 2510	0	2869 ± 1434
Alaska	500	200	34	24	128	131	866 ± 525	0	185 ± 93
Arizona	185000	60000	9	61	159	15336	101217 ± 61344	0	207887 ± 103944
Arkansas	12000	4000	10	9	122	870	5745 ± 3482	0	1567 ± 783
California	1750000	760000	9	59	158	148050	977127 ± 592198	0	1941095 ± 970547
Colorado	123000	75000	11	64	159	13314	87874 ± 53257	0	154930 ± 77465
Connecticut	19000	9500	16	13	145	2691	17757 ± 10762	0	4372 ± 2186
Delaware	5000	3000	19	10	143	852	5621 ± 3406	0	896 ± 448
Florida	114000	30000	7	43	149	6816	44986 ± 27265	0	83741 ± 41871
Georgia	78000	23000	9	23	147	5970	39404 ± 23881	0	30515 ± 15257
Hawaii	2000	1000	9	57	130	145	956 ± 580	0	1835 ± 918
Idaho	574000	320000	11	65	157	60731	400822 ± 242922	0	717725 ± 358863
Illinois	98000	46000	39	16	146	33772	222897 ± 135089	0	27711 ± 13855
Indiana	172000	62000	29	24	150	43943	290021 ± 175770	0	72733 ± 36366
Iowa	210000	130000	34	20	152	67738	447069 ± 270951	0	79692 ± 39846
Kansas	122000	85000	21	36	151	24608	162414 ± 98432	0	84371 ± 42185
Kentucky	77000	50000	14	3	134	9262	61132 ± 37050	0	3970 ± 1985
Louisiana	19000	5000	10	9	125	1379	9103 ± 5517	0	2483 ± 1241
Maine	32000	17000	20	10	144	5674	37450 ± 22697	0	5674 ± 2837
Maryland	53000	28000	22	8	144	10330	68179 ± 41320	0	7513 ± 3756
Massachusetts	13500	7500	22	8	138	2554	16856 ± 10216	0	1857 ± 929
Michigan	361000	148000	24	29	158	80977	534445 ± 323906	0	195693 ± 97847


Minnesota	470000	290000	39	17	145	166961	1101942 ± 667844	0	145556 ± 72778
Mississippi	15000	7000	10	12	135	1236	8160 ± 4946	0	2967 ± 1484
Missouri	95000	50000	42	11	131	32560	214898 ± 130241	0	17055 ± 8528
Montana	14000	7000	19	42	150	2425	16007 ± 9701	0	10722 ± 5361
Nebraska	58000	20000	26	29	150	13233	87337 ± 52932	0	29520 ± 14760
Nevada	28000	10000	10	65	157	2569	16958 ± 10277	0	33402 ± 16701
New Hampshire	15000	7500	19	10	149	2583	17050 ± 10333	0	2719 ± 1360
New Jersey	7500	4000	25	6	138	1603	10581 ± 6412	0	769 ± 385
New Mexico	322000	140000	9	61	164	28173	185944 ± 112693	0	381905 ± 190953
New York	610000	330000	17	13	151	96041	633869 ± 384163	0	146886 ± 73443
North Carolina	44000	21000	11	10	152	4440	29306 ± 17761	0	8073 ± 4037
North Dakota	20000	10000	38	15	142	6604	43590 ± 26418	0	5214 ± 2607
Ohio	270000	125000	38	15	145	89967	593780 ± 359867	0	71026 ± 35513
Oklahoma	54000	20000	21	45	140	9409	62100 ± 37636	0	40325 ± 20162
Oregon	121000	65000	11	50	150	12245	80815 ± 48979	0	111315 ± 55658
Pennsylvania	543000	310000	24	6	147	118770	783881 ± 475079	0	59385 ± 29692
Rhode Island	1100	500	25	5	141	235	1549 ± 939	0	94 ± 47
South Carolina	16000	7000	8	18	144	1108	7315 ± 4433	0	4988 ± 2494
South Dakota	90000	25000	24	31	150	18635	122989 ± 74539	0	48140 ± 24070
Tennessee	50000	35000	12	4	140	5415	35737 ± 21659	0	3610 ± 1805
Texas	425000	230000	11	58	155	44298	292368 ± 177193	0	467144 ± 233572
Utah	87000	42000	15	56	151	11918	78662 ± 47674	0	88991 ± 44496
Vermont	135000	61000	17	13	145	20068	132448 ± 80271	0	30692 ± 15346
Virginia	95000	49000	11	5	145	9285	61282 ± 37140	0	8441 ± 4220
Washington	252000	122000	11	56	160	26658	175941 ± 106631	0	271425 ± 135712
West Virginia	10000	5000	23	7	134	1900	12543 ± 7602	0	1157 ± 578
Wisconsin	1265000	700000	38	17	150	443940	2930005 ± 1775761	0	397210 ± 198605
Wyoming	6000	5000	19	43	150	1123	7413 ± 4493	0	5084 ± 2542
<i>total kg N<sub>2</sub>O y<sup>-1</sup></i>						<i>1709175</i>	<i>11280554 ± 6836699</i>	<i>0</i>	<i>6023128 ± 3011564</i>
<b>total Tg CO<sub>2</sub>e y<sup>-1</sup></b>						<b>0.51</b>	<b>3.36 ± 2.04</b>	<b>0</b>	<b>1.79 ± 0.90</b>

738 **Figure 1.** Sources of N<sub>2</sub>O and CH<sub>4</sub> on dairies. Thin arrows indicate movement of manure  
739 between locations. Thick arrows indicate relative emission rate. Hardstandings are not shown  
740 but have negligible emissions.



741  
742


743 **Figure 2.** Geographic distribution of sampling sites, marked as triangles (created in GeoMapApp  
744 v. 3.3.8, <http://www.geomapapp.org/>).



745




746 **Figure 3.** Comparison of modeled CH<sub>4</sub> emissions and field measurement means and standard  
747 errors for the largest CH<sub>4</sub> sources.




748  
749

750 **Figure 4.** Comparison of modeled CH<sub>4</sub> emissions for slurry using the current slurry *MCF* used  
751 by the European Union and the revised *MCFs* for deep pit storage and slurry storage calculated  
752 in this study (Table 4).



753  
754

755 **Figure 5.** Comparison of modeled N<sub>2</sub>O emissions of solid manure piles and anaerobic lagoons  
756 calculated using current and revised  $EF_{N_2O}$  values.



757