

UC Santa Barbara

UC Santa Barbara Previously Published Works

Title

Complete Genome Sequences of Two Phylogenetically Distinct Nitrospina Strains Isolated from the Atlantic and Pacific Oceans

Permalink

<https://escholarship.org/uc/item/5gz5182w>

Journal

Microbiology Resource Announcements, 11(5)

ISSN

2576-098X

Authors

Bayer, Barbara
Kellom, Matthew
Valois, Frederica
et al.

Publication Date

2022-05-19

DOI

10.1128/mra.00100-22

Peer reviewed



Complete Genome Sequences of Two Phylogenetically Distinct *Nitrospina* Strains Isolated from the Atlantic and Pacific Oceans

Barbara Bayer,^{a*} Matthew Kellom,^b Frederica Valois,^c John B. Waterbury,^c Alyson E. Santoro^a

^aDepartment of Ecology, Evolution, and Marine Biology, University of California, Santa Barbara, California, USA

^bDOE Joint Genome Institute, Lawrence Berkeley National Laboratory, Berkeley, California, USA

^cBiology Department, Woods Hole Oceanographic Institution, Woods Hole, Massachusetts, USA

ABSTRACT The complete genome sequences of two chemoautotrophic nitrite-oxidizing bacteria of the genus *Nitrospina* are reported. *Nitrospina gracilis* strain Nb-211 was isolated from the Atlantic Ocean, and *Nitrospina* sp. strain Nb-3 was isolated from the Pacific Ocean. We report two highly similar ~3.07-Mbp genome sequences that differ by the presence of ferric iron chelator (siderophore) biosynthesis genes.

Nitrospina species are aerobic, chemoautotrophic, nitrite-oxidizing bacteria that have so far only been found in marine habitats (1), where they play an important role in the nitrogen cycle (2). *Nitrospina gracilis* Nb-211 (ATCC 25379), first described in 1971 (3), was isolated from surface waters (13-m depth) of the Atlantic Ocean approximately 200 miles from the mouth of the Amazon River. *Nitrospina* sp. strain Nb-3 was isolated from the Pacific Ocean off the coast of Peru and has not been validly described; however, its 16S rRNA gene sequence was published in 1994 (4). Both strains belong to the *Nitrospinaceae* family within the *Nitrospinae/Nitrospinota* (Joint Genome Institute/Genome Taxonomy Database [JGI/GTDB]) phylum.

For genomic sequencing, cultures were grown in 2-L glass bottles in artificial seawater medium containing 2 mM nitrite and incubated in the dark without agitation as described previously (5). Cells were collected via centrifugation (1 h, 15,000 × *g*, 10°C), and DNA was extracted from the cell pellets using a cetyltrimethylammonium bromide (CTAB)-phenol-chloroform protocol (6). Draft genomes were generated at the DOE JGI using the Pacific Biosciences (PacBio) sequencing technology (7). Genomic DNA was sheared to 10 kb using g-TUBE columns (Covaris) and subjected to library preparation using the SMRTbell Express template prep 2.0 kit. The PacBio SMRTbell library was purified and size-selected using AMPure PB beads and sequenced on the PacBio Sequel platform, which generated 123,391 subreads (5,002.7 ± 3,337.2 bp) totaling 617,291,847 bp for strain Nb-211 and 89,053 subreads (6,935.6 ± 5,348.8 bp) totaling 617,633,853 bp for strain Nb-3. Reads of >5 kb were assembled with HGAP (v. smrtlink/8.0.0.80529, HGAP 4 [1.0]) using default settings (8). The input read coverage was 188.4× for strain Nb-211 and 189.5× for strain Nb-3. The final draft genome sequences consisted of one scaffold each, with a total size of 3,069,626 bp for strain Nb-211 and 3,075,869 bp for strain Nb-3 (Table 1). We confirmed complete circularization with the Circlator pipeline v.1.5.5 (9), which uses nucmer v.3.1 (10) to check for alignment between assembled contigs at opposite ends of the assembly, identifying a 50,007-bp alignment with 100% identity for strain Nb-211 and 50,012 bp with 99.99% identity for strain Nb-3.

Both genomes were annotated using the Integrated Microbial Genomes (IMG) Annotation Pipeline (IMGAP) v.5.0.22/3. The genome of *Nitrospina gracilis* Nb-211 contains 2,939 coding DNA sequences (CDS), and that of *Nitrospina* sp. Nb-3 contains 2,905 CDS. Both genome sequences share an average nucleotide identity (ANI) of

Editor Frank J. Stewart, Montana State University

This is a work of the U.S. Government and is not subject to copyright protection in the United States. Foreign copyrights may apply.

Address correspondence to Barbara Bayer, bbayer@ucsb.edu, or Alyson E. Santoro, asantoro@ucsb.edu.

*Present address: Barbara Bayer, Department of Microbiology and Ecosystem Science, University of Vienna, Vienna, Austria.

The authors declare no conflict of interest.

Received 9 February 2022

Accepted 12 April 2022

Published 2 May 2022

TABLE 1 Genome features of *Nitrospina gracilis* Nb-211 and *Nitrospina* sp. Nb-3

| Strain | GenBank accession no. | JGI taxon ID | Genome size (bp) | G+C content (%) | No. of DNA scaffolds | No. of total genes | No. of protein coding genes | No. of rRNA operons | No. of tRNA operons | CRISPR count |
|----------------------------|-----------------------|--------------|------------------|-----------------|----------------------|--------------------|-----------------------------|---------------------|---------------------|--------------|
| <i>N. gracilis</i> Nb-211 | JAKJKD010000001 | 2917506613 | 3,069,626 | 57.43 | 1 | 2,939 | 2,879 | 1 | 50 | 2 |
| <i>Nitrospina</i> sp. Nb-3 | JAKJKC010000001 | 2929071401 | 3,075,869 | 56.21 | 1 | 2,939 | 2,846 | 1 | 49 | 0 |

85.5%, well below the intraspecies threshold of 96.5% (11). Strain Nb-3 shares 99.98% ANI with the previously published draft genome sequence of *Nitrospina gracilis* strain 3/211 (12), indicating that the latter likely derives from the culture originally designated strain Nb-3 (4). Strain Nb-3 encodes a putative iron chelator (siderophore) biosynthesis gene cluster, which is absent in strain Nb-211, potentially reflecting adaptations to differences in iron availability in the respective ocean basins the strains were isolated from. Strain Nb-211 was isolated near the Amazon River, which is a source of iron to the Atlantic Ocean (13), while strain Nb-3 was isolated from the relatively iron-deplete North Pacific (14).

Data availability. The whole-genome shotgun sequencing project of *Nitrospina gracilis* Nb-211 has been deposited at DDBJ/ENA/GenBank under BioProject number [PRJNA708439](https://ncbi.nlm.nih.gov/bioproject/PRJNA708439) and accession number [JAKJKD000000000](https://ncbi.nlm.nih.gov/nucl/JAKJKD000000000). The whole-genome shotgun sequencing project of *Nitrospina* sp. Nb-3 has been deposited at DDBJ/ENA/GenBank under BioProject number [PRJNA783628](https://ncbi.nlm.nih.gov/bioproject/PRJNA783628) and accession number [JAKJKC000000000](https://ncbi.nlm.nih.gov/nucl/JAKJKC000000000). The NCBI Sequence Read Archive (SRA) accession numbers for the raw reads are [SRR17430281](https://ncbi.nlm.nih.gov/sra/SRR17430281) for strain Nb-211 and [SRR17430190](https://ncbi.nlm.nih.gov/sra/SRR17430190) for strain Nb-3.

ACKNOWLEDGMENTS

This work was supported by a Simons Foundation Early Career Investigator Award (3435889) and a U.S. National Science Foundation award OCE-1924512 to A.E.S. B.B. was supported by the Austrian Science Fund (FWF) project number J4426-B. The work conducted by the U.S. Department of Energy Joint Genome Institute, a DOE Office of Science User Facility, is supported under contract number DE-AC02-05CH11231. These data were generated for JGI proposal number 506203 to B.B. and A.E.S.

We thank Christa Pennacchio, who managed the project at JGI.

REFERENCES

- Daims H, Lückner S, Wagner M. 2016. A new perspective on microbes formerly known as nitrite-oxidizing bacteria. *Trends Microbiol* 24:699–712. <https://doi.org/10.1016/j.tim.2016.05.004>.
- Kuypers MMM, Marchant HK, Kartal B. 2018. The microbial nitrogen-cycling network. *Nat Rev Microbiol* 16:263–276. <https://doi.org/10.1038/nrmicro.2018.9>.
- Watson SW, Waterbury JB. 1971. Characteristics of two marine nitrite oxidizing bacteria, *Nitrospina gracilis* nov. gen. nov. sp. and *Nitrococcus mobilis* nov. gen. nov. sp. *Archives of Microbiology* 77:203–230.
- Teske A, Alm E, Regan JM, Toze S, Rittmann BE, Stahl DA. 1994. Evolutionary relationships among ammonia- and nitrite-oxidizing bacteria. *J Bacteriol* 176:6623–6630. <https://doi.org/10.1128/jb.176.21.6623-6630.1994>.
- Bayer B, Saito MA, McIlvin MR, Lückner S, Moran DM, Lankiewicz TS, Dupont CL, Santoro AE. 2021. Metabolic versatility of the nitrite-oxidizing bacterium *Nitrospira marina* and its proteomic response to oxygen-limited conditions. *ISME J* 15:1025–1039. <https://doi.org/10.1038/s41396-020-00828-3>.
- JGI. 2012. Bacterial genomic DNA isolation using CTAB. <https://jgi.doe.gov/wp-content/uploads/2014/02/JGI-Bacterial-DNA-isolation-CTAB-Protocol-2012.pdf>.
- Eid J, Fehr A, Gray J, Luong K, Lyle J, Otto G, Peluso P, Rank D, Baybayan P, Bettman B, Bibillo A, Bjornson K, Chaudhuri B, Christians F, Cicero R, Clark S, Dalal R, deWinter A, Dixon J, Foquet M, Gaertner A, Hardenbol P, Heiner C, Hester K, Holden D, Kearns G, Kong X, Kuse R, Lacroix Y, Lin S, Lundquist P, Ma C, Marks P, Maxham M, Murphy D, Park I, Pham T, Phillips M, Roy J, Sebra R, Shen G, Sorenson J, Tomaney A, Travers K, Trulson M, Veceli J, Wegener J, Wu D, Yang A, Zaccarin D, et al. 2009. Real-time DNA sequencing from single polymerase molecules. *Science* 323:133–138. <https://doi.org/10.1126/science.1162986>.
- Chin CS, Alexander DH, Marks P, Klammer AA, Drake J, Heiner C, Clum A, Copeland A, Huddleston J, Eichler EE, Turner SW, Korlach J. 2013. Nonhybrid, finished microbial genome assemblies from long-read SMRT sequencing data. *Nat Methods* 10:563–569. <https://doi.org/10.1038/nmeth.2474>.
- Hunt M, De Silva N, Otto TD, Parkhill J, Keane JA, Harris SR. 2015. Circlator: automated circularization of genome assemblies using long sequencing reads. *Genome Biol* 16:294. <https://doi.org/10.1186/s13059-015-0849-0>.
- Kurtz S, Phillippy A, Delcher AL, Smoot M, Shumway M, Antonescu C, Salzberg SL. 2004. Versatile and open software for comparing large genomes. *Genome Biol* 5:R12. <https://doi.org/10.1186/gb-2004-5-2-r12>.
- Varghese NJ, Mukherjee S, Ivanova N, Konstantinidis KT, Mavrommatis K, Kyrpides NC, Pati A. 2015. Microbial species delineation using whole genome sequences. *Nucleic Acids Res* 43:6761–6771. <https://doi.org/10.1093/nar/gkv657>.
- Lückner S, Nowka B, Rattei T, Spieck E, Daims H. 2013. The genome of *Nitrospina gracilis* illuminates the metabolism and evolution of the major marine nitrite oxidizer. *Front Microbiol* 4:27. <https://doi.org/10.3389/fmicb.2013.00027>.
- Subramaniam A, Yager PL, Carpenter EJ, Mahaffey C, Björkman K, Cooley S, Kustka AB, Montoya JP, Sañudo-Wilhelmy SA, Shipe R, Capone DG. 2008. Amazon River enhances diazotrophy and carbon sequestration in the tropical North Atlantic Ocean. *Proc Natl Acad Sci U S A* 105:10460–10465. <https://doi.org/10.1073/pnas.0710279105>.
- Moore JK, Braucher O. 2007. Observations of dissolved iron concentrations in the World Ocean: implications and constraints for ocean biogeochemical models. *Biogeosciences* 4:1241–1277. <https://doi.org/10.5194/bgd-4-1241-2007>.