

# **UCLA**

## **UCLA Previously Published Works**

### **Title**

An update on the pharmacotherapeutic options and treatment strategies for systemic sclerosis.

### **Permalink**

<https://escholarship.org/uc/item/5h8669kz>

### **Journal**

Expert Opinion on Pharmacotherapy, 21(16)

### **Authors**

McMahan, Zsuzsanna

Volkman, Elizabeth

### **Publication Date**

2020-11-01

### **DOI**

10.1080/14656566.2020.1793960

Peer reviewed



Published in final edited form as:

*Expert Opin Pharmacother.* 2020 November ; 21(16): 2041–2056. doi:10.1080/14656566.2020.1793960.

## An update on the pharmacotherapeutic options and treatment strategies for systemic sclerosis

Zsuzsanna H. McMahan<sup>a</sup>, Elizabeth R. Volkman<sup>b</sup>

<sup>a</sup>Division of Rheumatology, Johns Hopkins University, Baltimore, MD, USA;

<sup>b</sup>Division of Rheumatology, University of California, Los Angeles, CA, USA

### Abstract

**Introduction:** Systemic sclerosis (SSc) is a multi-dimensional connective tissue disease of unknown etiology. Given the immense clinical complexity of SSc, the treatment of this condition is not standardized and considerable heterogeneity exists in SSc management approaches. The purpose of this article is to highlight novel therapeutic strategies and new medications under development for the treatment of systemic sclerosis (SSc).

**Areas covered:** Herein, the authors focus primarily on recently completed clinical trials and phase 3 and 4 clinical trials of therapeutic agents that show promise in SSc. This review is organized by the clinical complications that occur in SSc, for which novel treatment strategies are under study.

**Expert opinion:** Combining therapies to address the individual manifestations of SSc is a cornerstone to the comprehensive management of this condition. Therapeutic strategies must take into account the organs involved, the level of disease activity in each area, and the disease stage. Controlling the complex biological network, progressive vasculopathy and fibrosis, as well as manifestations of end-organ dysfunction are all critical considerations when determining the best treatment approach for SSc.

### Keywords

Scleroderma; systemic sclerosis; treatment; therapy; update

## 1. Introduction

Systemic sclerosis (SSc) is a multi-dimensional connective tissue disease of unknown etiology. The pathological hallmarks of this condition (e.g. fibrosis, inflammation, autoimmunity, vasculopathy) uniquely converge to cause varying degrees of organ system dysfunction and damage. The skin is the most common organ system affected in SSc,

---

**CONTACT** Zsuzsanna H. McMahan zcmcaha1@jhmi.edu Division of Rheumatology, Johns Hopkins University, Baltimore, MD 21224, USA.

Declaration of interest

ER Volkman has received grant support from Corbus Pharmaceuticals and Forbius, and consulting fees from Forbius and Boehringer Ingelheim. ZH McMahan has received salary support from Corbus Pharmaceuticals for her work as a subinvestigator on the Lenabasum study. The authors have no other relevant affiliations or financial involvement with any organization or entity with a financial interest in or financial conflict with the subject matter or materials discussed in the manuscript apart from those disclosed.

followed by the gastro-intestinal (GI) tract and then the lungs. While SSc often presents with the onset of Raynaud phenomenon, the evolution of the disease vastly differs based on a number of defined (i.e. SSc cutaneous subtype, sex, auto-antibody profile) and yet to be defined (i.e. genetics, epigenetics, environmental factors) variables.

### 1.1. Determining when therapy is indicated

Given the immense clinical complexity of SSc, the treatment of this condition is not standardized and considerable heterogeneity exists in SSc management approaches. General treatment algorithms have been proposed [1]; however, emerging and experimental therapies are changing the land-scape of the treatment paradigm for SSc [2,3]. Selection of therapy is often guided by the extent and severity of organ involvement (Figure 1). For example, if a patient has diffuse cutaneous sclerosis and interstitial lung disease (ILD), preferred interventions would target both organ systems simultaneously.

Beyond the extent and severity of organ involvement, other factors affect treatment decisions, such as the duration of disease and its current activity. For instance, the approach to managing an SSc patient with long-standing ILD and stable lung function may contrast with the approach adopted to treat an SSc patient with new-onset ILD presenting with a decline in lung function. The latter scenario suggests active disease and would likely require a more aggressive treatment strategy, while the former scenario may require a close monitoring approach.

In addition, specific patient characteristics may also contribute to treatment decisions. Factors such as sex and auto-antibody status may confer a heightened or lessened risk for the progression of disease within certain organ systems. As an example, male patients with SSc have an increased risk of progression of ILD compared with female patients [4], while patients who possess the centromere antibody seem to have a decreased risk of ILD progression compared with patients who do not possess these antibodies [5]. Table 1 summarizes some of the key factors that often affect treatment decisions in SSc patients beyond the organ system affected.

The present review describes the current, as well as late-stage, investigational therapies for SSc.

## 2. Novel strategies for treating complications of SSC

While the review is organized by organ system, we encourage a holistic management approach that targets treating the patient as a whole, considering the factors outlined in Table 1. Where applicable, we will point out certain therapies that may treat multiple dimensions of this disease, as well as therapies that may be combined in potentially synergistic ways. The review will conclude with a review of the areas where more drug discovery and development are needed. This review specifically focuses on late-stage clinical trials, and therefore preclinical and earlier stages of investigations are not in the scope of this review but have been recently summarized else-where [3,6].

## 2.1. Cutaneous sclerosis

Nearly all patients with SSc have cutaneous involvement. The extent of cutaneous involvement varies immensely with some patients presenting with rapidly evolving diffuse cutaneous sclerosis (dcSSc) and others presenting with stable limited cutaneous sclerosis (lcSSc). In patients with dcSSc, skin thickening occurs in the hands and feet and extends proximally beyond the elbows or knees and often involves the trunk; whereas in patients with lcSSc, skin thickening is confined to the distal extremities, or may only affect the fingers (i.e. scler-odactyly) [7]. The cutaneous manifestations of SSc not only cause functional disability but they substantially contribute to pain, psychological distress, and body image dissatisfaction [8].

**2.1.1. Existing therapies for cutaneous sclerosis**—While the natural history of cutaneous sclerosis in dcSSc often involves a gradual improvement over time [9], randomized-controlled trials (RCTs) have demonstrated that treatment with immunosuppression can lead to a greater reduction in the extent of cutaneous sclerosis [10]. The mRSS is a measure of skin thickness and is often used as the primary outcome in clinical trials of patients with dcSSc. For instance, in Scleroderma Lung Study (SLS) I (12 months of oral cyclophosphamide [CYC] versus 12 months of placebo for SSc-ILD), patients with dcSSc randomized to CYC had an improvement in their mRSS of  $-5.3$  at 12 months; whereas the dcSSc patients randomized to placebo group had an improvement in mRSS of only  $-1.7$  at 12 months ( $P = 0.008$ ) [10]. The mRSS (score range of 0 [no skin thickening] to 51 [most severe]) is a measure of skin thickness and is often used as the primary outcome in clinical trials of patients with dcSSc. A number of studies have demonstrated that it is sensitive to change in the context of SSc clinical trials; however, it should ideally be performed by the same examiner in a trial who has experience with this assessment to reduce inter-observer variability [11].

In addition to CYC, studies have demonstrated that treatment with mycophenolate mofetil (MMF) leads to improvements in cutaneous sclerosis [12,13]. In SLS II (12 months of CYC, followed by 12 months of placebo versus 24 months of MMF), change in mRSS was a key secondary outcome and improved in a clinically meaningful manner in both treatment groups (CYC:  $-5.35$ ; MMF:  $-4.90$ ) [11,12]. In a post-hoc analysis comparing the MMF arm of SLS II, with the placebo arm of SLS I, patients in the MMF arm experienced a greater reduction in the extent of cutaneous sclerosis at 24 months ( $-4.9$  versus  $-2.4$ , respectively) in all patients, and in patients with dcSSc ( $-6.3$  versus  $-3.9$ , respectively), and this difference was statistically significant [14]. In this post-hoc analysis, mRSS was also a key secondary outcome.

In light of the evidence from the aforementioned studies, MMF and CYC are often considered first-line therapies for the treatment of cutaneous sclerosis in patients with dcSSc, with CYC typically reserved for patients with severe cutaneous sclerosis who do not respond to MMF. Although the most recent EULAR treatment guidelines for SSc [15] include methotrexate as a first-line treatment for cutaneous sclerosis in SSc, the evidence for this approach is poor as there have been no RCTs comparing methotrexate with placebo for the treatment of cutaneous sclerosis in SSc. Hematopoietic stem cell transplant (HSCT) is a

potentially viable option for patients with rapidly evolving dcSSc refractory to treatment with immunosuppression. Given the risks (e.g. treatment-related side effects and early treatment-related mortality) associated with this procedure, this option is typically reserved for patients with rapidly progressive cutaneous sclerosis and underlying organ involvement refractory to treatment with immunosuppression. In the Autologous Stem Cell Transplantation International Scleroderma (ASTIS) trial (autologous HSCT versus monthly intravenous CYC for 12 months), the mean improvement in mRSS from baseline to 24 months was greater in the HSCT group (−19.9) than in the control group (−9.8) ( $P < .001$ ) [16]. In the Scleroderma: Cyclophosphamide Or Transplantation (SCOT) trial (autologous HSCT versus monthly intravenous CYC for 12 months) [17], numerically more patients in the HSCT arm experienced a clinically meaningful improvement in mRSS compared with patients in the CYC arm. There was also a long-term survival benefit associated with HSCT in the SCOT trial; however, the results of the survival analysis should be interpreted with caution since the comparator arm only received CYC for 12 months, and typically patients with SSc receive immunosuppression beyond 1 year.

**2.1.2. Emerging therapies for cutaneous sclerosis**—In clinical practice, a considerable proportion of patients with dcSSc do not respond adequately or are intolerant to treatment with CYC or MMF. New treatment options are needed for patients who possess a progressive, treatment-resistant phenotype of cutaneous sclerosis. Promising therapeutic agents, such as tocilizumab [18] and abatacept [19], have failed to meet the primary endpoint of change in mRSS in RCTs, leaving an unmet clinical need (Table 2).

The novel agonist to the cannabinoid receptor type 2 (CB2), lenabasum, was found to reduce the expression of genes and proteins in key inflammatory and fibrotic pathways implicated in the pathogenesis of SSc [20]. In a phase II study of 43 patients with dcSSc, treatment with lenabasum was associated with an improvement in the American College of Rheumatology (ACR) Combined Response Index in diffuse cutaneous Systemic Sclerosis (CRISS) score (33% achieved a positive response in the lenabasum arm compared with 0% in the placebo arm) and there was a trend for a significant treatment effect on mRSS ( $P = 0.085$ ) [21]. The CRISS is a composite response index derived from patients with dcSSc and is comprised of the following endpoints with differential weighting: mRSS, forced vital capacity (FVC)%-predicted, Health Assessment Questionnaire Disability Index (HAQ-DI), Patient Global Assessment, and Physician Global Assessment [22]. In the open-label extension to the Phase II study, a clinically meaningful improvement in mRSS was observed under treatment with lenabasum (−8.4 points at 6 months, −9.8 points at 12 months, and −10.7 points at 18 months), although given that there was no placebo arm in the extension study, it is unclear whether this represents a continued treatment effect versus the natural history of the disease [23]. The drug appears to be well-tolerated, and the phase III study (NCT03398837) has completed enrollment ( $N = 365$ ), and the expected study completion date is in 2020.

The anti-fibrotic, pirfenidone, is currently under investigation in a phase II trial for SSc-ILD (SLS III), in which a key secondary endpoint is a change in mRSS (NCT03221257). An open-label phase II study of pirfenidone in SSc-ILD suggested an acceptable safety and tolerability profile of this agent [24]. In SLS III, pirfenidone is combined with upfront MMF therapy to determine whether combining an anti-fibrotic with MMF leads to an improvement

in skin and lung outcomes compared with MMF alone. Another novel anti-fibrotic, nintedanib was recently approved by the FDA for the treatment of SSc-ILD as described further below; however, this phase III study failed to detect a significant treatment effect for nintedanib on mRSS [25].

The B-cell depleting agent, rituximab, is frequently used in clinical practice to treat patients with dcSSc who fail to respond to conventional immunosuppressive therapies. Large RCTs of rituximab for SSc have not been performed; however, one relatively small RCT demonstrated that rituximab treatment led to a greater improvement in mRSS at 6 months compared with CYC [26]. In addition, small, open-label studies have demonstrated favorable effects on skin thickening [27,28]. A case-control analysis of 63 SSc patients from European Scleroderma Trial and Research (EUSTAR) cohort demonstrated that the patients with dcSSc who received rituximab experienced a greater reduction in MRSS compared with matched controls ( $N = 25$ ;  $-24.0 \pm 5.2\%$  vs  $-7.7 \pm 4.3\%$ ;  $p = 0.03$ ) [29]. A small observational study of 18 patients with SSc demonstrated that combining rituximab with MMF is safe and leads to significant improvements in mRSS [30]. As discussed further below, there is also evidence that rituximab may possess disease-modifying effects on ILD in SSc, suggesting that this agent may play a role in the management of patients with treatment-resistant skin and lung disease. A phase II study combining rituximab with belimumab and MMF is ongoing (NCT03222492).

## 2.2. Interstitial lung disease

The majority of patients with SSc have interstitial abnormalities identified on high-resolution computed tomography (HRCT) imaging [31]. Patients with both lcSSc and dcSSc can develop ILD, and when present, the development of ILD in SSc increases the risk of mortality by at least threefold [32]. Furthermore, data from observational studies in both the US [33] and Europe [34] suggest that ILD is likely the leading cause of SSc-related death. While progression rates of ILD in SSc vary [35], patients typically experience progression within the first 4–5 years of their presentation [33]; therefore, treatment should be initiated early in affected patients to prevent ILD progression, especially in patients who possess features that increase their likelihood of ILD progression (e.g. Scl-70 antibody presence, male sex, African American race) [36].

**2.2.1. Existing therapies for SSc-ILD**—The treatment paradigm for SSc-ILD has historically involved the initiation of immunosuppressive therapy in patients who exhibit signs of or risk factors for the progression of ILD. In SLS I (oral CYC versus placebo), treatment with CYC led to significant improvements in the FVC%-predicted [10], total lung capacity (TLC)%-predicted [10], radiographic fibrosis [37], and quality of life [38] at 12 months. However, a year after cessation of CYC, there was no difference in the FVC%-predicted and TLC%-predicted between patients randomized to CYC versus placebo [39]. Moreover, during a 12-year follow-up period, there was no difference in long-term morbidity and mortality outcomes between patients randomized to CYC versus placebo in SLS I [40], suggesting that 1 year of treatment does not lead to a sustained improvement in health outcomes in SSc-ILD, and maintenance therapy is important. In SLS II [12], treatment with MMF for 24 months led to a significant improvement in FVC%-predicted [12], quantitative

radiographic fibrosis [37], and patient-reported outcomes [41]. There was no significant difference in efficacy outcomes between patients randomized to MMF versus CYC in SLS II, although MMF was better tolerated and had a more favorable safety profile compared with CYC in this study [12]. For instance, there were 15 versus 4 premature study drug withdrawals in the CYC and MMF arms, respectively, and over twice as many deaths in the CYC arm (11 CYC; 5 MMF), with most deaths attributable to progressive of ILD. When the MMF arm of SLS II was compared with the placebo arm of SLS I (with adjustment for baseline disease severity), patients treated with MMF demonstrated a significant improvement in FVC % predicted, DLCO% predicted, and dyspnea compared with the placebo group [14]. The collective findings from SLS I and II suggest that: (1) the duration of treatment for SSc-ILD should be longer than 1 year to yield a sustained benefit; (2) treatment with MMF and CYC both lead to short-term improvements in lung function, radiographic fibrosis, and quality of life; and (3) MMF appears to be safer and better tolerated than CYC.

**2.2.2. Emerging therapies for SSc-ILD**—Similar to cutaneous sclerosis, a substantial portion of patients with SSc-ILD experience progression of ILD despite treatment with CYC or MMF. HSCT may ameliorate ILD in carefully selected patients with dcSSc [16,17]. In the ASTIS study, in which 86% and 87% of the patients randomized to HSCT and CYC, respectively, had ILD, more patients in the HSCT arm experienced an improvement in the FVC compared with the CYC arm [16]. Similarly, in the SCOT study, in which 100% and 95% of the patients randomized to HSCT and CYC, respectively, had ILD, fewer patients randomized to HSCT experienced respiratory failure in the HSCT arm (N = 5) compared with the CYC arm (N = 13) at 1 year [17].

The tyrosine kinase inhibitor, nintedanib, was recently found to slow the rate of progression of ILD in the largest RCT (SENSCIS) ever conducted in SSc (N = 576) [25]. In this study, patients were randomized to 12 months of nintedanib versus 12 months of placebo, and the entry criteria permitted patients to continue treatment with MMF if they were taking a stable dose of MMF for at least 6 months prior to screening. Approximately half of all patients in each study arm were taking MMF at baseline. The rate of decline of FVC (mL/year) was greatest in the placebo arm not on background MMF at baseline (−119.3), and lowest in the nintedanib arm taking MMF at baseline (−40.2), and similar in the nintedanib alone arm (−63.9) and placebo arm on background MMF at baseline (−66.5). In contrast with MMF and CYC [12,41], treatment with nintedanib did not lead to an improvement in self-reported dyspnea as measured by the St. George's Respiratory Questionnaire (SGRQ), nor mRSS [25]. It is unclear why patients randomized to nintedanib did not experience an improvement in the SGRQ relative to placebo; however, it may be because responses to this questionnaire are influenced by extra-pulmonary manifestations of SSc not targeted by nintedanib (e.g. cutaneous sclerosis, arthritis). In terms of safety, patients on combination therapy (MMF + nintedanib) appeared to have a similar adverse event profile compared with patients on nintedanib alone; however, the majority of patients on nintedanib (76%) reported diarrhea (compared with 32% in the placebo arm), and this should be taken into consideration, particularly when treating patients who have co-morbid SSc-related lower gastrointestinal tract involvement.



Although it is difficult to draw meaningful conclusions about the benefits of upfront combination therapy (MMF + nintedanib) for SSc-ILD since patients were not randomized to MMF in the SENSICIS trial, the results support the hypothesis that using nintedanib as add-on therapy to MMF may be advantageous from a lung function standpoint. To this end, SLS III (NCT03221257) was conceived and designed to test the hypothesis that combining an anti-fibrotic therapy (e.g. pirfenidone) with a cytotoxic, immunosuppressive agent (e.g. mycophenolate) upfront may lead to an improved and faster treatment response compared with using an immunosuppressive agent alone. In SLS III, patients with SSc-ILD who are treatment naïve or early in their course of treatment with MMF (treatment with MMF for 6 months prior to enrollment) are randomized to either treatment with MMF alone or treatment with MMF plus pirfenidone for 18 months. In the prior open-label study phase II study on pirfenidone for SSc, MMF was combined with pirfenidone in 63.5% of patients with good tolerability [24].

The monoclonal antibody against the interleukin (IL)-6 receptor, tocilizumab, may play a role in the treatment of ILD in SSc, particularly in patients with early dcSSc with inflammatory features. Two RCTs have investigated the safety and efficacy of tocilizumab in patients with early dSSc with elevated acute phase reactant proteins, and while the primary outcome for these studies was mRSS, the FVC was a key secondary endpoint [18,42]. In the phase II faSScinate trial, fewer patients in the tocilizumab arm experienced a decline in FVC at 48 weeks than in the placebo group ( $p = 0.0373$ ) [42]. Furthermore, the FVC treatment effect appeared to be sustained in the open-label extension period [18]. The phase III trial of tocilizumab for SSc did not meet the primary endpoint of mRSS [43].

As with cutaneous sclerosis treatment, rituximab is frequently used to treat SSc patients with progressive ILD resistant to CYC and MMF. Evidence for using rituximab in SSc-ILD is largely based on observational studies [27–30] and one small RCT of short duration, which demonstrated an improvement in lung function over 6 months in patients randomized to rituximab versus CYC [26]. In addition, a nested case-control analysis of the EUSTAR cohort found that the SSc-ILD patients ( $N = 9$ ) treated with rituximab experienced stability in the FVC; whereas the matched-control patients experienced a significant decline in FVC over a median of 6 months [29].

To date, only one small RCT has been published on rituximab for SSc-ILD, and this study found that treatment with rituximab was associated with a significant improvement in lung function compared with placebo [44]. In this study, the majority of the rituximab-treated patients also experienced a reduction in ground glass opacities (but not reticulations) on HRCT at 18 months (based on visual assessment) [44]. A larger double-blind, RCT comparing IV CYC (600 mg/m<sup>2</sup> body surface area monthly for 6 months) versus rituximab (1 g at baseline and at 2 weeks) for connective tissue disease-related ILD, including SSc-ILD, is currently underway in the UK (NCT01862926). The primary outcome for this study is the change in FVC at 48 weeks [45] (Table 3).

### 2.3. Raynaud phenomenon

Approximately 90–95% of patients with SSc have Raynaud phenomenon (RP), and it is often the presenting feature of SSc. It is characterized by peripheral vasospasm that leads to



a transient discoloration (erythema, cyanosis, and/or pallor) and/or numbness in the digits immediately following specific triggers, such as exposure to cold temperatures, relative changes in the temperature of the surrounding environment, and/or exposure to stress. The complications of RP are particularly severe in SSc, as the vasospasm compounds the already reduced blood flow in vessels affected by SSc vasculopathy [46]. Up to about 50% of patients with SSc are affected by RP may experience ischemic complications such as digital ulcers, pits, or gangrene [47]. This subset of patients requires more aggressive therapy with medication to prevent the loss of digital tissue. Recent studies have focused on determining whether specific drugs are more effective in preventing or healing digital ulcers, as some medications are known to be more effective for one than the other.

**2.3.1. Existing therapies for SSc-Raynaud phenomenon and its ischemic complications**—The current standard of care for the management of patients experiencing RP-related digital ischemia involves minimizing environmental triggers and initiating medications that maximize peripheral blood flow (e.g. calcium channel blockers, phosphodiesterase inhibitors, endothelial receptor antagonists). On-demand therapy with phosphodiesterase inhibitors (i.e. sildenafil) for the treatment of primary or secondary RP was recently shown not to have clinically relevant efficacy, given the significant heterogeneity in patient responses [48]. Few studies have focused on the prevention of RP-associated ulcers. In the past decade, these areas of study have become areas of increasing interest (Table 4).

**2.3.2. Emerging therapies and new applications of existing therapies for SSc-associated Raynaud's**

**2.3.2.1. Phosphodiesterase inhibitors.:** The specific roles of phosphodiesterase inhibitors in healing RP-associated digital ulcers in SSc continues to be an active area of investigation, as this class of medications improves digital blood flow, and has been shown to have significant efficacy in secondary Raynaud's [49,50]. The SEDUCE study group recently evaluated the effects of sildenafil on ischemic digital ulcer healing in SSc in a randomized placebo-controlled trial [51]. Eighty-three patients with a total of 192 digital ulcers were included in the intention-to-treat analysis (89 in the sildenafil group, 103 in the placebo group), however the primary endpoint was not reached.

In order to expand therapeutic options for the treatment of SSc-related RP attacks, tadalafil was studied in a prospective double-blind placebo-controlled crossover study. Patients were randomized to receive a fixed dose of tadalafil at 20 mg daily or placebo for a 4 week period, and the RP condition score, frequency of RP episodes, and duration of RP episodes between treatment groups was compared. While tadalafil was well-tolerated, there was no significant difference in treatment response between the tadalafil arm and the placebo arm [52].

However, a subsequent study examined the benefits of tadalafil as an add-on therapy for the healing and prevention of digital ulcers. This double-blind, randomized, cross-over trial evaluated the efficacy of tadalafil as add-on therapy in patients with treatment-resistant RP. Patients with SSc and MCTD who were on vasodilators, but still having 4 or more episodes

of RP attacks per week were included and randomized to receive either tadalafil 20 mg per day or placebo. Significant improvements from baseline were observed in several outcome measures during tadalafil therapy compared to placebo, including mean daily frequency of RP episodes, mean daily duration of RP, and mean daily RP condition score. In the tadalafil group, all 24 digital lesions healed compared to only 3 of 13 in the placebo group, and only one new ulcer was reported during tadalafil therapy compared to 13 in patients on placebo therapy [53]. Collectively, these data suggest that phosphodiesterase-inhibitors play an important role in the healing and prevention of digital ulcers in SSc-related RP.

**2.3.2.2. Prostacyclins and prostaglandins.:** Prostacyclins and prostaglandins continue to serve as key therapies used to treat challenging cases of RP in the context of SSc. Iloprost, for example, serves as a standard treatment of existing digital ulcers [54–56]. It is currently indicated for patients with severe disabling Raynaud’s unresponsive to other therapies for the prevention of ischemic pain, peripheral ulcers, and necrosis of the digits to prevent amputation. It may be administered by one of three schedules: (a) 3-day schedule as an inpatient for RP/SSc; (b) 5-day schedule as an outpatient for RP/SSc; or (c) continuous infusion for the treatment of patient with active or extensive digital ulcers, severe digital ischemia, or for patients who cannot tolerate higher rates of the infusion. Details of the complex infusion protocols can be found here [57].

Treprostanil, a synthetic analog of prostacyclin (PGI<sub>2</sub>), has also been studied as a therapy for patients with SSc-associated Raynaud’s and/or digital ulcers, in both oral and topical formulations. Oral treprostanil initially showed promise in patients with SSc and digital ischemia in a phase 1 study, where effective absorption and temporal associations with improved cutaneous perfusion and temperature were noted [58]. The recurrence of digital ulcers in patients with SSc after discontinuation of oral treprostanil was also noted in a multicentered retrospective study, also suggesting some benefit [59]. However, the association between vascular biomarkers and digital ulcerations in SSc was recently evaluated using the DISTOL-1 randomized controlled trial cohort, and a lack of strong response to any vascular, angiogenic, or inflammatory markers suggested that these pathways are not primary drivers in the development of digital ulcer clinical outcomes in an SSc population [60].

The effects of topical treprostanil have also been a focus of recent work. Treprostanil iontophoresis in patients with SSc was also studied to determine its ability to improve digital blood flow during local (hand) cooling [61]. It showed promising as digital treprostanil iontophoresis shifted skin blood flow upward during local cooling on the hand and during the initial rewarming phase in patients with SSc. The safety profile of treprostanil hydrogel iontophoresis was also recently examined in a 2-stage randomized, placebo-controlled single ascending-dose study among healthy volunteers and patients with SSc-related digital ulcers and was found to be fairly well-tolerated, with 2 minimal local adverse effects reported among 5 SSc patients with digital ulcers [62].

A phase 2 multi-center, double-blind, RCT is currently examining the effects of intravenous iloprost on RP in patients with SSc. Forty-one patients were enrolled and randomized to receive either intravenous iloprost or placebo infusion, and the primary endpoint is the

change in weekly frequency of symptomatic RP attacks at 21 days. Results of this study are pending ([NCT03867097](#)), and the phase 3 trial is enrolling ([NCT04040322](#)).

**2.3.2.3. Endothelin receptor antagonists.:** Endothelin-1 is a well-recognized promoter of vasculopathy in SSc. Prior studies have demonstrated that bosentan, a dual endothelin receptor antagonist, successfully prevents RP-related digital ulcers in SSc. In an earlier clinical trial, RAPIDS-1, which enrolled SSc patients with or without digital ulcers at baseline, bosentan significantly reduced the number of new digital ulcers compared to placebo. The purpose of subsequent RAPIDS-2 trial is to evaluate the effects of bosentan compared to placebo on ulcer prevention and healing over a 24-week treatment period. Patients received bosentan 62.5 mg twice daily for 4 weeks and then 125 mg twice daily for 20 weeks or the equivalent placebo, and time to complete healing of the cardinal ulcer and the total number of new digital ulcers per patient over 24 weeks was assessed [63]. The investigators found that bosentan treatment was associated with a 30% reduction in the number of new ulcers compared with placebo (mean  $\pm$  standard error:  $1.9 \pm 0.2$  vs  $2.7 \pm 0.3$  new ulcers;  $p = 0.04$ ), and that the effect was greater in patients who entered the trial with more ulcers. There was no significant difference between treatments in healing rate of the cardinal ulcer [63]. This data suggests that bosentan may play an important role in digital ulcer prevention in SSc, but its role in digital ulcer healing is unclear.

As bosentan demonstrated success in its positive effects on digital ulcer preventions, an international group of investigators evaluated the efficacy of a newer dual endothelin receptor antagonist, macitentan, in reducing the number of new digital ulcers in SSc patients. Two double-blind placebo-controlled trials were conducted and enrolled patients with SSc and active digital ulcers, and patients were randomized to receive oral doses of 3 mg macitentan, 10 mg of macitentan, or placebo once daily and stratified according to the number of digital ulcers at baseline. In both trials, patients on placebo had a lower mean number of new digital ulcers and had fewer adverse events than patients in both treatment groups. As a result, the investigators recommended against using macitentan for the prevention of digital ulcers in this patient population [64] ([NCT01474109](#); [NCT01474122](#)).

**2.3.2.4. Topical agents.:** The data supporting the efficacy of a variety of topical agents in the management of RP have been variable. A systematic review and meta-analysis recently examined the effects of local topical nitrates in primary and secondary RP with respect to parameters of digital blood flow, and clinical severity was recently completed [65]. Seven placebo-controlled trials were included (346 patients) in the meta-analysis, with 4 trials involving nitroglycerin ointments, 2 involving the nitroglycerin gel vehicle MQX-503, and 1 involving a compounded nitrite. The results of the meta-analysis demonstrated a moderate-to-large treatment effect in RP (standardized mean difference [SMD] = 0.70; 95% CI, 0.35–1.05;  $P < .0001$ ). Subgroup analyses showed a large treatment effect in secondary RP (SMD = 0.95; 95% CI, 0.25–1.65;  $P = .008$ ) and moderate effect in primary RP (SMD = 0.45; 95% CI, 0.05–0.85;  $P = .03$ ) [65].

**2.3.2.5. Other therapies under study.:** A relatively novel therapy that shows promise in RP involves the local injection of botulinum toxin type A (Btx-A) in the hands of patients with SSc. Btx-A is a neurotoxin that acts as a neuromuscular blocking agent by blocking the

release of acetylcholine from presynaptic nerve terminals, thereby interfering with vascular smooth muscle contraction and enhancing local circulation [66]. In a recent double-blind, RCT, Btx-A was administered (50 units in 2.5 ml sterile saline) in one randomly selected hand and sterile saline (2.5 ml) in the opposite hand [67]. Follow-up at 1 and 4 months post-injection included laser Doppler imaging of hands, patient-reported outcomes, and physical examination. At 1-month follow-up, a significantly greater reduction in average blood flow was observed in Btx-A-treated hands compared to placebo-treated hands ( $p = 0.024$ ). Change in blood flow at a 4-month follow-up was not significantly different between groups. Ultimately, the investigators concluded that there may be a role of Btx-A in treating a subset of patients with RP, and that further studies defining the patients who are most likely to benefit from this intervention are warranted. A phase 3 study is currently recruiting in France to assess whether or not a single injection schedule of BTX-A in both hands improves RP secondary to SSc better than a placebo at 4, 12, and 24 weeks after the treatment. (NCT03717961).

Riociguat, a stimulator of soluble guanylate cyclase which has downstream effects stimulating vasodilation, was recently studied to determine its efficacy and safety in SSc-associated digital ulcers [68]. In this multi-center randomized, double-blind, placebo-controlled pilot study, SSc patients with at least one visible active or painful digital ulcer were enrolled and randomized (1:1 placebo or riociguat maximum of 2.5 mg three times daily) for an 8-week titration period, followed by an 8-week stable dosing period. An optional 16-week open-label extension phase for patients with active DU/reoccurrence of DUs within 1 month of the end of the main treatment phase followed. Ultimately, treatment with riociguat did not reduce the number of digital ulcer burden compared to placebo at 16 weeks.

The effects of statins on endothelial dysfunction and RP in SSc are currently under study by a group in Pittsburgh [69]. In this double-blind, randomized, placebo-controlled trial of atorvastatin 40 mg once daily vs placebo, 24 patients with early diffuse SSc (<3 years of SSc symptoms and RP) were enrolled if they had been on stable RP medications for at least 4 weeks. Improvement in microvascular endothelial function measured by reactive hyperemia index (RHI) was the primary outcome, and secondary outcomes included change in macrovascular endothelial function by brachial flow-mediated (FMD) dilation, and RP severity using the RP condition score (RCS) and visual analog scale (RP-VAS). In the atorvastatin treatment group, 60% (6/10) of patients improved their RHI, compared to 29% (4/14) in the placebo group ( $p = 0.12$ ). No difference in change in peak FMD% was noted between groups. The RCS decreased 2 points in the statin group compared to no change in the placebo ( $p = 0.12$ ; Table 2). While the results demonstrated a non-significant improvement in microvascular endothelial function and RCS scores with the treatment of atorvastatin, the number of patients enrolled into the trial was small and thus it may have been underpowered to capture a significant difference between groups.

Although acetylsalicylic acid (ASA) has been available for over 100 years, it has not been systematically studied in SSc, and it may potentially play a role in preventing SSc-related vascular injury. An ongoing study in Brazil aims to evaluate the effectiveness of ASA on microcirculation alterations in SSc patients. In this phase 4 placebo-controlled clinical trial,

70 patients will be randomized to take either 100 mg daily ASA or placebo for 4 weeks. Outcome measures will include periungual panoramic capillary microscopy, videocapillaroscopy and laser Doppler imaging, as well as a panel of vascular biomarkers. This study is still recruiting and results are not yet available.

## 2.4. Pulmonary hypertension

Pulmonary arterial hypertension (PAH) affects approximately 7–12% of patients with SSc, and it is recognized as a leading cause of SSc-related death [70,71]. Right ventricular failure has been associated with poor survival in SSc-PAH, and modest responses to existing therapies leave SSc patients with PAH with a higher relative mortality compared to PAH from other causes [72,73]. Monotherapy with prostacyclins, phosphodiesterase inhibitors, and endothelin receptor antagonists were the standard of care for many years. However, given the significant morbidity and mortality still associated with SSc-PAH, ongoing studies are now exploring alternative strategies, including the novel approach of combining these therapies (Table 5). [74] In addition, because approximately 50–70% of the pulmonary vasculature needs to be affected or obstructed before resting mPAP is elevated, investigators are also now exploring the treatment of exercise pulmonary hypertension (abnormal hemodynamic response to exercise), which is likely a marker of early pulmonary vascular disease. Determining whether the initiation of therapy at this earlier stage of PAH is beneficial for patients is another key focus of ongoing investigation [74].

**2.4.1. Novel combinations of existing therapies for SSc-pulmonary arterial hypertension**—Data from the Pulmonary Hypertension Assessment and Recognition of Outcomes in Scleroderma (PHAROS) registry suggest that patients with SSc-related PAH have a significantly shorter time to clinical worsening when treated with phosphodiesterase-5 inhibitor (PDE5i) monotherapy compared to patients treated with endothelin receptor antagonists (ERA) and PDE5i combination therapy [75]. Such data have fueled interest in the study of combination therapy for PAH in SSc. Upfront combination therapy with phosphodiesterase inhibitors and endothelial receptor antagonists was recently studied in the a treatment-naïve group of patients with SSc-PAH. In this multi-center, open-label, clinical trial, patients were treated for 36 weeks with tadalafil 20 mg daily and ambrisentan 5 mg daily, with medication up-titration occurring at week 4 as tolerated. Investigators then compared the change in RV mass on cardiac MRI (standard volumetric cine images) from baseline to 36-week follow-up within subjects. They found that measures of right and left ventricular systolic and diastolic function were improved after treatment. In addition, combination therapy was associated with significant improvements in RV and LV function as measured by cardiac MRI [76], suggesting that combining these medications is a promising therapeutic strategy.

Although trials comparing the efficacy of combination therapy vs. monotherapy for pulmonary hypertension in SSc are limited, one study compared the efficacy and safety of monotherapy with phosphodiesterase inhibitors (sildenafil) versus initial combination therapy with phosphodiesterase inhibitors and endothelial receptor antagonists (sildenafil and bosentan) for treatment of SSc-PAH. In this single-center, double-blind, RCT, 34 patients with SSc-PAH (Pulmonary Artery Systolic Pressures >35 mmHg by echocar-

diography), with a forced vital capacity >60% were randomized to receive either sildenafil plus placebo or to combination therapy with sildenafil plus bosentan for 24 weeks. Ultimately, there were no significant changes between groups in pulmonary artery pressures at 24 weeks from baseline, or in secondary end points including change in 6 Minute Walk Distance or Time To Clinical Worsening (TTCW is defined as the first occurrence of all-cause deaths, PAH related hospitalization, worsening of symptoms defined as a decrease of >15% in 6 min walk distance and worsening of NYHA functional class); however, the investigators recommended larger, better powered studies, as combination therapy was well-tolerated in these patients [77] ([NCT03053739](#)).

Because of the interest in treating early PAH, investigators have also explored the role of treating exercise-induced pulmonary hypertension (PH) in SSc. In cardiology, stress tests are increasingly utilized to better characterize hemodynamic changes, and a strong rationale now exists to suggest that a reduction in pulmonary vascular reserve may be an early signal of subclinical pulmonary hypertension [78]. Some studies now suggest that among SSc patients with normal resting mPAP, an excessive increase in mPAP during exercise coupled with an impairment in vascular dispensability may be indicative of an early stage of pulmonary vasculopathy, associated with reduced survival similar to patients with PH measured at rest [79,80].

A pilot study recently evaluated whether treatment of exercise-induced PH open-label daily ambrisentan positively affects changes in a 24-week interval in hemodynamics and exercise capacity in patients with SSc. Exercise-induced PH was defined as a mean pulmonary artery pressure of >30 mm Hg with maximum exercise and a transpulmonary gradient (TPG) of >15 mm Hg. Patients had normal hemodynamics at rest and were treated with 5–10 mg of ambrisentan daily. From baseline to 24 weeks, significant changes were identified in mean exercise pulmonary vascular resistance, mean 6 minute walk distance, mean exercise cardiac output, mean pulmonary artery pressure, and total pulmonary resistance. Placebo-controlled studies are now needed to confirm that these findings are attributable to a drug effect and to define the optimal therapeutic regimen for these patients [81].

The application of statin therapy is novel in the management of PAH. Statins have a vasoprotective effect, and may therefore add value to the management of PAH, a condition in which vascular dysfunction is prominent. The effects of rosuvastatin on ameliorating vascular dysfunction in SSc-related pulmonary hypertension were recently examined in a study of 40 patients with SSc randomized to receive either rosuvastatin or placebo. All participants completed transthoracic echocardiography, 6 minute walk tests, were categorized by WHO functional class, and had tolerability and safety monitored. The results of this phase-3 trial are not yet reported ([NCT00984932](#)).

#### **2.4.2. Emerging therapies for SSc-pulmonary hypertension secondary to ILD**

—A new study is now seeking to understand patients with interstitial lung disease (ILD) and scleroderma who develop pulmonary hypertension (PH) and how they fit into the treatment schema in SSc. Investigators from National Jewish and the University of Pittsburgh are using pressure-volume loops to derive right ventriculo-vascular coupling, pulmonary impedance, and invasive cardiopulmonary exercise testing to compare the efficacy of chronic macitentan



therapy in improving right ventricular hemodynamics, exercise capacity, and symptoms in scleroderma ILD-PH patients with and without PVL. The trial is currently ongoing ([NCT03726398](#)).

#### **2.4.3. Emerging therapies for PAH secondary to connective tissue disease—**

As connective tissue disease-associated PAH has a relatively poor prognosis relative to PAH from other causes, recent trials have focused on identifying novel therapies to improve outcomes for such patients. The safety and efficacy of riociguat were recently evaluated in an exploratory analysis using the 12-week, phase III Pulmonary Arterial hypertension sGC-stimulator Trial (PATENT)-1, and the long-term extension PATENT-2 data. Investigators specifically examined the results of the study in the subset of patients enrolled who had PAH-associated with SSc or another defined connective tissue disease. In the prospectively planned analysis, it was determined that riociguat was well-tolerated and associated with positive trends in several endpoints, including 6-minute walk distance, and that these findings were sustained at 2 years in this patient subgroup, though further studies need to be done [82].

A similar approach was taken to examine the effects of selexipag using the GRIPHON study population, specifically looking at the subset of patients with PAH-associated connective tissue disease. Of the 334 patients with PAH from connective tissue disease in this population, 170 patients had SSc. Selexipag was associated with delayed progression of PAH, as measured by a reduction in PAH related morbidity/mortality events, and was well-tolerated among the PAH-connective tissue disease population, including patients with SSc [83].

### **2.5. Gastrointestinal disease**

The majority of patients with SSc experience gastrointestinal tract involvement. The gastrointestinal (GI) complications of SSc are variable, and include dysmotility, sphincter dysfunction, vascular malformations, and dysbiosis, each of which is managed differently. As there are no measures of immune-mediated disease activity in the GI tract, differentiating disease activity from damage is a major challenge. Furthermore, since no proven disease-modifying therapies exist for SSc-related GI involvement, the management of SSc-GI complications is currently focused on addressing and ameliorating patient symptoms. Patient symptoms may result from a variety of factors including underlying GI dysmotility and microbiota dysbiosis [84,85]. Recent studies have therefore focused on evaluating the role of probiotics in SSc, as well as identifying novel medications that can be used to enhance GI transit in this population.

#### **2.5.1. Emerging therapies for SSc-GI disease**

**2.5.1.1. Probiotics.:** The role of probiotics in patients with GI complications and SSc is unclear. The effects of probiotics on GI symptoms and on the immune system in patients with SSc with moderate-to-severe total scores on the UCLA GIT 2.0 were recently reported [86]. Patients were randomly assigned to receive a daily dose of probiotics (several strains of *Lactobacillus*) or placebo for 8 weeks. Seventy-three patients were randomized to receive probiotics or placebo. While there was no difference in UCLA GIT 2.0 scores, the probiotic



group showed a significant decrease in the proportion of Th17 cells compared to placebo, suggesting a possible immunomodulatory effect in SSc.

In another probiotic RCT, 37 patients with SSc with a moderate to severe total score on the UCLA GIT 2.0 were randomized to receive probiotics (containing *Lactobacillus* and *Bifidobacterium*) and 36 patients were randomized to placebo. Both groups were followed for 8 weeks, completed UCLA GIT 2.0 questionnaires and HAQ-DI surveys. Because probiotics are thought to act by modulating the microbiome and the immune response, the authors also collected serum samples to explore changes in the circulating levels of Th1, Th2, Th17, and regulatory T cells. At week 8, there was no significant difference in total or subdomain UCLA GIT 2.0 scores between the two groups, in the HAQ-DI score, or in the proportion of Th1, Th2, and regulatory T cells; however, the probiotic group did have a significantly decreased proportion of Th17 cells compared to placebo ( $p = 0.003$ ). Overall, the authors concluded that probiotics did not improve GI symptoms in SSc patients [86], however the use of the UCLA GIT total score as an outcome measure may not have been a sensitive enough tool to detect improvement in symptoms of distention, bloating, and diarrhea.

Another study, which focused on a more homogenous SSc population, recently determined that the addition of probiotics may enhance existing therapeutic GI strategies [87]. In this open-label pilot trial, investigators sought to evaluate how treatment with probiotics, antibiotics, or a combination of both compare in the management of GI symptoms in 40 SSc patients with small intestinal bacterial overgrowth (SIBO) assessed by hydrogen breath test. Patients were assigned to one of the three treatment groups (*Saccharomyces boulardii*, metronidazole, or combination therapy) for 24 weeks, and patient-reported outcomes were collected (NIH-GI PROMIS). Interestingly, at the end of the 2 month period, SIBO was eradicated in 55% of the combination therapy group, 33% of the probiotic group, and 25% of the antibiotic treatment group. In addition, the probiotic group and combination therapy groups had decreased diarrhea, abdominal pain, and gas, bloating, and flatulence. These symptomatic improvements were not identified in the antibiotic group. Reductions in expired hydrogen at 45 to 60 min were 48% and 44% in the combination group, 18% and 20% in the antibiotic group, and 53% and 60% in the probiotic groups in the first and second months, respectively ( $p < 0.01$ ). These data support a role for probiotics to alleviate clinical symptoms in a subset of patients with SSc.

**2.5.1.2. Promotility agents.:** The safety and efficacy of prucalopride, a 5HT<sub>4</sub> receptor agonist was recently demonstrated in a cross-over 2 × 2 study [67]. Patients with mild to moderately severe SSc-GI symptoms were enrolled ( $n = 40$ ) and randomized 1:1 to prucalopride 2 mg/day or no treatment for 1 month and vice versa after a 2-week washout period. Prucalopride was significantly associated with more intestinal evacuations and improved orocecal transit times, as well as significant improvements in the UCLA GIT 2.0 constipation, reflux, and bloating scores, suggesting that it may be effective in treating dysmotility symptoms in SSc patients.

Interestingly, buspirone, an oral 5-HT<sub>1A</sub> receptor agonist, may improve the dysfunction of the lower esophageal sphincter in patients with SSc. In an open-label trial, the effects of

buspirone on esophageal motor function and symptoms in SSc patients with esophageal involvement were evaluated. All 30 patients enrolled had symptomatic esophageal involvement, despite PPI use, and underwent high-resolution manometry and CT chest for assessment of motor function and esophageal dilatation, respectively. Visual analog scales were used to score GI symptom severity. In the 22 patients who completed the trial, lower esophageal sphincter resting pressure significantly increased after buspirone administration. Scores for heartburn and regurgitation significantly decreased at 4 weeks compared to baseline. These findings suggest that buspirone could improve symptoms in patients with SSc who report reflux symptoms despite under-going standard treatment [88].

While several other agents have been proposed in case series to target dysmotility in SSc GI disease (e.g. IVIG, pyr-idostigmine), large prospective placebo-controlled trials are lacking and should be a focus of future research.

**2.5.1.3. Other considerations.:** Among patients with severe GI disease, it is important to ensure nutritional goals are met, as the prevalence of malnutrition in SSc is estimated to be between 15% and 58% [89]. A validated screening tool to assess for malnutrition is the Malnutrition Universal Screening Tool (a.k.a. 'MUST'), and it has been recommended by some experts that all SSc patients should be screened [90,91]. This tool includes body mass index (BMI), unintentional weight loss in the preceding 3–6 month period, and an acute disease effect score to estimate the risk of malnutrition. In patients who screen positive, referrals to a nutritionist and gastroenterologist should be considered. Supplemental enteral or parenteral nutrition may be necessary to sustain nutrition and weight in patients with severe disease [89].

### 3. Conclusion

An expanding pipeline of investigational therapeutic agents now exists for treating the unique clinical manifestations of SSc. This review highlighted products in development in phase 3 and phase 4 clinical trials. However, our growing understanding of the pathogenesis of SSc supports the clinical application of several novel therapies, including anti-fibrotic therapies and biologic agents. While these interventions have the potential to change the clinical course of the disease, determining the optimal drug and/or combination of drugs will remain an ongoing focus of future work. In addition, the early identification of organ involvement and application of targeted therapy, prior to the development of severe organ damage, will remain an important priority. We remain optimistic that current drug development and clinical trials will continue to yield promising therapeutic strategies for improving health outcomes for all patients with SSc.

### 4. Expert opinion

In the last decade, the number of therapeutic options available to treat the unique clinical dimensions of SSc has dramatically increased. As highlighted in this review, these agents target different aspects of the immune system and elicit differential effects on various organ systems within and between individual patients. Data from all of the clinical trials conducted to date suggest that SSc is a heterogeneous disease both in terms of manifestations and

response to therapies. This heterogeneity is perhaps the most challenging component of drug discovery and development in this field.

It is our opinion that future therapeutic studies should consider combining therapies that target different aspects of the immune system (e.g. Anti-inflammatory plus anti-fibrotic). Similar to other complex connective tissue diseases that employ combination therapy (e.g. systemic lupus erythematosus and rheumatoid arthritis), patients with SSc may also benefit from this approach. Research studies are therefore needed to understand the optimal timing to add on certain therapies and to determine whether this should be done upfront or whether treatments should be added sequentially over specified time frames.

In addition, improved clinical and biological phenotyping may help homogenize study cohorts to increase the likelihood of detecting significant treatment effects. Our clinical experience has shown us that certain patients derive benefit from IL-6 inhibition, even though the phase III RCT of the IL-6 inhibitor, tocilizumab, did not meet its primary endpoint. To us, this suggests that we need to do a better job of enriching study cohorts to include the patients who may derive the most benefit from the therapeutic intervention under study. While this may constrain local enrollment into studies, improved global recruitment efforts, as demonstrated in the SENSICIS trial [25], may overcome this limitation.

Another hurdle in SSc therapeutics is identifying and defining objective treatment-responsive endpoints. While we have several valid endpoints to study ILD in SSc, endpoints for the other clinical manifestations of SSc are lacking. The GI tract is the perfect example of this. How can we determine whether biologics or anti-fibrotics modify the course of GI disease in SSc if no valid, objective SSc-GI endpoints exist? To this end, we are currently conducting studies to investigate whether the GI microbiome may be a marker of GI disease activity in SSc. However, additional dedicated research efforts primarily aimed at developing and validating endpoints in SSc are greatly needed.

Finally, a significant unmet therapeutic need in SSc is prevention. The collective efforts to halt inflammation, fibrosis, and vascular changes in SSc are typically initiated after there is already evidence of end-organ damage (i.e. loss of lung function, renal insufficiency, etc.). Furthering our knowledge of the pathobiology of SSc may help uncover treatment targets that could curtail the progression of early SSc. The goal here would be to intervene early with an agent that could selectively target key mediators of profibrotic and proinflammatory pathways to reduce the likelihood of a patient developing specific clinical manifestations of SSc, such as ILD.

In summary, despite the fact that the number of clinical trials in SSc has increased in recent years, there is still a great deal more work to do in SSc research to: (a) understand the optimal timing to initiate therapy in SSc; (b) discover how to combine therapies; (c) improve our ability to phenotype patients; (d) define new disease-specific endpoints; and (e) determine the best preventative treatment strategies. We anticipate that research conducted in the next decade will help address these unanswered questions and propel this field forward in new and exciting ways.

## Funding

Z McMahan is supported by the National Institutes of Health/National Institute of Arthritis and Musculoskeletal and Skin Diseases via grant K23 AR071473, the Scleroderma Research Foundation and the Jerome L Greene Foundation. ER Volkman is supported by National Institutes of Health/The National Heart, Lung, and Blood Institute via grant K23 HL150237-01 and the Rheumatology Research Foundation.

### Reviewer disclosures

One referee has received speaker's fees, travel grants, research funding, or compensation for consultancies or board memberships from Abbvie, Actelion, Baxalta/Shire, Bristol-Myers Squibb, Boehringer Ingelheim, Celgene, Chugai, Genzyme, Gilead Biosciences, Hexal/Sandoz, Janssen-Cilag, Merck Sharp and Dohme, Novartis, Pfizer, Roche, Sanofi Pasteur and UCB Pharma. Peer reviewers on this manuscript have no other relevant financial relationships or otherwise to disclose.

## References

Papers of special note have been highlighted as either of interest (●) or of considerable interest (●●) to readers.

1. Fernández-Codina A, Walker KM, Pope JE, et al. Treatment algorithms for systemic sclerosis according to experts. *Arthritis Rheumatol*. 2018 11;70(11):1820–1828. [PubMed: 29781586]
2. Volkman ER, Varga J. Emerging targets of disease-modifying therapy for systemic sclerosis. *Nat Rev Rheumatol*. 2019 4;15(4):208–224. [PubMed: 30796362]
3. Zhang YDJ. Therapeutic molecular targets of SSc-ILD. *J Scleroderma Related Disord*. 2020;5(25):17–30.
4. Peoples C, Medsger TA Jr., Lucas M, et al. Gender differences in systemic sclerosis: relationship to clinical features, serologic status and outcomes. *J Scleroderma Relat Disord*. 2016 May-Aug;1(2):177–240.
5. Steen VD. The many faces of scleroderma. *Rheum Dis Clin North Am*. 2008 2;34(1):1–15;v. [PubMed: 18329529]
6. Sierra-Sepulveda A, Esquinca-Gonzalez A, Benavides-Suarez SA, et al. Systemic sclerosis pathogenesis and emerging therapies, beyond the fibroblast. *Biomed Res Int*. 2019;2019:4569826. [PubMed: 30809542]
7. Volkman ER, Furst DE. Management of systemic sclerosis-related skin disease: a review of existing and experimental therapeutic approaches. *Rheum Dis Clin North Am*. 2015 8;41(3):399–417. [PubMed: 26210126]
8. Medsger TA Jr. Natural history of systemic sclerosis and the assessment of disease activity, severity, functional status, and psychologic well-being. *Rheum Dis Clin North Am*. 2003 5;29(2):255–73, vi. [PubMed: 12841294]
9. Sultan N, Pope JE, Clements PJ. Scleroderma trials study G. The health assessment questionnaire (HAQ) is strongly predictive of good outcome in early diffuse scleroderma: results from an analysis of two randomized controlled trials in early diffuse scleroderma. *Rheumatology (Oxford)*. 2004 4;43(4):472–478. [PubMed: 14679295]
10. Tashkin DP, Elashoff R, Clements PJ, et al. Cyclophosphamide versus placebo in scleroderma lung disease. *N Engl J Med*. 2006 6 22;354(25):2655–2666. [PubMed: 16790698]
11. Khanna D, Furst DE, Clements PJ, et al. Standardization of the modified Rodnan skin score for use in clinical trials of systemic sclerosis. *J Scleroderma Relat Disord*. 2017 Jan-Apr;2(1):11–18. [PubMed: 28516167]
12. Tashkin DP, Roth MD, Clements PJ, et al. Mycophenolate mofetil versus oral cyclophosphamide in scleroderma-related interstitial lung disease (SLS II): a randomised controlled, double-blind, parallel group trial. *Lancet Respir Med*. 2016 9 4;4(9):708–719. [PubMed: 27469583] • This landmark RCT demonstrated that MMF was not only as effective as CYC at improving lung function and radiographic fibrosis in ILD, but was safer and better tolerated than CYC.

13. Namas R, Tashkin DP, Furst DE, et al. Efficacy of mycophenolate mofetil and oral cyclophosphamide on skin thickness: post hoc analyses from two randomized placebo-controlled trials. *Arthritis Care Res (Hoboken)*. 2018 3;70(3):439–444. [PubMed: 28544580]
14. Volkmann ER, Tashkin DP, Li N, et al. Mycophenolate mofetil versus placebo for systemic sclerosis-related interstitial lung disease: an analysis of scleroderma lung studies I and II. *Arthritis Rheumatol*. 2017 7;69(7):1451–1460. [PubMed: 28376288]
15. Kowal-Bielecka O, Fransen J, Avouac J, et al. Update of EULAR recommendations for the treatment of systemic sclerosis. *Ann Rheum Dis*. 2017 8;76(8):1327–1339. [PubMed: 27941129]
16. van Laar JM, Farge D, Sont JK, et al. Autologous hematopoietic stem cell transplantation vs intravenous pulse cyclophosphamide in diffuse cutaneous systemic sclerosis: a randomized clinical trial. *JAMA*. 2014 6 25;311(24):2490–2498. [PubMed: 25058083] • This was the first large, RCT to evaluate the safety and efficacy of HSCT in SSc and demonstrated improved long-term mortality in dsSSc patients randomized to HSCT compared with CYC.
17. Sullivan KM, Goldmuntz EA, Keyes-Elstein L, et al. Myeloablative autologous stem-cell transplantation for severe scleroderma. *N Engl J Med*. 2018 1 4;378(1):35–47. [PubMed: 29298160] • This was the second large, RCT to evaluate the safety and efficacy of HSCT in SSc and also demonstrated improved long-term mortality in dsSSc patients randomized to HSCT compared with CYC.
18. Khanna D, Denton CP, Lin CJF, et al. Safety and efficacy of subcutaneous tocilizumab in systemic sclerosis: results from the open-label period of a phase II randomised controlled trial (faSScinate). *Ann Rheum Dis*. 2018 2;77(2):212–220. [PubMed: 29066464]
19. Khanna D, Spino C, Johnson S, et al. Abatacept in early diffuse cutaneous systemic sclerosis: results of a phase II investigator-initiated, multicenter, double-blind, randomized, placebo-controlled trial. *Arthritis Rheumatol*. 2020 1;72 (1):125–136. [PubMed: 31342624]
20. Gonzalez EG, Selvi E, Balistreri E, et al. Synthetic cannabinoid ajulemic acid exerts potent antifibrotic effects in experimental models of systemic sclerosis. *Ann Rheum Dis*. 2012 9;71 (9):1545–1551.. [PubMed: 22492781]
21. Spiera R, Hummers L, Chung L, et al. Safety and efficacy of lenabasum (JBT-101) in diffuse cutaneous systemic sclerosis subjects treated for one year in an open-label extension of trial JBT101-SSC-001. *EULAR Annual Meeting (Amsterdam, Netherlands)*; 2018.
22. Khanna D, Berrocal VJ, Giannini EH, et al. The American college of rheumatology provisional composite response index for clinical trials in early diffuse cutaneous systemic sclerosis. *Arthritis Rheumatol*. 2016 2;68(2):299–311. [PubMed: 26808827]
23. R Dn S, Bloom B, White B, et al. Baseline subject demographics and disease characteristics in a phase 3 study of safety and efficacy of lenabasum in diffuse cutaneous systemic sclerosis. *Atlanta GA: ACR Annual Meeting*; 2019.
24. Khanna D, Albera C, Fischer A, et al. An open-label, phase II study of the safety and tolerability of pirfenidone in patients with scleroderma-associated interstitial lung disease: the LOTUSS trial. *J Rheumatol*. 2016 9;43(9):1672–1679. [PubMed: 27370878]
25. Distler O, Highland KB, Gahlemann M, et al. Nintedanib for systemic sclerosis-associated interstitial lung disease. *N Engl J Med*. 2019 6 27;380(26):2518–2528. [PubMed: 31112379] • This was the largest RCT ever conducted in SSc and demonstrated that treatment with nintedanib slows the rate of decline of lung function in patients with SSc-ILD compared with placebo.
26. Sircar G, Goswami RP, Sircar D, et al. Intravenous cyclophosphamide vs rituximab for the treatment of early diffuse scleroderma lung disease: open label, randomized, controlled trial. *Rheumatology (Oxford)*. 2018 12;57(12):2106–2113. [PubMed: 30053212]
27. Daoussis D, Melissaropoulos K, Sakellaropoulos G, et al. A multicenter, open-label, comparative study of B-cell depletion therapy with Rituximab for systemic sclerosis-associated interstitial lung disease. *Semin Arthritis Rheum*. 2017 4;46(5):625–631.. [PubMed: 27839742]
28. Smith V, Van Praet JT, Vandooren B, et al. Rituximab in diffuse cutaneous systemic sclerosis: an open-label clinical and histopatho-logical study. *Ann Rheum Dis*. 2010 1;69(1):193–197. [PubMed: 19103636]

29. Jordan S, Distler JH, Maurer B, et al. Effects and safety of rituximab in systemic sclerosis: an analysis from the European Scleroderma Trial and Research (EUSTAR) group. *Ann Rheum Dis*. 2015 6;74 (6):1188–1194. [PubMed: 24442885]
30. Fraticelli P, Fischetti C, Salaffi F, et al. Combination therapy with rituximab and mycophenolate mofetil in systemic sclerosis. A single-centre case series study. *Clin Exp Rheumatol*. 2018 Jul-Aug;36(Suppl 113(4)):142–145. [PubMed: 30277864]
31. Steele R, Hudson M, Lo E, et al. Clinical decision rule to predict the presence of interstitial lung disease in systemic sclerosis. *Arthritis Care Res (Hoboken)*. 2012 4;64(4):519–524. [PubMed: 22213733]
32. Sanchez-Cano D, Ortego-Centeno N, Callejas JL, et al. Interstitial lung disease in systemic sclerosis: data from the spanish scleroderma study group. *Rheumatol Int*. 2018 3;38(3):363–374.
33. Steen VD, Conte C, Owens GR, et al. Severe restrictive lung disease in systemic sclerosis. *Arthritis Rheum*. 1994 9;37(9):1283–1289. [PubMed: 7945490]
34. Tyndall AJ, Bannert B, Vonk M, et al. Causes and risk factors for death in systemic sclerosis: a study from the EULAR Scleroderma Trials and Research (EUSTAR) database. *Ann Rheum Dis*. 2010 10;69(10):1809–1815.. [PubMed: 20551155]
35. Volkmann ER, Tashkin DP, Kuwana M, et al. Progression of Interstitial Lung Disease in Systemic Sclerosis: the Importance of Pneumoproteins Krebs von den Lungen 6 and CCL18. *Arthritis Rheumatol*. 2019 12;71(12):2059–2067.. [PubMed: 31233287]
36. Fischer A, Strek ME, Cottin V, et al. Proceedings of the American College of Rheumatology/ Association of Physicians of Great Britain and Ireland connective tissue disease-associated interstitial lung disease summit: a multidisciplinary approach to address challenges and opportunities. *Arthritis Rheumatol*. 2019 2;71(2):182–195.. [PubMed: 30604506]
37. Goldin JG, Kim GHJ, Tseng CH, et al. Longitudinal changes in quantitative interstitial lung disease on computed tomography after immunosuppression in the scleroderma lung study II. *Ann Am Thorac Soc*. 2018 11;15(11):1286–1295. [PubMed: 30265153]
38. Khanna D, Yan X, Tashkin DP, et al. Impact of oral cyclophosphamide on health-related quality of life in patients with active scleroderma lung disease: results from the scleroderma lung study. *Arthritis Rheum*. 2007 5;56(5):1676–1684. [PubMed: 17469162]
39. Tashkin DP, Elashoff R, Clements PJ, et al. Effects of 1-year treatment with cyclophosphamide on outcomes at 2 years in scleroderma lung disease. *Am J Respir Crit Care Med*. 2007 11 15;176 (10):1026–1034. [PubMed: 17717203]
40. Volkmann ER, Tashkin DP, Sim M, et al. Cyclophosphamide for systemic sclerosis-related interstitial lung disease: a comparison of scleroderma lung study I and II. *J Rheumatol*. 2019 10;46 (10):1316–1325. [PubMed: 30770517]
41. Volkmann ER, Tashkin D, Roth M, LeClair H, et al. Mycophenolate mofetil and cyclophosphamide improve health-related quality of life in patients with systemic sclerosis who participated in SLS II. *Arthritis Rheumatol*. 2019;71(suppl 10).
42. Khanna D, Denton CP, Jhreis A, et al. Safety and efficacy of subcutaneous tocilizumab in adults with systemic sclerosis (faSScinate): a phase 2, randomised, controlled trial. *Lancet*. 2016 6 25;387(10038):2630–2640. [PubMed: 27156934]
43. D Lc K, Kuwana M, Allamore Y, et al. Efficacy and safety of tocilizumab for the treatment of systemic sclerosis: results from a phase 3 randomized controlled trial. *Arthritis Rheumatol*. 2018;70(suppl 10).
44. Daoussis D, Lioussis SN, Tsamandas AC, et al. Experience with rituximab in scleroderma: results from a 1-year, proof-of-principle study. *Rheumatology (Oxford)*. 2010 2;49(2):271–280. [PubMed: 19447770]
45. Saunders P, Tsipouri V, Keir GJ, et al. Rituximab versus cyclophosphamide for the treatment of connective tissue disease-associated interstitial lung disease (RECITAL): study protocol for a randomised controlled trial. *Trials*. 2017 6 15;18(1):275. [PubMed: 28619061]
46. Wigley FM, Flavahan NA. Raynaud's Phenomenon. *N Engl J Med*. 2016 8 11;375(6):556–565. [PubMed: 27509103]



47. Nihtyanova SI, Brough GM, Black CM, et al. Clinical burden of digital vasculopathy in limited and diffuse cutaneous systemic sclerosis. *Ann Rheum Dis.* 2008 1;67(1):120–123. [PubMed: 17660220]
48. Roustit M, Giai J, Gaget O, et al. On-demand sildenafil as a treatment for Raynaud phenomenon: a series of n-of-1 trials. *Ann Intern Med.* 2018 11;169(10):694–703. [PubMed: 30383134]
49. Roustit M, Blaise S, Allanore Y, et al. Phosphodiesterase-5 inhibitors for the treatment of secondary Raynaud's phenomenon: systematic review and meta-analysis of randomised trials. *Ann Rheum Dis.* 2013 10;72(10):1696–1699. [PubMed: 23426043]
50. Andrigueti FV, Ebbing PCC, Arismendi MI, et al. Evaluation of the effect of sildenafil on the microvascular blood flow in patients with systemic sclerosis: a randomised, double-blind, placebo-controlled study. *Clin Exp Rheumatol.* 2017 Sep-Oct;35(Suppl 106(4)):151–158. [PubMed: 28281457]
51. Hachulla E, Hatron PY, Carpentier P, et al. Efficacy of sildenafil on ischaemic digital ulcer healing in systemic sclerosis: the placebo-controlled SEDUCE study. *Ann Rheum Dis.* 2016 6;75(6):1009–1015. [PubMed: 25995322]
52. Schioppa E, Hsu VM, Impens AJ, et al. Randomized placebo-controlled crossover trial of tadalafil in Raynaud's phenomenon secondary to systemic sclerosis. *J Rheumatol.* 2009 10;36(10):2264–2268. [PubMed: 19755613]
53. Shenoy PD, Kumar S, Jha LK, et al. Efficacy of tadalafil in secondary Raynaud's phenomenon resistant to vasodilator therapy: a double-blind randomized cross-over trial. *Rheumatology (Oxford).* 2010 12;49(12):2420–2428. [PubMed: 20837499]
54. Wigley FM, Seibold JR, Wise RA, et al. Intravenous iloprost treatment of Raynaud's phenomenon and ischemic ulcers secondary to systemic sclerosis. *J Rheumatol.* 1992 9;19(9):1407–1414. [PubMed: 1279170]
55. Wigley FM, Wise RA, Seibold JR, et al. Intravenous iloprost infusion in patients with Raynaud phenomenon secondary to systemic sclerosis. A multicenter, placebo-controlled, double-blind study. *Ann Intern Med.* 1994 2 1;120(3):199–206. [PubMed: 7506013]
56. McHugh NJ, Csuka M, Watson H, et al. Infusion of iloprost, a prostacyclin analogue, for treatment of Raynaud's phenomenon in systemic sclerosis. *Ann Rheum Dis.* 1988 1;47(1):43–47. [PubMed: 2449871]
57. Milio G, Corrado E, Genova C, et al. Iloprost treatment in patients with Raynaud's phenomenon secondary to systemic sclerosis and the quality of life: a new therapeutic protocol. *Rheumatology (Oxford).* 2006 8;45(8):999–1004. [PubMed: 16484290]
58. Shah AA, Schioppa E, Hummers LK, et al. Open label study of escalating doses of oral treprostinil diethanolamine in patients with systemic sclerosis and digital ischemia: pharmacokinetics and correlation with digital perfusion. *Arthritis Res Ther.* 2013 4;15(2):R54. [PubMed: 23597147]
59. Shah AA, Schioppa E, Chatterjee S, et al. The recurrence of digital ulcers in patients with systemic sclerosis after discontinuation of oral treprostinil. *J Rheumatol.* 2016 9;43(9):1665–1671. [PubMed: 27307535]
60. Mecoli CA, Perin J, Van Eyk JE, et al. Vascular biomarkers and digital ulcerations in systemic sclerosis: results from a randomized controlled trial of oral treprostinil (DISTOL-1). *Clin Rheumatol.* 2020 4;39(4):1199–1205. [PubMed: 31858338]
61. Gaillard-Bigot F, Roustit M, Blaise S, et al. Treprostinil iontophoresis improves digital blood flow during local cooling in systemic sclerosis. *Microcirculation.* 2016 4;23(3):266–270. [PubMed: 26833587]
62. Guigui A, Mazet R, Blaise S, et al. Treprostinil hydrogel iontophoresis in systemic sclerosis-related digital skin ulcers: a safety study. *J Clin Pharmacol.* 2020 2;60:758–767. [PubMed: 32080862]
63. Matucci-Cerinic M, Denton CP, Furst DE, et al. Bosentan treatment of digital ulcers related to systemic sclerosis: results from the RAPIDS-2 randomised, double-blind, placebo-controlled trial. *Ann Rheum Dis.* 2011 1;70(1):32–38. [PubMed: 20805294]
64. Khanna D, Denton CP, Merkel PA, et al. Effect of macitentan on the development of new ischemic digital ulcers in patients with systemic sclerosis: DUAL-1 and DUAL-2 randomized clinical trials. *JAMA.* 2016 5 10;315(18):1975–1988. [PubMed: 27163986]




65. Curtiss P, Schwager Z, Cobos G, et al. A systematic review and meta-analysis of the effects of topical nitrates in the treatment of primary and secondary Raynaud's phenomenon. *J Am Acad Dermatol*. 2018 6;78(6):1110–18 e3. [PubMed: 29408338]
66. Segreto F, Marangi GF, Cerbone V, et al. The role of botulinum toxin a in the treatment of Raynaud phenomenon. *Ann Plast Surg*. 2016 9;77(3):318–323. [PubMed: 26808752]
67. Bello RJ, Cooney CM, Melamed E, et al. The therapeutic efficacy of botulinum toxin in treating scleroderma-associated Raynaud's phenomenon: a randomized, double-blind, placebo-controlled clinical trial. *Arthritis Rheumatol*. 2017 8;69(8):1661–1669. [PubMed: 28426903]
68. Nagaraja V, Spino C, Bush E, et al. A multicenter randomized, double-blind, placebo-controlled pilot study to assess the efficacy and safety of riociguat in systemic sclerosis-associated digital ulcers. *Arthritis Res Ther*. 2019 9;21(1):202. [PubMed: 31481106]
69. R Lm D, Goundappa B, Medsger T, et al. The effect of atorvastatin on microvascular endothelial function and Raynaud phenomenon in early diffuse scleroderma: results of the “tamer” study. *Arthritis Rheumatol*. 2019;71(suppl 10).
70. Mukerjee D, St George D, Coleiro B, et al. Prevalence and outcome in systemic sclerosis associated pulmonary arterial hypertension: application of a registry approach. *Ann Rheum Dis*. 2003 11;62 (11):1088–1093. [PubMed: 14583573]
71. Steen VD, Medsger TA. Changes in causes of death in systemic sclerosis, 1972–2002. *Ann Rheum Dis*. 2007 7;66(7):940–944. [PubMed: 17329309]
72. Chung L, Farber HW, Benza R, et al. Unique predictors of mortality in patients with pulmonary arterial hypertension associated with systemic sclerosis in the REVEAL registry. *Chest*. 2014 12;146(6):1494–1504. [PubMed: 24992469]
73. Fisher MR, Mathai SC, Champion HC, et al. Clinical differences between idiopathic and scleroderma-related pulmonary hypertension. *Arthritis Rheum*. 2006 9;54(9):3043–3050. [PubMed: 16947776]
74. Mullin CJ, Mathai SC. New insights into the recognition, classification and management of systemic sclerosis-associated pulmonary hypertension. *Curr Opin Rheumatol*. 2017 11;29(6):561–567. [PubMed: 28817465]
75. Lammi MR, Mathai SC, Saketkoo LA, et al. Association between initial oral therapy and outcomes in systemic sclerosis-related pulmonary arterial hypertension. *Arthritis Rheumatol*. 2016 3;68(3):740–748. [PubMed: 26479414]
76. Hassoun PM, Zamanian RT, Damico R, et al. Ambrisentan and tadalafil up-front combination therapy in scleroderma-associated pulmonary arterial hypertension. *Am J Respir Crit Care Med*. 2015 11 1;192(9):1102–1110. [PubMed: 26360334] • This open-label trial was one of the first studies to demonstrate the hemodynamic benefits from upfront combination vasodilatory therapy in SSc-PAH.
77. Dwivedi PSS, Dhir V, Rohit MK, et al. A randomized double blind placebo controlled trial to compare the efficacy of initial combination therapy vs monotherapy for pulmonary arterial hypertension in systemic sclerosis. *Ann Rheum Dis*. 2018;77:1531.
78. Herve P, Lau EM, Sitbon O, et al. Criteria for diagnosis of exercise pulmonary hypertension. *Eur Respir J*. 2015 9;46(3):728–737. [PubMed: 26022955]
79. Stamm A, Saxer S, Lichtblau M, et al. Exercise pulmonary haemo-dynamics predict outcome in patients with systemic sclerosis. *Eur Respir J*. 2016 12;48(6):1658–1667. [PubMed: 27824602]
80. Yang S, Wu J, Lei S, et al. Abnormal pulmonary artery systolic pressure response after exercise in systemic sclerosis patients: A PRISMA-compliant meta-analysis. *Medicine (Baltimore)*. 2019 2;98(6):e14342. [PubMed: 30732161]
81. Saggarr R, Khanna D, Shapiro S, et al. Brief report: effect of ambrisentan treatment on exercise-induced pulmonary hypertension in systemic sclerosis: a prospective single-center, open-label pilot study. *Arthritis Rheum*. 2012 12;64(12):4072–4077. [PubMed: 22777623]
82. Humbert M, Coghlan JG, Ghofrani HA, et al. Riociguat for the treatment of pulmonary arterial hypertension associated with connective tissue disease: results from PATENT-1 and PATENT-2. *Ann Rheum Dis*. 2017 2;76(2):422–426. [PubMed: 27457511]
83. Gaine S, Chin K, Coghlan G, et al. Selexipag for the treatment of connective tissue disease-associated pulmonary arterial hypertension. *Eur Respir J*. 2017 8;50:2.

84. Catania MV, Sortino MA, Rampello L, et al. Adenosine deaminase increases release of excitatory amino acids through a mechanism independent of adenosine depletion. *Neuropharmacology*. 1991 2;30(2):153–159. [PubMed: 1709478]
85. Frech TM, Khanna D, Maranian P, et al. Probiotics for the treatment of systemic sclerosis-associated gastrointestinal bloating/distention. *Clin Exp Rheumatol*. 2011 Mar-Apr;29(2 Suppl 65):S22–5. [PubMed: 21586214]
86. Marighela TF, Arismendi MI, Marville V, et al. Effect of probiotics on gastrointestinal symptoms and immune parameters in systemic sclerosis: a randomized placebo-controlled trial. *Rheumatology (Oxford)*. 2019 11 1;58(11):1985–1990. [PubMed: 31056685]
87. Garcia-Collinot G, Madrigal-Santillan EO, Martinez-Bencomo MA, et al. Effectiveness of *saccharomyces boulardii* and metronidazole for small intestinal bacterial overgrowth in systemic sclerosis. *Dig Dis Sci*. 2020 4;65(4):1134–1143. [PubMed: 31549334]
88. Karamanolis GP, Panopoulos S, Denaxas K, et al. The 5-HT<sub>1A</sub> receptor agonist buspirone improves esophageal motor function and symptoms in systemic sclerosis: a 4-week, open-label trial. *Arthritis Res Ther*. 2016 9 1;18:195. [PubMed: 27586891]
89. Bharadwaj S, Tandon P, Gohel T, et al. Gastrointestinal manifestations, malnutrition, and role of enteral and parenteral nutrition in patients with scleroderma. *J Clin Gastroenterol*. 2015 8;49(7):559–564. [PubMed: 25992813]
90. Baron M, Bernier P, Côté LF, et al. Screening and therapy for malnutrition and related gastrointestinal disorders in systemic sclerosis: recommendations of a North American expert panel. *Clin Exp Rheumatol*. 2010 Mar-Apr;28(2 Suppl 58):S42–6.
91. Stratton RJ, Hackston A, Longmore D, et al. Malnutrition in hospital outpatients and inpatients: prevalence, concurrent validity and ease of use of the 'malnutrition universal screening tool' ('MUST') for adults. *Br J Nutr*. 2004 11;92(5):799–808. [PubMed: 15533269]

**Article highlights**

- A variety of novel therapeutic strategies for the management of SSc now exist and are grounded in solid scientific research.
- These therapeutic strategies primarily target individual organ manifestations of SSc.
- Immunosuppressants with potential disease-modifying effects continue to be important components of the therapeutic armamentarium for SSc.
- Patients with refractory disease may require combination therapy with anti-fibrotics and/or other agents.
- Defining mechanistically based SSc subgroups based on biological and clinical profiles will reduce heterogeneity in clinical trial cohorts and may enhance our ability to detect treatment effects.
- Preventing both the onset and progression of individual organ manifestation remains an important unmet clinical need in SSc.



**Figure 1.** Drugs targeting specific complications of systemic sclerosis in the skin, lungs, peripheral vasculature, pulmonary arteries, and gastrointestinal tract.

Author Manuscript

Author Manuscript

Author Manuscript

Author Manuscript

**Table 1.** Examples of some factors that often drive treatment decisions in SSc beyond the organ system involved.

Factor	Example
Sex	Male patients with SSc-ILD have an increased risk of progression of ILD and may warrant more aggressive upfront therapy for ILD.
Disease duration	Patients with SSc of <5 years disease duration may require a more aggressive approach for the management of ILD and/or cutaneous sclerosis since these dimensions of the disease are more likely to progress within this time frame.
Scl-70 antibody positivity	Patients with newly diagnosed SSc-ILD who have normal lung function may benefit from treatment for ILD because their ILD is more likely to progress rapidly.
Centromere antibody positivity	Patient with newly diagnosed SSc-ILD who have normal lung function may <i>not</i> benefit from treatment for ILD because their ILD is less likely to progress rapidly.
Overlap syndromes	Patients with overlap syndromes, such as rheumatoid arthritis, may benefit from therapy which targets both inflammatory arthritis and SSc
Age	Older patients with co-morbidities may have an increased risk of infectious complications when multiple immunosuppressant therapies are used in combination

**Table 2.**

Existing and phase 2, 3 investigational therapies for dcSSc.

Agent	Trial Phase (NCT)	Skin endpoint	Efficacy outcomes
Cyclophosphamide	3 (NCT00004563)	Secondary endpoint: mRSS	Significant improvement at 12 months compared with placebo arm
Mycophenolate mofetil (MMF)	2 (NCT00883129)	Secondary endpoint: mRSS	Clinically meaningful improvement at 24 months with no difference between cyclophosphamide arm
Abatacept	2 (NCT02161406)	Primary endpoint: mRSS	No significant skin treatment effect compared with placebo arm
Tocilizumab	3 (NCT02453256)	Primary endpoint: mRSS	No significant skin treatment effect compared with placebo arm
Belimumab + Rituximab + MMF	2 (NCT03844061)	Primary endpoint: CRIS	Trial ongoing
Brentuximab Vedotin	1/2 (NCT03222492)	Exploratory endpoints: mRSS, CRIS	Trial ongoing
Pirfenidone	2 (NCT03221257)	Secondary endpoint: mRSS	Trial ongoing
Nintedanib	3 (NCT02597933)	Secondary endpoint: mRSS	No significant skin treatment effect compared with placebo arm
Intravenous immunoglobulin (IVIg)	2 (NCT04138485)	Primary endpoint: CRIS; Secondary endpoint: mRSS	Trial ongoing

**Table 3.**

Existing and phase 2, 3 investigational therapies for SSC-ILD.

Agent	Trial Phase (NCT)	Skin endpoint	Efficacy outcomes
Cyclophosphamide (CYC)	3 (NCT00004563)	Primary endpoint: FVC%-predicted	Significant improvement at 12 months
Mycophenolate mofetil (MMF)	2 (NCT00883129)	Primary endpoint: FVC%-predicted	Significant improvement at 24 months
Tocilizumab	3 (NCT02453256)	Secondary endpoint: FVC%-predicted	Trial ongoing
Pirfenidone	2 (NCT03221257)	Primary endpoint: FVC%-predicted	Trial ongoing
Nintedanib	3 (NCT02597933)	Primary endpoint: Rate of decline of FVC	Significant treatment effect at 12 months
Rituximab + MMF	3 (NCT02990286)	Primary endpoint: FVC%-predicted	Trial ongoing
Rituximab vs. CYC	2/3 (NCT01862926)	Primary endpoint: Absolute change in FVC	Trial ongoing
Bortezomib + MMF	2 (NCT02370693)	Secondary endpoint: Change in FVC	Trial ongoing



**Table 4.**

Existing and phase 3 and 4 investigational therapies for SSC Raynaud phenomenon.

Agent	Raynaud's phenomenon			Efficacy outcomes
	Trial Phase (NCT)	Vascular endpoint		
Sildenafil vs. Placebo	3 (NCT01295736)	Primary endpoints: time to healing of ischemic digital ulcers		No significant differences in primary endpoint at 90 days
Tadalafil vs. Placebo	3 (reference 47)	Primary endpoints: Raynaud's Phenomenon (RP) condition score, frequency of RP episodes, duration of RP episodes		No significant differences in primary endpoint
Add-on tadalafil vs. add-on placebo	3 (NCT00626665)	Primary endpoints: frequency and duration of RP attacks, evolution of trophic digital lesions		Significant improvement relative to baseline and placebo
Bosentan vs. placebo	3 (NCT00077584)	Primary endpoints: time to complete healing of the cardinal ulcer and the total number of new digital ulcers		Significant improvement in the reduction of number of new ulcers compared to placebo
Macitentan vs. placebo	3 (NCT01474109)	Primary endpoints: Incidence rate of new digital ulcers		No significant differences in primary endpoint at 16 weeks
Macitentan vs. placebo	3 (NCT01474122)	Primary endpoints: Incidence Rate of New Digital Ulcers		No significant differences in primary endpoint primary endpoints at 16 weeks
Intravenous Iloprost vs. Placebo	3 (NCT04040322)	Primary endpoint: Frequency of symptomatic RP attacks		Trial ongoing
Botox-A vs. Placebo	3 (NCT03717961)	Primary endpoint: Absolute change from baseline in the number of RP attacks per week at 4 weeks		Trial ongoing
Aspirin vs. placebo	4 (NCT03558854)	Primary endpoints: Serum level of thromboxane A		Trial ongoing

**Table 5.**

Existing and phase 3 and 4 investigational therapies for SSC pulmonary hypertension.

Agent	Pulmonary hypertension			Efficacy outcomes
	Trial Phase (NCT)	Endpoint (s)		
Upfront tadalafil and ambrisentan	4 (NCT01042158)	Primary endpoint: Assessment of change in right ventricular mass via standard volumetric cine images of the right heart		Significant therapeutic effect at 36 weeks
Sildenafil vs. Sildenafil plus bosentan	4 (NCT03053739)	Primary endpoint: Change in Pulmonary artery pressures measured by echocardiography		No significant change in primary endpoints at 24 weeks
Ambrisentan (open label)	4 (NCT01051960)	Primary endpoint: Change in multipoint exercise total pulmonary resistance (TPR)/from baseline		Significant treatment effect at 24 weeks
Macitentan (open label)	3 (NCT03726398)	Primary endpoint: Change in exercise pulmonary vascular resistance (PVR)		Trial ongoing
Rosuvastatin vs. placebo	3 (NCT00984932)	Primary endpoints: exercise capacity measured by the SMWT, mPAP and WHO functional class change		Recruitment completed, Results pending