

Lawrence Berkeley National Laboratory

LBL Publications

Title

Pion Pair Production in Photon-Photon Collisions at Spear

Permalink

<https://escholarship.org/uc/item/5hj9s1wn>

Authors

Roussarie, A

Burke, D L

Abrams, G S

et al.

Publication Date

1981-05-01

Copyright Information

This work is made available under the terms of a Creative Commons Attribution License, available at <https://creativecommons.org/licenses/by/4.0/>



Lawrence Berkeley Laboratory

UNIVERSITY OF CALIFORNIA

Physics, Computer Science & Mathematics Division

OCT 9 1981

Submitted to Physics Letters B


PION PAIR PRODUCTION IN PHOTON-PHOTON COLLISIONS
AT SPEAR

Lawrence Berkeley Laboratory and Department of Physics,
University of California, Berkeley, CA, and Stanford
Linear Accelerator Center, Stanford University,
Stanford, CA

May 1981

TWO-WEEK LOAN COPY

*This is a Library Circulating Copy
which may be borrowed for two weeks.
For a personal retention copy, call
Tech. Info. Division, Ext. 6782.*



LBL-13196
-2

DISCLAIMER

This document was prepared as an account of work sponsored by the United States Government. While this document is believed to contain correct information, neither the United States Government nor any agency thereof, nor the Regents of the University of California, nor any of their employees, makes any warranty, express or implied, or assumes any legal responsibility for the accuracy, completeness, or usefulness of any information, apparatus, product, or process disclosed, or represents that its use would not infringe privately owned rights. Reference herein to any specific commercial product, process, or service by its trade name, trademark, manufacturer, or otherwise, does not necessarily constitute or imply its endorsement, recommendation, or favoring by the United States Government or any agency thereof, or the Regents of the University of California. The views and opinions of authors expressed herein do not necessarily state or reflect those of the United States Government or any agency thereof or the Regents of the University of California.

PION PAIR PRODUCTION IN PHOTON-PHOTON COLLISIONS AT SPEAR*

A. Roussarie,[¶] D.L. Burke, G.S. Abrams, M.S. Alam,[†] C.A. Blocker,
 A. Blondel, A.M. Boyarski, M. Breidenbach, W.C. Carithers, W. Chinowsky,
 M.W. Coles,^{††} S. Cooper,^{††} W.E. Dieterle, J.B. Dillon, J. Dorenbosch,[‡]
 J.M. Dorfan, M.W. Eaton, G.J. Feldman, M.E.B. Franklin, G. Gidal,
 G. Goldhaber, G. Hanson, K.G. Hayes,[‡] T. Himel,[‡] D.G. Hitlin,^{‡‡}
 R.J. Hollebeek, W.R. Innes, J.A. Jaros, P. Jenni,[‡] A.D. Johnson, J.A. Kadyk,
 A.J. Lankford, R.R. Larsen, M. Levi, V. Lüth, R.E. Millikan, M.E. Nelson,
 C.Y. Pang, J.F. Patrick, M.L. Perl, B. Richter, D.L. Scharre,
 R.H. Schindler,[‡] R.F. Schwitters,[‡] J.L. Siegrist, J. Strait, H. Taureg,[‡]
 M. Tonutti,^{||} G.H. Trilling, E.N. Vella, R.A. Vidal, I. Videau,[§] J.M. Weiss,
 and H. Zaccone.[¶]

Stanford Linear Accelerator Center
 Stanford University, Stanford, California 94305

and

Lawrence Berkeley Laboratory and Department of Physics
 University of California, Berkeley CA 94720

ABSTRACT

We report a measurement of the cross section for the process
 $\gamma\gamma \rightarrow \pi^+\pi^-$ at invariant masses $500 \text{ MeV}/c^2 < m_{\pi\pi} < 2000 \text{ MeV}/c^2$. A
 value for the radiative width of the $f(1270)$ tensor meson
 $\Gamma_{f \rightarrow \gamma\gamma} = 3.6 \pm 0.3 \pm 0.5 \text{ keV}$ (helicity $\lambda=2$) has been obtained
 from a fit to the observed $\pi\pi$ mass spectrum.

*Work supported in part by Department of Energy, contracts
 DE-AC03-76SF00515 and W-7405-ENG-48, by the Universität Bonn,
 Federal Republic of Germany, the Centre d'Etudes Nucléaires de
 Saclay, France and the Miller Institute for Basic Research in
 Science, Berkeley, California.

Present addresses:

[¶]Vanderbilt University, Nashville, TN 37235.

^{††}DESY, Hamburg. Fed. Rep. of Germany.

[‡]EP Division, CERN, Geneva, Switzerland.

^{‡‡}California Institute of Technology, Pasadena, CA 91125.

[‡]Harvard University, Cambridge, MA 02138.

^{||}Universität Bonn, Fed. Rep. of Germany

[¶]CEN-Saclay, France.

[§]LPNHE, École Polytechnique, Palaiseau, France.

The process $\gamma\gamma \rightarrow \pi\pi$ is an ideal place to study the properties
 of $J^C = \text{even}^+$ resonances¹ and, at high masses, it becomes a testing
 ground for QCD². In this Letter we present a study of the process

$$e^+e^- \rightarrow e^+e^-\pi^+\pi^- \quad (1)$$

at invariant masses $500 \text{ MeV}/c^2 \leq m_{\pi\pi} \leq 2000 \text{ MeV}/c^2$. A strong signal
 at the well-known $f(1270)$ resonance occurs midway in this region.
 Below the resonance the cross section for reaction (1) is found to
 be large ($\sim 100\text{nb}$) while at higher masses it falls rapidly.

The data presented here were taken with the SLAC-LBL Mark II
 detector³ at SPEAR with center-of-mass energies between $4.5 \text{ GeV}/c^2$
 and $6.5 \text{ GeV}/c^2$. The accumulated luminosity was $13,200 \text{ nb}^{-1}$. Events
 were selected that contained two detected tracks of opposite charge
 and no other detected charged particles. Separation of reaction (1)
 from the large background of two-photon initiated QED processes

$$e^+e^- \rightarrow e^+e^- + (e^+e^- \text{ or } \mu^+\mu^-) \quad (2)$$

has been done using the signals from the Mark II liquid argon system.
 To ensure that each track was within the fiducial acceptance of the
 liquid argon calorimeters; the laboratory polar angle of each track
 was required to satisfy $|\cos\theta_L| \leq 0.6$. The outgoing small angle
 beam electrons were not detected.

Backgrounds to process (1), other than the QED processes (2), include beam-gas events, higher multiplicity two-photon processes, and events from the e^+e^- annihilation channel. The beam-gas contamination was minimized by requiring the vertex of each event to be in the measured interaction region. Sideband subtractions were then made to correct the data for the remaining beam-gas background. Separation of the desired two-prong final state from beam-beam backgrounds was achieved by accepting only those events for which the net transverse momentum of the observed prongs, $|\Sigma \vec{p}_T|$, was less than 100 MeV/c. The two-photon production of the $\eta'(958)$ and of the four pion final state ($\rho^0\rho^0$) produce a cluster of events near the rho mass. Since these processes have previously been observed in the same detector^{3a,4}, a direct computation of the feed-down from these sources that survives the p_T cut has been made with a Monte Carlo. Remaining backgrounds were determined by fitting the observed $|\Sigma \vec{p}_T|$ distributions in various mass bins with a combination of the shape expected from the photon-photon processes (1) and (2) and the shape expected from events produced in the annihilation channel. The two distributions were determined by Monte Carlo simulations. The detected two-prong mass spectrum is shown in Fig. 1 along with the various contributions as they have been calculated or measured. The non-QED backgrounds are only a few percent of the total two-prong sample, but they become 5-30% in the raw $\pi\pi$ signal that we will present below.

The QED curve shown in Fig. 1 was computed by Monte Carlo⁵ and normalized by the measured luminosity of the experiment. Because this normalization is uncertain by $\pm 6\%$, the extraction of the $\pi\pi$ signal by a direct subtraction of the expected QED from the observed spectrum is impossible. To enhance the $\pi\pi$ signal relative to the QED background, we make use of the relatively large probability that a low momentum pion will interact in the magnet coil which is positioned between the central drift chamber and the liquid argon calorimeters. Unlike the electromagnetic interactions of electrons and muons, the hadronic nature of these pion interactions leads to a substantial probability that a pion will not deposit any energy in the liquid argon along the track trajectory. Accordingly, we identify a particle as a pion if it is assigned less energy by the liquid argon tracking programs than would be expected from a minimum ionizing particle. The event selection criteria include geometrical cuts to guarantee that all accepted events contain two-prongs within the fiducial acceptance of the calorimeters. Sources of pions, muons, and electrons that have been unambiguously identified without the use of the liquid argon signals have been used to calibrate the resulting pion identification probability and the electron and muon misidentification probabilities. The results are given in Fig. 2a. The pion efficiency for momenta below 400 MeV/c was checked using a large sample of two-prong beam-gas events. These events are relatively free of contamination by electrons and muons. Events containing protons, identified by time of flight, were removed. The

agreement of the pion efficiency obtained from these events with that in Fig. 2a was found to be better than 10%.

The pion identification criteria have been applied to each event in the two-prong sample (Fig. 1). The histogram in Fig. 2b shows the number of these events that contain at least one identified pion. The points with errors in this figure are the numbers of events that would have satisfied these criteria if the two-prong sample consisted of only electron and muon pairs (reactions 2 above). To compute these points the QED Monte Carlo⁵ was used to determine the expected ratio of detected electron pairs to detected muon pairs and then each event in the two-prong sample was weighted by the probability that it is a misidentified electron pair plus the corresponding probability that it is a misidentified muon pair. A definite excess of events at $m_{\pi\pi} \gtrsim 750 \text{ MeV}/c^2$ is seen from which we extract the $\pi^+\pi^-$ signal. This analysis has been carried out in a completely parallel manner with events that have been required to consist of two identified pions. This latter sample contains negligible QED background at $m_{\pi\pi} \gtrsim 700 \text{ MeV}/c^2$.

The two-photon cross section $\sigma_{\gamma\gamma \rightarrow \pi^+\pi^-}$ is related to the measured cross section $\sigma_{ee \rightarrow ee\pi\pi}$ by⁶

$$\frac{d\sigma_{ee \rightarrow ee\pi\pi}}{ds_{\pi\pi} d\Omega^*} = \left(\frac{\alpha}{\pi}\right)^2 \frac{f(E_b, s_{\pi\pi})}{s_{\pi\pi}} \frac{d\sigma_{\gamma\gamma \rightarrow \pi\pi}}{d\Omega^*}, \quad (3)$$

where $s_{\pi\pi}$ is the $\gamma\gamma$ invariant mass squared, E_b is the beam energy, and $d\Omega^* = d(\cos\theta^*) d\phi^*$ is the $\gamma\gamma$ c.m. solid angle element. With the restriction $|\cos\theta_L| \leq 0.6$ there is not much information in the c.m. angular distribution and, so, shown in Fig. 3 is the acceptance corrected cross section $\sigma_{\gamma\gamma \rightarrow \pi\pi}(m_{\pi\pi})$ integrated over the range $|\cos\theta^*| \leq 0.35$. The agreement between the results extracted with the two levels of pion identification indicates that the $\pi\pi$ final state has been accurately separated from the QED background. Also included in Fig. 3 is a data point at high mass that was measured using the Mark II muon and liquid argon systems to directly reject muon and electron pairs. This point has been corrected for all backgrounds with previously described techniques, and the pion efficiencies and electron and muon misidentification probabilities have been studied with the test samples previously described. The data tend to fall below the Born cross section⁷ at masses $\gtrsim 1.5 \text{ GeV}/c^2$. The prediction of a recent QCD calculation² also shown in the figure, indicates that data from higher beam energies will be needed in order to reach the interesting region at masses above the tail of the $f(1270)$.

The two-prong sample (Fig. 1) exhibits a large enhancement over the QED background in the mass region of the $f(1270)$ meson. We have previously reported⁸ that this structure is not consistent with the d-wave Breit-Wigner expected from parameters of the f determined in hadronic reactions⁹. Since we now have a more complete measurement of the $\pi\pi$ mass spectrum it is possible to investigate the hypothesis

that the Breit-Wigner for the resonant amplitude is modified by interference with the non-f background¹⁰. Although it contains a large QED component, the spectrum in Fig. 1 gives the best accuracy for this study. We use the detected spectrum of identified $\pi\pi$ events (with $|\cos\theta_L| < 0.6$) to normalize the QED curve in Fig. 1; this normalization is determined so that the excess of two-prong events over QED in the region $700 \text{ MeV}/c^2 \leq m_{\pi\pi} \leq 950 \text{ MeV}/c^2$ is consistent with the measured number of $\pi\pi$ events in the same mass region. Shown in Fig. 4 is the resulting non-QED spectrum in the vicinity of the f. Errors in the measured $\pi\pi$ yield result in a $\pm 4\%$ uncertainty in the QED normalization which is taken into account in the fitting procedure described below.

We fit the spectrum in Fig. 4 with the sum

$$\pi\pi(m) = f(m) + \alpha.C(m) + 2\beta.\cos\delta(m).\sqrt{\alpha.C(m).f(m)} + f'(m) \quad (4)$$

In this expression $f(m)$ is the contribution to the rate appropriate for a relativistic d-wave Breit-Wigner with resonance mass and width equal to $1271 \text{ MeV}/c^2$ and $180 \text{ MeV}/c^2$ respectively. All relevant photon flux factors have been included, and we assume that the f is only produced with helicity ± 2 . Two extreme cases for the shape of the non-f $\pi\pi$ production $C(m)$, were considered. The first was the simple Born term at all masses, and the second consisted of using the Born term at masses below $1 \text{ GeV}/c^2$, but replacing it with the QCD prediction at higher masses. (See Fig. 3.) The third term in Eq.(4) is an interference between the first two terms. We take the non-f amplitude to be real (like the Born term) so that the relative

phase of these two terms is equal to the resonant phase shift $\delta(m)$, modulo π . Since we are not able to do a partial wave analysis for the $\pi\pi$ spectrum, the parameter β has been included and we do fits for various values $-1 \leq \beta \leq 1$. The last term in Eq.(4) represents the contribution made by the process $\gamma\gamma \rightarrow f'(1515) \rightarrow K^+K^-$ where the kaons have been misidentified as pions. If we use the SU(3) expectation for the ratio $\Gamma_{f' \rightarrow \gamma\gamma} / \Gamma_{f \rightarrow \gamma\gamma}$ ($= 0.14$ assuming an m^3 phase space correction) and take the branching ratio $f' \rightarrow K^+K^-$ to be 50%, then a Monte Carlo calculation, which includes losses due to kaon decays, gives this term to be 3.8% of the f. The yield f' events is shown in Fig. 4 for the $\gamma\gamma$ partial width of the f obtained below.

The fitted parameters are the normalization of the non-f background (α) and the $\gamma\gamma$ partial width of the f(1270). Although α varies by a factor of ≈ 1.5 as the QED normalization is varied, the $\gamma\gamma$ partial width of the f depends very little on either the QED normalization, the value of β , or the choice for the continuum $C(m)$. The fits give (for helicity $\lambda=\pm 2$)

$$\Gamma_{f \rightarrow \gamma\gamma} = 3.6 \pm 0.3(\text{statistical}) \pm 0.5(\text{systematic}) \text{ keV}, \quad (5)$$

where the systematic error includes contributions from uncertainties in the luminosity measurement and acceptance calculation, as well as contributions from the fitting procedure. A value of the partial width $6.8 \pm 0.6 \pm 1.0 \text{ keV}$ is obtained if the f is assumed to be produced only with helicity $\lambda=0$. We display on Fig.4 the results of fits with $\beta=0, 0.5, \text{ and } 1.0$ and with $C(m)$ given by the Born term. If hypothesis (4), with $C(m)$ given by the Born term, is correct, then the data

require $\beta \geq 0.10$ (95% CL). If the background is parameterized by the QCD prediction above $m_{\pi\pi} = 1.0 \text{ GeV}/c^2$ then the fitted values of $\Gamma_{f \rightarrow \gamma\gamma}$ increase by $\approx 10\%$ (included in the systematic error quoted above), and the data require a somewhat larger value of β . The contribution of the f to the $\pi\pi$ spectrum (i.e., $\pi\pi(m) - \alpha.C(m) - f'(m)$ in eq. 4) as determined by this latter fit is given in Fig. 3 for $\beta=0.5$.

Our result (5) can be compared with previously reported values of $\Gamma_{f \rightarrow \gamma\gamma} = 2.3 \pm 0.5 \pm 0.35 \text{ keV}$ obtained at PETRA¹¹ and $\Gamma_{f \rightarrow \gamma\gamma} = 9.5 \pm 3.9 \pm 2.4 \text{ keV}$ from another SPEAR experiment¹². Theoretical predictions of the radiative width of the f are numerous¹ and vary from 1 keV to 20 keV. Our measurement clearly favors the lower end of this range.

We wish to acknowledge several enlightening discussions with Professors S. Brodsky and F. Gilman of the SLAC theory group.

REFERENCES

1. F. J. Gilman, SLAC-PUB-2461, 1980. Presented at the International Conference on Two-Photon Interactions, Lake Tahoe, Calif. 1979.
M. Greco, LNF-80/28 (P), 1980. Presented at the International Workshop on $\gamma\gamma$ Collisions, Amiens, France, 1980.
2. S. J. Brodsky and G. P. Lepage, presented at the 20th International Conference on High Energy Physics, Madison, Wisconsin, 1980; and SLAC-PUB-2587 (1980).
3. a) G. S. Abrams et al., Phys. Rev. Lett. 43 (1979) 477.
b) R. H. Schindler, SLAC-REPORT-219 (1979), Ph. D. Thesis unpublished.
4. D. L. Burke et al., SLAC-PUB-2693 (1981), to be published in Phys. Lett. B.
5. J. Smith, J.A.M. Vermaseren and G. Grammar, Phys. Rev. D15 (1971) 1532.
6. G. Bonneau, M. Gourdin, and F. Martin, Nucl. Phys. B54 (1972) 573.
J. H. Field, Nucl. Phys. B168 (1980) 477.
7. S. J. Brodsky, T. Kinoshita and H. Terazawa, Phys. Rev. D4 (1971) 1532.
8. A. Roussarie, presented at the 20th International Conference on High Energy Physics, Madison, Wisconsin, 1980; and SLAC-PUB-2599 (1980).
9. Particle Data Group, Rev. Mod. Phys. 52 (1980) 51.
10. D. Faiman, H. J. Lipkin, and H. R. Rubinstein, Phys. Lett. 59B (1975) 269.
11. PLUTO Collab., Ch. Berger et al., Phys. Lett. 94B (1980) 254.
12. C. J. Biddick et al., Phys. Lett. 97B (1980) 320.

Figure Captions

Fig. 1. The observed two-prong mass spectrum. The curves represent various contributions as measured and/or computed.

Fig. 2. (a) Pion identification efficiency and muon and electron misidentification probabilities. The muons are cosmic rays and the electrons come from photons that convert in the beam pipe.
 (b) Raw $\pi^+\pi^-$ signal from events with at least one identified pion. The points with errors give the yields that would be expected from QED in the absence of pion pair production. Also shown is the $\pi^+\pi^-$ identification probability.

Fig. 3. The cross section $\sigma_{\gamma\gamma \rightarrow \pi\pi}$ integrated over the range of c.m. angles $|\cos\theta^*| \leq 0.35$. The curves are: — Born, — QCD, and --- f(1270).

Fig. 4. The non-QED spectrum near the f(1270) mass. The number of misidentified $\gamma\gamma \rightarrow f'(1515) \rightarrow K^+K^-$ expected from SU(3) is shown for a branching ratio $BR(f' \rightarrow K^+K^-) = 50\%$. The remaining curves are described in the text.




Fig. 1




Fig. 2




Fig. 3




Fig. 4

This report was done with support from the Department of Energy. Any conclusions or opinions expressed in this report represent solely those of the author(s) and not necessarily those of The Regents of the University of California, the Lawrence Berkeley Laboratory or the Department of Energy.

Reference to a company or product name does not imply approval or recommendation of the product by the University of California or the U.S. Department of Energy to the exclusion of others that may be suitable.

TECHNICAL INFORMATION DEPARTMENT
LAWRENCE BERKELEY LABORATORY
UNIVERSITY OF CALIFORNIA
BERKELEY, CALIFORNIA 94720