UC Irvine

UC Irvine Previously Published Works

Title

Stratospheric ozone destruction by man-made chlorofluoromethanes.

Permalink

<https://escholarship.org/uc/item/5hk1w60k>

Journal

Science (New York, N.Y.), 185(4157)

ISSN

0036-8075

Authors

Cicerone, RJ
Stolarski, RS
Walters, S

Publication Date

1974-09-01

DOI

10.1126/science.185.4157.1165

Copyright Information

This work is made available under the terms of a Creative Commons Attribution License, available at <https://creativecommons.org/licenses/by/4.0/>

Peer reviewed

References and Notes

1. Key references include W. L. Bragg, R. W. James, C. H. Bosanquet, *Phil. Mag.* **44**, 433 (1922); R. J. Havighurst, *Phys. Rev.* **29**, 1 (1927); E. O. Wollan, *ibid.* **38**, 15 (1931). Corresponding studies by electron diffraction were first reported by L. S. Bartell and L. O. Brockway [*Phys. Rev.* **90**, 833 (1953)].
2. E. Schrödinger, *Ann. Phys. (Leipzig)* **79**, 361 (1926); D. R. Hartree, *Proc. Cambridge Phil. Soc.* **24**, 89 (1928); *ibid.*, p. 111; *ibid.*, p. 426.
3. E. W. Müller, *Z. Naturforsch. A* **5**, 473 (1950).
4. —, *J. Appl. Phys.* **27**, 474 (1956).
5. See, for example, A. V. Crews, J. Wall, J. Langmore, *Science* **168**, 1338 (1970); F. P. Ottensmeyer, E. E. Schmidt, A. J. Olbrecht, *ibid.* **179**, 175 (1973).
6. D. Gabor, *Nature (Lond.)* **161**, 777 (1948); *Proc. Roy. Soc. London Ser. A* **197**, 454 (1949).
7. L. S. Bartell, in *Techniques of Chemistry: Physical Methods in Chemistry*, A. Weissberger and B. W. Rossiter, Eds. (Interscience, New York, ed. 4, 1972), vol. 1, part 3D, pp. 125–158.
8. E. Clementi and D. L. Raimondi, *J. Chem. Phys.* **38**, 2686 (1963).
9. This work was supported by a grant from the National Science Foundation.

6 May 1974

Stratospheric Ozone Destruction by Man-Made

Chlorofluoromethanes

Abstract. Calculations indicate that chlorofluoromethanes produced by man can greatly affect the concentrations of stratospheric ozone in future decades. This effect follows the release of chlorine from these compounds in the stratosphere. Present usage levels of chlorofluoromethanes can lead to chlorine-catalyzed ozone destruction rates that will exceed natural sinks of ozone by 1985 or 1990.

Chlorofluoromethanes (CF_xCl_y), principally CF_2Cl_2 and CFCl_3 , are being produced as aerosol propellants and refrigerants in large and growing amounts, and their atmospheric concentrations are increasing (1–3). These compounds have been considered valuable as tracers of atmospheric motions because they are relatively inert chemically with atmospheric lifetimes exceeding 10 years (4). Unlike CCl_4 (1), they seem to have no natural sources or sinks in the troposphere; their lifetimes are controlled by diffusion into the stratosphere where they can be photodissociated by ultraviolet light (5). Molina and Rowland (5) have noted that this stratospheric sink for CF_xCl_y also represents a potential sink for stratospheric O_3 . This is so because the photodissociation of CF_xCl_y releases chlorine atoms which can catalytically destroy O_3 through reactions like those of the nitrogen oxides (NO_x) with O_3 (5–8). Because of the great importance of stratospheric O_3 (9), we have reexamined this potential effect and its likely time evolution. We find that current CF_xCl_y usage levels and trends can lead to chlorine-catalyzed O_3 destruction rates exceeding all natural sinks of stratospheric O_3 by the early 1980's. Stratospheric changes will continue long after ground-level emissions cease. For example, if emissions were curtailed now, the resultant O_3 destruction would maximize around 1990 and would remain significant for several decades. Our calculations also indicate that the CIX concentrations (the sum of the concentrations of Cl , ClO , and HCl) will increase significantly in the stratosphere but not in the troposphere.

Hence, critical monitoring of this problem will require measurements in the stratosphere; tropospheric observations alone will not suffice.

After release, CF_xCl_y molecules diffuse upward to be photolyzed by solar radiation (chiefly 175- to 220-nm ultraviolet wavelengths) in the stratosphere. On the basis of several atmospheric measurements (1–3) and known chemical properties of the chlorofluoromethanes, stratospheric photolysis ap-

Fig. 1. Globally integrated O_3 destruction rates for three model time histories of CF_xCl_y emission. Corresponding rates appear for the natural O_3 sinks due to pure oxygen (labeled "Chapman") reactions and NO_x reactions. The width of the NO_x band is due mostly to a lack of knowledge of the natural NO_x abundance and altitude distribution. All quantities were computed on the basis of present-day atmospheric background and have been integrated to 60-km altitude. The stratospheric O_3 production rate, integrated globally, is about $6 \times 10^{31} \text{ sec}^{-1}$. Above 50 km the hydrogen oxides can also destroy O_3 .

pears to be the major sink for CF_xCl_y (5). Several other possible sinks have been suggested but appear to be negligible (10). After release of the first chlorine atom by a solar photon, chemical reactions will probably remove the remaining chlorine and fluorine atoms from the $\text{CF}_x\text{Cl}_{y-1}$ radical, temporarily forming phosgene-type molecules (5). Once free in the stratosphere, chlorine atoms can catalyze the recombination of O_3 and atomic oxygen. The key reaction are (5–8)

Eventually, the chlorine released from CF_xCl_y should reach the ground through downward diffusion and tropospheric rainout (6–8).

We have quantitatively estimated globally averaged rates of O_3 destruction due to the chlorine atoms released in the initial photodissociation from man-made CF_xCl_y . Figure 1 displays the altitude integral of these rates for three presumed time histories of CF_2Cl_2 and CFCl_3 (11): model 1, exponentially increasing with a doubling time of 3.5 years, the current pattern; model 2, exponential increase from 1960 to 1975, then constant at the 1975 rate; and model 3, exponential increase from 1960 to 1975, then immediate cessation. Figure 1 also presents globally integrated O_3 destruction rates due to the chemical reactions of oxygen alone (labeled "Chapman") and for the NO_x catalytic cycles. These two O_3 sinks are currently believed to control stratospheric O_3 concentrations (8, 12). Models 1 and 2 yield rapidly increasing O_3 destruction rates that will equal the natural sinks by about 1982 and 1986, respectively. Model 3 shows that, if CF_xCl_y emissions were curtailed now, the ensuing O_3 destruction rates would maximize around 1990 at a rate comparable to major natural cycles and would persist for several decades. Even larger effects may be possible because in our calculations we consider only the first chlorine atom released from each CF_xCl_y molecule; in reality, all four halogen atoms may be freed (13).

Important parameters in calculations leading to Fig. 1 appear in Table 1. To account for vertical transport in our time-dependent, one-dimensional (altitude) model we adopted and smoothed the relatively high eddy diffusion coefficient (K) profile of Wofsy and McElroy (14). To compute photodissociation coefficients, J , for CF_2Cl_2 and CFCl_3 we used available laboratory

photoabsorption data (15). We evaluated the O_3 -destroying effect of the chlorine oxides (ClO_x) by using the photochemical reaction scheme of Stolarski and Cicerone (6) after adding the reactions $ClO + NO \rightarrow Cl + NO_2$ and $H_2 + Cl \rightarrow HCl + H$ (16). In the calculations we assumed an atmospheric background of O , O_3 , CH_4 , OH , NO , and H_2 that was constant in time. A constant O_3 profile representing present conditions is probably not consistent with the large O_3 -destroying rates of ClO_x of Fig. 1, but through its use we can compare the effect of the CF_xCl_y to the present natural O_3 sinks. Other procedures and input data are discussed in (17).

The accumulation (mixing ratio) of CF_xCl_y molecules is shown as a function of altitude and time in Fig. 2, predicted from emission history model 2. Corresponding profiles are also shown for CIX. Because the relevant photochemical processes are fast relative to the transport processes, one may calculate the amounts of Cl , ClO , and HCl present at each altitude from the CIX concentration and the equations of photochemical equilibrium (6–8). The time evolution of the CIX profiles show large increases above 20 km and virtually no change below 15 km. The precise CIX concentration predicted for a given altitude and time depends on several computational parameters, especially K , but the profile shapes clearly indicate the effects of a mid-stratospheric source (see the Q_{Cl} values in Table 1), slow transport below 25 km, and eventual loss to the ground.

The 1985 chlorine production rate, Q_{Cl} for 1985 in Table 1, is 5×10^6 $cm^{-2} sec^{-1}$, integrated over altitude. Under steady-state conditions this figure will grow to 2×10^7 $cm^{-2} sec^{-1}$ if only one chlorine atom is taken from

Fig. 2. Calculated fractional concentrations [mixing ratios (volume-volume)] of atmospheric CF_xCl_y (dashed curves) and those for CIX (solid curves). In the calculations we employed emission history model 2 with a rapid temporal accumulation of CF_xCl_y , resulting in a similar increase of stratospheric CIX. Ground-level CF_xCl_y concentrations computed for the present are in good agreement with measured values (1–3). Also, the computed initial CIX profile is not inconsistent with available measurements (17).

each CF_xCl_y and model 2 (constant) emissions are assumed. This man-made CIX source is comparable to the projected NO_x input due to large supersonic transport (SST) fleets, and it appears that ClO_x destroys O_3 more efficiently than NO_x (6). McElroy *et al.* (12) find that O_3 depletion from SST NO_x is sensitive to the injection altitude. The O_3 effect due to added CIX will be difficult to evaluate, but current models (6–8) indicate that CIX added above 25 km is much more important than that added below.

We have not attempted to predict O_3 concentrations or their changes due to increasing amounts of CIX from CF_xCl_y usage. Predictions of O_3 concentrations and trends due to perturbations are still the subject of considerable debate (18). However, Fig. 1 indicates that, regardless of the precise magnitude of the effect of increasing destruction rates on O_3 content, chlorine atom produc-

tion from present or exponentially increasing CF_xCl_y concentrations will eventually convert the O_3 layer from NO_x to ClO_x control. Uncertainties in the initial CIX profile, conversion rates to chlorine, or catalytic efficiency do not alter the basic conclusion, only the time of passage to ClO_x control. Studies should be pursued with more complex interactive photochemical models to assess the impact of these large predicted increases in O_3 destruction on the O_3 content. In addition to continued careful monitoring of O_3 , other key stratospheric trace constituents must be measured. For example, the stratospheric concentrations of OH radicals and atomic oxygen, although central to questions of stratospheric chemistry, have not, to our knowledge, been measured. The present stratospheric abundance of gaseous chlorine, natural or otherwise, must also be determined.

RALPH J. CICERONE

RICHARD S. STOLARSKI

STACY WALTERS

Space Physics Research Laboratory,
University of Michigan,
Ann Arbor 48105

References and Notes

1. J. E. Lovelock, *Nature (Lond.)* **230**, 379 (1971); C. W. Su and E. D. Goldberg, *ibid.* **245**, 27 (1973).
2. J. E. Lovelock, R. J. Maggs, R. J. Wade, *ibid.* **241**, 194 (1973).
3. P. E. Wilkiss, R. A. Lamontagne, R. E. Larson, J. W. Swinnerton, C. R. Dickson, T. Thompson, *ibid.* **245**, 45 (1973). More recent measurements by J. E. Lovelock [*ibid.* **249**, 434 (1974)] and P. E. Wilkiss (personal communication) show further increases in atmospheric concentrations.
4. L. Machta, paper presented as part of the Proceedings of the 2nd International Union of Theoretical and Applied Mechanics—International Union of Geodesy and Geophysics Symposium on Turbulent Diffusion in Environmental Pollution, Charlottesville, Virginia, 1973; see also (1, 2, 5).
5. M. J. Molina and F. S. Rowland, *Nature (Lond.)* **249**, 810 (1974).
6. R. S. Stolarski and R. J. Cicerone, *Can. J. Chem.* **52**, 1610 (1974).
7. S. C. Wofsy and M. B. McElroy, *ibid.*, p. 1582.
8. P. Crutzen, *ibid.*, p. 1569; see also (5).
9. P. Cutchis, *Science* **184**, 13 (1974).
10. Experiments by O. C. Taylor, N. E. Hester, and E. A. Cardiff (personal communication) show that soils and plants do not act as sinks for CF_2Cl_2 and $CFCl_3$. Other studies (N. E. Hester, personal communication) show that these compounds are remarkably stable in simulated photochemical smog. Also, possible reaction with atmospheric $O(^1D)$ atoms, even at a fast (3×10^{-10} $cm^3 sec^{-1}$) rate, would not be a significant sink because of small $O(^1D)$ concentrations. The search for natural sinks should continue.
11. Worldwide rates of production of CF_2Cl_2 and $CFCl_3$ are currently about 0.45×10^6 and 0.27×10^6 kg, respectively (R. L. McCarthy, personal communication). Between 1960 and the present, this production grew exponentially with a doubling time of 3.5 years; see also (2, 4).
12. M. B. McElroy, S. C. Wofsy, J. E. Penner, J. C. McConnell, *J. Atmos. Sci.* **31**, 287 (1974).
13. F. S. Rowland (personal communication) tells us that all the chlorine atoms are likely to be rapidly removed from the CF_xCl_{y-1} radical

Table 1. Input data at several altitudes: K , the eddy diffusion coefficient; the NO concentration; J , the photodissociation coefficients for $CFCl_3$ and CF_2Cl_2 , respectively; Q_{Cl} , the predicted release rate of chlorine atoms assuming a model 2 emission time history. The J values were computed for a solar zenith angle of 60° and were divided by 2 to represent a diurnal average. Note: 1.5(4) means 1.5×10^4 .

Altitude (km)	K ($cm^2 sec^{-1}$)	NO concentration (cm^{-3})	J_{CFCl_3} (sec^{-1})	$J_{CF_2Cl_2}$ (sec^{-1})	Q_{Cl} ($cm^{-2} sec^{-1}$)	
					1980	1985
15	1.5(4)	1.9(8)	0	0	0	0
20	2.5(3)	5.0(8)	3.0(–10)	2.7(–11)	0.14	0.24
25	7.6(3)	7.2(8)	2.1(–8)	2.2(–9)	1.2	2.06
30	3.5(4)	8.0(8)	3.6(–7)	4.0(–8)	2.4	4.14
35	6.5(4)	5.8(8)	2.0(–6)	2.3(–7)	1.5	2.68
40	1.0(5)	5.4(8)	4.6(–6)	5.0(–7)	0.4	0.82
45	1.6(5)	4.7(8)	6.4(–6)	7.5(–7)	0.12	0.23
50	2.5(5)	3.3(8)	7.6(–6)	9.5(–7)	0.03	0.06

- by reactions in the atmosphere. Although the fluorine atoms released by similar reactions can also destroy O_3 catalytically, the reaction $CH_4 + F \rightarrow CH_3 + HF$ will terminate the fluorine chains more quickly than $CH_4 + Cl \rightarrow CH_3 + HCl$ does for chlorine. Further, reinitiation of fluorine chains will not proceed via $HF + OH \rightarrow F + H_2O$, whereas $HCl + OH \rightarrow Cl + H_2O$ does reinitiate the chlorine chains.
14. S. C. Wofsy and M. B. McElroy, *J. Geophys. Res.* **78**, 2619 (1973).
 15. J. Doucet, P. Sauvageau, C. Sandorfy, *J. Chem. Phys.* **58**, 3708 (1973); M. J. Molina and F. S. Rowland, unpublished data between 186 and 226 nm. Solar intensities were taken from R. T. Brinkmann, A. Green, C. Barth, *Calif. Inst. Technol. Jet Propulsion Lab. Tech. Rep.* 32-951 (1966). Absorption by O_2 was computed from the band absorption data of R. D. Hudson and S. Mahle [*J. Geophys. Res.* **77**, 2902 (1972)].
 16. Both D. D. Davis and D. D. Garvin (personal communication) have suggested that $ClO + NO \rightarrow Cl + NO_2$, which couples ClO_x and NO_x cycles, is important for stratospheric chlorine chemistry. This reaction and $H_2 + Cl + HCl + H$ were subsequently included in model studies by Wofsy and McElroy (7).
 17. We computed photolysis rates for $CFCl_3$ and CF_2Cl_2 versus altitude and time through an implicit numerical solution of the diffusion equation with 10-day time steps and altitude steps ΔZ of 1 km. These loss rates for CF_2Cl_2 then served as chlorine production rates, Q , in the photochemical, vertical transport model for CIX. Table 1 shows an example of Q along with selected input data. The OH profile was taken from (12); the reaction rate for $HCl + OH \rightarrow Cl + H_2O$ was $2 \times 10^{-12} \exp(-335/T)$ (where T is the absolute temperature) (J. G. Anderson, personal communication). The remainder of the photochemical scheme and input data were adopted from (6). For the CIX calculations the boundary conditions were: zero flux at the upper boundary (80 km), ground upward flux of $2 \times 10^{10} \text{ cm}^{-2} \text{ sec}^{-1}$ [see (7)], and diffusion to the ground at a velocity of 0.4 cm sec^{-1} . These conditions yield ground-level concentrations of 1 part per billion (ppb), consistent with data from C. Junge [*Tellus* **9**, 528 (1957)], an initial stratospheric concentration of 0.55 ppb, below the 0.8 ppb estimated for HCl at 16 km by C. B. Farmer (paper presented at the 3rd Conference on the Climatic Impact Assessment Program, Cambridge, Massachusetts, 27 February 1974). A 10-day rainout period (0 to 5 km), tapering to 100 days at 10 km, was assumed for HCl. The initial CIX profile, though not critical for the present problem, implies a column-integrated O_3 destruction rate of $2 \times 10^{12} \text{ cm}^{-2} \text{ sec}^{-1}$.
 18. H. Foley and M. Ruderman, *J. Geophys. Res.* **78**, 4441 (1973); H. Johnston, G. Whitten, J. Birks, *ibid.*, p. 6107; E. Hestvedt, *Can. J. Chem.* **52**, 1592 (1974); M. A. Ruderman, *Science* **184**, 1079 (1974); see also (12).
 19. We thank M. J. Molina and F. S. Rowland for several helpful discussions and for making available to us their not yet published laboratory data; we also thank D. D. Davis for ideas and criticism. Support provided by NASA through grant NGR 23-005-616 and by the Climatic Impact Assessment Program, Department of Transportation, through NSF grant GA-43326.
 - 21 June 1974

Methane Production in the Interstitial Waters of Sulfate-Depleted Marine Sediments

Abstract. *Methane in the interstitial waters of anoxic Long Island Sound sediments does not reach appreciable concentrations until about 90 percent of seawater sulfate is removed by sulfate-reducing bacteria. This is in agreement with laboratory studies of anoxic marine sediments sealed in jars, which indicate that methane production does not occur until dissolved sulfate is totally exhausted. Upward diffusion of methane or its production in sulfate-free microenvironments, or both, can explain the observed coexistence of measurable concentrations of methane and sulfate in the upper portions of anoxic sediments.*

Amounts of methane in the marine environment in excess of that resulting from equilibration with the atmosphere most frequently occur in anoxic waters such as those found in fjords (1, 2) and in the interstitial water of anoxic sediments (3-6). In the absence of sources associated with shipping and industrial activity or natural seeps from oil and gas reservoirs (7), most of this methane results from anaerobic bacterial decomposition of organic matter. Methane bacteria are readily found in anoxic environments, where they are terminal organisms in the microbial food chain (8); moreover, there is some evidence that methane is not produced until dissolved sulfate has been previously removed by sulfate-reducing bacteria (5, 6).

The interstitial waters of Recent organic-rich marine sediments are ideal for studying the relation between methane and dissolved sulfate distributions

because of the large concentration changes over short depth intervals that result from high bacterial activity. In this report we present the results of a study of methane and dissolved sulfate in the interstitial waters of Long Island Sound sediments which suggest that significant production of methane does not begin until dissolved sulfate concentrations approach zero. Results of laboratory studies of time-dependent changes in the chemistry of anoxic marine sediments indicate that sulfate reduction and methane production are mutually exclusive metabolic processes.

Gravity cores were collected at three stations in Long Island Sound. Station TH is located approximately 2 km south of the coastal town of Guilford, Connecticut, at a water depth of approximately 7 m. Stations BS and SC are located in two shallow harbors near Guilford. The water depths at these two stations ranged from 1.5 to 4 m be-

cause of tidal fluctuations. Interstitial waters were sampled without coming into contact with air by transferring sediments from sealed core liners to a filter-press type squeezer through an interlock flushed with CO_2 or He (9). Dissolved methane was measured by liquid stripping techniques developed for measuring dissolved gases in seawater by Swinnerton *et al.* (10) and applied to interstitial water measurements by Reeburgh (11). Dissolved sulfate was measured by gravimetric analysis as $BaSO_4$. Blank corrections for precipitation of nonsulfate material led to uncertainties of $\pm 0.5 \text{ mmole liter}^{-1}$ in dissolved sulfate concentrations.

Concentrations of methane and dissolved sulfate in the interstitial waters are plotted as a function of depth in Fig. 1. Differences in the depth of complete sulfate reduction at the three stations are probably a result of variations in sedimentation rates and the content of organic matter in the sediments. The leveling off of methane concentrations at station SC at about $1.0 \text{ mmole liter}^{-1}$ is consistent with reaching saturation with respect to methane. The solubility of methane calculated from the Setchenow relation by using solubility data from Winkler (12) and Atkinson and Richards (2) ranges from 1.1 to $2.3 \text{ mmole liter}^{-1}$ (25 to 51 ml liter^{-1}) for the temperature range (4° to $28^\circ C$) and salinity range (27 to 31 per mil) encountered in these sediments.

Reeburgh (4) has presented evidence of methane saturation in Chesapeake Bay sediments, which results in the formation of trapped methane bubbles that strip other dissolved gases such as N_2 and Ar from interstitial waters in the sediments. We infer from the low concentrations of dissolved N_2 and Ar at station SC that this process may be taking place there (13).

The data in Fig. 1 show that in the interstitial waters of Long Island Sound sediments high methane concentrations do not occur unless sulfate concentrations have been appreciably lowered. Only where dissolved sulfate concentrations approach zero do concentrations of methane attain saturation.

Four alternative hypotheses can be used to explain these results. The first is that methane is produced at roughly the same rate throughout the sediment column by methane bacteria but is consumed by sulfate-reducing bacteria through the reaction