

UCLA

UCLA Previously Published Works

Title

Sex hormones in the modulation of irritable bowel syndrome

Permalink

<https://escholarship.org/uc/item/5j29q385>

Journal

World Journal of Gastroenterology, 20(10)

ISSN

1007-9327

Authors

Mulak, Agata
Taché, Yvette
Larauche, Muriel


Publication Date

2014

DOI

10.3748/wjg.v20.i10.2433

Peer reviewed



WJG 20th Anniversary Special Issues (4): Irritable bowel syndrome

Sex hormones in the modulation of irritable bowel syndrome

Agata Mulak, Yvette Taché, Muriel Larauche

Agata Mulak, Department of Gastroenterology and Hepatology, Wrocław Medical University, 50-556 Wrocław, Poland
Yvette Taché, Muriel Larauche, CURE: Digestive Diseases Research Center and Oppenheimer Family Center for Neurobiology of Stress, Digestive Diseases Division, Department of Medicine, University of California Los Angeles, VA Greater Los Angeles Healthcare System, Los Angeles, CA 90073, United States
Author contributions: Mulak A designed and wrote the paper; Taché Y designed, wrote and reviewed the paper; Larauche M wrote and reviewed the paper.

Supported by The Veterans Administration Research Career Scientist Award (to Taché Y); National Institute of Health grants No. P50 DK-64539 (to Taché Y); and No. K01-DK088937 (to Larauche M)

Correspondence to: Agata Mulak, MD, PhD, Department of Gastroenterology and Hepatology, Wrocław Medical University, Borowska 213, 50-556 Wrocław, Poland. agata.mulak@wp.pl
Telephone: +48-71-7332120 Fax: +48-71-7332129

Received: September 25, 2013 Revised: December 10, 2013

Accepted: January 14, 2014

Published online: March 14, 2014

Abstract

Compelling evidence indicates sex and gender differences in epidemiology, symptomatology, pathophysiology, and treatment outcome in irritable bowel syndrome (IBS). Based on the female predominance as well as the correlation between IBS symptoms and hormonal status, several models have been proposed to examine the role of sex hormones in gastrointestinal (GI) function including differences in GI symptoms expression in distinct phases of the menstrual cycle, in pre- and post-menopausal women, during pregnancy, hormonal treatment or after oophorectomy. Sex hormones may influence peripheral and central regulatory mechanisms of the brain-gut axis involved in the pathophysiology of IBS contributing to the alterations in visceral sensitivity, motility, intestinal barrier function, and immune activation of intestinal mucosa. Sex differences in stress response of the hypothalamic-pituitary-adrenal axis and autonomic nervous system, neuroimmune interac-

tions triggered by stress, as well as estrogen interactions with serotonin and corticotropin-releasing factor signaling systems are being increasingly recognized. A concept of "microgenderome" related to the potential role of sex hormone modulation of the gut microbiota is also emerging. Significant differences between IBS female and male patients regarding symptomatology and comorbidity with other chronic pain syndromes and psychiatric disorders, together with differences in efficacy of serotonergic medications in IBS patients confirm the necessity for more sex-tailored therapeutic approach in this disorder.

© 2014 Baishideng Publishing Group Co., Limited. All rights reserved.

Key words: Brain-gut axis; Irritable bowel syndrome; Microbiota; Pain modulation; Sex hormones

Core tip: Recent clinical and experimental findings support the modulatory actions of sex hormones exerted at different levels of the brain-gut-microbiota axis in irritable bowel syndrome (IBS). Sex hormones may influence peripheral and central regulatory mechanisms contributing to the alterations in visceral sensitivity, motility, permeability, and immune activation of intestinal mucosa. A new concept of "microgenderome" is emerging based on the observations that the gender bias present in numerous diseases may be reinforced by the commensal microbiota of the host. Significant sex differences in epidemiology, symptomatology, and treatment outcome in IBS indicate the necessity for sex-tailored therapeutic approach in this disorder.

Mulak A, Taché Y, Larauche M. Sex hormones in the modulation of irritable bowel syndrome. *World J Gastroenterol* 2014; 20(10): 2433-2448 Available from: URL: <http://www.wjgnet.com/1007-9327/full/v20/i10/2433.htm> DOI: <http://dx.doi.org/10.3748/wjg.v20.i10.2433>

INTRODUCTION

Sex hormones, in particular estrogens, play a significant role in the physiology and pathology of the gastrointestinal (GI) tract including regulation of motor and sensory function^[1,2]. Irritable bowel syndrome (IBS) is a GI sensory-motor disorder characterized by abdominal pain or discomfort associated with a change in bowel habits^[3]. The role of gonadal hormones in symptomatology and pathophysiology of IBS is being increasingly recognized based on the female predominance as well as the correlation between IBS symptoms and hormonal status during menstrual cycle phases, pregnancy or menopause^[4]. Sex differences in stress and pain response are considered as crucial factors in the pathogenesis of functional GI disorders^[4,5]. Sex hormones influence peripheral and central regulatory mechanisms of the brain-gut axis involved in the pathophysiology of IBS contributing to alterations in visceral sensitivity, motility, permeability, and immune activation of intestinal mucosa^[4,6,7]. Among numerous interactions of sex hormones with other neurotransmitters, estrogen interactions with serotonin and corticotropin-releasing factor (CRF) signaling systems play a pivotal role^[8,9]. Estrogens can also modulate neuroimmune interactions triggered by stress *via* the brain-gut axis^[10]. Recently, the gut microbiota has been also recognized as an important element in the bi-directional communication along the brain-gut axis through neural, immune, and endocrine pathways^[11,12]. In the present article we will review recent clinical and experimental findings supporting the modulatory effect of sex hormones, in particular estrogens, on different levels of the brain-gut-microbiota axis in IBS and their clinical implications regarding the symptomatology, pathophysiology and treatment of IBS.

SEX AND GENDER DIFFERENCES IN IBS PREVALENCE AND SYMPTOMATOLOGY

In Western countries the female-to-male ratio among non-patient population of IBS sufferers is 2:1^[13]. Within the patient population in primary or tertiary care settings females outnumber male patients by 3:1 to 5:1, respectively^[13-15]. However, in many Eastern countries such as India, China and South Korea, the female predominance among IBS patients is not observed^[16]. Likewise, results of recent meta-analysis studies in South Asia, South America, or Africa confirmed that IBS prevalence was not significantly higher in women, compared to men^[17]. Therefore gender-related and socio-cultural differences in health care-seeking behavior are suggested to also account in IBS symptoms reporting^[18]. Further epidemiologic studies from different world regions are needed to elucidate the complex interactions between genetic, environmental, psychological and/or cultural factors that may contribute to sex differences in IBS symptoms^[19,20].

Additionally, as the prevalence of IBS subtypes varied according to gender^[17], the dominating subtype of IBS in different countries may also affect the male-

to-female ratio. Women with IBS, compared to male patients, are more likely to report constipation, bloating, severe abdominal pain, and feeling of incomplete evacuation, while men with IBS more frequently complain of diarrhea-associated symptoms^[17,19,21]. In fact, earlier studies indicate that women have slower colonic transit in comparison with men^[22,23]. The results of the recent study by Tang *et al.*^[24] conducted in Chinese population confirmed significant differences between female and male IBS patients in their rating of abdominal pain/discomfort with regard to severity and duration, but not frequency of pain attacks. Interestingly, sex differences in dietary coping with GI symptoms have also been reported^[25]. Female IBS patients seem to be more willing to implement nutritional behavior changes alleviating the GI problems than men, although both men and women could benefit similarly from these changes^[25].

Cain *et al.*^[26] reported higher GI symptoms (pain, distension, bloating, intestinal gas) in postmenopausal women than in men, but the greatest differences in the overall symptom reporting between men and women were associated with somatic symptoms such as joint and muscle pain. This gender-related difference was most prominent when postmenopausal women were compared to men. Gender differences were much weaker for psychological and emotional symptoms except for fatigue, sleep disturbances and stress^[24,26]. Noteworthy, there is a wide spectrum of chronic pain disorders frequently overlapping with IBS namely fibromyalgia, migraine headache, chronic pelvic pain, interstitial cystitis, and chronic fatigue syndrome. These diseases are also characterized by female predominance with a correlation between their symptoms and hormonal status^[27-29]. In addition, women with IBS, more frequently show comorbidity with affective or mood symptoms including anxiety and depression as compared to women without IBS^[30]. There are also reports indicating that women with IBS exhibit more anxiety and depressive symptoms compared to men with IBS^[24,26]. Sex differences in the prevalence of concomitant somatic symptoms, as well as anxiety and depression may significantly contribute to the greater impairment of quality of life in female patients and affect treatment results^[24,31].

Sex-related difference in IBS prevalence emerges around the time of puberty and increases during the early adult years. Women are suffering from IBS most commonly in the late teenage to mid forties, additionally suggesting the role of reproductive hormones in the pathophysiology of the disorder. The incidence of IBS in women steadily declines with age and approaches the rate observed among men by the 7th decade of life^[32]. By contrast, the prevalence of IBS among males is fairly constant within the age range of 20-70 years^[32].

CORRELATION BETWEEN IBS SYMPTOMS AND HORMONAL STATUS

Menstrual cycle

The menstrual cycle in women is divided into three

Table 1 Correlation between hormonal status and irritable bowel syndrome symptoms expression^[3,6]

Status	Hormone levels	IBS and pain related symptoms expression	Ref.
Late luteal phase (premenstrual)	Rapid decline in estrogen and progesterone levels	Exacerbation of bowel symptoms	[33,34]
Menstruation (menses)	Lowest levels of estrogen and progesterone	Increased bloating Exacerbation of bowel symptoms Increased abdominal pain/discomfort Lower rectal sensitivity threshold	[34,35,37,38,40]
Dysmenorrhea	Disturbances in hormonal interactions at different regulatory levels (lower progesterone level)	Exacerbation of bowel symptoms	[41]
Pregnancy	Physiological hyperestrogenemia and hyperprogesteronemia	Reduced pain sensitivity and alleviation of many chronic pain syndromes Exacerbation of constipation (prolonged gastrointestinal transit)	[27,51,73]
Menopause	Decline in ovarian hormones	Decrease in IBS incidence High prevalence of constipation and somatic discomfort syndromes	[26,53,54]
Oral contraceptives	Estrogen and progestin administration	Reduced abdominal symptoms at menses	[55]
Hormone replacement therapy	Estrogen (and progesterone) supplementation	Increased prevalence of IBS in postmenopausal women during HRT Prolongation of IBS symptoms to a later age	[58]
Oophorectomy	Ovarian hormone deficiency	Exacerbation or occurrence of gastrointestinal symptoms after gynecological surgery	[60]
Men with IBS	Lower level of luteinizing hormone in middle-aged men Elevated level of sex hormone-binding globulin in young men	Generally more prevalent diarrhea (compared to women with IBS)	[66,70]
Transsexual women (male-to-female subjects)	Estrogen/anti-androgen treatment	Development of chronic pain including visceral pain	[72]

IBS: Irritable bowel syndrome; HRT: Hormone replacement therapy.

phases: the follicular (proliferative) phase, ovulation, and the luteal (secretory) phase. Estrogen levels are increasing during the midfollicular phase and then drop precipitously after ovulation. This is followed by a secondary rise in estrogen levels during the midluteal phase with a decrease before menstruation. The secondary rise in estradiol parallels the rise of serum progesterone and 17-hydroxyprogesterone levels^[53].

Dynamic changes in ovarian hormones during menstrual cycle can modulate GI contractility, transit, secretion, visceral sensitivity, and immune function at multiple target sites, including those located in the periphery and the brain^[5]. Clinical studies indicate that declining or low ovarian hormone levels in women (such as during menses) may contribute to the occurrence or exacerbation of GI symptoms, including abdominal pain or discomfort, altered bowel habits and bloating that varies across the menstrual cycle phases (Table 1)^[3,4-37]. Rectal sensitivity thresholds have been shown to be significantly lower in IBS patients at menses relative to those at other cycle phases indicating that IBS symptoms experience may be modified by ovarian hormone status^[38]. Also in animal studies it has been shown that both visceral and somatic sensitivity vary over the rat estrous cycle and that high levels of ovarian hormones (proestrus/estrus stages) are associated with enhanced sensitivity^[39]. Therefore the menstrual cycle provides a natural model to explore the effects of ovarian hormones on the bowel function.

Approximately one third of otherwise asymptomatic women experience GI symptoms at the time of menstruation^[3,4]. About 40% of women with IBS report

influence of the menstrual cycle on their symptoms^[35]. Whitehead *et al.*^[37] found that in women with functional bowel disorders (FBDs), including IBS, bowel symptoms seem to be affected by menstruation to a greater degree than in women without FBDs, suggesting that IBS women may respond differently to the fluctuations in the ovarian hormones. Variation in GI symptoms during the menstrual cycle can be related to motor disturbances and/or a change in perception of colonic motor events, as well as alterations in colonic epithelial barrier and mucosal immunity^[10,40].

IBS female patients are more likely to report dysmenorrhea and premenstrual distress syndrome than those who do not suffer from IBS^[41-43]. Moreover, IBS patients with dysmenorrhea report noticeably more GI symptoms than non-dysmenorrheic women^[41]. In a 10-year follow-up-study conducted in Iceland it has been shown that IBS female patients with dysmenorrhea were twice more likely to have increased symptoms compared to IBS patients without dysmenorrhea^[43].

A significant connection between IBS and endometriosis has also been reported^[43,44]. Additionally, polycystic ovary syndrome (PCOS), the most common female endocrine disorder affecting up to 10% of reproductive-age women characterized by chronic anovulation and hyperandrogenism, is associated with the increased prevalence of IBS^[45-47]. Interestingly, IBS coexisting with PCOS was associated with a higher BMI and percent body fat when compared to PCOS alone^[45]. The relationships between obesity, hormonal status and IBS require further investigation, particularly in the context

of obesity being linked with increased inflammatory mediators and in the light of recent reports on the GI dysbiosis^[46,48].

Pregnancy

Pregnancy is characterized by high ovarian hormones levels as well as an increase in opioid-mediated antinociception^[3,49]. Little is known, however, regarding IBS symptoms and pregnancy. Many chronic pain syndromes frequently associated with IBS, like migraine headache for example, are alleviated during the time of pregnancy^[27]. Similarly, in rodents high ovarian hormones levels during pregnancy reduce somatic and visceral pain sensitivity^[50]. During the time of the physiological hyperestrogenemia and hyperprogesteronemia a prolonged GI transit is also observed^[51]. Additionally, numerous psychological variables affecting the autonomic nervous system (ANS) may trigger or modulate symptoms reported in pregnant women^[52].

Menopause

Data concerning the impact of the menopause transition on IBS patients remain inconsistent. Although the decline in ovarian hormones may induce or exacerbate GI symptoms, generally, in postmenopausal period, the incidence of IBS decreases significantly^[53-55]. However, according to some recent data, IBS symptoms severity may increase after menopause as well^[43]. Cain *et al.*^[26] found that various GI symptoms were reported more frequently by postmenopausal women compared with men, but these differences were not significant when controlled for age. In one study, gas and excessive flatulence were more prevalent in post- than premenopausal healthy women^[53].

Hormone supplementation

Premenopausal healthy women taking oral contraceptives (OCs), monophasic or triphasic preparations, report a typical increase in GI symptoms at menses^[55]. However, women with IBS taking OCs, which contain both estrogen and progestin, appeared to have reduced levels of abdominal symptoms compared with IBS women not taking OCs^[55]. At the same time, the pattern of GI and non-GI symptoms over the menstrual cycle was similar in female patients with IBS, regardless of OCs use or the predominant bowel pattern^[55]. Noteworthy, in women with dysmenorrhea that may coexist with IBS, OCs often reduce the symptoms^[15]. Recently, Bird *et al.*^[56] reported an increased risk for development of IBS with drospirenone. Drospirenone is a synthetic progestin approved in combination with ethinyl-estradiol as an OC. Although it was designed as an antimineralocorticoid steroid, it exhibits antiandrogen activity^[56]. In another study evaluating the effect of hormone supplementation on IBS symptoms, the therapeutic efficacy of gonadotropin-releasing hormone agonist (leuprolid) in female patients with menstrual cycle-related symptoms has been reported^[57]. However, the use of this medication is limited by its side effects^[57].

Based on the recent meta-analysis, there are insufficient data to determine the exact effect of hormone supplementation during menopause on IBS symptoms^[19]. In postmenopausal women, hormone replacement therapy (HRT) has been reported to be associated with the increased prevalence of IBS. HRT may prolongs IBS symptoms to a later age or even induce changes in GI function in females not previously affected^[58]. One of the confounding factors may be related to the fact that women with IBS are more likely to report various pre- and postmenopausal symptoms, and thus may be prescribed HRT to a greater degree. However, Ruigómez *et al.*^[58] have shown that both current and past users of HRT presented an increased risk of IBS compared to non-users, even after adjusting for comorbidity and consultation pattern. This increased risk was irrespective of treatment duration, regimen or route of administration of HRT^[58].

Gynecological surgery

There are few data concerning the prevalence of oophorectomy or hysterectomy in IBS female patients, mostly because these surgical procedures are excluding factors in the studies of IBS patients. However, it has been reported that the rate of hysterectomy is about twice higher in women with IBS compared to controls^[59]. It is conceivable that IBS patients, because of the chronic abdominal pain, are more likely to be qualified for various surgical procedures (not only gynecological, but also GI surgery like cholecystectomy and appendectomy)^[59]. In fact, in a number of women, GI symptoms emerge for the first time after gynecological surgery^[60]. Preclinical studies however remain controversial. There is indeed evidence in mice that ovariectomy generates a slow developing and persistent hyperalgesic state localized to the abdomen, lower limbs and abdominal viscera, which is reversed by estrogen supplementation^[61,62]. In contrast, in rats, ovariectomy decreased the magnitude of the visceromotor response to colorectal distension compared with cycling rats^[63] and abolished restraint stress-induced visceral hypersensitivity^[64]. The sensitivity to colorectal distension and the influence of stress on visceral pain were restored by estrogen replacement at a dose comparable to the proestrus level^[65].

Male sex hormones

Most of the explanations of sex-related differences in IBS have focused on the concept that women might be more susceptible, while less attention has been given to the concept that male hormones may be protective against pain disorders including IBS^[66]. Androgens, higher in males than females, appear to protect against the development of chronic pain disorders in humans, and testosterone exerts an analgesic effect in experimental pain models, in both men and women^[67-69]. Differences in androgen levels, their receptors as well as sites of action may play a role in the sex difference in the risk of developing chronic pain disorders. There are only few reports concerning the role of sex hormones in male

patients with IBS^[67,70,71]. Houghton *et al*^[66] found that testosterone levels, although similar in the patient and control groups, correlated negatively with perceptual thresholds of rectal distension and overall well-being in IBS patients. In the same study it was found that middle-aged male IBS patients tended to have lower levels of luteinizing hormone compared with male control subjects^[66]. Kim *et al*^[70] have also reported that the sex hormone status in young male patients is different from that of older male patients and that an elevated sex hormone-binding globulin level might play a key role in the pathophysiology of IBS in young men. Interestingly, a highly significant reduction in male-trait scores in men with IBS has been confirmed^[71]. Another unique model to study the relationship between sex hormones and chronic pain was proposed by Aloisi *et al*^[72] who evaluated the results of sex-crossed hormone administration in transgender subjects. About half of the female-to-male subjects treated with testosterone reported a significant improvement of the chronic pain (*e.g.*, headache) present before the treatment. Conversely, about one-third of the male-to-female subjects receiving estrogen/anti-androgen treatment developed chronic pain including headaches, breast and musculoskeletal pain, and in some cases visceral pain as well^[72]. These findings support experimental and clinical data suggesting that sex steroid hormones play a crucial role in pain perception and modulation.

SEX HORMONE MODULATION OF THE BRAIN-GUT AXIS AT THE CENTRAL NERVOUS SYSTEM LEVEL

Estrogens

The abundant distribution of estrogen receptors (ERs) at all levels of the brain-gut axis, including the central nervous system (CNS), spinal cord, and the enteric nervous system supports the multiplicity of neuronal action^[73]. There are two subtypes of ERs: ER- α and ER- β . Estrogens, similarly to progesterone and testosterone, exert their function by binding to either specific intracellular (nuclear) receptors that act as ligand-dependent transcription factors (classical mechanisms) or membrane-bound receptors (mERs) that stimulate several signal transduction pathways (non-classical mechanisms). The family of nuclear receptors mediate rather slow genomic action of estradiol resulting in enhancement or repression of gene transcription and thus protein synthesis alterations. In contrast, mERs are involved in the rapid action of estrogens related to the activation of various protein-kinase cascades and phosphorylation of proteins, but estrogenic rapid signaling can also occur by recruiting intracellular pathways that can act *via* the genome through phosphorylated cyclic adenosine monophosphate (cAMP) response element protein (pCREB) and intermediate early genes^[74]. In addition to the well described G protein-coupled receptor (GPR30), multiple mERs have recently been discovered, such as the classical nuclear ER- α and ER- β , ER- α 44, ER-X and mER-G α q^[74-76].

ERs are spread throughout the brain, including the amygdala, hypothalamus, pituitary, hippocampus, cerebral cortex, mid-brain, and brain stem, providing neuro-anatomical support for potential numerous target sites of estrogen actions on neurocognitive processes^[73,77]. Based on the results of brain imaging studies, greater responsiveness of emotional arousal circuits in relation to visceral pain has been implicated as inducing central mechanisms of pain amplification in IBS, with female subjects showing greater response than male subjects^[78]. Recent results confirmed sex differences in emotion-related cognitive processes and functioning of brain networks including the prefrontal regions, cingulate, insula, and amygdala in IBS and healthy control subjects^[79].

Estrogens may act in the CNS through multiple pathways modulating production and action of neurotransmitters, influencing electrical excitability and synaptic function, and changing the morphological features of neural elements involved in the function^[77,80,81]. Estrogens have been documented to exert differential, sometimes opposite effects on pain. Clinical and experimental data indicate that both analgesic and hyperalgesic responses can be induced by estrogens depending upon the experimental conditions^[67]. Estrogens were shown to enhance neuronal system activities during development and in adult life, for instance through the hippocampal neuronal circuits involving acetylcholine, glutamate and brain-derived neurotrophic factor^[82]. Noteworthy, elevated levels of estrogens in fertile women have been associated with the increased number of μ -opioid receptors in the brain regions related to pain processing^[68]. There is accumulating evidence that estrogens have a significant impact on neuronal plasticity-related process and ameliorate recovery after chronic stress (Table 2)^[73].

Estrogens may also contribute to the important sex differences in the stress-related hypothalamic-pituitary-adrenal (HPA) axis response that have been documented in a number of clinical and experimental studies^[83]. The menstrual cycle phases, menopausal status and pregnancy have been shown to affect the HPA axis as well as ANS functions^[5]. Women between puberty and menopause usually show lower HPA axis and autonomic responses to psychological stressors than men of the same age^[84]. However, the HPA axis response to psychological stressors is higher in the luteal phase, when post-stress free cortisol level approaches that for men^[84]. CRF is a key mediator of the HPA axis and the brain-gut axis response to stress at both central and peripheral levels^[9,85,86]. The co-localization of ER- α with CRF receptors in the hypothalamus represents one of the possible neuroendocrine interactions between CRF signaling pathways and estrogens^[87]. Importantly, activation of both receptors ER- α and ER- β has been shown to stimulate CRF gene expression in the hypothalamic paraventricular nucleus (PVN)^[83,88]. Additionally, estrogens induce also an increase in glucocorticoid receptor expression in the amygdala^[88]. In the recent functional magnetic resonance imaging study, it was demonstrated that significant sex differences in brain activity in stress

Table 2 Sex hormone modulation of the brain-gut-microbiota axis

Level of the brain-gut-microbiota axis	Estrogen	Progesterone	Testosterone
Central nervous system	Analgesic or hyperalgesic effect ^[67] Excitatory action on neurons ^[72] Estrogen-induced increase in the number of μ -opioid receptors ^[68] Enhancement of serotonergic postsynaptic responsiveness in the brain ^[8] Central interaction with CRF signaling pathways-modulation of stress responsiveness ^[87,89] : Stimulation of CRF gene expression in PVN ^[83] Increase in glucocorticoid receptor expression in the amygdala ^[83] Influence on neuronal plasticity-related processes ^[73]	Activation of the γ -aminobutyric acid (GABA) receptors, major inhibitory receptors in the brain ^[77] Neuroprotective action in the hippocampus ^[80]	Analgesic effect ^[72] Inhibition of stress-induced ACTH release ^[103]
Autonomic nervous system	Attenuation of sympathetic responsiveness ^[108]	Reduced cholinergic responsiveness ^[5]	Regulation of parasympathetic tone ^[110]
Enteric nervous system/ Gut immune system	Expression of estrogen receptors in enteric neurons, regulation of neurogenic reflexes ^[73] Activation of colonic NK1 receptors in stress-induced visceral hypersensitivity ^[64] Augmentation of mast cells secretion ^[118] Effects on both pro- and anti-inflammatory pathways ^[113] Peripheral interaction with CRF signaling pathways, modulation of colonic motor and sensory responses to stress ^[87] Regulation of 5-HT3 receptor expression in rat colon ^[120] Regulation of secretory and absorptive function of gastrointestinal epithelial cells ^[128] Enhanced expression of trans-membrane tight junction protein in non-inflamed colon ^[124] Decreased production of proinflammatory cytokines in experimental colitis in female rats ^[125,126]	Inhibition of gastrointestinal motility ^[130] Inhibition of visceral signaling following colonic inflammation ^[100] Inhibition of mast cells degranulation ^[131] Immunosuppressive action related to inhibition of NF κ B activation in macrophages ^[133]	Stimulation of smooth muscle contractions ^[135] Decreased production of proinflammatory mediators inducing visceral hyperalgesia ^[69,136] No effect on mast cells degranulation ^[137] Decreased TLR4 expression of macrophages and monocytes ^[138]
Gut microbiota	ER- β expression affects the gut microbiota composition ^[143] Microbial β -glucuronidase activity determines estrogens deconjugation enabling their reabsorption <i>via</i> enterohepatic circulation ^[146] Direct effect on bacterial metabolism, growth and expression of virulence factors ^[132] Bacterial hydroxysteroid dehydrogenase regulates the balance between active and inactive steroids ^[132]	Direct effect on bacterial metabolism, growth, and expression of virulence factors ^[132]	Reversible 17 β reduction of androgens may regulate testosterone level ^[148] Commensal microbiota-dependent testosterone production protects against autoimmune disease in mice ^[149] Testosterone production protects against autoimmune disease in mice ^[149]

ACTH: Adrenocorticotrophic hormone; CRF: Corticotropin-releasing factor; NF κ B: Nuclear factor κ B; PVN: Paraventricular nucleus; TLR4: Toll-like receptor 4.

response circuitry were dependent on women’s menstrual cycle phase^[89]. In addition, chronic treatment with estrogens modulates brain circuitry responsive to stress^[90]. Furthermore, administration of estradiol and progesterone directly to the amygdala in rats increases pain response to visceral stimulation suggesting that an amygdala-dependent mechanism may be responsible, at least in part, for the exacerbation of visceral symptomatology in females^[91]. A recent meta-analysis by Tillisch *et al*^[92] points to the amygdala, a brain region known to facilitate HPA axis output, as one of the most consistently activated areas following rectal stimulation in IBS patient compared with controls. Of significance, the activation of the amygdala by corticosterone eliminates spontaneously occurring differences in visceral and somatic pain perception in cycling female rats, resulting in visceral hypersensitivity during metestrus/diestrus, and increased somatic sensitivity during both metestrus/diestrus as

well as proestrus/estrus^[39]. This observation could explain the lowered pain thresholds and higher incidence of somatic pain observed in women with IBS^[39].

Childhood trauma (early adverse life event, EAL) is associated with changes in HPA axis responsiveness in IBS^[93]. Dysregulation of the HPA axis in IBS patients has been related to blunted adrenocorticotrophic hormone (ACTH) levels and enhanced cortisol response to visceral stimulation^[94]. However, little is known on sex-differences in EAL-induced visceral pain. Interestingly, sexually dimorphic effects of unpredictable EAL on visceral pain behavior in a rodent model has been demonstrated^[95]. Female rats exposed neonatally to different pairings of an odor and shock developed visceral hypersensitivity in adulthood, while in contrast, in male rats, visceral sensitivity was not significantly different after EAL. Visceral sensitivity following unpredictable EAL was reversed by ovariectomy and reestablished by estradiol

replacement. These data suggest estrogen-mediated pivotal mechanisms in maintaining visceral hypersensitivity^[95].

The serotonergic system represents another potential contribution to sex differences in pain modulation^[3,8]. In the CNS, serotonin (5-HT) generally has been associated with descending pain inhibition, whereas in the periphery, 5-HT is an inflammatory mediator and is generally pronociceptive and prokinetic. Estrogens enhance serotonergic postsynaptic responsiveness in the brain^[8]. Additionally, estrogens enhance 5-HT synthesis in most part of the brain by increasing expression of the enzyme tryptophan hydroxylase and decreasing the expression of the serotonin re-uptake transporter^[96]. The serotonergic and reproductive endocrine systems are also prominently involved in both the regulation of mood and behavioral states. In addition, interactions between these systems have profound implications for the etiology and treatment of anxiety disorders^[97]. A growing body of evidence also indicates sex-dependent differences in serotonin-related genetic polymorphisms in IBS patients^[98], particularly with regard to anxiety and depressive disorders more common in women with IBS^[99].

Progesterone

The role of progesterone in sex-related differences in pain modulation is less clear. Progesterone activates intracellular receptors to regulate genomic processes, and also affects cell membrane receptors, especially in neurons^[100]. Membrane progesterone receptors present in the hippocampus were suggested to contribute the neuroprotective action of the hormone^[80]. At the CNS level, progesterone action seems to be dependent on the activation of the γ -aminobutyric acid receptors that are major inhibitory receptors in the brain^[101].

Androgens

While estrogens are commonly indicated as CNS stimulant, androgen receptor-mediated actions are often related to CNS inhibition, which may underlie the lower incidence of many forms of chronic pain in men^[72]. Androgens participate in the regulation of the HPA axis response to chronic stress and the autonomic circuitry^[102]. Optical and electron microscopic immunocytochemical studies in rodents have revealed that the distribution of androgen receptors is overlapping with that of ER- α , ER- β , as well as progesterone receptors in three major autonomic regions in the brain: the rostral ventrolateral medulla, nucleus of the solitary tract and PVN^[80]. In male rats, testosterone inhibits the acute restraint stress-induced ACTH release^[103] that, ultimately, may impact on other brain stress-related CRF-mediated influence on colonic motility and visceral pain^[9].

SEX HORMONE EFFECTS ON THE AUTONOMIC NERVOUS SYSTEM

Estrogens

Estrogens influence also nociceptive pathways at the

level of primary afferent nerves and spinal cord projections^[104]. A direct involvement of ERs in nociceptive transmission is possible *via* their activation of enkephalin synthesizing cells in the superficial laminae of the spinal cord^[105]. Additionally, estrogens modulate the responsiveness of primary vagal afferents neurons to substance P and the activation of glutamate receptors involved in the afferent pain pathways^[40]. Spinal estrogen receptors ER- α and ER- β have been also shown to contribute to the facilitation of N-methyl-D-aspartate-dependent colon-to-urethra cross-organ reflex sensitization, which is presumed to underlie pelvic viscerovisceral referred pain^[106].

Autonomic dysregulation in response to a visceral stressor is an objective physiologic correlate in IBS^[107]. Tillisch *et al*^[108] reported gender differences in the ANS reactivity to colorectal distension in IBS patients, with men demonstrating increased sympathetic nervous system activation and decreased parasympathetic activation compared to women. There are also data indicating menstrual cycle-linked differences in the ANS tone that are likely to result from estrogen exposure, and its attenuating influence on sympathetic responsiveness^[109]. Furthermore, many other chronic pain syndromes, frequently coexisting with IBS can be also related to autonomic disturbances^[28].

Progesterone and androgens

Progesterone has been shown to reduce cholinergic responsiveness^[5]. However, little is known about the effect of testosterone on the ANS. Recently, it has been noticed that testosterone deficiency is accompanied by a decrease in basal parasympathetic tone and reduced baroreflex sensitivity in men with heart failure^[110].

SEX HORMONE ACTIONS AT THE ENTERIC NERVOUS AND GUT IMMUNE SYSTEMS

Estrogens

Within the enteric neurons of the colon, where both CRF receptor subtype 1 (CRF₁) and ERs are expressed, interactions between CRF signaling pathways and estrogens participate in the stimulation of the colonic motor function^[90]. Additionally, a local paracrine/autocrine pro-inflammatory action by CRF₁ receptor activation was reported in several models of intestinal inflammation both *in vitro* and *in vivo*, as well as the up-regulation of CRF and CRF₁ expression in immune cells of the human colonic lamina propria in response to inflammation^[86].

There is compelling evidence suggesting an up-regulated gut immune function in patients with IBS, particularly with post-infectious IBS^[111]. Gastrointestinal inflammation seems to be strongly modulated by stress, especially in IBS patients being characterized by enhanced stress responsiveness^[15]. Important sex-related differences in IBS patients related to neuroimmune inter-

actions have been suggested^[71,112]. Female sex is an independent risk factor for developing postinfectious IBS^[3]. The following observations support sex differences in immune response: females produce stronger cellular as well as humoral immune reaction, have a greater resistance to bacterial infections, and are more likely to develop autoimmune diseases compared to men, symptoms of which depend on hormonal status^[113]. Estrogens may influence both pro- and anti-inflammatory pathways. The effect of estrogens in inflammatory responses has been found extremely complex and dependent on the estrogen level, the cell type, specific inflammatory factors, the type of tissue that is inflamed, the time course of the inflammatory response (*e.g.*, acute vs chronic), and the time point at which estrogen exposure occurs^[113]. In an experimental model, estrogens contributed also to the colonic neurokinin-1 receptor-mediated effects of stress-induced visceral hypersensitivity to colorectal distension^[64].

Mast cells represent another crucial link in sex-dependent neuroimmune interactions as they co-express CRF and sex hormone receptors^[114-116]. The number of colonic mucosal mast cells was found to be higher in female compared to male IBS patients^[10]. Mediators released by activated mast cells, characterized by extensive anatomical and functional communication with intrinsic and extrinsic nervous system of the gut, evoke visceral hypersensitivity and increase mucosal permeability^[117]. Notably, mast cells are involved in many other disorders, frequently overlapping with IBS such as fibromyalgia, interstitial cystitis, chronic fatigue syndrome and migraine, all of which occur more often in women, are exacerbated during ovulation and reduced during pregnancy^[10]. These sex-related differences in the prevalence and severity of chronic pain disorders could be related to the fact that mast cells express progesterone and estrogen receptors^[10]. Estradiol has been shown to augment mast cells secretion, whereas tamoxifen (an estradiol receptor antagonist) inhibits this function^[118].

The serotonergic system at the peripheral level may also contribute to sex differences in modulation of GI motility, secretion and sensitivity^[3,8]. Fluctuations in estrogen levels during ovarian cycle cause predictable changes in 5-HT system in women^[8]. Moreover, 5-HT concentration varies with sex and menstrual status in patients with diarrhea-predominant IBS^[119]. Experimental studies indicate that colonic 5-HT₃ receptor gene expression is increased in ovariectomized rats exposed to restraint stress and restored with hormone replacement after ovariectomy^[120]. Recently, Galligan *et al.*^[121] proposed serotonin transporter (*SERT*) gene knockout (KO) rats as a new interesting model for studying interactions between serotonin, sex, and visceral sensation. *SERT* KO female rats display an increased colonic extracellular serotonin associated with visceral hypersensitivity and hyperexcitability of colon projecting sensory neurons, which is not observed in male *SERT* KO rats^[121]. Gender difference has been also shown in *SERT* activity and serotonin concentration in platelets of IBS patients^[122].

Estrogen-dependent modulation of the intestinal barrier function is another component in sex-related differences in IBS. It has been well established that stress involving the activation of CRF₁ receptors alters intestinal barrier that appears to be a prerequisite for the development of visceral hypersensitivity in both human and rodents^[6,123]. In the colon, ERs signaling enhances expression of trans-membrane tight junction proteins in non-inflamed conditions^[124], and decreases production of proinflammatory cytokines in experimental colitis^[125,126]. In human, acute experimental stress evokes a differential gender-dependent increase in intestinal macromolecular permeability^[127]. A significant increase in albumin permeability in healthy women, but not in men, could explain enhanced female susceptibility to IBS^[127].

Additionally, ERs are localized on the epithelial cells throughout the GI mucosa and may affect secretory and absorptive functions^[37,128,129]. The fluid retention that occurs in females during the cycle may be associated with the extra-nuclear action of estrogen that can stimulate calcium entry into colonic epithelial cells as well as suppress c-AMP-dependent chloride secretion in the distal colonic epithelium in females only, both in rats and humans^[128].

Progesterone and androgens

At the peripheral level progesterone has been suggested to influence both visceral sensitivity and motility *via* prostaglandins^[100]. Overexpression of progesterone receptors in colonic muscle in women with slow transit constipation is associated with lower levels of prostaglandin PGF_{2α} and tromboxane A that cause muscle contraction, and higher levels of PGs that cause muscle relaxation (such as PGE₂)^[130]. Progesterone may also inhibit estrogen-dependent mast cells degranulation^[131]. Progesterone receptors have been identified in epithelial cells, granulocytes, macrophages, and lymphocytes^[132]. Progesterone is known to exert an immunosuppressive action as it inhibits the activation of nuclear factor (NF)κB and increases the expression of the suppressor of cytokine signaling protein (COS1) in macrophages^[133].

Testosterone and its active metabolite 5α-dihydrotestosterone are potent modulators of colonic motility by stimulating smooth muscle contractions through non-genomic calcium sensitization pathways^[134]. In the urethral calculus model of visceral pain, Aloisi *et al.*^[135] did not find any significant effect of testosterone on visceral pain. However, there is growing body of evidence that androgens may contribute to the modulation of visceral pain by decreasing pro-inflammatory mediators that participate in the development of hyperalgesia^[69,136]. Apart from female sex hormone receptor expression, mast cells also express androgen receptor, however, testosterone treatment had no effect on mast cell degranulation^[137]. Testosterone decreases also the expression of macrophage and monocyte Toll-like receptor 4, which is involved in the activation of the innate system response to pathogen challenge^[138].

INTERACTIONS BETWEEN SEX HORMONES AND THE GUT MICROBIOTA

The increasing knowledge of the role of microbiota in health and disease state has shed a light on the critical role of the enteric microbiota, both commensal and pathogenic organisms, in regulation the brain-gut axis. This has consequently led to the coining of a new term: the brain-gut-enteric microbiota axis^[11]. The bi-directional communication between the gut bacteria and the brain occurs through neural, immune, and endocrine pathways^[12,48,139,140] which may be modulated by sex hormones, in particular estrogens. In fact, it has been recently reported that the microbiome-gut-brain axis during early life regulates the hippocampal serotonergic system in a sex-dependent manner^[141]. Clarke *et al.*^[141] found that male germ free mice, unlike females, display a significant elevation in the hippocampal concentration of 5-HT and its metabolite, compared with conventionally colonized control animals.

Numerous studies have reported the effects of sex hormones on the dimorphic sex differences in the response to microbial and viral infections^[132]. Besides the role of sex hormones in the modulation of the immune system, they have a direct effect over bacterial metabolism, growth, and expression of virulence factors. For instance during pregnancy, the proportion of certain bacteria species associated with plaque microbiota is altered with a noticeable increase in the ratio of anaerobic to facultative bacteria^[142]. Of significance, recent studies indicate that steroid nuclear receptor expression including ER- β can determine the intestinal microbiota composition^[143].

Moreover, the gut microbiota may also affect estrogens metabolisms and their systemic level^[144]. Conjugated estrogens are excreted in the bile and pass into the distal ileum, where they are variably deconjugated and may be reabsorbed from the gut lumen and enter the circulation *via* the portal vein^[145]. It has been shown in men and postmenopausal women that the intestinal microbiota richness and function, associated for example with β -glucuronidase activity, influence levels of non-ovarian estrogens *via* enterohepatic circulation^[146]. Bacteria are capable of metabolizing sex hormones through the activity of various enzymes such as hydroxysteroid dehydrogenase that regulate the balance between active and inactive steroids^[130]. In particular, fecal bacteria can perform hydrolytic, reductive and oxidative reactions of estrogens and androgens^[147]. Reversible 17 β reduction of androgens carried out by the gut microbiota is suggested to play a role in the regulation of testosterone level^[148]. The results of a landmark study published recently by Markle *et al.*^[149] provide astonishing conclusions indicating that sex differences in the gut microbiome drive hormone-dependent regulation of autoimmunity. In the study performed in the non-obese diabetic mouse model of type 1 diabetes, they showed that male puberty in mice leads to changes in the gut microbiota

that reinforce testosterone production, which is protective against the development of T and B cell functions linked to autoimmune disease^[149]. In mice, the properties of the male-associated microbiota can be transferred to younger females and exert testosterone-mediated protection from autoimmune disease upon recipients. The observations that early-life microbial exposures determine sex hormone levels and modify sex-mediated immune regulation may have crucial implications for the pathophysiology of IBS. A new concept of “microgenderome” is emerging based on the recent observations that the gender bias present in numerous diseases is not entirely a host-intrinsic factor, but may be exercised and/or reinforced by the commensal microbiota of the host^[150]. Undoubtedly, further studies are needed to elucidate the role of microgenderome in IBS.

THERAPEUTIC IMPLICATIONS

The modulator role of sex hormones on the bi-directional interactions within the brain-gut-microbiota axis may have significant therapeutic implications in IBS. However, although gender differences in responses to treatment modalities exist, the approach to IBS patients in both genders is quite similar so far. Clinical observations confirm that alosetron, a 5-HT₃ receptor antagonist, is more effective in improving urgency and loose stools in IBS-diarrhea predominant women than men^[120,151,152]. The basis for this noticeable sex difference in therapeutic efficacy of alosetron could be associated with sex-related differences in 5-HT₃ receptor expression, lower alosetron clearance in women, and/or greater 5-HT synthesis in certain brain regions in IBS male patients compared with female IBS patients^[153]. Sex difference in genetic polymorphism of the 5-HT transporter (SERT) promoter region has been also suggested and may induce the different expression of affective symptoms in women compared with men^[154]. Additionally, the potential role of interaction between gonadal hormones and the cytochrome P450 pathway may be considered in sex-related differences in drug clearance^[155]. Differences in adipose tissue compartment in women compared to men may affect this process as well^[140].

Women with IBS are more susceptible to anxiety and depression and other stress-related disorders. However, in the study comparing the efficacy of treatment with paroxetine alone or combined with psychotherapy, no gender effect was reported^[156]. Preliminary observations suggesting that IBS female patients may better respond to hypnotherapy^[157] is yet to be confirmed. The recent results of randomized controlled trial have shown that gender, age, disease duration and IBS type have no influence on the long-term success of gut-directed hypnotherapy^[158].

Regarding the role of sex hormones in the pathogenesis of IBS, therapeutic approaches aiming to suppress ovarian steroidogenesis have been also considered. In fact, gonadotropin-releasing hormone agonist (leuprolide)




Figure 1 Sex hormones in the mutual brain-gut-microbiota interactions. Sex hormones influence peripheral and central regulatory mechanisms involved in the pathophysiology of irritable bowel syndrome contributing to the alterations in stress response, visceral sensitivity and motility, intestinal barrier function, and immune activation of intestinal mucosa. Sex hormones have also a direct effect on the gut microbiota. ENS: Enteric nervous system.

was reported to be effective in IBS female patients with menstrual cycle-related symptoms^[57]. Nevertheless, many unpleasant side effects of leuprolide similar to climacteric-like syndrome significantly limit its application^[57].

Noteworthy, interactions between gonadal hormones and pain modulation are bi-directional, as pain therapies in different experimental and clinical conditions have been found to affect the gonads as well^[159,160]. For example, morphine treatment increased estrogen receptor, androgen receptor and *TRPV1* genes expression in the ovary, whereas in the testis the opiate reduced ER- α and ER- β mRNA expression not affecting androgen receptor and *TRPV1* expression^[160].

A pivotal interdependence between the composition and stability of the gut microbiota and GI function as well as stress-related behavioral changes indicate a great therapeutic potential of probiotics, prebiotics and antibiotics in IBS^[161,162]. So far, no gender specificity in probiotics efficacy in IBS patients has been reported^[163]. Nevertheless, in the light of the microgenderome concept and sex-dependent differences in the immune regulation driven by gut microbiome^[150], gender specificity

in microflora manipulation seem to be essential and is expected to be extensively explored in the near future.

CONCLUSION

The results of epidemiological studies and clinical observations confirm significant sex and gender differences in the IBS prevalence and symptomatology. Furthermore, a growing number of clinical and experimental data strongly support a crucial role of sex hormones in the regulatory mechanisms of the brain-gut-microbiota axis involved in the pathophysiology of IBS (Figure 1). Some discrepancies in the results, especially related to the influence of estrogens, may result from different experimental conditions or heterogeneous groups of patients (*e.g.*, different age, menstrual status), but they also reflect the very complex nature of sex hormone actions. Estrogens can induce dual effects, both analgesic or hyperalgesic, as well as pro- or anti-inflammatory. Noteworthy, alterations in estrogen-induced visceral sensitivity seem to depend not only on the gonadal hormones levels, but more so on sudden changes in their levels, their sus-

tained genomic effects, and complex interactions with other neurotransmitters. Concomitant alterations in the number (up- or down-regulation) and sensitivity of ERs may play a crucial role in these processes as well. Thus, the physiological fluctuation in sex hormones may evoke for example different responses in IBS female patients compared to healthy women. Furthermore, a growing body of evidence indicates a protective role of androgens in pain modulation and anti-inflammatory properties of testosterone that may inhibit the development of visceral hyperalgesia. That could contribute to the higher susceptibility of women to IBS. A better understanding of the role of sex hormones in the modulation of the brain-gut-microbiota axis should enable a more effective and sex-tailored therapeutic approach in IBS.

REFERENCES

- Hogan AM, Collins D, Baird AW, Winter DC. Estrogen and its role in gastrointestinal health and disease. *Int J Colorectal Dis* 2009; **24**: 1367-1375 [PMID: 19655153 DOI: 10.1007/s00384-009-0785-0]
- Ji Y, Tang B, Traub RJ. The visceromotor response to colorectal distention fluctuates with the estrous cycle in rats. *Neuroscience* 2008; **154**: 1562-1567 [PMID: 18550290 DOI: 10.1016/j.neuroscience.2008.04.070]
- Longstreth GF, Thompson WG, Chey WD, Houghton LA, Mearin F, Spiller RC. Functional bowel disorders. *Gastroenterology* 2006; **130**: 1480-1491 [PMID: 16678561]
- Heitkemper MM, Chang L. Do fluctuations in ovarian hormones affect gastrointestinal symptoms in women with irritable bowel syndrome? *Gen Med* 2009; **6** Suppl 2: 152-167 [PMID: 19406367 DOI: 10.1016/j.genm.2009.03.004]
- Filligim RB, King CD, Ribeiro-Dasilva MC, Rahim-Williams B, Riley JL. Sex, gender, and pain: a review of recent clinical and experimental findings. *J Pain* 2009; **10**: 447-485 [PMID: 19411059 DOI: 10.1016/j.jpain.2008.12.001]
- Larauche M, Mulak A, Taché Y. Stress and visceral pain: from animal models to clinical therapies. *Exp Neurol* 2012; **233**: 49-67 [PMID: 21575632 DOI: 10.1016/j.expneurol.2011.04.020]
- Mulak A, Bonaz B. Irritable bowel syndrome: a model of the brain-gut interactions. *Med Sci Monit* 2004; **10**: RA55-RA62 [PMID: 15260348]
- Rybaczylk LA, Bashaw MJ, Pathak DR, Moody SM, Gilders RM, Holzschu DL. An overlooked connection: serotonergic mediation of estrogen-related physiology and pathology. *BMC Womens Health* 2005; **5**: 12 [PMID: 16368009]
- Taché Y, Bonaz B. Corticotropin-releasing factor receptors and stress-related alterations of gut motor function. *J Clin Invest* 2007; **117**: 33-40 [PMID: 17200704]
- Cremon C, Gargano L, Morselli-Labate AM, Santini D, Cogliandro RF, De Giorgio R, Stanghellini V, Corinaldesi R, Barbara G. Mucosal immune activation in irritable bowel syndrome: gender-dependence and association with digestive symptoms. *Am J Gastroenterol* 2009; **104**: 392-400 [PMID: 19174797 DOI: 10.1038/ajg.2008.94]
- Rhee SH, Pothoulakis C, Mayer EA. Principles and clinical implications of the brain-gut-enteric microbiota axis. *Nat Rev Gastroenterol Hepatol* 2009; **6**: 306-314 [PMID: 19404271 DOI: 10.1038/nrgastro.2009.35]
- Cryan JE, Dinan TG. Mind-altering microorganisms: the impact of the gut microbiota on brain and behaviour. *Nat Rev Neurosci* 2012; **13**: 701-712 [PMID: 22968153 DOI: 10.1038/nrn3346]
- Heitkemper M, Jarrett M, Bond EF, Chang L. Impact of sex and gender on irritable bowel syndrome. *Biol Res Nurs* 2003; **5**: 56-65 [PMID: 12886671]
- Longstreth GF, Wolde-Tsadik G. Irritable bowel-type symptoms in HMO examinees. Prevalence, demographics, and clinical correlates. *Dig Dis Sci* 1993; **38**: 1581-1589 [PMID: 8359067]
- Toner BB, Akman D. Gender role and irritable bowel syndrome: literature review and hypothesis. *Am J Gastroenterol* 2000; **95**: 11-16 [PMID: 10638553]
- Gwee KA, Lu CL, Ghoshal UC. Epidemiology of irritable bowel syndrome in Asia: something old, something new, something borrowed. *J Gastroenterol Hepatol* 2009; **24**: 1601-1607 [PMID: 19788601 DOI: 10.1111/j.1440-1746.2009.05984.x]
- Lovell RM, Ford AC. Effect of gender on prevalence of irritable bowel syndrome in the community: systematic review and meta-analysis. *Am J Gastroenterol* 2012; **107**: 991-1000 [PMID: 22613905 DOI: 10.1038/ajg.2012.131]
- Husain N, Chaudhry IB, Jafri F, Niaz SK, Tomenson B, Creed F. A population-based study of irritable bowel syndrome in a non-Western population. *Neurogastroenterol Motil* 2008; **20**: 1022-1029 [PMID: 18492027 DOI: 10.1111/j.1365-2982.2008.01143.x]
- Adeyemo MA, Spiegel BM, Chang L. Meta-analysis: do irritable bowel syndrome symptoms vary between men and women? *Aliment Pharmacol Ther* 2010; **32**: 738-755 [PMID: 20662786 DOI: 10.1111/j.1365-2036.2010.04409.x]
- Heitkemper MM, Jarrett ME. Update on irritable bowel syndrome and gender differences. *Nutr Clin Pract* 2008; **23**: 275-283 [PMID: 18595860 DOI: 10.1177/0884533608318672]
- Chang L, Toner BB, Fukudo S, Guthrie E, Locke GR, Norton NJ, Sperber AD. Gender, age, society, culture, and the patient's perspective in the functional gastrointestinal disorders. *Gastroenterology* 2006; **130**: 1435-1446 [PMID: 16678557]
- Meier R, Beglinger C, Dederding JP, Meyer-Wyss B, Fumagalli M, Rowedder A, Turberg Y, Brignoli R. Influence of age, gender, hormonal status and smoking habits on colonic transit time. *Neurogastroenterol Motil* 1995; **7**: 235-238 [PMID: 8574912]
- Lampe JW, Fredstrom SB, Slavin JL, Potter JD. Sex differences in colonic function: a randomised trial. *Gut* 1993; **34**: 531-536 [PMID: 8387940]
- Tang YR, Yang WW, Wang YL, Lin L. Sex differences in the symptoms and psychological factors that influence quality of life in patients with irritable bowel syndrome. *Eur J Gastroenterol Hepatol* 2012; **24**: 702-707 [PMID: 22382707 DOI: 10.1097/MEG.0b013e328351b2c2]
- Faresjö A, Johansson S, Faresjö T, Roos S, Hallert C. Sex differences in dietary coping with gastrointestinal symptoms. *Eur J Gastroenterol Hepatol* 2010; **22**: 327-333 [PMID: 19550348 DOI: 10.1097/MEG.0b013e32832b9c53]
- Cain KC, Jarrett ME, Burr RL, Rosen S, Hertig VL, Heitkemper MM. Gender differences in gastrointestinal, psychological, and somatic symptoms in irritable bowel syndrome. *Dig Dis Sci* 2009; **54**: 1542-1549 [PMID: 18979200]
- Mulak A, Waszczuk E, Paradowski L. Irritable bowel syndrome as an interdisciplinary clinical problem. *Adv Clin Exp Med* 2008; **17**: 667-675
- Warnock JK, Clayton AH. Chronic episodic disorders in women. *Psychiatr Clin North Am* 2003; **26**: 725-740 [PMID: 14563106]
- Heitkemper M, Jarrett M. Irritable bowel syndrome: does gender matter? *J Psychosom Res* 2008; **64**: 583-587 [PMID: 18501258 DOI: 10.1016/j.jpsychores.2008.02.020]
- Palsson OS, Drossman DA. Psychiatric and psychological dysfunction in irritable bowel syndrome and the role of psychological treatments. *Gastroenterol Clin North Am* 2005; **34**: 281-303 [PMID: 15862936]
- Surdea-Blaga T, Băban A, Dumitrascu DL. Psychosocial determinants of irritable bowel syndrome. *World J Gastroenterol* 2012; **18**: 616-626 [PMID: 22363132 DOI: 10.3748/wjg.v18.i7.616]
- García Rodríguez LA, Ruigómez A, Wallander MA, Johans-

- son S, Olbe L. Detection of colorectal tumor and inflammatory bowel disease during follow-up of patients with initial diagnosis of irritable bowel syndrome. *Scand J Gastroenterol* 2000; **35**: 306-311 [PMID: 10766326]
- 33 **Farage MA**, Neill S, MacLean AB. Physiological changes associated with the menstrual cycle: a review. *Obstet Gynecol Surv* 2009; **64**: 58-72 [PMID: 19099613 DOI: 10.1097/OGX.0b013e3181932a37]
- 34 **Moore J**, Barlow D, Jewell D, Kennedy S. Do gastrointestinal symptoms vary with the menstrual cycle? *Br J Obstet Gynaecol* 1998; **105**: 1322-1325 [PMID: 9883927]
- 35 **Chang L**, Lee OY, Naliboff B, Schmulson M, Mayer EA. Sensation of bloating and visible abdominal distension in patients with irritable bowel syndrome. *Am J Gastroenterol* 2001; **96**: 3341-3347 [PMID: 11774947]
- 36 **Mulak A**, Taché Y. Sex difference in irritable bowel syndrome: do gonadal hormones play a role? *Gastroenterol Pol* 2010; **17**: 89-97
- 37 **Whitehead WE**, Cheskin LJ, Heller BR, Robinson JC, Crowell MD, Benjamin C, Schuster MM. Evidence for exacerbation of irritable bowel syndrome during menses. *Gastroenterology* 1990; **98**: 1485-1489 [PMID: 2338190]
- 38 **Houghton LA**, Lea R, Jackson N, Whorwell PJ. The menstrual cycle affects rectal sensitivity in patients with irritable bowel syndrome but not healthy volunteers. *Gut* 2002; **50**: 471-474 [PMID: 11889064]
- 39 **Gustafsson JK**, Greenwood-Van Meerveld B. Amygdala activation by corticosterone alters visceral and somatic pain in cycling female rats. *Am J Physiol Gastrointest Liver Physiol* 2011; **300**: G1080-G1085 [PMID: 21454447 DOI: 10.1152/ajpgi.00349.2010]
- 40 **Ouyang A**, Wrzos HF. Contribution of gender to pathophysiology and clinical presentation of IBS: should management be different in women? *Am J Gastroenterol* 2006; **101**: S602-S609 [PMID: 17177863]
- 41 **Kane SV**, Sable K, Hanauer SB. The menstrual cycle and its effect on inflammatory bowel disease and irritable bowel syndrome: a prevalence study. *Am J Gastroenterol* 1998; **93**: 1867-1872 [PMID: 9772046]
- 42 **Altman G**, Cain KC, Motzer S, Jarrett M, Burr R, Heitkemper M. Increased symptoms in female IBS patients with dysmenorrhea and PMS. *Gastroenterol Nurs* 2006; **29**: 4-11 [PMID: 16552294]
- 43 **Olafsdottir LB**, Gudjonsson H, Jonsdottir HH, Björnsson E, Thjodleifsson B. Natural history of irritable bowel syndrome in women and dysmenorrhea: a 10-year follow-up study. *Gastroenterol Res Pract* 2012; **2012**: 534204 [PMID: 22474441 DOI: 10.1155/2012/534204]
- 44 **Meurs-Szojda MM**, Mijatovic V, Felt-Bersma RJ, Hompes PG. Irritable bowel syndrome and chronic constipation in patients with endometriosis. *Colorectal Dis* 2011; **13**: 67-71 [PMID: 19832874 DOI: 10.1111/j.1463-1318.2009.02055.x]
- 45 **Mathur R**, Ko A, Hwang LJ, Low K, Azziz R, Pimentel M. Polycystic ovary syndrome is associated with an increased prevalence of irritable bowel syndrome. *Dig Dis Sci* 2010; **55**: 1085-1089 [PMID: 19697132 DOI: 10.1007/s10620-009-0890-5]
- 46 **Karagiannides I**, Pothoulakis C. Neuropeptides, mesenteric fat, and intestinal inflammation. *Ann N Y Acad Sci* 2008; **1144**: 127-135 [PMID: 19076372 DOI: 10.1196/annals.1418.009]
- 47 **Sverrisdóttir YB**, Mogren T, Kataoka J, Janson PO, Stener-Victorin E. Is polycystic ovary syndrome associated with high sympathetic nerve activity and size at birth? *Am J Physiol Endocrinol Metab* 2008; **294**: E576-E581 [PMID: 18198350 DOI: 10.1152/ajpendo.00725.2007]
- 48 **Grenham S**, Clarke G, Cryan JF, Dinan TG. Brain-gut-microbe communication in health and disease. *Front Physiol* 2011; **2**: 94 [PMID: 22162969 DOI: 10.3389/fphys.2011.00094]
- 49 **Cogan R**, Spinnato JA. Pain and discomfort thresholds in late pregnancy. *Pain* 1986; **27**: 63-68 [PMID: 3785964]
- 50 **Gintzler AR**. Endorphin-mediated increases in pain threshold during pregnancy. *Science* 1980; **210**: 193-195 [PMID: 7414330]
- 51 **Chiloiro M**, Darconza G, Piccioli E, De Carne M, Clemente C, Riezzo G. Gastric emptying and orocecal transit time in pregnancy. *J Gastroenterol* 2001; **36**: 538-543 [PMID: 11519832]
- 52 **Jarrett M**, Heitkemper M, Cain KC, Tuftin M, Walker EA, Bond EF, Levy RL. The relationship between psychological distress and gastrointestinal symptoms in women with irritable bowel syndrome. *Nurs Res* 1998; **47**: 154-161 [PMID: 9610649]
- 53 **Triadafilopoulos G**, Finlayson M, Grellet C. Bowel dysfunction in postmenopausal women. *Women Health* 1998; **27**: 55-66 [PMID: 9796084]
- 54 **Gonenne J**, Esfandyari T, Camilleri M, Burton DD, Stephens DA, Baxter KL, Zinsmeister AR, Bharucha AE. Effect of female sex hormone supplementation and withdrawal on gastrointestinal and colonic transit in postmenopausal women. *Neurogastroenterol Motil* 2006; **18**: 911-918 [PMID: 16961694]
- 55 **Heitkemper MM**, Cain KC, Jarrett ME, Burr RL, Hertig V, Bond EF. Symptoms across the menstrual cycle in women with irritable bowel syndrome. *Am J Gastroenterol* 2003; **98**: 420-430 [PMID: 12591063]
- 56 **Bird ST**, Liu W, Brophy JM, Bressler B, Delaney JA, Etminan M. Irritable bowel syndrome and drospirenone-containing oral contraceptives; a comparative-safety study. *Curr Drug Saf* 2012; **7**: 8-15 [PMID: 22663950]
- 57 **Palomba S**, Orio F, Manguso F, Russo T, Falbo A, Lombardi G, Doldo P, Zullo F. Leuprolide acetate treatment with and without coadministration of tibolone in premenopausal women with menstrual cycle-related irritable bowel syndrome. *Fertil Steril* 2005; **83**: 1012-1020 [PMID: 15820814]
- 58 **Ruigómez A**, García Rodríguez LA, Johansson S, Wallander MA. Is hormone replacement therapy associated with an increased risk of irritable bowel syndrome? *Maturitas* 2003; **44**: 133-140 [PMID: 12590009]
- 59 **Longstreth GF**, Yao JF. Irritable bowel syndrome and surgery: a multivariable analysis. *Gastroenterology* 2004; **126**: 1665-1673 [PMID: 15188159]
- 60 **Sperber AD**, Morris CB, Greemberg L, Bangdiwala SI, Goldstein D, Sheiner E, Rusabrov Y, Hu Y, Katz M, Freud T, Neville A, Drossman DA. Development of abdominal pain and IBS following gynecological surgery: a prospective, controlled study. *Gastroenterology* 2008; **134**: 75-84 [PMID: 18166349 DOI: 10.1053/j.gastro.2007.10.041]
- 61 **Sanoja R**, Cervero F. Estrogen-dependent abdominal hyperalgesia induced by ovariectomy in adult mice: a model of functional abdominal pain. *Pain* 2005; **118**: 243-253 [PMID: 16202534]
- 62 **Sanoja R**, Cervero F. Estrogen modulation of ovariectomy-induced hyperalgesia in adult mice. *Eur J Pain* 2008; **12**: 573-581 [PMID: 17959401]
- 63 **Ji Y**, Murphy AZ, Traub RJ. Estrogen modulates the visceromotor reflex and responses of spinal dorsal horn neurons to colorectal stimulation in the rat. *J Neurosci* 2003; **23**: 3908-3915 [PMID: 12736360]
- 64 **Bradesi S**, Eutamene H, Garcia-Villar R, Fioramonti J, Bueno L. Stress-induced visceral hypersensitivity in female rats is estrogen-dependent and involves tachykinin NK1 receptors. *Pain* 2003; **102**: 227-234 [PMID: 12670663]
- 65 **Taché Y**, Million M, Nelson AG, Lamy C, Wang L. Role of corticotropin-releasing factor pathways in stress-related alterations of colonic motor function and viscerosensitivity in female rodents. *Genet Med* 2005; **2**: 146-154 [PMID: 16290887]
- 66 **Houghton LA**, Jackson NA, Whorwell PJ, Morris J. Do male sex hormones protect from irritable bowel syndrome? *Am J Gastroenterol* 2000; **95**: 2296-2300 [PMID: 11007231]
- 67 **Aloisi AM**. Gonadal hormones and sex differences in pain

- reactivity. *Clin J Pain* 2003; **19**: 168-174 [PMID: 12792555]
- 68 Cairns BE, Gazerani P. Sex-related differences in pain. *Maturnitas* 2009; **63**: 292-296 [PMID: 19595525]
- 69 Vodo S, Bechi N, Petroni A, Muscoli C, Aloisi AM. Testosterone-induced effects on lipids and inflammation. *Mediators Inflamm* 2013; **2013**: 183041 [PMID: 23606790 DOI: 10.1155/2013/183041]
- 70 Kim BJ, Rhee PL, Park JH, Chang DK, Kim YH, Son HJ, Kim JJ, Rhee JC, Lee H. Male sex hormones may influence the symptoms of irritable bowel syndrome in young men. *Digestion* 2008; **78**: 88-92 [PMID: 18974649 DOI: 10.1159/000166600]
- 71 Miller V, Whitaker K, Morris JA, Whorwell PJ. Gender and irritable bowel syndrome: the male connection. *J Clin Gastroenterol* 2004; **38**: 558-560 [PMID: 15232357]
- 72 Aloisi AM, Bachicco V, Costantino A, Stefani R, Ceccarelli I, Bertaccini A, Meriggola MC. Cross-sex hormone administration changes pain in transsexual women and men. *Pain* 2007; **132** Suppl 1: S60-S67 [PMID: 17379410]
- 73 Ter Horst GJ, Wichmann R, Gerrits M, Westenbroek C, Lin Y. Sex differences in stress responses: focus on ovarian hormones. *Physiol Behav* 2009; **97**: 239-249 [PMID: 19275910 DOI: 10.1016/j.physbeh.2009.02.036]
- 74 Sinchak K, Wagner EJ. Estradiol signaling in the regulation of reproduction and energy balance. *Front Neuroendocrinol* 2012; **33**: 342-363 [PMID: 22981653 DOI: 10.1016/j.yfrne.2012.08.004]
- 75 Gintzler AR, Liu NJ. Importance of sex to pain and its amelioration; relevance of spinal estrogens and its membrane receptors. *Front Neuroendocrinol* 2012; **33**: 412-424 [PMID: 23036438 DOI: 10.1016/j.yfrne.2012.09.004]
- 76 Wierman ME. Sex steroid effects at target tissues: mechanisms of action. *Adv Physiol Educ* 2007; **31**: 26-33 [PMID: 17327579]
- 77 Pfaffl MW, Lange IG, Meyer HH. The gastrointestinal tract as target of steroid hormone action: quantification of steroid receptor mRNA expression (AR, ERalpha, ERbeta and PR) in 10 bovine gastrointestinal tract compartments by kinetic RT-PCR. *J Steroid Biochem Mol Biol* 2003; **84**: 159-166 [PMID: 12710999]
- 78 Naliboff BD, Berman S, Chang L, Derbyshire SW, Suyenobu B, Vogt BA, Mandelkern M, Mayer EA. Sex-related differences in IBS patients: central processing of visceral stimuli. *Gastroenterology* 2003; **124**: 1738-1747 [PMID: 12806606]
- 79 Labus JS, Gupta A, Coveleskie K, Tillisch K, Kilpatrick L, Jarcho J, Feier N, Bueller J, Stains J, Smith S, Suyenobu B, Naliboff B, Mayer EA. Sex differences in emotion-related cognitive processes in irritable bowel syndrome and healthy control subjects. *Pain* 2013; **154**: 2088-2099 [PMID: 23791896 DOI: 10.1016/j.pain.2013.06.024]
- 80 McEwen BS, Akama KT, Spencer-Segal JL, Milner TA, Waters EM. Estrogen effects on the brain: actions beyond the hypothalamus via novel mechanisms. *Behav Neurosci* 2012; **126**: 4-16 [PMID: 22289042 DOI: 10.1037/a0026708]
- 81 Kelly MJ, Rønnekleiv OK. A selective membrane estrogen receptor agonist maintains autonomic functions in hypoestrogenic states. *Brain Res* 2013; **1514**: 75-82 [PMID: 23535448 DOI: 10.1016/j.brainres.2013.03.020]
- 82 Scharfman HE, MacLusky NJ. Differential regulation of BDNF, synaptic plasticity and sprouting in the hippocampal mossy fiber pathway of male and female rats. *Neuropharmacology* 2014; **76** Pt C: 696-708 [PMID: 23660230]
- 83 Chen XN, Zhu H, Meng QY, Zhou JN. Estrogen receptor-alpha and -beta regulate the human corticotropin-releasing hormone gene through similar pathways. *Brain Res* 2008; **1223**: 1-10 [PMID: 18597742 DOI: 10.1016/j.brainres.2008.05.043]
- 84 Kajantie E, Phillips DI. The effects of sex and hormonal status on the physiological response to acute psychosocial stress. *Psychoneuroendocrinology* 2006; **31**: 151-178 [PMID: 16139959]
- 85 Larauche M, Gourcerol G, Wang L, Pambukchian K, Brunnhuber S, Adelson DW, Rivier J, Million M, Taché Y. Cortagine, a CRF1 agonist, induces stresslike alterations of colonic function and visceral hypersensitivity in rodents primarily through peripheral pathways. *Am J Physiol Gastrointest Liver Physiol* 2009; **297**: G215-G227 [PMID: 19407218 DOI: 10.1152/ajpgi.00072.2009]
- 86 Yuan PQ, Wu SV, Elliott J, Anton PA, Chatzaki E, Million M, Taché Y. Expression of corticotropin releasing factor receptor type 1 (CRF1) in the human gastrointestinal tract and upregulation in the colonic mucosa in patients with ulcerative colitis. *Peptides* 2012; **38**: 62-69 [PMID: 22948128 DOI: 10.1016/j.peptides.2012.07.028]
- 87 Bao AM, Hestiantoro A, Van Someren EJ, Swaab DF, Zhou JN. Colocalization of corticotropin-releasing hormone and oestrogen receptor-alpha in the paraventricular nucleus of the hypothalamus in mood disorders. *Brain* 2005; **128**: 1301-1313 [PMID: 15705605]
- 88 Peiffer A, Lapointe B, Barden N. Hormonal regulation of type II glucocorticoid receptor messenger ribonucleic acid in rat brain. *Endocrinology* 1991; **129**: 2166-2174 [PMID: 1915097]
- 89 Goldstein JM, Jerram M, Abbs B, Whitfield-Gabrieli S, Makris N. Sex differences in stress response circuitry activation dependent on female hormonal cycle. *J Neurosci* 2010; **30**: 431-438 [PMID: 20071507 DOI: 10.1523/JNEUROSCI.3021-09.2010]
- 90 Shansky RM, Morrison JH. Stress-induced dendritic remodeling in the medial prefrontal cortex: effects of circuit, hormones and rest. *Brain Res* 2009; **1293**: 108-113 [PMID: 19361488 DOI: 10.1016/j.brainres.2009.03.062]
- 91 Myers B, Schulkin J, Greenwood-Van Meerveld B. Sex steroids localized to the amygdala increase pain responses to visceral stimulation in rats. *J Pain* 2011; **12**: 486-494 [PMID: 21167789 DOI: 10.1016/j.jpain.2010.10.007]
- 92 Tillisch K, Mayer EA, Labus JS. Quantitative meta-analysis identifies brain regions activated during rectal distension in irritable bowel syndrome. *Gastroenterology* 2011; **140**: 91-100 [PMID: 20696168 DOI: 10.1053/j.gastro.2010.07.053]
- 93 Videlock EJ, Adeyemo M, Licudine A, Hirano M, Ohning G, Mayer M, Mayer EA, Chang L. Childhood trauma is associated with hypothalamic-pituitary-adrenal axis responsiveness in irritable bowel syndrome. *Gastroenterology* 2009; **137**: 1954-1962 [PMID: 19737564 DOI: 10.1053/j.gastro.2009.08.058]
- 94 Chang L, Sundaresh S, Elliott J, Anton PA, Baldi P, Licudine A, Mayer M, Vuong T, Hirano M, Naliboff BD, Ameen VZ, Mayer EA. Dysregulation of the hypothalamic-pituitary-adrenal (HPA) axis in irritable bowel syndrome. *Neurogastroenterol Motil* 2009; **21**: 149-159 [PMID: 18684212 DOI: 10.1111/j.1365-2982.2008.01171.x]
- 95 Chaloner A, Greenwood-Van Meerveld B. Sexually dimorphic effects of unpredictable early life adversity on visceral pain behavior in a rodent model. *J Pain* 2013; **14**: 270-280 [PMID: 23348370 DOI: 10.1016/j.jpain.2012.11.008]
- 96 Gupta S, McCarron KE, Welch KM, Berman NE. Mechanisms of pain modulation by sex hormones in migraine. *Headache* 2011; **51**: 905-922 [PMID: 21631476 DOI: 10.1111/j.1526-4610.2011.01908.x]
- 97 Linthorst AC. Interactions between corticotropin-releasing hormone and serotonin: implications for the aetiology and treatment of anxiety disorders. *Handb Exp Pharmacol* 2005; **(169)**: 181-204 [PMID: 16594259]
- 98 Niesler B, Kapeller J, Fell C, Atkinson W, Möller D, Fischer C, Whorwell P, Houghton LA. 5-HTTLPR and STin2 polymorphisms in the serotonin transporter gene and irritable bowel syndrome: effect of bowel habit and sex. *Eur J Gas-*

- troenterol Hepatol* 2010; **22**: 856-861 [PMID: 19561511 DOI: 10.1097/MEG.0b013e32832e9d6b]
- 99 **Mizuno T**, Aoki M, Shimada Y, Inoue M, Nakaya K, Takahashi T, Itoyama Y, Kanazawa M, Utsumi A, Endo Y, Nomura T, Hiratsuka M, Mizugaki M, Goto J, Hongo M, Fukudo S. Gender difference in association between polymorphism of serotonin transporter gene regulatory region and anxiety. *J Psychosom Res* 2006; **60**: 91-97 [PMID: 16380315]
- 100 **Ji Y**, Tang B, Traub RJ. Modulatory effects of estrogen and progesterone on colorectal hyperalgesia in the rat. *Pain* 2005; **117**: 433-442 [PMID: 16154701]
- 101 **Wang M**. Neurosteroids and GABA-A Receptor Function. *Front Endocrinol (Lausanne)* 2011; **2**: 44 [PMID: 22654809 DOI: 10.3389/fendo.2011.00044]
- 102 **Pasquali R**. The hypothalamic-pituitary-adrenal axis and sex hormones in chronic stress and obesity: pathophysiological and clinical aspects. *Ann N Y Acad Sci* 2012; **1264**: 20-35 [PMID: 22612409 DOI: 10.1111/j.1749-6632.2012.06569.x]
- 103 **Viau V**, Meaney MJ. The inhibitory effect of testosterone on hypothalamic-pituitary-adrenal responses to stress is mediated by the medial preoptic area. *J Neurosci* 1996; **16**: 1866-1876 [PMID: 8774455]
- 104 **Chaban V**. Estrogen and Visceral Nociception at the Level of Primary Sensory Neurons. *Pain Res Treat* 2012; **2012**: [PMID: 21984961]
- 105 **Amandusson A**, Hermanson O, Blomqvist A. Colocalization of oestrogen receptor immunoreactivity and preproenkephalin mRNA expression to neurons in the superficial laminae of the spinal and medullary dorsal horn of rats. *Eur J Neurosci* 1996; **8**: 2440-2445 [PMID: 8950107]
- 106 **Peng HY**, Chen GD, Lai CY, Hsieh MC, Hsu HH, Wu HC, Lin TB. PI3K modulates estrogen-dependent facilitation of colon-to-urethra cross-organ reflex sensitization in ovariectomized female rats. *J Neurochem* 2010; **113**: 54-66 [PMID: 20067569 DOI: 10.1111/j.1471-4159.2010.06577.x]
- 107 **Cheng P**, Shih W, Alberto M, Presson AP, Licudine A, Mayer EA, Naliboff BD, Chang L. Autonomic response to a visceral stressor is dysregulated in irritable bowel syndrome and correlates with duration of disease. *Neurogastroenterol Motil* 2013; **25**: e650-e659 [PMID: 23822743 DOI: 10.1111/nmo.12177]
- 108 **Tillisch K**, Mayer EA, Labus JS, Stains J, Chang L, Naliboff BD. Sex specific alterations in autonomic function among patients with irritable bowel syndrome. *Gut* 2005; **54**: 1396-1401 [PMID: 15923667]
- 109 **Matsumoto T**, Ushiroyama T, Morimura M, Moritani T, Hayashi T, Suzuki T, Tatsumi N. Autonomic nervous system activity in the late luteal phase of eumenorrhic women with premenstrual symptomatology. *J Psychosom Obstet Gynaecol* 2006; **27**: 131-139 [PMID: 17214447]
- 110 **Rydlewska A**, Maj J, Katkowski B, Biel B, Ponikowska B, Banasiak W, Ponikowski P, Jankowska EA. Circulating testosterone and estradiol, autonomic balance and baroreflex sensitivity in middle-aged and elderly men with heart failure. *Aging Male* 2013; **16**: 58-66 [PMID: 23581721 DOI: 10.3109/13685538.2013.768979]
- 111 **Spiller R**, Garsed K. Postinfectious irritable bowel syndrome. *Gastroenterology* 2009; **136**: 1979-1988 [PMID: 19457422 DOI: 10.1053/j.gastro.2009.02.074]
- 112 **Ong TS**. Biomechanical properties of the Ong capsular lens and results of four years of clinical experience. *J Am Intra-ocul Implant Soc* 1984; **10**: 44-48 [PMID: 6706815]
- 113 **Straub RH**. The complex role of estrogens in inflammation. *Endocr Rev* 2007; **28**: 521-574 [PMID: 17640948]
- 114 **Gué M**, Del Rio-Lacheze C, Eutamene H, Théodorou V, Fioramonti J, Bueno L. Stress-induced visceral hypersensitivity to rectal distension in rats: role of CRF and mast cells. *Neurogastroenterol Motil* 1997; **9**: 271-279 [PMID: 9430796]
- 115 **Larauche M**. Novel insights in the role of peripheral corticotropin-releasing factor and mast cells in stress-induced visceral hypersensitivity. *Neurogastroenterol Motil* 2012; **24**: 201-205 [PMID: 22316289 DOI: 10.1111/j.1365-2982.2011.01867.x]
- 116 **Zierau O**, Zenclussen AC, Jensen F. Role of female sex hormones, estradiol and progesterone, in mast cell behavior. *Front Immunol* 2012; **3**: 169 [PMID: 22723800 DOI: 10.3389/fimmu.2012.00169]
- 117 **Barbara G**, Stanghellini V, De Giorgio R, Corinaldesi R. Functional gastrointestinal disorders and mast cells: implications for therapy. *Neurogastroenterol Motil* 2006; **18**: 6-17 [PMID: 16371078]
- 118 **Vliagoftis H**, Dimitriadou V, Boucher W, Rozniecki JJ, Correia I, Raam S, Theoharides TC. Estradiol augments while tamoxifen inhibits rat mast cell secretion. *Int Arch Allergy Immunol* 1992; **98**: 398-409 [PMID: 1384869]
- 119 **Houghton LA**, Brown H, Atkinson W, Morris J, Fell C, Whorwell PJ, Lockhart S, Keevil B. 5-hydroxytryptamine signalling in irritable bowel syndrome with diarrhoea: effects of gender and menstrual status. *Aliment Pharmacol Ther* 2009; **30**: 919-929 [PMID: 19691669 DOI: 10.1111/j.1365-2036.2009.04121.x]
- 120 **Li TJ**, Yu BP, Dong WG, Luo HS, Xu L, Li MQ. Ovarian hormone modulates 5-hydroxytryptamine 3 receptors mRNA expression in rat colon with restraint stress-induced bowel dysfunction. *World J Gastroenterol* 2004; **10**: 2723-2726 [PMID: 15309727]
- 121 **Galligan JJ**, Patel BA, Schneider SP, Wang H, Zhao H, Novotny M, Bian X, Kabeer R, Fried D, Swain GM. Visceral hypersensitivity in female but not in male serotonin transporter knockout rats. *Neurogastroenterol Motil* 2013; **25**: e373-e381 [PMID: 23594365 DOI: 10.1111/nmo.12133]
- 122 **Franke L**, Schmidtman M, Riedl A, van der Voort I, Uebelhack R, Mönnikes H. Serotonin transporter activity and serotonin concentration in platelets of patients with irritable bowel syndrome: effect of gender. *J Gastroenterol* 2010; **45**: 389-398 [PMID: 19997855 DOI: 10.1007/s00535-009-0167-y]
- 123 **Piche T**, Barbara G, Aubert P, Bruley des Varannes S, Dainese R, Nano JL, Cremon C, Stanghellini V, De Giorgio R, Galmiche JP, Neunlist M. Impaired intestinal barrier integrity in the colon of patients with irritable bowel syndrome: involvement of soluble mediators. *Gut* 2009; **58**: 196-201 [PMID: 18824556 DOI: 10.1136/gut.2007.140806]
- 124 **Braniste V**, Leveque M, Buisson-Brenac C, Bueno L, Fioramonti J, Houdeau E. Oestradiol decreases colonic permeability through oestrogen receptor beta-mediated up-regulation of occludin and junctional adhesion molecule-A in epithelial cells. *J Physiol* 2009; **587**: 3317-3328 [PMID: 19433574 DOI: 10.1113/jphysiol.2009.169300]
- 125 **Houdeau E**, Moriez R, Leveque M, Salvador-Cartier C, Waget A, Leng L, Bueno L, Bucala R, Fioramonti J. Sex steroid regulation of macrophage migration inhibitory factor in normal and inflamed colon in the female rat. *Gastroenterology* 2007; **132**: 982-993 [PMID: 17324399]
- 126 **Verdú EF**, Deng Y, Bercik P, Collins SM. Modulatory effects of estrogen in two murine models of experimental colitis. *Am J Physiol Gastrointest Liver Physiol* 2002; **283**: G27-G36 [PMID: 12065288]
- 127 **Alonso C**, Guilarte M, Vicario M, Ramos L, Rezzi S, Martínez C, Lobo B, Martín FP, Pigrau M, González-Castro AM, Gallart M, Malagelada JR, Azpiroz F, Kochhar S, Santos J. Acute experimental stress evokes a differential gender-determined increase in human intestinal macromolecular permeability. *Neurogastroenterol Motil* 2012; **24**: 740-746; e348-e349 [PMID: 22625665 DOI: 10.1111/j.1365-2982.2012.01928.x]
- 128 **Levin ER**. Membrane oestrogen receptor alpha signalling to cell functions. *J Physiol* 2009; **587**: 5019-5023 [PMID: 19687123 DOI: 10.1113/jphysiol.2009.177097]

- 129 **Kawano N**, Koji T, Hishikawa Y, Murase K, Murata I, Kohno S. Identification and localization of estrogen receptor alpha- and beta-positive cells in adult male and female mouse intestine at various estrogen levels. *Histochem Cell Biol* 2004; **121**: 399-405 [PMID: 15138841]
- 130 **Cong P**, Pricolo V, Biancani P, Behar J. Abnormalities of prostaglandins and cyclooxygenase enzymes in female patients with slow-transit constipation. *Gastroenterology* 2007; **133**: 445-453 [PMID: 17681165]
- 131 **Vasiadi M**, Kempuraj D, Boucher W, Kalogeromitros D, Theoharides TC. Progesterone inhibits mast cell secretion. *Int J Immunopathol Pharmacol* 2006; **19**: 787-794 [PMID: 17166400]
- 132 **García-Gómez E**, González-Pedrajo B, Camacho-Arroyo I. Role of sex steroid hormones in bacterial-host interactions. *Biomed Res Int* 2013; **2013**: 928290 [PMID: 23509808 DOI: 10.1155/2013/928290]
- 133 **Su L**, Sun Y, Ma F, Lü P, Huang H, Zhou J. Progesterone inhibits Toll-like receptor 4-mediated innate immune response in macrophages by suppressing NF-kappaB activation and enhancing SOCS1 expression. *Immunol Lett* 2009; **125**: 151-155 [PMID: 19607861 DOI: 10.1016/j.imlet.2009.07.003]
- 134 **González-Montelongo MC**, Marín R, Gómez T, Marrero-Alonso J, Díaz M. Androgens induce nongenomic stimulation of colonic contractile activity through induction of calcium sensitization and phosphorylation of LC20 and CPI-17. *Mol Endocrinol* 2010; **24**: 1007-1023 [PMID: 20207835 DOI: 10.1210/me.2009-0472]
- 135 **Aloisi AM**, Affaitati G, Ceccarelli I, Fiorenzani P, Lerza R, Rossi C, Pace MC, Chiefari M, Aurilio C, Giamberardino MA. Estradiol and testosterone differently affect visceral pain-related behavioural responses in male and female rats. *Eur J Pain* 2010; **14**: 602-607 [PMID: 19948419 DOI: 10.1016/j.ejpain.2009.10.016]
- 136 **Sommer C**, Kress M. Recent findings on how proinflammatory cytokines cause pain: peripheral mechanisms in inflammatory and neuropathic hyperalgesia. *Neurosci Lett* 2004; **361**: 184-187 [PMID: 15135924]
- 137 **Chen W**, Beck I, Schober W, Brockow K, Effner R, Buters JT, Behrendt H, Ring J. Human mast cells express androgen receptors but treatment with testosterone exerts no influence on IgE-independent mast cell degranulation elicited by neuromuscular blocking agents. *Exp Dermatol* 2010; **19**: 302-304 [PMID: 19758318 DOI: 10.1111/j.1600-0625.2009.00969.x]
- 138 **Rettew JA**, Huet-Hudson YM, Marriott I. Testosterone reduces macrophage expression in the mouse of toll-like receptor 4, a trigger for inflammation and innate immunity. *Biol Reprod* 2008; **78**: 432-437 [PMID: 18003947]
- 139 **Dinan TG**, Cryan JF. Regulation of the stress response by the gut microbiota: implications for psychoneuroendocrinology. *Psychoneuroendocrinology* 2012; **37**: 1369-1378 [PMID: 22483040 DOI: 10.1016/j.psyneuen.2012.03.007]
- 140 **Dinan TG**, Cryan JF. Melancholic microbes: a link between gut microbiota and depression? *Neurogastroenterol Motil* 2013; **25**: 713-719 [PMID: 23910373 DOI: 10.1111/nmo.12198]
- 141 **Clarke G**, Grenham S, Scully P, Fitzgerald P, Moloney RD, Shanahan F, Dinan TG, Cryan JF. The microbiome-gut-brain axis during early life regulates the hippocampal serotonergic system in a sex-dependent manner. *Mol Psychiatry* 2013; **18**: 666-673 [PMID: 22688187 DOI: 10.1038/mp.2012.77]
- 142 **Kornman KS**, Loesch WJ. Effects of estradiol and progesterone on *Bacteroides melanogenicus* and *Bacteroides gingivalis*. *Infect Immun* 1982; **35**: 256-263 [PMID: 6119293]
- 143 **Menon R**, Watson SE, Thomas LN, Allred CD, Dabney A, Azcarate-Peril MA, Sturino JM. Diet complexity and estrogen receptor β status affect the composition of the murine intestinal microbiota. *Appl Environ Microbiol* 2013; **79**: 5763-5773 [PMID: 23872567 DOI: 10.1128/AEM.01182-13]
- 144 **Adlercreutz H**, Pulkkinen MO, Hämäläinen EK, Korpela JT. Studies on the role of intestinal bacteria in metabolism of synthetic and natural steroid hormones. *J Steroid Biochem* 1984; **20**: 217-229 [PMID: 6231418]
- 145 **Raftogianis R**, Creveling C, Weinsilboum R, Weisz J. Estrogen metabolism by conjugation. *J Natl Cancer Inst Monogr* 2000; (27): 113-124 [PMID: 10963623]
- 146 **Flores R**, Shi J, Fuhrman B, Xu X, Veenstra TD, Gail MH, Gajer P, Ravel J, Goedert JJ. Fecal microbial determinants of fecal and systemic estrogens and estrogen metabolites: a cross-sectional study. *J Transl Med* 2012; **10**: 253 [PMID: 23259758 DOI: 10.1186/1479-5876-10-253]
- 147 **Lombardi P**, Goldin B, Boutin E, Gorbach SL. Metabolism of androgens and estrogens by human fecal microorganisms. *J Steroid Biochem* 1978; **9**: 795-801 [PMID: 713557]
- 148 **Donova MV**, Egorova OV. Microbial steroid transformations: current state and prospects. *Appl Microbiol Biotechnol* 2012; **94**: 1423-1447 [PMID: 22562163 DOI: 10.1007/s00253-012-4078-0]
- 149 **Markle JG**, Frank DN, Mortin-Toth S, Robertson CE, Feazel LM, Rolle-Kampczyk U, von Bergen M, McCoy KD, Macpherson AJ, Danska JS. Sex differences in the gut microbiome drive hormone-dependent regulation of autoimmunity. *Science* 2013; **339**: 1084-1088 [PMID: 23328391 DOI: 10.1126/science.1233521]
- 150 **Flak MB**, Neves JF, Blumberg RS. Immunology. Welcome to the microgenderome. *Science* 2013; **339**: 1044-1045 [PMID: 23449586 DOI: 10.1126/science.1236226]
- 151 **Koch KM**, Palmer JL, Noordin N, Tomlinson JJ, Baidoo C. Sex and age differences in the pharmacokinetics of alosetron. *Br J Clin Pharmacol* 2002; **53**: 238-242 [PMID: 11874386]
- 152 **Voß U**, Lewerenz A, Nieber K. Treatment of irritable bowel syndrome: sex and gender specific aspects. *Handb Exp Pharmacol* 2012; (214): 473-497 [PMID: 23027463 DOI: 10.1007/978-3-642-30726-3_21]
- 153 **Nakai A**, Diksic M, Kumakura Y, D'Souza D, Kersey K. The effects of the 5-HT₃ antagonist, alosetron, on brain serotonin synthesis in patients with irritable bowel syndrome. *Neurogastroenterol Motil* 2005; **17**: 212-221 [PMID: 15787942]
- 154 **Kano M**, Mizuno T, Kawano Y, Aoki M, Kanazawa M, Fukudo S. Serotonin transporter gene promoter polymorphism and alexithymia. *Neuropsychobiology* 2012; **65**: 76-82 [PMID: 22222552]
- 155 **Monostory K**, Dvorak Z. Steroid regulation of drug-metabolizing cytochromes P450. *Curr Drug Metab* 2011; **12**: 154-172 [PMID: 21395541]
- 156 **Creed F**, Fernandes L, Guthrie E, Palmer S, Ratcliffe J, Read N, Rigby C, Thompson D, Tomenson B. The cost-effectiveness of psychotherapy and paroxetine for severe irritable bowel syndrome. *Gastroenterology* 2003; **124**: 303-317 [PMID: 12557136]
- 157 **Whorwell PJ**, Prior A, Faragher EB. Controlled trial of hypnotherapy in the treatment of severe refractory irritable bowel syndrome. *Lancet* 1984; **2**: 1232-1234 [PMID: 6150275]
- 158 **Moser G**, Trägner S, Gajowniczek EE, Mikulits A, Michalski M, Kazemi-Shirazi L, Kulnigg-Dabsch S, Führer M, Ponocny-Seliger E, Dejaco C, Miehsler W. Long-term success of GUT-directed group hypnosis for patients with refractory irritable bowel syndrome: a randomized controlled trial. *Am J Gastroenterol* 2013; **108**: 602-609 [PMID: 23419384 DOI: 10.1038/ajg.2013.19]
- 159 **Brennan MJ**. The effect of opioid therapy on endocrine function. *Am J Med* 2013; **126**: S12-S18 [PMID: 23414717 DOI: 10.1016/j.amjmed.2012.12.001]
- 160 **Vodo S**, Arcelli D, Fiorenzani P, Meriggiola MC, Butkevich I, Di Canio C, Mikhailenko V, Aloisi AM. Gonadal ER α / β , AR and TRPV1 gene expression: modulation by pain and morphine treatment in male and female rats. *Physiol Behav* 2013; **110-111**: 80-86 [PMID: 23287630 DOI: 10.1016/j.physbeh.2012.12.014]

- 161 **Saulnier DM**, Ringel Y, Heyman MB, Foster JA, Bercik P, Shulman RJ, Versalovic J, Verdu EF, Dinan TG, Hecht G, Guarner F. The intestinal microbiome, probiotics and prebiotics in neurogastroenterology. *Gut Microbes* 2013; **4**: 17-27 [PMID: 23202796 DOI: 10.4161/gmic.22973]
- 162 **Mulak A**. Good bacteria for good mood: probiotics and the brain-gut axis interactions. *Pol Arch Med Wewn* 2013; **123**: 71-72 [PMID: 23449192]
- 163 **O'Mahony L**, McCarthy J, Kelly P, Hurley G, Luo F, Chen K, O'Sullivan GC, Kiely B, Collins JK, Shanahan F, Quigley EM. Lactobacillus and bifidobacterium in irritable bowel syndrome: symptom responses and relationship to cytokine profiles. *Gastroenterology* 2005; **128**: 541-551 [PMID: 15765388]

P- Reviewers: Abraham P, Poli-Neto OB **S- Editor:** Gou SX
L- Editor: A **E- Editor:** Wu HL





百世登

Baishideng®

Published by **Baishideng Publishing Group Co., Limited**

Flat C, 23/F., Lucky Plaza,

315-321 Lockhart Road, Wan Chai, Hong Kong, China

Fax: +852-65557188

Telephone: +852-31779906

E-mail: bpgoffice@wjgnet.com

<http://www.wjgnet.com>



ISSN 1007-9327



9 771007 932045