

UC Berkeley

UC Berkeley Previously Published Works

Title

Expansion of Signal Transduction Pathways in Fungi by Extensive Genome Duplication

Permalink

<https://escholarship.org/uc/item/5jw144n1>

Journal

Current Biology, 26(12)

ISSN

0960-9822

Authors

Corrochano, Luis M
Kuo, Alan
Marcet-Houben, Marina
[et al.](#)

Publication Date

2016-06-01

DOI

10.1016/j.cub.2016.04.038

Peer reviewed



Published in final edited form as:

Curr Biol. 2016 June 20; 26(12): 1577–1584. doi:10.1016/j.cub.2016.04.038.

Expansion of signal transduction pathways in fungi by extensive genome duplication

A full list of authors and affiliations appears at the end of the article.

Summary

Plants and fungi use light and other signals to regulate development, growth, and metabolism. The fruiting bodies of the fungus *Phycomyces blakesleeanus* are single cells that react to environmental cues, including light, but the mechanisms are largely unknown [1]. The related fungus *Mucor circinelloides* is an opportunistic human pathogen that changes its mode of growth upon receipt of signals from the environment to facilitate pathogenesis [2]. Understanding how these organisms respond to environmental cues should provide insights into the mechanisms of sensory perception and signal transduction by a single eukaryotic cell, and their role in pathogenesis. We sequenced the genomes of *P. blakesleeanus* and *M. circinelloides*, and show that they have been shaped by an extensive genome duplication or, most likely, a whole genome duplication (WGD), which is rarely observed in fungi [3-6]. We show that the genome duplication has expanded gene families, including those involved in signal transduction, and that duplicated genes have specialized, as evidenced by differences in their regulation by light. The transcriptional response to light varies with the developmental stage and is still observed in a photoreceptor mutant of *P. blakesleeanus*. A phototropic mutant of *P. blakesleeanus* with a heterozygous mutation in the photoreceptor gene *madA* demonstrates that photosensor dosage is important for the magnitude of signal transduction. We conclude that the genome duplication provided the means to improve signal transduction for enhanced perception of environmental signals. Our results will help to understand the role of genome dynamics in the evolution of sensory perception in eukaryotes.

Correspondence to: L.M. Corrochano. corrochano@us.es.

Author contributions

A.K., M.M.-H., S.P., A.S., J.M.V.-E. and J.G. contributed equally to this work. L.M.C. coordinated the project. E.L.B., S.B., S.T.-M. and L.M.C. conceived the project. I.V.G. directed all genome-sequencing and analysis efforts at JGI. J.R.-R., V.G.T., A.M.-D., C.S., M.S., E.M.L., F.S.-F. and V.G. provided DNA and RNA samples. A.K., A.S., E.L., J.-F.C., W.S., J.M., J.S., J.G., H.T. and D.B. sequenced, assembled and annotated the genomes. M.M.-H., A.S., T.G. and I.V.G. did genome comparisons and phylogenomics analysis. M.I.A., J.A., E.P.B., I.B., G.B., L.P.C., D.C., E.C.O., L.M.C., A.D., M.E., A.P.E., V.G., F.G., G.G., J.H., B.H., B.A.H., A.I., E.A.I., B.F.L., J.L.L., S.C.L., W.L., A.T.M., H.R.M., A.M., E.M.-T., J.A.O., R.A.O., M. Olmedo, M. Orejas, L.O.-C., A.G.P., J.R.-R., J.R.-H., R.R.-V., C.S., M.S., E.S., F.S.-F., D.S., K.S., V.G.T., N.J.T., M.R.T., S.T.-M., R.P.V., A.W. and J.S.Y. analyzed the genomes. J.M.V.-E., E.M.L., S.L.-G., F.S.-F., R.R.-V., S.T.-M., V.G., L.M.C. and A.H.-E performed the transcriptomic analysis. S.P., S.B., K.M. and A.I. performed the genetic characterization of *madI*. L.M.C., B.A.H., S.T.-M., A.I., A.H.-E., T.G. and I.V.G. wrote the paper with input from all authors.

Data deposition

Genome assemblies and annotations have been deposited in DDBJ/EMBL/GenBank under the accessions AMYC00000000 (*P. blakesleeanus*) and AMYB00000000 (*M. circinelloides*), and are accessible via the JGI Mycocosm portal (<http://jgi.doe.gov/fungi>). The mtDNAs have been deposited in DDBJ/EMBL/GenBank under the accessions KR809878 (*P. blakesleeanus*) and KR809877 (*M. circinelloides*). Phylomes can be accessed at phylomeDB (<http://phylomedb.org>) with phylomeIDs 252, 253, 254, 255 and 256. Gene expression data have been deposited in the GEO database with accessions GSE64369 (*P. blakesleeanus*) and GSE58264 (*M. circinelloides*).

Results and Discussion

Genome Duplications in the Evolution of the Mucoromycotina Fungi

Gene duplication has expanded the number of genes for photoreception in *P. blakesleeanus* and *M. circinelloides* (Figure 1A, 1B) [7-9], and we hypothesize that gene duplications and specialization may have provided new proteins to expand their sensory repertoire. We thus sequenced the 53.9 Megabase (Mb) *P. blakesleeanus* and the 36.6 Mb *M. circinelloides* genomes, and their respective mitochondrial DNAs (mtDNAs) (Supplemental Experimental Procedures; Table S1A-D; Figure S1). The increase in genome size in *P. blakesleeanus* is, in part, due to repetitive DNA, including transcribed transposable elements (Table S1E-G). 16,528 (*P. blakesleeanus*) and 11,719 (*M. circinelloides*) protein-coding genes were annotated and compared to proteins from other fungi (Figure 1C). Comparison of the two genomes with that of *Rhizopus delemar* and other fungi suggests that a whole-genome duplication (WGD) occurred early in the Mucoromycotina lineage. The fungal kingdom contains the subkingdom Dikarya and a number of early divergent lineages including the Mucoromycotina with *P. blakesleeanus*, *M. circinelloides*, and *R. delemar* [10]. Two WGDs have been reported in fungi: in the Saccharomycotina, a lineage of the Dikarya [4-6], and in *R. delemar* [3].

Ancient WGDs are difficult to detect because gene loss and rearrangements result in the absence of regions of synteny. However, genomes from Mucoromycotina species have more members per gene family than genomes from Dikarya fungi (2.9-3.6 versus 1.6–2.2) and a large fraction of gene families with more members than average (50%-68% versus 6.9%-22% for each Dikarya species) (Supplemental Experimental Procedures; Table S2A). Moreover, Mucoromycotina genomes have more duplicated regions than other fungal genomes, with 4-13 genes on average (Tables S2B and S2C). We confirmed the presence of duplicated regions after WGDs in the genomes of *S. cerevisiae* and *R. delemar* as expected, and also large amounts of duplicated DNA in several Dikarya fungi (Tables S2B and S2C). The *Puccinia graminis* and *Laccaria bicolor* genomes have expanded lineage-specific gene families proposed to be involved in pathogenesis and symbiosis [11, 12]. These duplicated regions contain large fractions of lineage-specific genes (Table S2C) supporting the proposal that they arose after species-specific segmental duplications. Additional WGD signatures can be observed in families of three genes from genome pairs. These types of duplicates are more frequent in the genomes of Mucoromycotina species than in non-Mucoromycotina species, suggesting that the former harbors traces of past WGDs (Table S2D).

To gain further insight into past genome expansions in the Mucoromycotina, we reconstructed the complete collection of evolutionary histories (*i.e.* the phylome) for genes within Mucoromycotina fungi with 13 other fungal genomes. The gene trees were analyzed to detect and date duplication events [13] (Figure 1C). This method has been used to characterize the WGD that took place in the *S. cerevisiae* lineage [14]. In addition to the WGD described in *R. delemar* (0.43 duplications per gene), we detected a larger duplication peak (0.70-0.96 duplications per gene) in the lineage preceding the Mucoromycotina species consistent with a WGD preceding the diversification of this lineage. This early WGD explains gene duplications in the oxidative phosphorylation complex in Mucorales [15] and

segmental duplications in *Lichtheimia corymbifera* where previously a species-specific WGD had been rejected [16].

The best explanation for our observations is a WGD predating the diversification of the Mucorales followed by a WGD in the *R. delemar* lineage and, subsequently, rampant gene loss as observed in yeast [5]. The alternative explanation (lineage-specific gene duplications) is less parsimonious. Although segmental duplications can create paralogous regions with shared synteny, we consider it unlikely that numerous such events affecting large regions of the genome would have coincided in time or affected multiple lineages in parallel.

Expansion of Gene Families in Mucoromycotina fungi

Duplicated genes in the four Mucoromycotina species contained an abundance of Gene Ontology (GO) terms for protein kinase activities (GO:4674, GO:4672, GO:4713, GO:4707; P-value at least 1×10^{-6}), fructose 2,6-bisphosphate metabolic process (GO:6003; P= 1.3×10^{-6}), ATP-binding (GO:5524; P= 1.8×10^{-46}), and protein transport (GO:15031; P= 1.6×10^{-17}), suggesting duplications of genes encoding signaling pathways and transport components (Supplemental Experimental Procedures; Figure 2; Data S1). This is supported by further analysis of the abundance of signal transduction genes. We have limited this analysis to the genomes of *P. blakesleeanus*, *M. circinelloides*, and *R. delemar* as they were the only Mucoromycotina genomes sequenced when we started the project.

Heterotrimeric G-protein signaling is central to the life cycle and virulence of fungi [17, 18]. All the gene families encoding the subunits of heterotrimeric G proteins are expanded. The G α subunit family contains 10, 12 and 12 genes in *P. blakesleeanus*, *M. circinelloides*, and *R. delemar*, respectively, compared to an average of three in the Ascomycota or seven in the Dikarya (Figure 2A). A single G β subunit gene has been found in the Ascomycota as well as in the basidiomycetes *Ustilago maydis* and *Cryptococcus neoformans* [19, 20]. In *P. blakesleeanus* we identified five G β genes, in *M. circinelloides* three, and in *R. delemar* four. Similarly, the Mucoromycotina genomes have three or four genes for the G γ subunit compared to an average of one in the Dikarya (Figure 2A). Theoretically, a very large number of G protein heterotrimers could be built from the multiple subunits. An estimate of 21, 21 and 10 genes for G protein coupled receptors (GPCR) in *P. blakesleeanus*, *M. circinelloides* and *R. delemar*, respectively (Data S1, sheet 1), suggests moderate expansion compared to 10-12 genes in Dikarya [21].

The number of other signal transduction genes has increased compared to the Dikarya, including genes for protein kinases, TRAFAC class GTPases, and regulators of GTPases of the Ras superfamily (Data S1, sheets 2-4). The expansion of kinase families is 3-4 fold, as there are 63, 70 and 82 CAMK genes in *P. blakesleeanus*, *M. circinelloides* and *R. delemar* respectively, compared to 22 and 21 in *Neurospora crassa* and *U. maydis*. Other families show larger expansion, e.g. 11-18 genes for casein kinase 1 in Mucoromycotina compared to 2-3 in Dikarya (Figure 2A; Data S1 sheet 4).

Some but not all families in a given category are expanded, e.g. for photoreception [22], the genes for components of the WC photoreceptor complex (WC-1 and WC-2) are duplicated, but not the cryptochrome gene; genes for casein kinase 1 are duplicated, but not those

encoding the sensor histidine kinases (Figure 2A). Genes for calcium or pH sensing show non-uniform duplication: there are multiple calmodulin genes and three calcineurin catalytic subunit genes, but only a single calcineurin regulatory subunit; in the pH pathway there are 3-4 genes for the PacC transcription factor and two genes for PalA, yet one gene for PalB or PalC as in the ascomycete *Aspergillus nidulans* [23] (Data S1, sheet 1). Cyclin families are expanded, but there is only a single mitotic cyclin, compared to several in the Dikarya (Figure 2A). Not all gene families have expanded. For example, the genes encoding proteins that participate in genome defense through RNAi have not duplicated (Figure 2A) [24].

The genome duplication has multiplied genes involved in cell wall biosynthesis, in particular chitin synthases and chitin deacetylases (Figure 2A). These enzymes may have specialized to modulate the growth response of the sporangiophore after environmental stimuli. Functional specialization after gene duplication should have played a key role in *M. circinelloides* and other pathogenic Mucoromycotina fungi. Mutants of the photoreceptor gene *wc-1* of *Fusarium oxysporum* and *C. neoformans* show decreased virulence [25, 26]. *M. circinelloides* *wc* genes have specialized their sensory role after gene duplication [9] suggesting that some WC proteins may serve as pathogenicity factors (Torres-Martínez, unpublished results). Calcineurin is a virulence factor in several fungi, including *M. circinelloides* [27]. The number of calcineurin A catalytic subunit genes has increased, with three genes in *M. circinelloides* compared to one in other fungi (Data S1, sheet 1). One of them, *cnaA*, is involved in virulence [27], confirming gene specialization to facilitate pathogenesis after duplication.

The expansion of some gene families can be accounted for by a WGD and retention of the resulting paralogs. The large number of chitin deacetylases or G α subunits genes (Figure 2A, Data S1, sheet 1), however, cannot be explained by a WGD alone, suggesting additional segmental duplications. Having more proteins for signal transduction and cell wall biosynthesis should have helped Mucoromycotina fungi to improve environmental sensing and responses, including the perception of potential hosts for pathogenic fungi. Elucidation of the biological role of duplicated genes will require further characterization.

Duplicated Genes Differ in their Transcriptional Response to Light

To investigate whether duplicated genes have specialized we asked if duplicated genes from *P. blakesleeanus* and *M. circinelloides* responded differently to light. The transcriptome of cultures kept in the dark or after exposure to 30 min of blue light (2.3×10^3 J/m²) indicates specialization (Figure 2B). Most of the genes encoding components of the photoreceptor complex (WCC) or the regulatory subunits of protein kinase A showed a similar expression pattern in *P. blakesleeanus* and *M. circinelloides* with only some genes regulated by light despite being transcriptionally active in both mycelia and sporangiophores (Figure 2B). In addition, genes encoding the photoreceptor WcoB, the G β subunits Gpb1 and Gpb3, and the kinases Pkac-1, Mps1-1, CK1-7 and CK1-8, showed opposite responses to light in *P. blakesleeanus* and *M. circinelloides*. Thus duplicated genes have evolved different patterns of expression in different organisms, as well as between specific tissues in the same organism.

A Refined Transcriptional Response to Light in *P. blakesleeanus*

The expansion in the number of photoreceptors and other signal transduction proteins may have allowed fine-tuning of the response to light, for example allowing tissue-specific transcriptional responses. We thus characterized the global transcriptional response in *P. blakesleeanus* mycelium and sporangiophores by RNAseq (Figure 3; Data S2). A total of 2,024 genes were responsive to light in the mycelium (1,421 induced and 603 repressed, about 12% of the protein-coding genes), compared to 1,212 genes in the sporangiophores (1,042 induced and 170 repressed). The transcriptional response to light was specific for each developmental stage because only 120 genes were light-regulated in both mycelium and sporangiophores (Figure 3A-B). The same analysis on a *madA madB* double mutant (strain L51) that is considered blind [28] detected only 159 light-regulated genes in the mycelium confirming the relevance of the Mad complex (Figure 3A). Surprisingly, the *madA madB* mutant showed a significant response to light in sporangiophores where 3,513 genes were regulated by light (Figure 3A). It is noteworthy that most responsive genes in the *madA madB* mutant sporangiophores were repressed by light (Figure 3A). This suggests the activity of light-dependent repressors in the absence of the Mad complex, as proposed by EMSA experiments that showed the binding of proteins in the dark to a light-regulated promoter in a *madA madB* mutant [29]. Seven gene clusters were enriched in regulatory genes in light-induced mRNAs from the mycelium of the wild type, and in ribosome biogenesis genes in the sporangiophores (Figure 3C). *P. blakesleeanus* and *M. circinelloides* have 879 and 650 genes encoding transcription factors (TFs), respectively (about 5% of the protein-coding genes), with an abundance of C₂H₂ Zn finger TFs (Fig. S2). Light regulates 9% of the *P. blakesleeanus* transcription factor genes (Data S1, sheets 5-7; Data S2, sheets 5-6), and we propose that the stage-specific transcriptional response to light relies on the expanded set of photoreceptors and light-dependent transcriptional regulators. Specialization of genes for signal transduction following WGD has been observed in vertebrate vision [30]. Strikingly, in both vertebrates and fungi the expansion of signal transduction genes after WGD has resulted in more elaborate sensory perception.

Reduced Sensitivity to Light in Strains with Wild Type and *madA* Mutant Nuclei

The photoreponse in the *P. blakesleeanus madA madB* mutant suggested the action of additional photoreceptors. Light perception in *madI* mutants is reduced 10 to 1000-fold, halfway between wild type and *madA* mutants, making MadI a candidate for a photoreceptor [31, 32] (Figure 4A). To identify *madI* we crossed two *madI* strains with a wild-type strain, and characterized the phototropism and molecular markers in the progeny [33]. Weak linkage was found for the *madI* mutation and three scaffolds, including the one carrying *madA* (Data S3). We therefore sequenced the genomes of two *madI* mutants (L151 and L153), along with another 17 *mad* mutant strains, and the sequences were scanned across the three scaffolds. We found that the two *madI* strains had an identical and unique mutation in *madA* [8] changing a conserved proline to leucine (Figure 4B-C). However, the *madI* strains also contained the wild type allele, indicating that they were heterozygous for this gene. To confirm these observations we sequenced *madA* in 63 *madI* x wild type progeny: eight only had the mutation in the *madA* gene, 35 were wild type, and 20 were heterozygotes. The two *madI* mutants had a high number of heterozygous sites across their genomes compared to other strains (Figure 4D). Analysis of all scaffolds in the L151 and

L153 genomes showed that the heterozygous single nucleotide polymorphisms (SNPs) are distributed throughout all chromosomes (Figure S3), suggesting that the two *madI* strains are heterokaryons or diploids, rather than being aneuploid or carrying a segmental duplication. The observation that two of the *madI* strains are heterozygous wild type/*madA* mutants shows that sensitivity to light is related to the dosage of the MadA photoreceptor.

Our characterization of the genomes of *P. blakesleeanus* and *M. circinelloides*, our comparative fungal genome analysis, and our gene function studies provide new insight into the occurrence and consequences of genome duplications in the evolution of fungi. Expansion and specialization of genes for signal transduction and cell-wall biosynthesis following genome duplication in the Mucoromycotina provided new proteins that have enabled these fungi to refine the way they perceive signals from the environment to regulate their growth and development. Our results provide new genomic tools to unravel the molecular mechanisms of sensory perception in early diverging fungi that will help to understand the evolution of sensory perception in eukaryotes.

Supplementary Material

Refer to Web version on PubMed Central for supplementary material.

Authors

Luis M. Corrochano¹, Alan Kuo², Marina Marcet-Houben^{3,4}, Silvia Polaino⁵, Asaf Salamov², José M. Villalobos-Escobedo⁶, Jane Grimwood^{2,7}, M. Isabel Álvarez⁸, Javier Avalos¹, Diane Bauer², Ernesto P. Benito^{8,9}, Isabelle Benoit¹⁰, Gertraud Burger¹¹, Lola P. Camino¹, David Cánovas¹, Enrique Cerdá-Olmedo¹, Jan-Fang Cheng², Angel Domínguez⁸, Marek Eliáš¹², Arturo P. Eslava⁸, Fabian Glaser¹³, Gabriel Gutiérrez¹, Joseph Heitman¹⁴, Bernard Henrissat^{15,16}, Enrique A. Iturriaga⁸, B. Franz Lang¹¹, José L. Lavín¹⁷, Soo Chan Lee¹⁴, Wenjun Li¹⁴, Erika Lindquist², Sergio López-García¹⁸, Eva M. Luque¹, Ana T. Marcos¹, Joel Martin², Kevin McCluskey¹⁹, Humberto R. Medina¹, Alejandro Miralles-Durán¹, Atsushi Miyazaki²⁰, Elisa Muñoz-Torres²¹, José A. Oguiza²², Robin A. Ohm², Margarita Orejas²³, Lucila Ortiz-Castellanos²⁴, Antonio G. Pisabarro²², Julio Rodríguez-Romero¹, José Ruiz-Herrera²⁴, Rosa Ruiz-Vázquez¹⁸, Catalina Sanz⁸, Wendy Schackwitz², Mahdi Shahriari⁸, Ekaterina Shelest²⁵, Fátima Silva-Franco¹⁸, Darren Soanes²⁶, Khajamohiddin Syed²⁷, Víctor G. Tagua¹, Nicholas J. Talbot²⁶, Michael R. Thon^{8,9}, Hope Tice², Ronald P. de Vries¹⁰, Ad Wiebenga¹⁰, Jagjit S. Yadav²⁷, Edward L. Braun²⁸, Scott E. Baker²⁹, Victoriano Garre¹⁸, Jeremy Schmutz^{2,7}, Benjamin A. Horwitz³⁰, Santiago Torres-Martínez¹⁸, Alexander Idnurm^{5,31}, Alfredo Herrera-Estrella⁶, Toni Gabaldón^{3,4,32}, and Igor V. Grigoriev²

Affiliations

¹Department of Genetics, University of Seville, Avda. Reina Mercedes s/n, 41012 Seville, Spain ²U.S. Department of Energy Joint Genome Institute, 2800 Mitchell Drive, Walnut Creek, CA 94598, USA ³Centre for Genomic Regulation (CRG), Doctor Aiguader 88, 08003 Barcelona, Spain ⁴Universitat Pompeu Fabra (UPF),

Doctor Aiguader 88, 08003 Barcelona, Spain ⁵School of Biological Sciences, University of Missouri-Kansas City, 5007 Rockhill Road, Kansas City, MO 64110, USA ⁶Laboratorio Nacional de Genómica para la Biodiversidad (LANGEBIO), Centro de Investigación y Estudios Avanzados, Kilómetro 9.6 Libramiento Norte, Carretera Irapuato-León, 36821 Irapuato, Guanajuato, México ⁷HudsonAlpha Institute of Biotechnology, 601 Genome Way Northwest, Huntsville, AL 35806, USA ⁸Departamento de Microbiología y Genética, Universidad de Salamanca, Plaza de los doctores de la Reina s/n, 37007 Salamanca, Spain ⁹Instituto Hispano-Luso de Investigaciones Agrarias (CIALE), Universidad de Salamanca, Río Duero 12, 37185 Salamanca, Spain ¹⁰CBS-KNAW Fungal Biodiversity Centre & Fungal Molecular Physiology, Utrecht University, Uppsalalaan 8, 3584 CT Utrecht, The Netherlands ¹¹Universite de Montreal, Pavillon Roger-Gaudry, Biochimie, CP 6128, Succursale Centre-Ville, Montreal QC, H3C 3J7, Canada ¹²Department of Biology and Ecology, Faculty of Science, University of Ostrava, Chittussiho 10, 710 00 Ostrava, Czech Republic ¹³Lokey Interdisciplinary Center for Life Sciences and Engineering, Technion - Israel Institute of Technology, Haifa 3200003, Israel ¹⁴Department of Molecular Genetics and Microbiology, Duke University Medical Center, Research Drive, Durham, NC 27710, USA ¹⁵Centre National de la Recherche Scientifique (CNRS), UMR7257, Université Aix-Marseille, 163 Avenue de Luminy, 13288 Marseille, France ¹⁶Department of Biological Sciences, King Abdulaziz University, P.O Box 80203, Jeddah 21589, Saudi Arabia ¹⁷Genome Analysis Platform, CIC bioGUNE, Bizkaia Technology Park, 48160 Derio, Bizkaia, Spain ¹⁸Departamento de Genética y Microbiología, Universidad de Murcia, 30071 Murcia, Spain ¹⁹Department of Plant Pathology, Kansas State University, 4024 Throckmorton Plant Sciences Center, Manhattan, KS 66506, USA ²⁰Department of Biological Sciences, Faculty of Science and Engineering, Ishinomaki Senshu University, Ishinomaki 986-8580, Japan ²¹Departamento de Biología Celular y Patología, Facultad de Medicina, Universidad de Salamanca, Avda. Campus Miguel de Unamuno, 37007 Salamanca, Spain ²²Department of Agrarian Production, Public University of Navarre, 31006 Pamplona, Spain ²³Instituto de Agroquímica y Tecnología de Alimentos, Consejo Superior de Investigaciones Científicas (IATA-CSIC), Avda. Catedrático Agustín Escardino 7, 46980 Paterna, Valencia, Spain ²⁴Departamento de Ingeniería Genética, Unidad Irapuato, Centro de Investigación y de Estudios Avanzados del Instituto Politécnico Nacional, Kilómetro 9.6 Libramiento Norte, Carretera Irapuato-León, 36821 Irapuato, Guanajuato, Mexico ²⁵Leibniz Institute for Natural Product Research and Infection Biology (Hans Knoell Institute), Beutenbergstr. 11a, 07745 Jena, Germany ²⁶Biosciences, College of Life & Environmental Sciences, University of Exeter, Stocker Road, Exeter EX4 4QD, UK ²⁷Department of Environmental Health, University of Cincinnati College of Medicine, 160 Panzeca Way, Cincinnati, OH 45267-0056, USA ²⁸Department of Biology, University of Florida, Gainesville, P.O. Box 118525, FL 32611-8525, USA ²⁹Pacific Northwest National Laboratory, 902 Battelle Blvd., Richland, WA 99352, USA ³⁰Department of Biology, Technion - Israel Institute of Technology, Haifa 32000,

Israel ³¹School of BioSciences, University of Melbourne, VIC 3010, Australia
³²Institució Catalana de Recerca i Estudis Avançats (ICREA), Barcelona, Spain

Acknowledgments

This article is dedicated to the memory of Max Delbrück (1906-1981). The work by the U.S. Department of Energy Joint Genome Institute, a DOE Office of Science User Facility, is supported by the Office of Science of the U.S. Department of Energy under Contract No. DE-AC02-05CH11231. The project was supported by European funds (European Regional Development Fund, ERDF), the Spanish Ministerio de Economía y Competitividad (BIO2005-25029-E, BIO2015-67148-R), and the Regional Government (Junta de Andalucía, P06-CVI-01650) to LMC; Conacyt (Mexico) (FORDECYT-2012-02-193512) to AHE; the US National Science Foundation (MCB-0920581) to AI; and the Czech Science Foundation (13-33039S) to M.E. We thank A. Gryganskyi, D. Armaleo, and P. Dyer for allowing us to use the genome sequences from *Lichtheimia corymbifera*, *Cladonia grayi*, and *Xanthoria parietina*, respectively, prior to publication. We thank M. M. Gil Sánchez for *P. blakesleeana* photographs in Fig. 1A.

References

1. Cerdá-Olmedo E. *Phycomyces* and the biology of light and color. *FEMS Microbiol Rev.* 2001; 25:503–512. [PubMed: 11742688]
2. Corrochano LM, Garre V. Photobiology in the Zygomycota: multiple photoreceptor genes for complex responses to light. *Fungal Genet Biol.* 2010; 47:893–899. [PubMed: 20466063]
3. Ma LJ, Ibrahim AS, Skory C, Grabherr MG, Burger G, Butler M, Elias M, Idnurm A, Lang BF, Sone T, et al. Genomic analysis of the basal lineage fungus *Rhizopus oryzae* reveals a whole-genome duplication. *PLoS Genet.* 2009; 5:e1000549. [PubMed: 19578406]
4. Wolfe KH, Shields DC. Molecular evidence for an ancient duplication of the entire yeast genome. *Nature.* 1997; 387:708–713. [PubMed: 9192896]
5. Dietrich FS, Voegeli S, Brachat S, Lerch A, Gates K, Steiner S, Mohr C, Pöhlmann R, Luedi P, Choi S, et al. The *Ashbya gossypii* genome as a tool for mapping the ancient *Saccharomyces cerevisiae* genome. *Science.* 2004; 304:304–307. [PubMed: 15001715]
6. Kellis M, Birren BW, Lander ES. Proof and evolutionary analysis of ancient genome duplication in the yeast *Saccharomyces cerevisiae*. *Nature.* 2004; 428:617–624. [PubMed: 15004568]
7. Sanz C, Rodríguez-Romero J, Idnurm A, Christie JM, Heitman J, Corrochano LM, Eslava AP. *Phycomyces* MADB interacts with MADA to form the primary photoreceptor complex for fungal phototropism. *Proc Natl Acad Sci USA.* 2009; 106:7095–7100. [PubMed: 19380729]
8. Idnurm A, Rodríguez-Romero J, Corrochano LM, Sanz C, Iturriaga EA, Eslava AP, Heitman J. The *Phycomyces madA* gene encodes a blue-light photoreceptor for phototropism and other light responses. *Proc Natl Acad Sci USA.* 2006; 103:4546–4551. [PubMed: 16537433]
9. Silva F, Torres-Martínez S, Garre V. Distinct *white collar-1* genes control specific light responses in *Mucor circinelloides*. *Mol Microbiol.* 2006; 61:1023–1037. [PubMed: 16879651]
10. McLaughlin DJ, Hibbett DS, Lutzoni F, Spatafora JW, Vilgalys R. The search for the fungal tree of life. *Trends Microbiol.* 2009; 17:488–497. [PubMed: 19782570]
11. Martin F, Aerts A, Ahren D, Brun A, Danchin EG, Duchaussoy F, Gibon J, Kohler A, Lindquist E, Pereda V, et al. The genome of *Laccaria bicolor* provides insights into mycorrhizal symbiosis. *Nature.* 2008; 452:88–92. [PubMed: 18322534]
12. Duplessis S, Cuomo CA, Lin YC, Aerts A, Tisserant E, Veneault-Fourrey C, Joly DL, Hacquard S, Amselem J, Cantarel BL, et al. Obligate biotrophy features unraveled by the genomic analysis of rust fungi. *Proc Natl Acad Sci USA.* 2011; 108:9166–9171. [PubMed: 21536894]
13. Huerta-Cepas J, Gabaldón T. Assigning duplication events to relative temporal scales in genome-wide studies. *Bioinformatics.* 2011; 27:38–45. [PubMed: 21075746]
14. Marcet-Houben M, Gabaldón T. Beyond the whole-genome duplication: phylogenetic evidence for an ancient interspecies hybridization in the baker's yeast lineage. *PLoS Biol.* 2015; 13:e1002220. [PubMed: 26252497]

15. Marcet-Houben M, Marceddu G, Gabaldón T. Phylogenomics of the oxidative phosphorylation in fungi reveals extensive gene duplication followed by functional divergence. *BMC Evol Biol.* 2009; 9:295. [PubMed: 20025735]
16. Schwartz VU, Winter S, Shelest E, Marcet-Houben M, Horn F, Wehner S, Linde J, Valiante V, Sammeth M, Riege K, et al. Gene expansion shapes genome architecture in the human pathogen *Lichtheimia corymbifera*: an evolutionary genomics analysis in the ancient terrestrial mucorales (Mucoromycotina). *PLoS Genet.* 2014; 10:e1004496. [PubMed: 25121733]
17. Lafon A, Han KH, Seo JA, Yu JH, d'Enfert C. G-protein and cAMP-mediated signaling in aspergilli: a genomic perspective. *Fungal Genet Biol.* 2006; 43:490–502. [PubMed: 16546420]
18. Li L, Wright SJ, Krystofova S, Park G, Borkovich KA. Heterotrimeric G protein signaling in filamentous fungi. *Annu Rev Microbiol.* 2007; 61:423–452. [PubMed: 17506673]
19. Müller P, Leibbrandt A, Teunissen H, Cubasch S, Aichinger C, Kahmann R. The G β -subunit-encoding gene *bpp1* controls cyclic-AMP signaling in *Ustilago maydis*. *Eukaryot Cell.* 2004; 3:806–814. [PubMed: 15190001]
20. Wang P, Perfect JR, Heitman J. The G-protein β subunit GPB1 is required for mating and haploid fruiting in *Cryptococcus neoformans*. *Mol Cell Biol.* 2000; 20:352–362. [PubMed: 10594037]
21. Xue C, Hsueh YP, Heitman J. Magnificent seven: roles of G protein-coupled receptors in extracellular sensing in fungi. *FEMS Microbiol Rev.* 2008; 32:1010–1032. [PubMed: 18811658]
22. Idnurm A, Verma S, Corrochano LM. A glimpse into the basis of vision in the kingdom Mycota. *Fungal Genet Biol.* 2010; 47:881–892. [PubMed: 20451644]
23. Peñalva MA, Tilburn J, Bignell E, Arst HN Jr. Ambient pH gene regulation in fungi: making connections. *Trends Microbiol.* 2008; 16:291–300. [PubMed: 18457952]
24. Nicolás FE, Moxon S, de Haro JP, Calo S, Grigoriev IV, Torres-Martínez S, Moulton V, Ruiz-Vázquez RM, Dalmay T. Endogenous short RNAs generated by Dicer 2 and RNA-dependent RNA polymerase 1 regulate mRNAs in the basal fungus *Mucor circinelloides*. *Nucleic Acids Res.* 2010; 38:5535–5541. [PubMed: 20427422]
25. Ruiz-Roldán MC, Garre V, Guarro J, Mariné M, Roncero MI. Role of the white collar 1 photoreceptor in carotenogenesis, UV resistance, hydrophobicity, and virulence of *Fusarium oxysporum*. *Eukaryot Cell.* 2008; 7:1227–1230. [PubMed: 18503005]
26. Idnurm A, Heitman J. Light controls growth and development via a conserved pathway in the fungal kingdom. *PLoS Biol.* 2005; 3:e95. [PubMed: 15760278]
27. Lee SC, Li A, Calo S, Heitman J. Calcineurin plays key roles in the dimorphic transition and virulence of the human pathogenic zygomycete *Mucor circinelloides*. *PLoS Pathog.* 2013; 9:e1003625. [PubMed: 24039585]
28. Lipson ED, Terasaka DT. Photogeotropism in *Phycomyces* double mutants. *Exp Mycol.* 1981; 5:101–111.
29. Sanz C, Benito EP, Orejas M, Alvarez MI, Eslava AP. Protein-DNA interactions in the promoter region of the *Phycomyces carB* and *carRA* genes correlate with the kinetics of their mRNA accumulation in response to light. *Fungal Genet Biol.* 2010; 47:773–781. [PubMed: 20580936]
30. Larhammar D, Nordström K, Larsson TA. Evolution of vertebrate rod and cone phototransduction genes. *Philos Trans R Soc Lond B Biol Sci.* 2009; 364:2867–2880. [PubMed: 19720650]
31. Campuzano V, Díaz-Mínguez JM, Eslava AP, Alvarez MI. A new gene (*madI*) involved in the phototropic response of *Phycomyces*. *Mol Gen Genet.* 1990; 223:148–151. [PubMed: 2259336]
32. Alvarez MI, Eslava AP, Lipson ED. Phototropism mutants of *Phycomyces blakesleeana* isolated at low light intensity. *Exp Mycol.* 1989; 13:38–48.
33. Chaudhary S, Polaino S, Shakya VP, Idnurm A. A new genetic linkage map of the zygomycete fungus *Phycomyces blakesleeana*. *PLoS One.* 2013; 8:e58931. [PubMed: 23516579]




Figure 1. Sensory perception and genome duplication in the Mucoromycotina

(A) The fruiting bodies, sporangiophores, of *Phycomyces blakesleeanus* grow out of the mycelium and reach several cm in length. The speed and direction of growth is controlled by signals from the environment including light, gravity, touch, wind, and the presence of nearby objects. The ball at the top of each fruiting body is the sporangium with spores. The direction of light is indicated by an arrow.

(B) The sporangiophores of *Mucor circinelloides* are small (about 5 mm) and show phototropism. The direction of light is indicated by an arrow.

(C) Evidence for a WGD in the Mucoromycotina. A fungal evolutionary tree with bootstrap support values lower than 95% indicated in black numbers at the branches. The average duplication per gene in each lineage is shown with a color that indicates the phylome used for the duplication density calculation. The branch where the proposed WGD took place is marked in red with a dot and an arrow. The graph (scale on the bottom) represents the percentage of genes in a given species that belong to one of the following categories: yellow, protein present in all species; light yellow, ancestral proteins that have homologs in the outgroups; brown, Fungi-specific proteins; green, Mucoromycotina-specific proteins which appear in all four species; light green, Mucoromycotina-specific proteins; grey, species-specific proteins. The red bars (scale on top) represent the total number of proteins encoded in each genome. See also Figure S1, Table S1, and Table S2.




Figure 2. Gene expansion and transcriptional specialization in the Mucoromycotina
 (A) Gene abundance in Mucoromycotina, *Neurospora crassa* and Dikarya fungi. The x-axis indicates number of genes, and the bars, from bottom to top, indicate numbers of predicted genes for the three Mucoromycotina (Pb, *P. blakesleeanus*; Mc, *M. circinelloides*; Rd, *R. delemar*), and average number of genes for *N. crassa* (Nc), and Dikarya.
 (B) Expression patterns in response to light of duplicated genes in *P. blakesleeanus* and *M. circinelloides*. Differential expression of genes was obtained for two *P. blakesleeanus* stages (mycelium and sporangiophore), and *M. circinelloides* mycelium. The three *wc-1* genes are *madA*, *wcoA*, and *wcoB*, and the four *wc-2* genes are *madB*, *wctB*, *wctC*, and *wctD*. Results are represented with the logarithm base ten of FDR, (FDR<0.05; fold-change>2). M= Mycelium, S= Sporangiophore. See also Figure S2 and Data S1.




Figure 3. The influence of light on gene expression in two developmental stages of *P. blakesleeanus*

(A) Differential expression (light/dark) in the mycelium or the sporangioophore of the wild-type and the *madA madB* mutant strain (L51). Differentially expressed genes with FDR 0.05 are shown in red.

(B) Overlap of genes induced and repressed in the wild type and mutant using RNA from mycelia (WT-ML and L51-ML) or sporangioophores (WT-SL and L51-SL).

(C) Category enrichment in differentially expressed genes (*FDR<0.05; **FDR<0.01). Each vertical block contains the up- and down-regulated categories. Color intensity represents the percentage of genes belonging to each category and includes only GO terms for Biological Processes. Clusters based on this percentage are displayed in different colors in the tree. See also Data S2.




Figure 4. The *P. blakesleeanus* *madI* strains are heterokaryons of wild type and *madA* nuclei
 (A) Phenotype of wild type, *madA*, and *madI* strains L151 and L153, with illumination from the right. The average bending response of *madI* strains is lower than in the wild type strain with the intensity that we used.
 (B) DNA sequencing chromatograms of a region of the *madA* gene from the two *madI* strains and progeny from the *madI* mutants crossed to wild-type UBC21. The progeny sequences represent the three different types observed in the strains obtained from crosses.
 (C) A segment of the sequence of MadA in different fungi with the conserved proline that is mutated in *madI* strains in bold.
 (D) Graph showing the ratio of heterozygous/non-heterozygous SNPs in the genomes of 19 *mad* mutants. See also Figure S3 and Data S3.