

UCLA

UCLA Previously Published Works

Title

Methamphetamine use drives decreases in viral suppression for people living with HIV released from a large municipal jail: Results of the LINK LA clinical trial

Permalink

<https://escholarship.org/uc/item/5mv5p2gj>

Authors

Goodman-Meza, David

Shoptaw, Steve

Weiss, Robert E

et al.

Publication Date

2019-09-01

DOI

10.1016/j.drugalcdep.2019.05.007

Peer reviewed



# HHS Public Access

Author manuscript

*Drug Alcohol Depend.* Author manuscript; available in PMC 2020 September 01.

Published in final edited form as:

*Drug Alcohol Depend.* 2019 September 01; 202: 178–184. doi:10.1016/j.drugalcdep.2019.05.007.

## Methamphetamine use drives decreases in viral suppression for people living with HIV released from a large municipal jail: Results of the LINK LA Clinical Trial

David Goodman-Meza<sup>a,\*</sup>, Steve Shoptaw<sup>b</sup>, Robert E. Weiss<sup>c</sup>, Terry Nakazono<sup>d</sup>, Nina T. Harawa<sup>b,e</sup>, Sae Takada<sup>f,g</sup>, Wendy H. Garland<sup>h</sup>, William E. Cunningham<sup>d,i</sup>

<sup>a</sup>Division of Infectious Diseases, David Geffen School of Medicine, University of California, Los Angeles, CA, USA

<sup>b</sup>Department of Family Medicine, David Geffen School of Medicine, University of California, Los Angeles, CA, USA

<sup>c</sup>Department of Biostatistics, Fielding School of Public Health, University of California, Los Angeles, CA, USA

<sup>d</sup>Department of Medicine, Division of General Internal Medicine and Health Services Research, David Geffen School of Medicine, University of California, Los Angeles, CA, USA

<sup>e</sup>Charles R. Drew University College of Medicine, Los Angeles, CA, USA

<sup>f</sup>National Clinician Scholars Program UCLA, Division of General Internal Medicine and Health Services Research, Department of Medicine, Los Angeles, CA, USA

<sup>g</sup>VA HSR and D Center for the Study of Healthcare Innovation, Implementation, and Policy, Los Angeles, CA, USA

<sup>h</sup>Division of HIV and STD Programs, Los Angeles County Department of Public Health, Los Angeles, CA, USA

<sup>i</sup>Department of Health Policy and Management, Fielding School of Public Health, University of California, Los Angeles, CA, USA

### Abstract

\***Corresponding author:** David Goodman-Meza, David Geffen School of Medicine at UCLA, 10833 Le Conte Ave (Room 37-121CHS), Los Angeles CA, 90095-1688.

Contributors

Study concept and design: Goodman-Meza, Cunningham, Weiss, Shoptaw. Statistical analysis: Nakazono and Weiss. Drafting of the manuscript: Goodman-Meza, Cunningham, Weiss, Nakazono, Harawa. Critical revision of the manuscript for important intellectual content: Goodman-Meza, Cunningham, Weiss, Shoptaw, Harawa, Garland. All authors reviewed and approved the final version of the manuscript.

**Publisher's Disclaimer:** This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final citable form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

Conflict of Interest  
No conflict declared.

**Background:** People living with HIV (PLWH) often experience decreases in HIV viral suppression (VS) after release from jail. The Linking Inmates to Care in LA (LINK LA) peer navigation intervention helped maintain VS 12 months after release from jail compared to standard of care. In this study, we analyzed correlates of substance use and tested whether substance use was an independent correlate of decreased VS in LINK LA participants.

**Methods:** We analyzed LINK LA data collected at baseline, 3, and 12 months. We defined high-risk drug use as any reported methamphetamine, cocaine, or opioid use in the 30 days prior to a study visit (or jail entry at baseline). We used generalized linear mixed models to test associations of sociodemographic variables with type of substance used, and we tested correlates of VS while controlling for time, the intervention, and their interaction.

**Results:** At baseline (n=356), 71% of participants reported high-risk drug use: 58% methamphetamine; 17% cocaine; 7% heroin; and 4% prescription opioids. Non-Hispanic Whites and those younger than 35 were most likely to use methamphetamine; Blacks were most likely to use cocaine; people who inject drugs were most likely to use opioids. Participants who used high-risk drugs had 53% lower adjusted odds than non-users of maintaining VS (AOR 0.47, 95% CI 0.31-0.70, p<0.001).

**Conclusion:** High-risk drug use, dominated by methamphetamine use, independently correlated with decreased VS among recently incarcerated PLWH. Improving HIV care continuum outcomes among populations leaving jail requires attention to efforts to address high-risk drug use.

## Keywords

Substance Use; Stimulant; Methamphetamine; HIV Viral Suppression; Jail

## 1. Introduction

In the United States, viral suppression (VS) improved dramatically in the general population of people living with HIV (PLWH) following the introduction of convenient, highly active, and highly tolerable antiretroviral therapy regimens (Nance et al., 2018). Yet, PLWH who are outside of the HIV care continuum and not virally suppressed continue to be the major source of HIV transmission (Escudero et al., 2017; Skarbinski et al., 2015) and preventable HIV-related morbidity and mortality (Kerr et al., 2014; 2012; Lesko et al., 2015). Disparities in engaging and retaining PLWH in the care continuum disproportionately affect PLWH who use substances (Korthuis and Edelman, 2018; Lazar et al., 2017), are involved in the criminal justice system (CJS) (Iroh et al., 2015), or are racial/ethnic minorities— especially Black and Hispanic PLWH (Lazar et al., 2017). Adequately engaging those who are not virally suppressed into the HIV care continuum is essential to reach goals of 90% of people leaving with HIV diagnoses, 90% of these in HIV medical care, and 90% of these virally suppressed (i.e., 90-90-90 goals) in the United States (Kay et al., 2016).

The transition period following release from correctional facilities can be marked by HIV viremia for PLWH due to medical care disruptions (Iroh et al., 2015). Linking Inmates to Care in LA (LINK LA) was a randomized controlled trial that was designed to test an intervention to improve linkage, retention, and antiretroviral therapy (ART) adherence among PLWH released from a large municipal jail with the goal of achieving or maintaining

VS (Cunningham et al., 2018). LINK LA enrolled a racially diverse group of PLWH (42% Black, 31% Hispanic, 27% White) prior to release from the nation's largest municipal jail, with 78% reporting either high-risk drug (stimulants or opioids) or binge alcohol use. LINK LA examined a peer navigation intervention that maintained VS in 49% of the intervention group from baseline to month 12, compared to a decrease from 52% to 30% in the control group, for a significant difference-in-difference of 22%. Although high-risk drug use declined over time for all participants (20% decline in the intervention arm, 16% in the control arm), there was no statistically significant effect of the intervention on high-risk drug use, and there was no interaction between the intervention and high-risk drug use.

In observational studies, a co-existing substance use disorder has often been associated with decreases in VS or HIV care engagement after release from incarceration (Ammon et al., 2018; Chitsaz et al., 2013; Harawa et al., 2017; Palepu et al., 2004). Of the many previous studies of the connections between drug use and HIV outcomes, a large proportion have focused on *injection* drug use as a key marker of the deleterious effects of drug use. However, many fewer studies have examined the links of HIV continuum outcomes with the specific types of drug used (Lesko et al., 2017; Westergaard et al., 2013; 2011).

For the present study, we had 3 main goals: 1) to examine the sociodemographic and clinical characteristics correlated with use of specific types of drugs (including binge alcohol use) among participants in the LINK LA study at baseline; 2) to examine the sociodemographic and clinical predictors of using specific drug types over the 12 months following release from Los Angeles (LA) County Jail; and 3) to evaluate the associations of specific substance use types with VS, independent of the intervention effects for the 12 months of follow-up among men and transgender women living with HIV after release from incarceration in the LINK LA study. We hypothesized that participants who reported use of any high-risk drugs would be less likely to maintain VS over time, over and above known positive effects of the peer navigation intervention on VS.

## 2. Methods

### 2.1. Parent Study

LINK LA was a randomized controlled trial conducted in Los Angeles (LA), California between December 2012 and October 2016. A detailed description of the study, intervention, and findings have been published (Cunningham et al., 2018). In brief, LINK LA enrolled PLWH who had been released from LA County Jail. Participants were 1) adult, 2) men or transgender women, 3) living with HIV, 4) residents of LA County, 5) English speaking, 6) eligible for or receiving antiretroviral therapy, and 7) selected for HIV transitional case management prior to enrollment. The study randomized participants to one of two groups: 1) the intervention group: received LINK LA peer navigation and a 12-session intervention based on Social Cognitive Theory imparted by peers over 6-months; 2) the control group: received standard transitional case management. Participants in both arms received referrals for post-release housing, substance abuse treatment, and HIV care. These referrals were provided by peer navigators to participants in the intervention group and by case managers to participants in the control group. Participants were interviewed in-person by trained research assistants before release (baseline) then at months 3, 6, and 12. All study

procedures were reviewed and overseen by the LA County Department of Public Health institutional review board (IRB) and the UCLA IRB.

## 2.2. Measures

**2.2.1. Substance Use.**—Substance use was assessed by a standardized questionnaire that included questions for alcohol (use and intoxication), heroin, methadone, prescription opioids misuse (e.g., oxycodone or other opioids), crack, cocaine, and methamphetamine or amphetamines, among others. At baseline, participants were asked if they ever used a substance; if they answered yes, the participants were asked if the substance was used within the 30 days prior to arrest and the duration use in years. At follow-up assessments, participants reported their use of the substance 30 days before the visits. There is extensive evidence to support the reliability and validity of self-reported substance use, when collected by trained interviewers, as in LINK LA (Dowling-Guyer et al., 2016; Johnson et al., 2000; Weiss et al., 1998). We created a composite variable, *high-risk drug use*, defined as any reported use of heroin, cocaine (powder or crack), methamphetamines or misuse of prescription opioids in the 30 days prior to being arrested (used at baseline) and 30-days prior to each follow-up assessment. We grouped the use of these substances, as they have the most consistently observed association with unsuppressed viral load in PLWH (Carrico et al., 2014; Qian et al., 2014; Rasbach et al, 2013). Binge alcohol use was defined as consuming 5 or more drinks on one occasion and also assessed for these periods.

**2.2.2. Covariates.**—Covariates included age, reported gender/HIV risk group, race and ethnicity, individual annual income, Short Form-12 (SF-12), time, intervention (navigation, control), CD4 count, and the interaction of time and intervention. Age was dichotomized as 34 years or less or 35 years or more based on age-related differences in patterns of substance use and of risk behaviors leading to HIV transmission. We grouped participants into mutually exclusive categories based on gender identity and/or HIV behavioral risk. The risk/gender categories included men who have sex with men (MSM), transgender women (TGW), people who inject drugs (PWID) regardless of sexual orientation and gender identity, and people in none of the previous categories (i.e., heterosexual cis-males who did not inject drugs). Race/ethnicity was defined as Hispanic, non-Hispanic Black, and non-Hispanic White. Individual annual income was dichotomized as \$10,000 or above and under \$10,000. Two SF-12 summary scores— the physical health composite and the mental health composite— were derived from the eight domains in the SF-12 and standardized using a T-score normed to the United States general population with mean of 50 and standard deviation of 10. Higher scores indicate better health (Ware et al., 1996). The SF-12 has been shown to be reliable and valid in PLWH and people with a substance use disorder (Ion et al, 2011; Pyne et al., 2011). CD4 count was dichotomized at 500 or more cells/ $\mu$ L and less than 500 cells/ $\mu$ L.

**2.2.3. HIV viral suppression.**—Viral load testing was performed at baseline, 3- and 12-months. Viral suppression (VS) was defined as an undetectable viral load or a quantitative value of less than 75 copies/mL. At baseline, the viral load was obtained from electronic medical records. At months 3 and 12, the viral load (HIV RNA) was obtained

from blood samples analyzed using RealTime HIV-1 assays (Abbott Laboratories). Blood samples for testing were obtained by a staff phlebotomist.

### 2.3. Statistical Analysis

We summarized baseline demographic characteristics and SF-12 composite scores by reported substance use as means and standard deviation or counts and percentages, where applicable. We modeled outcomes using longitudinal logistic random intercept models, which are particular cases of generalized linear mixed models (GLMMs). Outcomes are VS and reported substance use (methamphetamine, cocaine, and opioids each individually and as the composite “high-risk drug use”). The intervention, interview assessment (baseline, 3- and 12-months), and their interaction were used as predictors in all models. We first fit models with each additional covariate singularly. Possible covariates were age, gender/HIV risk group, race/ethnicity, income, SF-12 composite scores, and CD4 count. If the p-value for the corresponding coefficient was less than 0.05, the covariate was retained in a final multivariable model. To further determine the longitudinal association of each type of substance use with VS, we fit bivariate longitudinal random intercept models to estimate the correlation between participants’ frequency of reporting each type of substance on the logit scale and the participants’ VS over all three time-point assessments (Hara et al., 2013; Weiss, 2005). GLMMs robustly handle missing data via maximum likelihood estimation and treated the missing data as missing at random. We used SAS Proc Glimmix software version 9.4 (SAS Institute Inc., Cary, NC, USA) for all analyses.

## 3. Results

### 3.1. Baseline Characteristics by Reported Substance Use

LINK LA enrolled and randomized 356 participants between 2012 and 2015. Baseline characteristics by study arm were described previously (Cunningham et al., 2018). In brief, mean age was 39.5 (standard deviation [SD] 10.4), 85% were men and 15% TGW, 42% were Black and 31% Latino. The mean SF-12 mental health composite score was 38 (SD 12), and the mean physical health composite score was 51 (SD 10). The reported gender/HIV risk groups were 56% MSM, 12% PWID, 15% TGW, and 17% other. The percentages of each substance used at baseline by sociodemographic and clinical variables are presented in Table 1. Younger participants (34 years or younger) were more likely than older participants (35 years or older) to recently use binge alcohol, methamphetamines, and marijuana. Older participants were more likely to use cocaine than younger participants. White participants were the most likely to use any high-risk drugs and the most likely to use methamphetamines, followed by Hispanic then Black participants. Black participants were the most likely to use cocaine followed by Hispanic then White participants. By gender/HIV risk group, MSM were the most likely to use high-risk drugs and methamphetamines; TGW were the most likely to use cocaine; PWID were the most likely to use opioids.

### 3.2. Predictors of Substance Use Over Time

Results of longitudinal modeling of drug use over the course of the study are summarized in Table 2. For high-risk drug use, Black participants had lower odds than White participants of ongoing high-risk drug use, while Hispanic participants had similar odds as White

participants. MSM, TGW, and PWID all had higher odds of ongoing high-risk drug use than other participants. By individual drug, participants aged 34 years or less had more than twice the odds of methamphetamine use over time compared to those aged 35 years or older. Black participants had lower odds of methamphetamine use and higher odds of cocaine use than White participants. Again, MSM, TGW, and PWID all had higher odds of methamphetamine use than other participants. PWID were most likely to use any opioids.

### 3.3. Viral Suppression

Figure 1 shows the proportion of viral suppression in each arm and by high-risk drug use at study time points. In multivariable analysis, adjusted for the time-by-intervention interaction (Table 3), VS was independently predicted by baseline gender/HIV risk group, income, CD4 count, SF-12 mental health component score, and recent high-risk drug use. PWID was associated with higher odds of VS than participants in the other gender/HIV risk groups. Compared to those reporting a higher income, those reporting an income less than \$10,000 had lower odds of VS. Participants with a CD4 count lower than 500 cell/ $\mu$ L had lower odds of VS compared to those with a CD4 count over 500 cell/ $\mu$ L. SF-12 mental health composite score had a positive association with VS. As hypothesized, participants who used high-risk drugs at baseline had decreased odds of VS over time compared to those who did not use high-risk drugs (adjusted odds ratio 0.47, 95% confidence interval 0.31-0.70,  $p < 0.001$ ).

The longitudinal correlation of average VS with average frequency of high-risk drug use over the three follow-up visits was  $-0.55$  ( $p = 0.001$ ). When examining associations of individual substances with VS, this correlation appears to be driven by stimulant use. The longitudinal correlations of average VS with use of individual substances were: methamphetamine use,  $-0.34$  ( $p = 0.03$ ); cocaine use,  $-0.30$  ( $p = 0.07$ ); opioid use,  $-0.19$  ( $p = 0.32$ ); and binge alcohol use,  $0.33$  ( $p = 0.07$ ).

## 4. Discussion

The present study showed that the recent use of high-risk drugs, especially methamphetamines, was negatively associated with VS over 12 months in this population of PLWH released from a large municipal jail in Los Angeles. This longitudinal correlation was robust in analyses that adjusted for the significant effect of the intervention on VS over the same time period. The findings are important because efforts to end the HIV epidemic recognize the imperative of addressing the CJS, as gaps in care and loss of viral control occur frequently after release from correctional institutions (Iroh et al., 2015). Approximately 2.2 million Americans were estimated to be incarcerated in jails and prisons at the end of 2016, while an estimated 9 million are released from jails alone each year with a vastly disproportionate effect on Blacks and, to a lesser extent, Hispanics (Kaeble et al., 2018).

In the setting of the LINK LA clinical trial, which was conducted among PLWH released from the largest jail system in the US to test the effects of a theory-driven peer navigation intervention, we observed a close association of drug use with VS over and above the statistically significant effect of LINK LA on sustaining VS following release from custody. This longitudinal correlation highlights the importance of high-risk drug use as a barrier to

sustained VS in PLWH after release from incarceration, even in the presence of supportive interventions. It also underscores the community reentry period as a time when individuals are at increased risk of substance use-associated harms. During this period substance use can disorganize health-oriented behaviors like seeking HIV medical care, maintaining ART adherence, and securing social determinants of health, such as stable housing (Wooditch et al., 2018), that support sustained VS. Given the additional negative and cumulative influences of incarceration experiences on ART adherence over the lifetime (Milloy et al., 2011; Nosyk et al., 2014), future linkage-to-care interventions for PLWH post-incarceration may be enhanced by devising additional strategies to mitigate substance use.

In this context, an important finding was that the prevalence of high-risk drug use overall in LINK LA was 71%. The prevalence of methamphetamines in particular was 58%— much greater in this CJS sample than that found in other recent studies of substance use and the continuum of care among PLWH in the US (Gardner et al., 2016; Nolan et al., 2017). In a number of other studies of drug use and HIV care in criminal justice settings, cocaine and opiates, especially injected opiates such as heroin, have predominated (Althoff et al., 2013; Chitsaz et al., 2013; Vagenas et al., 2016). Pharmacotherapy with long-acting opioid antagonists prior to release from custody has been shown to be effective for reducing the disorganizing influences of opioid and alcohol use disorders in PLWH (Springer et al., 2018a; 2018b). However, in LINK LA, the prevalence of opiate use was fairly small, while methamphetamine use was common. Thus, it is unfortunate that no comparable pharmacotherapy for stimulant use is yet available (Bhatt et al., 2016; Castells et al., 2016) and surprising that PWIDs exhibited higher levels of viral suppression than other groups in our study. There is mixed support for agonist approaches to methamphetamine use disorder (Lee et al., 2018), but uptake may be limited in people involved with criminal justice systems. It is encouraging that contingency management is successfully being scaled up in treating stimulant use disorders in Veterans Affairs settings. This work provides key directions for research and programs targeting stimulant use during transitions into and out of jail settings (DePhilippis et al., 2018).

Our findings also point to discrete demographic and behavioral sub-groups in Los Angeles populations with CJS involvement who use distinct types of high-risk drugs, including methamphetamine, cocaine, heroin, and prescription opioids. Many of the differences in overall high-risk drug use were driven by the high prevalence of methamphetamine use among White, MSM, and TGW participants, whereas cocaine use was most common in Blacks. This mirrors substance use patterns found in vulnerable PLWH in the general populations of Los Angeles (Aralis et al., 2018; Brecht, 2018; Reback and Fletcher, 2014). The higher rates of high-risk drug use in the White study participants exemplifies a distinct type of criminal justice disparity, documented in previous research, that results in higher likelihoods of incarceration and longer sentences for Blacks than Whites arrested for similar charges (Iguchi et al, 2005; Rehavi and Starr, 2014; Steffensmeier et al., 2016). These disparities result in a concentration of Whites with more serious substance use in criminal justice settings. Previous research in vulnerable PLWH has also documented strong associations of type of substance use with access to the continuum of care for HIV (Sohler et al., 2007). Future interventions will need to accommodate racial differences in particular substance use patterns.



Mental health at baseline, as measured by the SF-12, was more than a full standard deviation below the US national norm, but better mental health scores independently predicted VS at 12 months. Mental health issues among PLWH in correctional settings are common. For example, in the EnhanceLink study— a cohort study of 878 PLWH leaving one of 20 jails— a high mental illness/emotional distress score as measured by Addiction Severity Index was found in 52.2% (Lincoln et al., 2015). In one study, PLWH who screened positive for depression prior to release from jail were more likely to anticipate needing income, housing, transportation, and assistance with adherence post-release than those who did not (Scheyett et al., 2010). In contrast to our findings, Loeliger et al. found that PLWH with higher severity scores of psychiatric need and untreated psychiatric illness during incarceration had higher levels of viral suppression than PLWH with a lower psychiatric need severity score over a 3-year period (Loeliger et al., 2018). Differences in measurements (SF-12 vs. psychiatric severity score) and study design (RCT vs. retrospective) make direct comparisons difficult. Yet, further study in measuring psychiatric need and treatment is necessary, as on face value treatment of mental health problems in PLWH prior to and after release should be helpful in reducing both continued substance use and increasing PLWH's chances of achieving and maintaining VS in the community (Meyer et al., 2011; Peters et al., 2017). Although not specific to PLWH, in a recent meta-analysis of 37 RCTs conducted in prisons worldwide to test several behavioral (non-pharmacological) therapies for common psychological conditions (e.g., depression and anxiety), cognitive behavioral therapy and mindfulness-based therapies had modest effects and should be tested in PLWH with a co-existing mental health disorder (Yoon et al., 2017). In PLWH, pharmacotherapy for alcohol and opioid use have recently shown promise in people with a co-existing mental illness (Robertson et al., 2018a; 2018b) reducing mental health hospitalizations and emergency department visits and improving psychotropic medication adherence after release from incarceration.

There were several limitations to this analysis. It was performed in the setting of a clinical trial with a behavioral intervention; hence, generalizability of findings to CJS settings without a linkage to care intervention may be limited. Nevertheless, both intervention and control participants were included in the analysis, and intervention assignment was controlled for. Due to small numbers of opioid users, we were underpowered to show significant associations with VS for this sub-group. Self-report bias was a possibility, as substance use data was collected via self-report and biologic testing was not performed. Additionally, we did not collect measures of diagnosis or severity of substance use disorders. We also acknowledge the potential for there to be many other factors between the relationship of high-risk drug use and viral suppression that include housing status, reincarceration, and therapy for drug use, among others. We did not control for these factors, as they are likely intermediates rather than confounders in the relationship between high-risk drug use and viral suppression. The strengths of this study include randomized controlled design with prospectively collected longitudinal, repeated measures, biological VS (HIV RNA) outcome data, and statistical methods to account for changes in substance use over time co-modeled with VS.

In summary, post-incarcerated PLWH who used high-risk drugs, especially methamphetamines, experienced decreased levels of VS over time, independent of peer

navigation services that maintained levels of VS. In this population, efforts to curtail stimulant use are urgently needed, given the high prevalence of and negative association with use. Future intervention research to improve VS and the overall health of post-incarcerated PLWH should address geographical and racial/ethnic differences in substances used.

## Acknowledgements

We would like to thank our participants and research assistants as well as our partners at the Los Angeles County Department of Public Health Mario Perez, Jennifer Sayles, Trista Bingham, Jane Rohde Bowers, Sonali Kulkarni, and Kim Bui, without whose contributions this study would not have been possible. We also thank Jimmy Ngo and J. Marcos Gracilazo for their tireless administrative support. Finally, we recognize at the Los Angeles County Sheriffs Department for their support for this study.

### Role of Funding Source

Dr. Goodman-Meza was supported by the a T32 NIMH Grant DA-2T32MH080634-11 (PI: Currier and Gorbach). Support for the primary collection of data was provided by a grant from NIH/NIDA, R01 DA030781 (PI, Cunningham). Additional support for time on this analysis (for Weiss, Shoptaw, Harawa, Cunningham) was provided by the Center for HIV Identification, Prevention, and Treatment (CHIPTS) NIMH grant P30-MH58107 (for Cunningham, Harawa) through UCLA Clinical and Translational Science Institute (CTSI) NIH/NCATS grants UL1-TR001881 (PI, Dubinett) and (for Weiss) the UCLA Center for AIDS Research (CFAR) grant 5P30AI028697, Core H. Dr. Cunningham's time was also supported in part by NIDA R01-DA039934 (MPIs, Schneider, Harawa, Kayo); NIMH R01-MH103076 (PI, Cunningham); NIA P30-AG021684 (RCMAR/CHIME; PI Mangione); NIMHD R01-MD011773 (PI, Cunningham); NINR Grants: R01-NR017334 (PI, Bogart), and R01-NR4014789 (PI, Ford and TL1-TR001883 (PI, Ettner). Dr. Takada's time was support by the VA Office of Academic Affiliations through the National Clinician Scholars Program. The contents do not represent the views of the U.S. Department of Veterans Affairs or the United States Government. The funders had no role in the design and conduct of the study; collection, management, analysis, and interpretation of the data; preparation, review, or approval of the manuscript; and decision to submit the manuscript for publication.

## References

- Ammon B, Iroh P, Tiruneh Y, Li X, Montague BT, Rich JD, Nijhawan AE, 2018 HIV care after jail: Low rates of engagement in a vulnerable population. *J. Urban Health* 4, 1–11.
- Bhatt M, Zielinski L, Baker-Beal L, Bhatnagar N, Mouravska N, Laplante P, Worster A, Thabane L, Samaan Z, 2016 Efficacy and safety of psychostimulants for amphetamine and methamphetamine use disorders: A systematic review and meta-analysis. *Syst. Rev* 5, 189. [PubMed: 27842569]
- Carrico AW, Shoptaw S, Cox C, Stall R, Li X, Ostrow DG, Vlahov D, Plankey MW, 2014 Stimulant use and progression to AIDS or mortality after the initiation of highly active antiretroviral therapy. *J. Acquir. Immune Defic. Syndr* 67, 508–513. [PubMed: 25271387]
- Castells X, Cunill R, Perez-Mana C, Vidal X, Capella D, 2016 Psychostimulant drugs for cocaine dependence. *Cochrane Database Syst. Rev* 9, CD007380. [PubMed: 27670244]
- Chitsaz E, Meyer JP, Krishnan A, Springer SA, Marcus R, Zaller N, Jordan AO, Lincoln T, Flanigan TP, Porterfield J, Altice FL, 2013 Contribution of substance use disorders on HIV treatment outcomes and antiretroviral medication adherence among HIV-infected persons entering jail. *AIDS Behav.* 17 Suppl 2, S118–1127. [PubMed: 23673792]
- Cunningham WE, Weiss RE, Nakazono T, Malek MA, Shoptaw SJ, Ettner SL, Harawa NT, 2018 Effectiveness of a peer navigation intervention to sustain viral suppression among HIV-positive men and transgender women released from jail: The LINK LA randomized clinical trial. *JAMA Intern. Med* 178, 542–553. [PubMed: 29532059]
- DePhilippis D, Petry NM, Bonn-Miller MO, Rosenbach SB, McKay JR, 2018 The national implementation of Contingency Management (CM) in the Department of Veterans Affairs: Attendance at CM sessions and substance use outcomes. *Drug Alcohol Depend.* 185, 367–373. [PubMed: 29524874]
- Dowling-Guyer S, Johnson ME, Fisher DG, Needle R, Watters J, Andersen M, Williams M, Kotranski L, Booth R, Rhodes F, Weatherby N, Estrada AL, Fleming D, Deren S, Tortu S, 2016 Reliability of


drug users' self-reported HIV risk behaviors and validity of self-reported recent drug use. *Assessment*. 1, 383–392.

- Gardner LI, Marks G, Shahani L, Giordano TP, Wilson TE, Drainoni ML, Keruly JC, Batey DS, Metsch LR, 2016 Assessing efficacy of a retention-in-care intervention among HIV patients with depression, anxiety, heavy alcohol consumption and illicit drug use. *AIDS*. 30, 1111–1119. [PubMed: 26760454]
- Hara M, Huang DYC, Weiss RE, Hser Y-I, 2013 Concurrent life-course trajectories of employment and marijuana-use: Exploring interdependence of longitudinal outcomes. *J. Subst. Abuse. Treat* 45, 426–432. [PubMed: 23890771]
- Harawa NT, Amani B, Rohde Bowers J, Sayles JN, Cunningham W, 2017 Understanding interactions of formerly incarcerated HIV-positive men and transgender women with substance use treatment, medical, and criminal justice systems. *Int. J. Drug Policy*. 48, 63–71 [PubMed: 28804052]
- Ion A, Cai W, Elston D, Pullenayegum E, Smaill F, Smieja M, 2011 A comparison of the MOS-HIV and SF-12v2 for measuring health-related quality of life of men and women living with HIV/ AIDS. *AIDS Res. Ther* 8, 5. [PubMed: 21272358]
- Iroh PA, Mayo H, Nijhawan AE, 2015 The HIV care cascade before, during, and after incarceration: A systematic review and data synthesis. *Am. J. Public Health*. 105, e5–16.
- Johnson ME, Fisher DG, Montoya I, Booth R, Rhodes F, Andersen M, Zhuo Z, Williams M, 2000 Reliability and validity of not-in-treatment drug users' follow-up self-reports. *AIDS Behav*. 4, 373–380.
- Kaebler D, Cowhig M, BJS Statisticians, 2018 Correctional populations in the United States, 2016 Bureau of Justice Statistics, Washington, D.C.
- Kay ES, Batey DS, Mugavero MJ, 2016 The HIV treatment cascade and care continuum: Updates, goals, and recommendations for the future. *AIDS Res. Ther* 13, 35. [PubMed: 27826353]
- Korthuis PT, Edelman EJ, 2018 Substance use and the HIV care continuum: Important advances. *Addict. Sci. Clin. Pract* 13.
- Lazar R, Kersanske L, Xia Q, Daskalakis D, Braunstein SL, 2017 Hospitalization rates among people with HIV/AIDS in New York City, 2013. *Clin. Infect. Dis* 65, 469–476. [PubMed: 28444155]
- Lee NK, Jenner L, Harney A, Cameron J, 2018 Pharmacotherapy for amphetamine dependence: A systematic review. *Drug Alcohol Depend*. 191, 309–337. [PubMed: 30173086]
- Lesko CR, Tong W, Moore RD, Lau B, 2017 Retention, antiretroviral therapy use and viral suppression by history of injection drug use among HIV-infected patients in an urban HIV clinical cohort. *AIDS Behav*. 21, 1016–1024. [PubMed: 27752872]
- Lincoln T, Simon-Levine D, Smith J, Donenberg GR, Springer SA, Zaller N, Altice FL, Moore K, Jordan AO, Draine J, Desabrais M, 2015 Prevalence and predictors of mental/emotional distress among HI V+ jail detainees at enrollment in an observational study. *J. Correct. Health Care*. 21, 125–139. [PubMed: 25788608]
- Loeliger KB, Meyer JP, Desai MM, Ciarleglio MM., Gallagher C, Altice FL, 2018 Retention in HIV care during the 3 years following release from incarceration: A cohort study. *PLoS Med*. 15, e1002667.
- Meyer JP, Chen NE, Springer SA, 2011 HIV Treatment in the criminal justice system: Critical knowledge and intervention gaps. *AIDS Res. Treat*. 2011, 1–10.
- Nance RM, Delaney JAC, Simoni JM, Wilson IB, Mayer KH, Whitney BM, Aunon FM, Safren SA, Mugavero MJ, Mathews WC, Christopoulos KA, Eron JJ, Napravnik S, Moore RD, Rodriguez B, Lau B, Fredericksen RJ, Saag MS, Kitahata MM, Crane HM, 2018 HIV viral suppression trends over time among HIV-infected patients receiving care in the United States, 1997 to 2015: A cohort study. *Ann. Intern. Med* 169, 376–384. [PubMed: 30140916]
- Nolan S, Walley AY, Heeren TC, Patts GJ, Ventura AS, Sullivan MM, Samet JH, Saitz R, 2017 HIV-infected individuals who use alcohol and other drugs, and virologic suppression. *AIDS Care*. 29, 1129–1136. [PubMed: 28513200]
- Palepu A, Tyndall MW, Chan K, Wood E, Montaner JSG, Hogg RS, 2004 Initiating highly active antiretroviral therapy and continuity of HIV care: The impact of incarceration and prison release on adherence and HIV treatment outcomes. *Antivir. Ther. (Lond.)* 9, 713–719. [PubMed: 15535408]

- Peters RH, Young MS, Rojas EC, Gorey CM, 2017 Evidence-based treatment and supervision practices for co-occurring mental and substance use disorders in the criminal justice system. *Am. J. Drug Alcohol Abuse.* 43, 475–488. [PubMed: 28375656]
- Pyne JM, Tripathi S, French M, McCollister K, Rapp RC, Booth BM, 2011 Longitudinal association of preference-weighted health-related quality of life measures and substance use disorder outcomes. *Addiction.* 106, 507–515. [PubMed: 21205046]
- Qian H-Z, Mitchell VJ, Bebawy S, Cassell EL, Perez G, McGowan CC, Sterling TR, Vermund SH, D'Aquila R, Hulgan T, 2014 Current drug use and lack of HIV virologic suppression: Point-of-care urine drug screen versus self-report. *BMC Infect. Dis* 14.
- Rasbach DA, Desruisseau AJ, Kipp AM, Stinnette S, Kheshti A, Shepherd BE, Sterling TR, Hulgan T, McGowan CC, Qian H-Z, 2013 Active cocaine use is associated with lack of HIV-1 virologic suppression independent of nonadherence to antiretroviral therapy: Use of a rapid screening tool during routine clinic visits. *AIDS Care.* 25, 109–117. [PubMed: 22670566]
- Scheyett A, Parker S, Golin C, White B, Davis CP, Wohl D, 2010 HIV-infected prison inmates: Depression and implications for release back to communities. *AIDS Behav.* 14, 300–307. [PubMed: 18709452]
- Sohler NL, Wong MD, Cunningham WE, Cabral H, Drainoni M-L, Cunningham CO, 2007 Type and pattern of illicit drug use and access to health care services for HIV-infected people. *AIDS Patient Care and STDs.* 21 Suppl. 1, S68–76. [PubMed: 17563292]
- Ware X, Kosinski M, Keller SD, 1996 A 12-Item short-form health survey: Construction of scales and preliminary tests of reliability and validity. *Med. Care.* 34, 220–233. [PubMed: 8628042]
- Weiss RD, Najavits LM, Greenfield SF, Soto JA, Shaw SR, Wyner D, 1998 Validity of substance use self-reports in dually diagnosed outpatients. *Am. J. Psychiatry.* 155, 127–128. [PubMed: 9433351]
- Weiss RE, 2005 Modeling longitudinal data. Springer New York, New York, NY.
- Westergaard RP, Hess T, Astemborski J, Mehta SH, Kirk GD, 2013 Longitudinal changes in engagement in care and viral suppression for HIV-infected injection drug users. *AIDS.* 27, 2559–2566. [PubMed: 23770493]
- Westergaard RP, Kirk GD, Richesson DR, Galai N, Mehta SH, 2011 Incarceration predicts virologic failure for HIV-infected injection drug users receiving antiretroviral therapy. *Clin. Infect. Dis* 53, 725–731. [PubMed: 21890777]
- Wooditch A, Mbaba M, Kiss M, Lawson W, Taxman F, Altice FL, 2018 Housing experiences among opioid-dependent, criminal justice-involved individuals in Washington, D.C. *J. Urban Health* 95, 61–70. [PubMed: 28550643]
- Yoon IA, Slade K, Fazel S, 2017 Outcomes of psychological therapies for prisoners with mental health problems: A systematic review and meta analysis. *J. Consult Clin. Psychol* 85, 783–802. [PubMed: 28569518]

### Highlights

- In Link LA, methamphetamine was the most prevalent drug used.
- Non-Hispanic Whites were most likely to report methamphetamine use.
- Non-Hispanic Blacks were more likely to report cocaine use.
- Any hard drug use was independently associated with decreased VS (viral suppression).
- Methamphetamine was the only substance independently associated with decreased VS.



**Figure 1.** Viral suppression by study arm and high-risk drug use over the course of the study.

**Table 1.** Baseline characteristics prior to release from jail for participants in LINK LA by type of substance used (n=356).

Characteristics	Total % (n)	High-Risk Drug Use <sup>b</sup> % (n)	Methamphetamine % (n)	Cocaine % (n)	Heroin % (n)	Prescription Opioids % (n)	Binge Alcohol % (n)
<b>Total</b>	100 (356)	71 (253)	58 (206)	17 (61)	7 (24)	4 (15)	25 (88)
<b>Study Arm</b>							
Intervention	51 (180)	67 (120)	58 (104)	14 (25)	8 (15)	6 (10)	23 (42)
Control	49 (176)	76 (133)	58 (102)	20 (36)	5 (9)	3 (5)	26 (46)
<b>Age</b>							
18-34 years	38 (137)	74 (101)	69 (95)***	10 (14)**	9 (13)	3 (4)	32 (44)*
35 or more	62 (219)	69 (152)	51 (111)	21 (47)	5 (11)	5 (11)	20 (44)
<b>Race/ethnicity</b>							
White, non-Hispanic	27 (95)	79 (75)*	74 (70)***	6 (6)***	7 (7)	7 (7)*	24 (23)
Black, non-Hispanic	42 (151)	63 (95)	40 (60)	26 (40)	4 (6)	1 (2)	23 (35)
Hispanic	31 (110)	75 (83)	69 (76)	14 (15)	10 (11)	5 (6)	27 (30)
<b>Risk/gender group</b>							
MSM	56 (201)	80 (160)***	70 (140)***	12 (24)**	4 (9)	3 (7)	22 (44)
PWID	12 (43)	70 (30)	51 (22)	23 (10)	21 (9)	7 (3)	30(13)
TGW	15 (53)	77 (41)	58 (31)	32 (17)	9 (5)	6 (3)	30 (16)
None of the above	17 (59)	37 (22)	22 (13)	18 (10)	2 (1)	3 (2)	25 (15)
<b>Educational attainment</b>							
Less than high school	37 (131)	74 (97)	62 (81)	17 (22)	11 (14)*	2 (3)	30 (39)
High school or more	63 (224)	69 (155)	55 (124)	17 (39)	4 (10)	5 (12)	21 (48)
<b>Annual income</b>							
\$10,000	42 (150)	69 (103)	55 (82)	19 (29)	8 (12)	1 (1)**	23 (35)
\$10,001 or more	58 (205)	73 (149)	60 (123)	16 (32)	6 (12)	7 (14)	25 (52)
<b>Insurance</b>							
Uninsured	55 (191)	70 (134)	58 (110)	18 (34)	8 (15)	4 (7)	28 (53)

Characteristics	Total % (n)	High-Risk Drug Use <sup>b</sup> % (n)	Methamphetamine % (n)	Cocaine % (n)	Heroin % (n)	Prescription Opioids % (n)	Binge Alcohol % (n)
Insured	45 (155)	72 (111)	57 (88)	17 (26)	5 (8)	5 (7)	22 (34)
<b>Homeless</b>							
Homeless	56 (198)	79 (156)***	64 (126)*	21 (41)*	10 (19)*	5 (10)	26 (52)
Not homeless	44 (158)	61 (97)	51 (80)	13 (20)	3 (5)	3 (5)	23 (36)
<b>Reincarceration</b>							
Reincarcerated	54 (192)	73 (141)	64 (123)	16 (31)	6 (12)	3 (6)	27 (51)
Number of times reincarcerated, mean (±SD)	1.3 (1.8)	1.3 (1.8)	1.4 (1.9)	1.3 (1.9)	1.3 (1.7)	1.1 (2.6)	1.2 (1.6)
Number of days per reincarceration, mean (±SD)	131 (147)	129 (150)	126 (151)	151 (138)	102 (86)	264 (222)	147 (177)
<b>Treatment visits</b>							
For substance use problems	15 (53)	81 (43)	60 (32)	28 (15)	6 (17)	6 (3)	23 (12)
For psychiatric/emotional problems	32 (112)	67 (75)	49 (55)	18 (20)	4 (5)	8 (9)	28 (31)
<b>Short Form-12</b>							
Mental health, mean (±SD)	38 (12)	38 (12)	37 (12)	38 (12)	34 (14)	35 (12)	34 (10)
Physical health, mean (±SD)	51 (10)	51 (10)	52 (9)	48 (11)	52 (9)	43 (13)	52 (10)
<b>CD4 count (cell/μL)<sup>d</sup></b>							
<500	51 (180)	72 (130)	55 (99)	18 (32)	7 (13)	6 (11)	27 (48)
500 or more	49 (175)	70 (122)	61 (106)	17 (29)	6 (11)	2 (4)	23 (40)
<b>Viral suppression at baseline<sup>c</sup></b>							
Suppressed	52 (184)	68 (126)	55 (101)	17 (32)	4 (8)	4 (8)	27 (50)
Not suppressed	48 (172)	74 (127)	61 (105)	17 (29)	9 (16)	4 (7)	22 (38)

Abbreviations: MSM, men who have sex with men; PWID, people who inject drugs; TGW, transgender women; SD, standard deviation.

Chi-square test was used to test for statistically significant differences between users and non-users within each substance use type.

\* p 0.05,

\*\* p 0.01,

\*\*\* p 0.001



Author Manuscript

Author Manuscript

Author Manuscript

Author Manuscript

<sup>a</sup>n=355 due to missing values

<sup>b</sup>*High-risk drug use* was defined as any reported use of heroin, cocaine (powder or crack), methamphetamines or misuse of prescription opioids in the 30 days prior to being arrested (used at baseline) and 30-days prior to each follow-up assessment.

<sup>c</sup>HIV viral suppression was defined as an undetectable viral load of less than 75 copies/mL.

**Table 2.**

Multivariable analysis of substance use over 12 months in LINK LA

Variable	High-Risk Drug Use <sup>a</sup>			Methamphetamine Use			Cocaine Use			Opioid Use		
	AOR	95% CI	P-value	AOR	95% CI	P-value	AOR	95% CI	P-value	AOR	95% CI	P-value
<b>Age group</b>												
34 or less				2.34	1.26-3.34	<0.01						
35 or more (reference)												
<b>Race/ethnicity</b>												
Hispanic	0.74	0.39-1.39	0.35	0.45	0.20-0.98	0.04	2.11	0.75-5.92	0.16			
Black, non-Hispanic	0.52	0.28-0.96	0.04	0.17	0.08-0.37	<0.001	5.44	2.04-14.5	<0.001			
White, non-Hispanic (reference)												
<b>Gender/HIV risk groups</b>												
MSM	8.45	4.14-17.3	<0.001	14.9	5.63-39.6	<0.001				3.39	0.81-14.2	0.10
PWID	4.86	1.89-12.5	0.01	4.66	1.36-16.0	0.01				18.6	3.34-104	<0.001
TGW	8.09	3.40-18.9	<0.001	9.54	3.07-29.6	<0.001				2.73	0.47-15.7	0.25
None of the above (reference)												

**Abbreviations:** AOR, adjusted odds ratio; CI, confidence interval; MSM, men who have sex with men; PWID, person who injects drugs; TGW, transgender women.

All models adjusted for the intervention, time, and intervention by time interaction. Variable that did not attain statistical significance (p<0.05) in univariate analysis have been omitted.

<sup>a</sup> *High-risk drug use* was defined as any reported use of heroin, cocaine (powder or crack), methamphetamines or misuse of prescription opioids in the 30 days prior to being arrested (used at baseline) and 30-days prior to each follow-up assessment.

**Table 3.**

Univariate and multivariable predictors of HIV viral suppression<sup>a</sup> over 12 months in LINK LA participants.

	Univariate GLMM models				Multivariate GLMM model			
		95% CI		P value		95% CI		P value
	OR	Lower	Upper	P value	AOR	Lower	Upper	P value
<b>Gender/HIV risk groups</b>								
MSM	1.04	0.58	1.89	0.89	1.25	0.68	2.30	0.47
TGW	0.56	0.26	1.19	0.13	0.70	0.32	1.51	0.36
PWID	2.03	0.89	4.61	0.09	2.29	1.00	5.24	0.05
None of the above (reference)								
<b>Income less than \$10,000</b>								
Yes	0.53	0.34	0.82	<0.01	0.57	0.37	0.89	0.01
No (reference)								
<b>CD4 less than 500 cell/μL</b>								
Yes	0.43	0.28	0.66	<0.001	0.44	0.28	0.67	<0.001
No (reference)								
<b>SF-12 (per unit increase)</b>								
Mental	1.03	1.02	1.05	<0.001	1.02	1.01	1.04	<0.01
Physical	1.03	1.01	1.04	<0.01				
<b>High risk drug use<sup>b</sup></b>								
Yes	0.66	0.41	1.06	0.09	0.47	0.31	0.70	<0.001
No (reference)								

**Abbreviations:** AOR, adjusted odds ratio; GLMM, generalized liner mixed models; MSM, men who have sex with men; OR, odds ratio; PWID, people who inject drugs; SF-12, 12-item Short Form Health Survey; TGW, transgender women.

Models were adjusted for the intervention, time and intervention by time interaction.

Variables that did not attain statistical significance ( $p < 0.10$ ) in univariate analysis have been omitted

<sup>a</sup>HIV viral suppression was defined as an undetectable viral load of less than 75 copies/mL.

<sup>b</sup>High-risk drug use was defined as any reported use of heroin, cocaine (powder or crack), methamphetamines or misuse of prescription opioids in the 30 days prior to being arrested (used at baseline) and 30-days prior to each follow-up assessment.