

UCLA

UCLA Previously Published Works

Title

Robust CRISPR/Cas9 mediated genome editing and its application in manipulating plant height in the first generation of hexaploid Ma bamboo (*Dendrocalamus latiflorus* Munro)

Permalink

<https://escholarship.org/uc/item/5pc5w2ct>

Journal

Plant Biotechnology Journal, 18(7)

ISSN

1467-7644

Authors

Ye, Shanwen
Chen, Gang
Kohnen, Markus V
[et al.](#)

Publication Date

2020-07-01

DOI

10.1111/pbi.13320

Peer reviewed



Robust CRISPR/Cas9 mediated genome editing and its application in manipulating plant height in the first generation of hexaploid Ma bamboo (*Dendrocalamus latiflorus* Munro)

Journal:	<i>Plant Biotechnology Journal</i>
Manuscript ID	PBI-00886-2019.R1
Wiley - Manuscript type:	Brief Communications
Date Submitted by the Author:	29-Oct-2019
Complete List of Authors:	Ye, ShanWen; Fujian Agriculture and Forestry University, Basic Forestry and Proteomics Center (BFPC), Fujian Provincial Key Laboratory of Haixia Applied Plant Systems Biology, College of Forestry Chen, Gang; Fujian Agriculture and Forestry University, Basic Forestry and Proteomics Center (BFPC), Fujian Provincial Key Laboratory of Haixia Applied Plant Systems Biology, College of Forestry Kohnen, Markus; Fujian Agriculture and Forestry University, Basic Forestry and Proteomics Center (BFPC), Fujian Provincial Key Laboratory of Haixia Applied Plant Systems Biology, College of Forestry Wang, WenJia; Fujian Agriculture and Forestry University, Basic Forestry and Proteomics Center (BFPC), Fujian Provincial Key Laboratory of Haixia Applied Plant Systems Biology, College of Forestry Cai, Changyang; Fujian Agriculture and Forestry University, Basic Forestry and Proteomics Center (BFPC), Fujian Provincial Key Laboratory of Haixia Applied Plant Systems Biology, College of Forestry Ding, WenSha; Fujian Agriculture and Forestry University, Basic Forestry and Proteomics Center (BFPC), Fujian Provincial Key Laboratory of Haixia Applied Plant Systems Biology, College of Forestry Wu, Chu; Fujian Agriculture and Forestry University, Basic Forestry and Proteomics Center (BFPC), Fujian Provincial Key Laboratory of Haixia Applied Plant Systems Biology, College of Forestry Gu, Lianfeng; Fujian Agriculture and Forestry University, Basic Forestry and Proteomics Center (BFPC), Fujian Provincial Key Laboratory of Haixia Applied Plant Systems Biology, College of Forestry, Zheng, Yushan; Fujian Agriculture and Forestry University, Basic Forestry and Proteomics Center (BFPC), Fujian Provincial Key Laboratory of Haixia Applied Plant Systems Biology, College of Forestry Ma, Xiangqing; Fujian Agriculture and Forestry University, Basic Forestry and Proteomics Center (BFPC), Fujian Provincial Key Laboratory of Haixia Applied Plant Systems Biology, College of Forestry Lin, Chentao; University of California Los Angeles Department of Molecular Cell and Developmental Biology Zhu, Qiang; Fujian Agriculture and Forestry University, Basic Forestry and Proteomics Center (BFPC), Fujian Provincial Key Laboratory of Haixia Applied Plant Systems Biology, College of Forestry
Keywords:	Bamboo, Genome editing, CRISPR/Cas9

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60



SCHOLARONE™
Manuscripts

1
2
3
4 **Title: Robust CRISPR/Cas9 mediated genome editing and its**
5
6 **application in manipulating plant height in the first generation of**
7
8 **hexaploid Ma bamboo (*Dendrocalamus latiflorus* Munro)**
9
10

11
12
13 Shanwen Ye¹, Gang Chen¹, Markus V. Kohnen¹, Wenjia Wang¹, Changyang Cai¹,
14 WenSha Ding¹, Chu Wu¹, Lianfeng Gu¹, Yushan Zheng¹, Xiangqing Ma¹, Chentao
15 Lin² and Qiang Zhu^{1,*}
16
17

18
19 ¹ *Basic Forestry and Proteomics Center (BFPC), Fujian Provincial Key Laboratory*
20 *of Haixia Applied Plant Systems Biology, College of Forestry, Fujian Agriculture and*
21 *Forestry University, Fujian, China;*
22

23
24
25 ²*Department of Molecular, Cell & Developmental Biology, University of California,*
26 *Los Angeles, California, USA*
27

28
29
30
31
32 Emails:

33
34 Shanwen Ye, yeshanwen@fafu.edu.cn;

35
36 Gang Chen, GangC@163.com;

37
38 Markus V. Kohnen, MKohnen@fafu.edu.cn;

39
40 Wenjia Wang, 3160808027@fafu.edu.cn;

41
42 Changyang Cai, caichangyang@fafu.edu.cn;

43
44 Wensha Ding, 1170428008@fafu.edu.cn;

45
46 Chu Wu, 1170538006@fafu.edu.cn;

47
48 Lianfeng Gu, lfgu@fafu.edu.cn;

49
50 Yushan Zheng, zys1960@163.com;

51
52 Xiangqing Ma, lxxmxq@126.com;

53
54 Chentao Lin, clin@mcdub.ucla.edu;

55
56 Qiang Zhu, zhuqiang@fafu.edu.cn (*Corresponding author).
57
58
59
60

1
2
3
4 **Dear Editor,**

5
6 Bamboo is a special grass to human due to its great economic and ecological values.
7
8 Around 2.5 billion people are directly producing and consuming bamboo, and its
9
10 international trade reached 68.8 billion US dollars in 2018 (Data from International
11
12 Bamboo and Rattan Organization). One major bamboo species in Asia is Ma bamboo
13
14 (*Dendrocalamus latiflorus Munro*), which is a hexaploid species with three
15
16 subgenomes (2n=72, AABBCC) (Guo et al., 2019). Despite its agronomic importance,
17
18 it is nearly impossible to modify bamboo traits by traditional breeding as it takes over
19
20 70 years to flower. Bamboo research largely lagged behind due to the lack of efficient
21
22 genetic manipulation tools.
23

24
25 The CRISPR (Clustered Regularly Interspaced Short Palindromic Repeat)/Cas9
26
27 provides straight-forward ways for genome editing in many plants (Yin et al., 2017),
28
29 but has never been applied in bamboo. Here, we reported the generation of bamboo
30
31 mutants with CRISPR/Cas9 technology by targeting one specific copy or all
32
33 homoeologous genes.
34

35
36 Since our recently established genetic transformation protocol is time-consuming
37
38 (~1.5 years) (Ye et al., 2017), we optimized the CRISPR/Cas9 system in bamboo
39
40 protoplast. We first improved the protoplast preparation methods, and could isolate
41
42 3.0×10^6 protoplasts/g fresh leaves. Next, we improved the PEG-mediated
43
44 transformation method, and reached efficiencies of 53.3% for a single plasmid and
45
46 29.8% for two co-transformed plasmids (**Figure 1A**), which is sufficient for
47
48 optimizing the CRISPR/Cas9 system. The maize *UBI* promoter was used to drive
49
50 Cas9 expression (Ye et al., 2017). Three polymerase III-dependent promoters from
51
52 rice (*OsU6a/OsU6b/OsU6c*) were selected to express the sgRNA cassettes (Ma et al.,
53
54 2015), as bamboo exhibits high genomic similarity with rice (Peng et al., 2013). To
55
56 check the effectiveness of CRISPR/Cas9 constructs, a frame-shift mutated *GFP*
57
58 (*mGFP*) containing an additional “guanine” thereby produces no fluorescence signal
59
60

1 was simultaneously co-transformed with CRISPR/Cas9 plasmids (**Figure 1B**).
2 Around 1.8% of the protoplasts transformed with the *UBI-Cas9/OsU6b-sgRNA*
3 construct showed strong signals within 72 h, indicating that the *mGFP* function was
4 restored by the CRISPR/Cas9 system through deleting the additional “guanine”
5 (**Figure 1C**). The *OsU6a* and *OsU6c* promoters work as well, however, with lower
6 efficiency than the *OsU6b* promoter, as positive signals were only occasionally
7 observed with more than 10 replicates. Taking together, the *UBI-Cas9/OsU6b-sgRNA*
8 construct effectively works in bamboo protoplast, and were used for the following
9 endogenous gene editing in Ma bamboo.

10
11 The putative phytoene synthase (*PSY1*) in bamboo, whose homolog in maize
12 functions in carotenoid biosynthesis (Zhu et al., 2016), was selected for the initial test.
13 Three bamboo *PSY1* alleles (*DlmPSY1-A*, *DlmPSY1-B*, *DlmPSY1-C*) were identified
14 and cloned by a homology cloning strategy (**Figure 1D**). To mutate all copies of
15 *DlmPSY1*, sgRNA1 targeting a conserved site among all *DlmPSY1* loci was designed
16 (**Figure 1D**). In addition, the sgRNA2 target site containing 2–3 single-nucleotide
17 polymorphisms (SNPs) in the spacer region among three *DlmPSY1* homoalleles and
18 was selected to test the tolerance of sgRNA mismatches (**Figure 1D**).

19
20 1,600 bamboo calluses induced from stem were transformed as described
21 previously (Ye et al., 2017). In total, 34 independent transgenic lines were confirmed
22 positive (2.1%) by PCR. Based on Sanger sequencing results, 22 (100%) and 10
23 (83.3%) independent T0 lines were edited in the sgRNA1 and sgRNA2 region,
24 respectively (**Figure 1E**), indicating that both constructs effectively induce
25 endogenous gene editing.

26
27 The editing profiles were further analyzed by sequencing. Eighteen lines (81.8%)
28 contained putative homozygote/biallelic mutations in all subgenomes at the sgRNA1
29 target site. In some lines, putative homozygote/biallelic mutations exist in one

1 subgenome while heterozygote or chimeric mutations appear in other subgenomes
2 (T0-10 and T0-26) (**Figure 1F**). Eight mutation types were identified from 590
3 independent clones (**Figure 1G**). The most frequent mutation type was deletion
4 (75%), of which 59.1% are small deletions (<2bp). The ratios of large fragment
5 deletions (≥ 14 bp), insertions, and combined indels were 15.9%, 2.21%, and 7.82%,
6 respectively (**Figure 1G**). Since bamboo propagates through asexual budding, those
7 homozygote/biallelic mutations will remain in the genome of their offspring clones
8 during breeding.

9
10 sgRNA2 that perfectly targets *DlmPSYI-A1*, but not *DlmPSYI-B1* or *DlmPSYI-C1*
11 was designed to study the recognition specificity (**Figure 1D**). Sequencing results
12 confirmed that 10 transgenic lines contain mutations in *DlmPSYI-A1*, but none in
13 *DlmPSYI-B1* and *DlmPSYI-C1* (**Figure 1E**). Two lines (20%) were putative
14 homozygous or biallelic mutations (T0-12 and T0-14), while 7 lines (70%) were
15 heterozygous/chimeric (T0-30 to T0-32 as representative examples, **Figure 1H**). The
16 ratios of deletions, insertions, and combined mutations were 86%, 9%, and 5%,
17 respectively (**Figure 1I**). The mutations were predominantly short nucleotide changes
18 (1–26bp), and 22.7% were 1bp nucleotide deletions (**Figure 1I**). Those data
19 demonstrated the successful application of the CRISPR/Cas9 system in mutating a
20 specific *DlmPSYI* allele.

21
22 Eighteen lines (81.8%) with homozygote/biallelic mutations in all subgenomes at
23 the sgRNA1 site exhibited albino phenotypes (**Figure 1J**), which appeared at an early
24 stage during tissue culture, and persisted at the plantlets stage (**Figure 1J**). Those
25 results suggest that genome editing takes place at an early stage in embryonic cells,
26 and led to the loss-of-function of all *DlmPSYI* alleles. Similar results were reported in
27 rice, wheat or cotton (Wang et al., 2018; Wang et al., 2014; Zhang et al., 2014). In
28 case of sgRNA2, although *DlmPSYI-A* was mutated, no visible phenotypic change
29 was observed due to the existence of the wild-type *DlmPSYI-B* and *DlmPSYI-C*

1 alleles.

2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

1 alleles.

2

3 Next, we applied this technology in bamboo molecular research. Bamboo is the

4 tallest grass in the world, while the underlying mechanism is unknown. Previously,

5 we identified several Gibberellin-responsive genes including *GRG1* (GA-responsive

6 gene 1, *PH01004823G0070*) that potentially acts in controlling bamboo height

7 (Zhang et al., 2018). Here two homozygote *grg1* mutants (efficiency 40%) in Ma

8 bamboo were produced using our optimized CRISPR/Cas9 technology. Mutation in

9 *GRG1* increased plant height (**Figure 1K**), mostly due to elongated internodes

10 (**Figure 1L-N**). Sequencing results confirmed that the *grg1* mutant has the putative

11 homozygous mutation in A1-subgenome, biallelic mutation in B1-subgenome, and

12 homozygous mutation in C1-subgenome (**Figure 1O**), indicating the loss-of-function

13 of *GRG1* in transgenic bamboo. To our knowledge, this is the first example on

14 controlling bamboo height through gene manipulation, which will contribute to

15 subsequent studies on the molecular mechanisms behind the fast growth of bamboo.

16
17 In summary, for the first time we engineered the hexaploid Ma bamboo through

18 CRISPR/Cas9 technology. The homozygote mutations were obtained in the first

19 generation of transgenic lines, which is extremely important for bamboo species due

20 to its long vegetative growth periods. We also confirmed the albino phenotype of

21 *dlmpsy1* mutant in bamboo and generated a bamboo mutant with altered plant height.

22 This demonstrates the applicability of CRISPR/Cas9 in bamboo and thereby boosts

23 future bamboo research and breeding.

24 25 26 27 28 29 30 31 32 33 34 35 36 37 38 39 40 41 42 43 44 45 46 47 48 49 50 51 52 53 54 55 56 57 58 59 60

27 Guo, Z.-H., Ma, P.-F., Yang, G.-Q., Hu, J.-Y., Liu, Y.-L., Xia, E.-H., Zhong, M.-C., Zhao, L., Sun, G.-L., Xu,

28 Y.-X., Zhao, Y.-J., Zhang, Y.-C., Zhang, Y.-X., Zhang, X.-M., Zhou, M.-Y., Guo, Y., Guo, C., Liu,

29 J.-X., Ye, X.-Y., Chen, Y.-M., Yang, Y., Han, B., Lin, C.-S., Lu, Y. and Li, D.-Z. (2019) Genome

30 sequences provide insights into the reticulate origin and unique traits of woody bamboos.

31 *Molecular Plant* DOI:<https://doi.org/10.1016/j.molp.2019.05.009>.

- 1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
- 1 Ma, X., Zhang, Q., Zhu, Q., Liu, W., Chen, Y., Qiu, R., Wang, B., Yang, Z., Li, H., Lin, Y., Xie, Y., Shen, R.,
2 Chen, S., Wang, Z., Guo, J., Chen, L., Zhao, X., Dong, Z. and Liu, Y.G. (2015) A Robust
3 CRISPR/Cas9 System for Convenient, High-Efficiency Multiplex Genome Editing in Monocot
4 and Dicot Plants. *Mol Plant* **8**, 1274-1284.
- 5 Peng, Z., Lu, Y., Li, L., Zhao, Q., Feng, Q., Gao, Z., Lu, H., Hu, T., Yao, N., Liu, K., Li, Y., Fan, D., Guo, Y., Li,
6 W., Weng, Q., Zhou, C., Zhang, L., Huang, T., Zhao, Y., Zhu, C., Liu, X., Yang, X., Wang, T.,
7 Miao, K., Zhuang, C., Cao, X., Tang, W., Liu, G., Liu, Y., Chen, J., Liu, Z., Yuan, L., Huang, X., Lu,
8 T., Fei, B., Ning, Z., Han, B. and Jiang, Z. (2013) The draft genome of the fast-growing
9 non-timber forest species moso bamboo (*Phyllostachys heterocycla*). *Nat Genet* **45**, 456-461.
- 10 Wang, P., Zhang, J., Sun, L., Ma, Y., Xu, J., Liang, S., Deng, J., Tan, J., Zhang, Q., Tu, L., Daniell, H., Jin, S.
11 and Zhang, X. (2018) High efficient multisites genome editing in allotetraploid cotton
12 (*Gossypium hirsutum*) using CRISPR/Cas9 system. *Plant Biotechnol J* **16**, 137-150.
- 13 Wang, Y., Cheng, X., Shan, Q., Zhang, Y., Liu, J., Gao, C. and Qiu, J.L. (2014) Simultaneous editing of
14 three homoeoalleles in hexaploid bread wheat confers heritable resistance to powdery
15 mildew. *Nat Biotechnol* **32**, 947-951.
- 16 Ye, S., Cai, C., Ren, H., Wang, W., Xiang, M., Tang, X., Zhu, C., Yin, T., Zhang, L. and Zhu, Q. (2017) An
17 Efficient Plant Regeneration and Transformation System of Ma Bamboo (*Dendrocalamus*
18 *latiflorus* Munro) Started from Young Shoot as Explant. *Frontiers in plant science* **8**, 1298.
- 19 Yin, K., Gao, C. and Qiu, J.L. (2017) Progress and prospects in plant genome editing. *Nature plants* **3**,
20 17107.
- 21 Zhang, H., Wang, H., Zhu, Q., Gao, Y., Wang, H., Zhao, L., Wang, Y., Xi, F., Wang, W., Yang, Y., Lin, C.
22 and Gu, L. (2018) Transcriptome characterization of moso bamboo (*Phyllostachys edulis*)
23 seedlings in response to exogenous gibberellin applications. *BMC Plant Biol* **18**, 125.
- 24 Zhang, H., Zhang, J., Wei, P., Zhang, B., Gou, F., Feng, Z., Mao, Y., Yang, L., Zhang, H., Xu, N. and Zhu,
25 J.K. (2014) The CRISPR/Cas9 system produces specific and homozygous targeted gene editing
26 in rice in one generation. *Plant Biotechnol J* **12**, 797-807.
- 27 Zhu, J., Song, N., Sun, S., Yang, W., Zhao, H., Song, W. and Lai, J. (2016) Efficiency and inheritance of
28 targeted mutagenesis in maize using CRISPR-Cas9. *J Genet Genomics* **43**, 25-36.

30 **Conflict of interests**

31 The authors declare that they have no conflict of interests.

33 **Fundings**

34 This work was supported by the National Natural Science Foundation of China grant
35 (No.31870660), Fujian Innovative Center for Germplasm Resources and Cultivation
36 of Woody plants (No.125/KLA15001E), Program for scientific and technological
37 innovation team in university of Fujian province (No.118/KLA18069A) to Q.Z. The

1 funding bodies were not involved in the design of the study or in any aspect of the
2 data collection, analysis and interpretation of data and in paper writing.

3 4 **Acknowledgements**

5 We thank Dr. Chella Thaniyarasu from Fujian Agriculture and Forestry University for
6 the critical reading of the manuscript.

7 8 **Authors' contributions**

9 Q.Z. conceived this project; L.F.G., Y.S.Z., X.Q.M. C.T.L. and Q.Z. designed
10 experiments, and interpreted the results; S.W.Y., G.C. and M.V.K. performed the
11 experiments and analyzed the data, W.J.W., C.Y.C., C.W., and D.W.S. helped to
12 perform the experiments and collect the data. All authors read and approved the
13 submission of this manuscript.

14 15 **Figure Legend**

16 **Figure 1. Genome editing in Ma bamboo using CRISPR/Cas9 technology**

17 **A.** Bamboo protoplast isolation and transformation. **a.** Microscopic image of
18 isolated bamboo protoplast transformed with *35S:tdTomato*. **b-d.** Images of
19 bamboo protoplasts co-expressing the fluorescence proteins *tdTomato* (**b**) and
20 GFP (**c**) driven by the 35S promoter, and their overlay (**d**).


21 **B.** CRISPR/Cas9 plasmids for bamboo protoplast. **Top:** CRISPR/Cas9 constructs
22 expressing the sgRNA directed against *mGFP* and driven by
23 *OsU6a/OsU6b/OsU6c* respectively; **Middle:** *mGFP*-expression construct,
24 *mGFP* contains one additional guanine (lower-green case) downstream of the
25 translational start site (red); **bottom:** GFP-expression construct. The sgRNA
26 was designed to produce the presumptive cleavage site at the third nucleotide
27 upstream of the PAM sequence (blue).

28 **C.** Representative bamboo protoplasts co-transfected with *mGFP* and
29 *OsU6b-sgRNA/UBI-Cas9* reproducibly emitting fluorescence signals (red

- 1
2
3
4 1 arrows).
- 5
6 2 **D.** *DlmPSYI* gene structure and sequences of the target sites. Gray boxes: exons;
7 3 black lines: introns; number in brackets: positions of start codon, stop codon
8 4 and sgRNA target sites (red and orange rectangles). The PAM regions (blue),
9 5 SNPs (red), and nucleotide sequences of the *sgRNAs* and *DlmPSYI* genes were
10 6 given.
- 11 7 **E.** Frequencies of the CRISPR/Cas9-induced mutations in two target sites of the
12 8 *DlmPSYI*.
- 13 9 **F.** Representative *DlmPSYI* mutations at the sgRNA1 site. T0-1, T0-2, T0-3,
14 10 T0-12 and T0-13 represent loss-of function mutants. T0-10 and T0-26 lines
15 11 contain heterozygote mutations in the C1-subgenome and chimeric mutations
16 12 in the A1-subgenome, respectively. Red: sgRNA target regions; blue: PAM
17 13 regions; green lowercase letters: nucleotide indels; dotted lines: omitted
18 14 nucleotides.
- 19 15 **G.** Frequencies of indels (left) and mutation types (right) at the sgRNA1 site of
20 16 *DlmPSYI*. *i*# and *d*#: # of bp inserted or deleted, respectively; $d \geq 14$: more
21 17 than 14 bp deletion; *i*+*d*: target sites with both deletions and insertions.
- 22 18 **H.** Representative *DlmPSYI* mutants at the sgRNA2 site. The represents
23 19 homozygote (T0-12), biallic (T0-14), and heterozygote (T0-30 to T0-32) at
24 20 A1-subgenome were shown.
- 25 21 **I.** Frequencies of indels (left) and mutation types (right) at the sgRNA2 site of
26 22 *DlmPSYI*. (legend: see G)
- 27 23 **J.** Phenotypes of representative *dlmpsyI* mutants. **a-c**, wild-type; **d-f**, *dlmpsyI*
28 24 mutant (T0-1).
- 29 25 **K-N.** Phenotypes of wild-type and the represented *grgI* mutant. Growth
30 26 phenotype (**K**) and internode elongation (**L**) of 5-month old wild-type (**a**) and
31 27 *grgI* (**b**) plants grown in the greenhouse. Plant heights (**M**) and internode lengths
32 28 (**N**) were quantified. **: $p < 0.01$.
- 33 29 **O.** Mutations of the *GRG1* gene were confirmed by Sanger sequencing. The
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

1
2
3
4 1 sgRNA target regions (red), PAM regions (blue), nucleotide insertions (green) and
5 2 their length (right side) are shown.
6
7 3
8
9 4
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

For Review Only










E

sgRNA	Numbers of transgenic lines	Numbers of plants analyzed	Numbers of plants with mutation	Mutation efficiency
sgRNA1	22	22	22	100%
sgRNA2	12	12	10	83.3%

F

WT	A1: GAGGAGGTCCGGCCAGCCTCCCGGCTCGA	
	B1: GAGGAGGTCCGGCCAGCCTCCCGGCTCGA	
	C1: GGGGAGGTCCGGCCAGCCTCCCGGCTCGA	
T0-1	A1: GAGGAGGTCCGGCCAGCCT - -CCCGGCTCGA	-2
	GAGGAGGTCCGGCCAGCCTcCCCGGCTCGA	+2
	B1: GAGGAGGTCCGGCCAGCCTcCCCGGCTCGA	+2
	GAGGAGGTCCGGCCAGCCT -CCCGGCTCGA	-1
	C1: GGGGAGGTCCGGCCAGCCT -CCCGGCTCGA	-1
	GGGGAGGTCCGGCCaCCT -CCCGGCTCGA	-2/+1
T0-2	A1: GAGGAGGTCCGGCCAGCCT - -CCCGGCTCGA	-2
	B1: GAGGAGGTCCGGCCAGCCTcCCCGGCTCGA	+2
	C1: GGGGAGGTCCGG - - - - -CTCGA	-14
	GGGGAGGTCCGGCCAGCCT -CCCGGCTCGA	-1
T0-3	A1: GAGGAGGTCCGGCCAGCCT -CCCGGCTCGA	-1
	B1: GAGGAGGTCCGGCCAGCCT -CCCGGCTCGA	-1
	GAGGAGGTCCGGCCAGCCTc - - - - -TTCTT	+2 /-27
	C1: GGGGAGGTCCGGCCAGCCT -CCCGGCTCGA	-1
T0-12	A1: GAGGAGGTCCGGCCAGCCT - -CCCGGCTCGA	-2
	B1: GAGGAGGTCCGGCCAGCCTcCCCGGCTCGA	+1
	GAGGAGGTCCGGCCAGCCT -CCCGGCTCGA	-1
	GAGGAGGTCCGGCCAGCCTaaCCCGGCTCGA	+2
	C1: GGGGAGGTCCGG - - - - -CTCGA	-14
	GGGGAGGTCCGGCCAGCCT -ccCGGCTCGA	-1
T0-13	A1: GAGGAGGTCCGG - - - - -CTCGA	-14
	B1: GAGGAGGTCCGGCCAGCCT -CCCGGCTCGA	-1
	GAGGAGGTCCGGC - - - - -CAAGGC	-21
	C1: GGGGAGGTCCGGCCAGC - - - - -GTGA	-12/+2
	GGGGAGGTCCGGCC - - - - -CCGGCTCGA	-8
T0-10	A1: GAGGAGGTCCGGCCAGCCT - -CCCGGCTCGA	-2
	B1: GAGGAGGTCCGGCCAGCCTcCCCGGCTCGA	+2
	GAGGAGGTCCGGCCAGCCT -CCCGGCTCGA	-1
	C1: GGGGAGGTCCGGCCAGCCTCCCGGCTCGA	WT
	GGGGAGGTCCGG - - - - -CTCGA	-14
T0-26	A1: GAGGAGGTCCGGCCAGCCTCCCGGCTCGA	WT
	GAGGAGGTCCGGCCAGCCT - -CCCGGCTCGA	-2
	GAGGAGGTCCGGCCAGCCTcCCCGGCTCGA	+2
	B1: GAGGAGGTCCGGCCAGCCT -CCCGGCTCGA	-1
	GAGGAGGTCCGGCCAGCCTcCCCGGCTCGA	+2
	C1: GGGGAGGTCCGG - - - - -CTCGA	-14
	GGGGAGGTCCGGCCAGCCT -CCCGGCTCGA	-1
T0-12	A1: GTTGCGCGGCACCTCCAG - - - - -CAACA	-17
T0-14	A1: GTTGCGCGGCACCTCCAGG - CTTGGCCTG	-1
	GTTGCGCGGCACCTCCAGG - - - - -ACGCC	-11
T0-30	A1: GTTGCGCGGCACCTCCAGGcCCTTGGCCTG	+1/WT
T0-31	A1: GTTGCGCGGCACCTC - - - - -CAGGT	-26/WT
T0-32	A1: GTTGCGCGGCACCTCCAGG - CTTGGCCT	-1 /WT



1
2
3
4 **Title: Robust CRISPR/Cas9 mediated genome editing and its**
5
6 **application in manipulating plant height in the first generation of**
7
8 **hexaploid Ma bamboo (*Dendrocalamus latiflorus* Munro)**
9
10

11
12
13 Shanwen Ye¹, Gang Chen¹, Markus V. Kohnen¹, Wenjia Wang¹, Changyang Cai¹,
14 WenSha Ding¹, Chu Wu¹, Lianfeng Gu¹, Yushan Zheng¹, Xiangqing Ma¹, Chentao
15 Lin² and Qiang Zhu^{1,*}
16
17

18
19 ¹ *Basic Forestry and Proteomics Center (BFPC), Fujian Provincial Key Laboratory*
20 *of Haixia Applied Plant Systems Biology, College of Forestry, Fujian Agriculture and*
21 *Forestry University, Fujian, China;*
22
23

24
25 ²*Department of Molecular, Cell & Developmental Biology, University of California,*
26 *Los Angeles, California, USA*
27
28

29
30
31
32 Emails:

33
34 Shanwen Ye, yeshanwen@fafu.edu.cn;

35
36 Gang Chen, GangC@163.com;

37
38 Markus V. Kohnen, MKohnen@fafu.edu.cn;

39
40 Wenjia Wang, 3160808027@fafu.edu.cn;

41
42 Changyang Cai, caichangyang@fafu.edu.cn;

43
44 Wensha Ding, 1170428008@fafu.edu.cn;

45
46 Chu Wu, 1170538006@fafu.edu.cn;

47
48 Lianfeng Gu, lfgu@fafu.edu.cn;

49
50 Yushan Zheng, zys1960@163.com;

51
52 Xiangqing Ma, lxxmxq@126.com;

53
54 Chentao Lin, clin@mcd.db.ucla.edu;

55
56 Qiang Zhu, zhuqiang@fafu.edu.cn (*Corresponding author).
57
58
59
60

1
2
3
4 **Dear Editor,**

5 Bamboo is a special grass to human due to its great economic and ecological values.
6
7 Around 2.5 billion people are directly producing and consuming bamboo, and its
8
9 international trade reached 68.8 billion US dollars in 2018 (Data from International
10
11 Bamboo and Rattan Organization). One major bamboo species in Asia is Ma bamboo
12
13 (*Dendrocalamus latiflorus Munro*), which is a hexaploid species with three
14
15 subgenomes (2n=72, AABBCC) (Guo et al., 2019). Despite its agronomic importance,
16
17 it is nearly impossible to modify bamboo traits by traditional breeding as it takes over
18
19 70 years to flower. Bamboo research largely lagged behind due to the lack of efficient
20
21 genetic manipulation tools.
22

23
24
25 The CRISPR (Clustered Regularly Interspaced Short Palindromic
26
27 ~~repeatRepeat~~)/Cas9 provides straight-forward ways for genome editing in many plants
28
29 (Yin et al., 2017), but has never been applied in bamboo. Here, we reported the
30
31 generation of bamboo mutants with CRISPR/Cas9 technology by targeting one
32
33 specific copy or all homoeologous genes.
34
35

36
37 Since our recently established genetic transformation protocol is time-consuming
38
39 (~1.5 years) (Ye et al., 2017), we optimized the CRISPR/Cas9 system in bamboo
40
41 protoplast. We first improved the protoplast preparation methods, and could isolate
42
43 3.0×10^6 protoplasts/g fresh leaves. Next, we improved the PEG-mediated
44
45 transformation method, and reached efficiencies of 53.3% for a single plasmid and
46
47 29.8% for two co-transformed plasmids (**Figure 1A**), which is sufficient for
48
49 optimizing the CRISPR/Cas9 system. The maize *UBI* promoter was used to drive
50
51 Cas9 expression (Ye et al., 2017). Three polymerase III-dependent promoters from
52
53 rice (*OsU6a/OsU6b/OsU6c*) were selected to express the sgRNA cassettes (Ma et al.,
54
55 2015), as bamboo exhibits high genomic synteny-similarity with rice (Peng et al.,
56
57 2013). To check the effectiveness of CRISPR/Cas9 constructs, a frame-shift mutated
58
59 *GFP* (*mGFP*) containing an additional “guanine” thereby produces no fluorescence
60

1 signal was simultaneously co-transformed with CRISPR/Cas9 plasmids (**Figure 1B**).
2 Around 1.8% of the protoplasts transformed with the *UBI-Cas9/OsU6b-sgRNA*
3 construct showed strong signals within 72 h, indicating that the *mGFP* function was
4 restored by the CRISPR/Cas9 system through deleting the additional “guanine”
5 (**Figure 1C**). The *OsU6a* and *OsU6c* promoters work as well, however, with lower
6 efficiency than the *OsU6b* promoter, as positive signals were only occasionally
7 observed with more than 10 replicates. Taking together, the *UBI-Cas9/OsU6b-sgRNA*
8 construct effectively works in bamboo protoplast, and were used for the following
9 endogenous gene editing in Ma bamboo.

10
11 The putative phytoene synthase (*PSY1*) in bamboo, whose homolog in maize
12 functions in carotenoid biosynthesis (Zhu et al., 2016), was selected for the initial test.
13 Three bamboo *PSY1* alleles (*DlmPSY1-A*, *DlmPSY1-B*, *DlmPSY1-C*) were identified
14 and cloned by a homology cloning strategy (**Figure 1D**). To mutate all copies of
15 *DlmPSY1*, sgRNA1 targeting a conserved site among all *DlmPSY1* loci was designed
16 (**Figure 1D**). In addition, the sgRNA2 target site containing 2–3 single-nucleotide
17 polymorphisms (SNPs) in the spacer region among three *DlmPSY1* homoalleles and
18 was selected to test the tolerance of sgRNA mismatches (**Figure 1D**).

19
20 1,600 bamboo calluses induced from stem were transformed Hygromycin-resistant
21 ~~bamboos were regenerated~~ as described previously (Ye et al., 2017). In total, 34
22 independent transgenic lines ~~regenerated from 1,600 calluses~~ were confirmed positive
23 (2.1%) by PCR. Based on Sanger sequencing results, 22 (100%) and 10 (83.3%)
24 independent T0 lines were edited in the sgRNA1 and sgRNA2 region, respectively
25 (**Figure 1E**), indicating that both constructs effectively induce endogenous gene
26 editing.

27
28 The editing profiles were further analyzed by sequencing. Eighteen lines (81.8%)
29 contained putative homozygote/biallelic mutations in all subgenomes at the sgRNA1

1 target site. In some lines, putative homozygote/biallelic mutations exist in one
2 subgenome while ~~heterozyote~~heterozygote or chimeric mutations appear in other
3 subgenomes (T0-10 and T0-26) (**Figure 1F**). Eight mutation types were identified
4 from 590 independent clones (**Figure 1G**). The most frequently ~~appeared~~
5 type ~~is~~-was deletion (75%), of which 59.1% are small deletions (<2bp). The ratios of
6 large fragment deletions (≥ 14 bp), insertions, and combined indels were 15.9%,
7 2.21%, and 7.82%, respectively (**Figure 1G**). Since bamboo propagates through
8 asexual budding, those homozygote/biallelic mutations will remain in the genome of
9 their offspring clones during breeding.

11 sgRNA2 that perfectly targets *DlmPSYI-A1*, but not *DlmPSYI-B1* or *DlmPSYI-C1*
12 was designed to study the recognition specificity (**Figure 1D**). Sequencing results
13 confirmed that 10 transgenic lines contain mutations in *DlmPSYI-A1*, but none in
14 *DlmPSYI-B1* and *DlmPSYI-C1* (**Figure 1E**). Two lines (20%) were putative
15 homozygous or biallelic mutations (T0-12 and T0-14), while 7 lines (70%) were
16 heterozygous/chimeric (T0-30 to T0-32 as representative examples, **Figure 1H**). The
17 ratios of deletions, insertions, and combined mutations were 86%, 9%, and 5%,
18 respectively (**Figure 1I**). The mutations were predominantly short nucleotide changes
19 (1–26bp), and 22.7% were 1bp nucleotide deletions (**Figure 1I**). Those data
20 demonstrated the successful application of the CRISPR/Cas9 system in mutating a
21 specific *DlmPSYI* allele.

23 Eighteen lines (81.8%) with homozygote/biallelic mutations in all subgenomes at
24 the sgRNA1 site exhibited albino phenotypes (**Figure 1J**), which appeared at an early
25 stage during tissue culture, and persisted at the plantlets stage (**Figure 1J**). Those
26 results suggest that genome editing takes place at an early stage in embryonic cells,
27 and led to the loss-of-function of all *DlmPSYI* alleles. Similar results were reported in
28 rice, wheat or cotton (Wang et al., 2018; Wang et al., 2014; Zhang et al., 2014). In
29 case of sgRNA2, although *DlmPSYI-A* was mutated, no visible phenotypic change

1 was observed due to the existence of the wild-type *DlmPSYI-B* and *DlmPSYI-C*
2 alleles.

3
4 Next, we applied this technology in bamboo molecular research. Bamboo is the
5 ~~highest-tallest~~ grass in the world, while the underlying mechanism is unknown.
6 Previously, we identified several Gibberellin-responsive genes including *GRG1*
7 (GA-responsive gene 1, *PH01004823G0070*) that potentially acts in controlling
8 bamboo height (Zhang et al., 2018). Here ~~atwo~~ homozygote *grg1* mutants (~~efficiency~~
9 ~~40%~~) in Ma bamboo ~~waswere~~ produced using our optimized CRISPR/Cas9
10 technology. Mutation in *GRG1* increased plant height (**Figure 1K**), mostly due to
11 elongated internodes (**Figure 1L-N**). Sequencing results confirmed that the *grg1*
12 mutant has the putative homozygous mutation in A1-subgenome, biallelic mutation in
13 B1-subgenome, and homozygous mutation in C1-subgenome (**Figure 1O**), indicating
14 the loss-of-function of *GRG1* in transgenic bamboo. To our knowledge, this is the
15 first example on controlling bamboo height through gene manipulation, which will
16 contribute to subsequent studies on the molecular mechanisms behind the fast growth
17 of bamboo.

18
19 In summary, for the first time we engineered the hexaploid Ma bamboo through
20 CRISPR/Cas9 technology. The homozygote mutations were obtained in the first
21 generation of transgenic lines, which is extremely important for bamboo species due
22 to its long vegetative growth periods. We also confirmed the albino phenotype of
23 *dmpsyl1* mutant in bamboo and generated a bamboo mutant with altered plant height.
24 This demonstrates the applicability of CRISPR/Cas9 in bamboo and thereby boosts
25 future bamboo research and breeding.

26 27 28 **References**

29 Guo, Z.-H., Ma, P.-F., Yang, G.-Q., Hu, J.-Y., Liu, Y.-L., Xia, E.-H., Zhong, M.-C., Zhao, L., Sun, G.-L., Xu,
30 Y.-X., Zhao, Y.-J., Zhang, Y.-C., Zhang, Y.-X., Zhang, X.-M., Zhou, M.-Y., Guo, Y., Guo, C., Liu,

- 1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
- 1 J.-X., Ye, X.-Y., Chen, Y.-M., Yang, Y., Han, B., Lin, C.-S., Lu, Y. and Li, D.-Z. (2019) Genome
2 sequences provide insights into the reticulate origin and unique traits of woody bamboos.
3 *Molecular Plant* DOI:<https://doi.org/10.1016/j.molp.2019.05.009>.
- 4 Ma, X., Zhang, Q., Zhu, Q., Liu, W., Chen, Y., Qiu, R., Wang, B., Yang, Z., Li, H., Lin, Y., Xie, Y., Shen, R.,
5 Chen, S., Wang, Z., Guo, J., Chen, L., Zhao, X., Dong, Z. and Liu, Y.G. (2015) A Robust
6 CRISPR/Cas9 System for Convenient, High-Efficiency Multiplex Genome Editing in Monocot
7 and Dicot Plants. *Mol Plant* **8**, 1274-1284.
- 8 Peng, Z., Lu, Y., Li, L., Zhao, Q., Feng, Q., Gao, Z., Lu, H., Hu, T., Yao, N., Liu, K., Li, Y., Fan, D., Guo, Y., Li,
9 W., Weng, Q., Zhou, C., Zhang, L., Huang, T., Zhao, Y., Zhu, C., Liu, X., Yang, X., Wang, T.,
10 Miao, K., Zhuang, C., Cao, X., Tang, W., Liu, G., Liu, Y., Chen, J., Liu, Z., Yuan, L., Huang, X., Lu,
11 T., Fei, B., Ning, Z., Han, B. and Jiang, Z. (2013) The draft genome of the fast-growing
12 non-timber forest species moso bamboo (*Phyllostachys heterocycla*). *Nat Genet* **45**, 456-461.
- 13 Wang, P., Zhang, J., Sun, L., Ma, Y., Xu, J., Liang, S., Deng, J., Tan, J., Zhang, Q., Tu, L., Daniell, H., Jin, S.
14 and Zhang, X. (2018) High efficient multisites genome editing in allotetraploid cotton
15 (*Gossypium hirsutum*) using CRISPR/Cas9 system. *Plant Biotechnol J* **16**, 137-150.
- 16 Wang, Y., Cheng, X., Shan, Q., Zhang, Y., Liu, J., Gao, C. and Qiu, J.L. (2014) Simultaneous editing of
17 three homoeoalleles in hexaploid bread wheat confers heritable resistance to powdery
18 mildew. *Nat Biotechnol* **32**, 947-951.
- 19 Ye, S., Cai, C., Ren, H., Wang, W., Xiang, M., Tang, X., Zhu, C., Yin, T., Zhang, L. and Zhu, Q. (2017) An
20 Efficient Plant Regeneration and Transformation System of Ma Bamboo (*Dendrocalamus*
21 *latiflorus* Munro) Started from Young Shoot as Explant. *Frontiers in plant science* **8**, 1298.
- 22 Yin, K., Gao, C. and Qiu, J.L. (2017) Progress and prospects in plant genome editing. *Nature plants* **3**,
23 17107.
- 24 Zhang, H., Wang, H., Zhu, Q., Gao, Y., Wang, H., Zhao, L., Wang, Y., Xi, F., Wang, W., Yang, Y., Lin, C.
25 and Gu, L. (2018) Transcriptome characterization of moso bamboo (*Phyllostachys edulis*)
26 seedlings in response to exogenous gibberellin applications. *BMC Plant Biol* **18**, 125.
- 27 Zhang, H., Zhang, J., Wei, P., Zhang, B., Gou, F., Feng, Z., Mao, Y., Yang, L., Zhang, H., Xu, N. and Zhu,
28 J.K. (2014) The CRISPR/Cas9 system produces specific and homozygous targeted gene editing
29 in rice in one generation. *Plant Biotechnol J* **12**, 797-807.
- 30 Zhu, J., Song, N., Sun, S., Yang, W., Zhao, H., Song, W. and Lai, J. (2016) Efficiency and inheritance of
31 targeted mutagenesis in maize using CRISPR-Cas9. *J Genet Genomics* **43**, 25-36.

33 **Conflict of interests**

34 The authors declare that they have no conflict of interests.

36 **Fundings**

37 This work was supported by the National Natural Science Foundation of China grant
38 (No.31870660), Fujian Innovative Center for Germplasm Resources and Cultivation
39 of Woody plants (No.125/KLA15001E), Program for scientific and technological

1 innovation team in university of Fujian province (No.118/KLA18069A) to Q.Z. The
2 funding bodies were not involved in the design of the study or in any aspect of the
3 data collection, analysis and interpretation of data and in paper writing.

4 5 **Acknowledgements**

6 We thank Dr. Chella Thaniyarasu from Fujian Agriculture and Forestry University for
7 the critical reading of the manuscript.

8 9 **Authors' contributions**

10 Q.Z. conceived this project; L.F.G., Y.S.Z., X.Q.M. C.T.L. and Q.Z. designed
11 experiments, and interpreted the results; S.W.Y., G.C. and M.V.K. performed the
12 experiments and analyzed the data, W.J.W., C.Y.C., C.W., and D.W.S. helped to
13 perform the experiments and collect the data. All authors read and approved the
14 submission of this manuscript.

15 16 **Figure Legend**

17 **Figure 1. Genome editing in Ma bamboo using CRISPR/Cas9 technology**

18 **A.** Bamboo protoplast isolation and transformation. **a.** Microscopic image of
19 isolated bamboo protoplast transformed with 35S:tdTomato. **b-d.** Images of
20 bamboo protoplasts co-expressing the fluorescence proteins *tdTomato* (**b**) and
21 GFP (**c**) driven by the 35S promoter, and their overlay (**d**).

22 **B.** CRISPR/Cas9 plasmids for bamboo protoplast. **Top:** CRISPR/Cas9 constructs
23 expressing the sgRNA directed against *mGFP* and driven by
24 *OsU6a/OsU6b/OsU6c* respectively; **Middle:** *mGFP*-expression construct,
25 *mGFP* contains one additional guanine (lower-green case) downstream of the
26 translational start site (red); **bottom:** GFP-expression construct. The sgRNA
27 was designed to produce the presumptive cleavage site at the third nucleotide
28 upstream of the PAM sequence (blue).

29 **C.** Representative bamboo protoplasts co-transfected with *mGFP* and

1
2
3
4 1 *OsU6b-sgRNA/UBI-Cas9* reproducibly emitting fluorescence signals (red
5
6 2 arrows).

7
8 3 **D.** *DlmPSYI* gene structure and sequences of the target sites. Gray boxes: exons;
9
10 4 black lines: introns; number in brackets: positions of start codon, stop codon
11
12 5 and sgRNA target sites (red and orange rectangles). The PAM regions (blue),
13
14 6 SNPs (red), and nucleotide sequences of the *sgRNAs* and *DlmPSYI* genes were
15
16 7 given.

17
18 8 **E.** Frequencies of the CRISPR/Cas9-induced mutations in two target sites of the
19
20 9 *DlmPSYI*.

21
22 10 **F.** Representative *DlmPSYI* mutations at the sgRNA1 site. T0-1, T0-2, T0-3,
23
24 11 T0-12 and T0-13 represent loss-of function mutants. T0-10 and T0-26 lines
25
26 12 contain heterozygote mutations in the C1-subgenome and chimeric mutations
27
28 13 in the A1-subgenome, respectively. Red: sgRNA target regions; blue: PAM
29
30 14 regions; green lowercase letters: nucleotide indels; dotted lines: omitted
31
32 15 nucleotides.

33
34 16 **G.** Frequencies of indels (left) and mutation types (right) at the sgRNA1 site of
35
36 17 *DlmPSYI*. i# and d#: # of bp inserted or deleted, respectively; $d \geq 14$: more
37
38 18 than 14 bp deletion; i+d: target sites with both deletions and insertions.

39
40 19 **H.** Representative *DlmPSYI* mutants at the sgRNA2 site. The represents
41
42 20 homozygote (T0-12), biallic (T0-14), and heterozygote (T0-30 to T0-32) at
43
44 21 A1-subgenome were shown.

45
46 22 **I.** Frequencies of indels (left) and mutation types (right) at the sgRNA2 site of
47
48 23 *DlmPSYI*. (legend: see G)

49
50 24 **J.** Phenotypes of representative *dlmpsy1* mutants. **a-c**, wild-type; **d-f**, *dlmpsy1*
51
52 25 mutant (T0-1).

53
54 26 **K-N.** Phenotypes of wild-type and the represented *grg1* mutant. Growth
55
56 27 phenotype (**K**) and internode elongation (**L**) of 5-month old wild-type (**a**) and
57
58 28 *grg1* (**b**) plants grown in the greenhouse. Plant heights (**M**) and internode lengths
59
60 29 (**N**) were quantified. **: $p < 0.01$.

1
2
3
4 1 **O.** Mutations of the *GRG1* gene were confirmed by Sanger sequencing. The
5 2 sgRNA target regions (red), PAM regions (blue), nucleotide insertions (green) and
6 3 their length (right side) are shown.
7 4
8 5
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

For Review Only