

Lawrence Berkeley National Laboratory

Recent Work

Title

THE SENSITIVITY OF THE FORWARD CROSS SECTION IN TRANSFER REACTIONS BETWEEN HEAVY IONS TO THE EDGE OF THE NUCLEAR FIELD

Permalink

<https://escholarship.org/uc/item/5pk6w9wn>

Authors

Ascutto, R.J.
Glendenning, Norman K.

Publication Date


1973-09-01

THE SENSITIVITY OF THE FORWARD CROSS SECTION
IN TRANSFER REACTIONS BETWEEN HEAVY IONS
TO THE EDGE OF THE NUCLEAR FIELD

R. J. Ascutto and Norman K. Glendenning

September 1973

Prepared for the U. S. Atomic Energy Commission
under Contract W-7405-ENG-48



TWO-WEEK LOAN COPY

This is a Library Circulating Copy
which may be borrowed for two weeks.
For a personal retention copy, call
Tech. Info. Division, Ext. 5545

DISCLAIMER

This document was prepared as an account of work sponsored by the United States Government. While this document is believed to contain correct information, neither the United States Government nor any agency thereof, nor the Regents of the University of California, nor any of their employees, makes any warranty, express or implied, or assumes any legal responsibility for the accuracy, completeness, or usefulness of any information, apparatus, product, or process disclosed, or represents that its use would not infringe privately owned rights. Reference herein to any specific commercial product, process, or service by its trade name, trademark, manufacturer, or otherwise, does not necessarily constitute or imply its endorsement, recommendation, or favoring by the United States Government or any agency thereof, or the Regents of the University of California. The views and opinions of authors expressed herein do not necessarily state or reflect those of the United States Government or any agency thereof or the Regents of the University of California.

THE SENSITIVITY OF THE FORWARD CROSS SECTION IN TRANSFER
REACTIONS BETWEEN HEAVY IONS TO THE EDGE OF THE NUCLEAR FIELD[†]

R. J. Ascutto

Wright Nuclear Structure Laboratory
Yale University
New Haven, Connecticut 06520

and

Norman K. Glendenning

Lawrence Berkeley Laboratory
University of California
Berkeley, California 94720

September 1973

The ratio of the cross section for two nucleon transfer in heavy ion reactions measured at forward angle and at the grazing angle depends very sensitively on the location of the edge of the imaginary potential compared to the real.

- - -


In this note we focus attention on the cross section in heavy ion reactions at angles smaller than the grazing angle, θ_g . Under classical sub-coulomb conditions, such scattering events would correspond to large impact parameters or distant collisions where the field is weak and causes little deflection. Of course, distant orbits cannot very effectively contribute to nucleon transfer reactions, especially when more than one nucleon is transferred, because of the exponential decay of the bound state wave functions. However, in the classical scattering of particles in a field, such as that which acts

[†]Work performed under the auspices of the U. S. Atomic Energy Commission.

between heavy ions, there exist also small impact parameter orbits which lead to forward scattering, for energies above the barrier, as was pointed out long ago by Ford and Wheeler [1]. Nuclei on such orbits can very easily transfer particles. They are, however, subject to absorption, because of their close encounter. However, this is precisely our interest in them. We claim that forward angle scattering in transfer reactions can be used as a very sensitive probe of the edge of the nuclear potential acting between heavy ions.

The situation is illustrated schematically in fig. 1. The distant orbit, a, is only slightly deflected by some small angle, θ , say. A closer orbit, b, since it feels a stronger repulsive field, is scattered to a larger angle. However, the two closer orbits c and d ($d < c$) feel a net attraction and scatter to the same angle θ as a. The orbits such as c and d are the ones of interest here.

In fig. 2 we show calculated cross sections for the reaction



leading to the ground and collective 2^+ state in ^{122}Sn , for $E = 100$ MeV, an energy significantly above the coulomb barrier (~ 60 MeV). The solid lines correspond to the optical model parameters listed in table 1. The ground state cross section shows the characteristic peak at the grazing angle while the 2^+ cross section exhibits a dip there due to destructive interference between direct and indirect modes of producing this state. This we have discussed separately [2]. The comparison we wish to make is with the dashed curve, which corresponds to a 5% increase in the radius of the imaginary potential. We see that this very modest change in potential results in a

difference in the ratio of forward to grazing angle cross section of more than a factor 10, which is a very large amplification factor. (If the forward angle cross section for the ground state is hard to measure experimentally, because of low cross section, we suggest a wide angle counter to integrate out the ripples and speed up data taking.)

Of course the question immediately comes to mind as to whether the effect of changing the radius of the imaginary potential relative to the real might not be more easily detected in the elastic cross section. This is not the case as shown in fig. 3 where we see a high precision experiment would be required to distinguish the two cross sections. Even then the multiple ambiguities in the optical model fit to elastic cross sections might not yield a unique determination. Nor would inelastic cross sections, or one nucleon transfer reactions be as appropriate for the measurement of this effect. For in such reactions, distant orbits, a, would make a larger contribution to the forward angle cross section, thus masking the effect produced on the close orbits c and d by the edge of the nuclear potential.

Since the de Broglie wave length of the relative motion between heavy ions in typical collisions is very short compared to the nuclear dimensions, there is a close correspondence between impact parameter and angular momentum. This makes it especially interesting to examine the S matrix elements as a function of L. We anticipate being able to relate rather directly, changes in such a plot to changes in the interaction region, such as potential radii or slope of the tail of bound state wave functions of transferred particles. Such a plot for the transfer reaction leading to the ground state of ^{122}Sn is shown in fig. 4, and corresponds to the cross section shown by a solid line in fig. 2.

The absolute value of the S matrix element is plotted versus integer values of L. These points are joined exhibiting a very smooth behavior of S. The sudden fall on the low side of the peak corresponds to the sudden onset of absorption.* The slower fall off on the high L side is governed by the tail of the transfer form factor. The points plotted on the figure indicate S matrix values in the case where the imaginary radius has been increased by 5% (dotted line of fig. 2). Here as anticipated we see the lower L region is effected but not the higher region.

It is also noteworthy that the peak in the S matrix is very little effected by the change in imaginary radius whereas, our calculations (not shown) indicate a considerable shift if the real radius is also increased. We conclude that the trajectories are principally determined by the real potential, while the imaginary potential produces little deflection.

The forward angle ripples in the differential cross section are, we believe, the result of interference between pairs of close impact orbits from opposite sides of the nucleus, such as c and d in fig. 1. This phenomenon is likely to be very general for heavy ion transfer reactions, especially where more than one nucleon is transferred, for it depends only on a sufficiently high localization in L. The frequency of the ripples is determined by the angular momentum L_0 corresponding to the peak in the S-matrix (angle between peaks is $\Delta\theta \approx \pi/L_0$). Indeed, if the S-matrix were a delta function $\delta(L - L_0)$, then the angular distribution would be given by $|Y_{L_0}(\theta)|^2$. The ripples will be damped according

* We use absorption only to mean loss to other channels than the elastic, inelastic and transfer channels explicitly treated in the calculation: We do not imply fusion!

to the width ΔL of S . So, for example, we see that the ripples are more damped in the dotted curve of fig. 2 corresponding to widths at half maximum of $\Delta L = 14$ and 11 , respectively, for the dotted and solid S -matrix shown in fig. 4.


To summarize, the ratio of the cross sections of two-nucleon transfer reactions, measured at forward angle and at the grazing angle, is a very sensitive probe of the relationship between the radius of the real and imaginary nuclear potentials. Such measurements should prove very useful in determining the details of the potential at the edge, which is of decisive importance in predicting cross sections of heavy ion reactions. The location of the edge of the imaginary potential relative to the real is also important in connection with the question of nuclear molecules. However, this kind of measurement should be made for the 0^+ state since it is very little effected by second-order processes. In contrast, second-order processes can strongly alter the forward cross-section, and are responsible for the forward peak in the 2^+ state [2].

References

1. K. W. Ford and J. A. Wheeler, Ann. of Physics (N.Y.) 7 (1959) 259.
2. R. J. Ascutto and N. K. Glendenning, Symposium on Heavy-Ion Transfer Reactions, Argonne (1973) 513; Phys. Letters 45B (1973) 85.


Table 1. Optical Potential Parameters: The radius of the nuclear part is $r_o (A_P^{1/3} + A_T^{1/3})$ and the charge radius $r_c A_T^{1/3}$.

V	W	r_o	a	r_c
-40	-15	1.31	.45	1.2



XBL 7310-1381

Fig. 1. Four classical orbits in the real part of the potential of table 1 are plotted, three of which, having impact parameters a , c and $-d$, ($d < c$) scatter to the same angle θ . The orbit b feels a stronger repulsion than a and therefore scatters to a larger angle. The circle marks the half value of the Woods-Saxon nuclear potential. The orbits are for 180 scattered by ^{120}Sn at 100 MeV lab energy. Dotted lines show pure Coulomb orbits.



XBL 739-1249

Fig. 2. Differential cross sections for the 0^+ and 2^+ state are compared for two values of the radius of the imaginary potential one of which (dotted curve) is 5% larger than the other. The shape of the 2^+ cross section has been strongly altered from the classical shape exhibited by the 0^+ by strong two-step processes which interfere destructively with the direct for stripping reactions as discussed in ref. 2. Cross sections are in mb./sr. and the two scales refer respectively to the 0^+ and 2^+ states.




Fig. 3. The elastic cross sections in the entrance channel corresponding to the reaction of fig. 2 for the two cases there, showing the weak sensitivity of the elastic cross section to $R_v - R_w$ compared to the two-nucleon transfer cross section. The optical model parameters are listed in table 1 for the solid line.




Fig. 4. The absolute value of the S-matrix elements for the 0^+ state are plotted as a function of L and the points (for integer L) joined. The solid lines of this and fig. 2 correspond as do the dots.

LEGAL NOTICE

This report was prepared as an account of work sponsored by the United States Government. Neither the United States nor the United States Atomic Energy Commission, nor any of their employees, nor any of their contractors, subcontractors, or their employees, makes any warranty, express or implied, or assumes any legal liability or responsibility for the accuracy, completeness or usefulness of any information, apparatus, product or process disclosed, or represents that its use would not infringe privately owned rights.

TECHNICAL INFORMATION DIVISION
LAWRENCE BERKELEY LABORATORY
UNIVERSITY OF CALIFORNIA
BERKELEY, CALIFORNIA 94720