

Lawrence Berkeley National Laboratory

LBL Publications

Title

Nuclear structure of proton-rich unstable nucleus ^{28}P studied by g-factor measurement

Permalink

<https://escholarship.org/uc/item/5r62m20n>

Journal

Chinese Physics C, 33(S1)

ISSN

1674-1137

Authors

Yong-Nan, Zheng

Dong-Mei, Zhou

Matsuta, K

et al.

Publication Date

2009-03-01

DOI

10.1088/1674-1137/33/s1/069

Peer reviewed

Nuclear structure of proton-rich unstable nucleus ^{28}P studied by g -factor measurement

To cite this article: Zheng Yong-Nan *et al* 2009 *Chinese Phys. C* **33** 215

View the [article online](#) for updates and enhancements.

Related content

- [Magnetic moment of extremely proton-rich nucleus \$^{23}\text{Al}\$](#)
T Nagatomo, K Matsuta, A Ozawa *et al*.
- [Magnetic moment of \$^{17}\text{Ne}\$](#)
L T Baby, C Bordeanu, M Hass *et al*.
- [Proton alignment in \$^{82}\text{Sr}\$ investigated by \$g\$ -factor measurements](#)
Fan Ping, Yuan Da-Qing, Zuo Yi *et al*.

Recent citations

- [Theab initioCalculation of Electric Field Gradient at the Site of P Impurity in -Al₃O₂](#)
Qiao-Li Zhang *et al*

Nuclear structure of proton-rich unstable nucleus ^{28}P studied by g -factor measurement^{*}

ZHENG Yong-Nan(郑永男)¹ ZHOU Dong-Mei(周冬梅)¹ K. Matsuta² M. Mihara² M. Fukuda²
D. Nishimura² J. Komurasaki² D. Ishikawa² R. Matsumiya² T. Nagatomo³ T. Izumikawa⁴
S. Takahashi⁵ H. Hirano⁵ T. Ohtsubo⁵ S. Momota⁶ Y. Nojiri⁷ A. Kitagawa⁷
M. Kanazawa⁷ M. Torikoshi⁷ S. Sato⁷ T. Minamisono⁸ J. R. Alonso⁹ G. F. Krebs⁹
T. J. M. Symons⁹ YUAN Da-Qing(袁大庆)¹ ZUO Yi(左翼)¹ FAN Ping(范平)¹ T. Suzuki¹⁰
ZHANG Xi-Zhen(张锡珍)¹ ZHU Sheng-Yun(朱升云)^{1;1)}

¹ (China Institute of Atomic Energy, P.O. Box 275-50, Beijing 102413, China)

² (Department of Physics, Osaka University, Toyonaka, Osaka 560-0043, Japan)

³ (RIKEN, Wako, Saitama 351-0198, Japan)

⁴ (RI Center, Niigata University, Niigata 951-8510, Japan)

⁵ (Department of Physics, Niigata University, Niigata 950-2181, Japan)

⁶ (Kochi University of Technology, Kochi 782-8502, Japan)

⁷ (National Institute of Radiological Sciences, Inage-ku, Chiba 263-8555, Japan)

⁸ (Fukui University of Technology, Fukui 910-8505, Japan)

⁹ (Lawrence Berkeley Laboratory, Berkeley, CA94720, USA)

¹⁰ (Department of Physics, Nihon University, Tokyo 156, Japan)

Abstract Nuclear structure of proton-rich unstable nucleus ^{28}P has been studied by measuring its g -factor for the first time. The g -factor of ^{28}P ($I^\pi = 3^+$, $T_{1/2} = 270.3$ ms) was measured by means of β -NMR technique combined with the new polarization technique for charge exchange reaction product in the intermediate energy heavy ion collisions. The obtained g -factor of $g = 0.1028(27)$ is very much quenched from the Schmidt value, but is well reproduced by the shell model ($+0.102$). In connection with the magnetic moment of the mirror partner and the β -ray transition probability, the orbital angular momenta and intrinsic spins of protons and neutrons have been determined as $\langle l_p \rangle = 0.43(29)$, $\langle l_n \rangle = 1.85(29)$, $\langle S_p \rangle = 0.28(4)$, and $\langle S_n \rangle = 0.44(4)$.

Key words ^{28}P , β -NMR, nuclear magnetic moment, exotic nuclei

PACS 21.10.Ky, 27.30.+t, 21.90.+f

1 Introduction

Proton-rich unstable nucleus ^{28}P ($I^\pi = 3^+$, $T_{1/2} = 270.3$ ms)^[1] has one proton in the $2s_{1/2}$ orbital outside the ^{27}Si core, with relatively small proton separation energy of 2.065 MeV, hence it may have proton halo structure, which may be responsible for the abrupt increase of the reaction cross section of ^{28}P on Si target at intermediate energies reported by Z. H. Liu et al.^[2]. The magnetic moment of this nucleus has always been of interest. No successful measure-

ment, however, has been done until now. We tried to measure it years ago using polarization transfer in the low energy nuclear reaction $^{28}\text{Si}(p,n)^{28}\text{P}$ and simply found it difficult. One difficulty may be the selection of implantation material. Si was known as a good implantation material for P isotopes, at that time, from the successful measurement of the magnetic moment of ^{29}P . However, in the case of ^{28}P , the Q moment may be very large compared with $Q(=0)$ for ^{29}P ($I^\pi = 1/2^+$), hence the spin relaxation from the quadrupole interactions may be significant. The

Received 3 September 2008

* Supported by National Natural Science Foundation of China (10435010, 10505032)

1) E-mail: zhuy@ciae.ac.cn

©2009 Chinese Physical Society and the Institute of High Energy Physics of the Chinese Academy of Sciences and the Institute of Modern Physics of the Chinese Academy of Sciences and IOP Publishing Ltd

magnetic moment of ^{28}P was estimated to be rather small, which give us very large $eqQ/\mu H$ ratio. In the present work, we have successfully measured the magnetic moment of ^{28}P using Pt as an implantation material, utilizing its small magnetic moment, to study structure of this nucleus. Combined with the magnetic moment of its mirror partner ^{28}Al , we can separate orbital angular momenta and intrinsic spins.

2 Experimental

The present experiment was performed at NIRS (National Institute of Radiological Sciences). The ^{28}P nuclei were produced through the charge exchange reaction bombarding 2 mm thick Be target with a primary beam of ^{28}Si at 100 AMeV, provided by the heavy ion synchrotron HIMAC. The ^{28}P nuclei were separated out with a fragment separator installed in the secondary beam line SB2. By selecting ejection angle $\theta = 0.7^\circ$ and momentum $\Delta p/p_0 = -2.0\%$, nuclear spin polarization was introduced in the ^{28}P nuclei. Thus polarized ^{28}P nuclei were slowed down and were implanted in a Pt foil cooled down to 15 K, placed under a strong magnetic field of 0.9 T ($FM = \pm 15$ kHz) or 1.0 T ($FM = \pm 50$ kHz), to preserve polarization. β -rays emitted from ^{28}P were detected by a pair of plastic-scintillation-counter telescopes placed above and below the Pt catcher relative to the polarization direction. To reduce background β -ray counts a 0.5 mm-thick Cu plate was used as an absorber. The β -NMR was detected by means of the β -ray asymmetry.

3 Results and discussion

Typical momentum distribution and the polarization of the secondary beam ^{28}P are shown in Fig. 1. Center momentum is significantly decelerated from the primary beam velocity, and the relatively large polarization is observed, which may show a strong influence from the pick-up process in the present charge exchange process.

The β -NMR spectra are shown in Fig. 2. From the spectra, g -factor of ^{28}P was determined to be $|g(^{28}\text{P})| = 0.1028(27)$, correcting 0.10% of the theoretical diamagnetism and including 0.27% of Knight shift^[3] in the error, from which the magnetic moment was deduced by the well known expression $\mu = gI$.

The obtained magnetic moment $|\mu(^{28}\text{P})| = 0.309$ (9) μ_N is very much quenched from the Schmidt value $+0.88$, but is well reproduced by the shell model value

$+0.306$ obtained by the OXBASH code^[4]. This agreement, at least, confirms the contribution of the valence $2s_{1/2}$ proton in main configuration, as is speculated from the reaction cross-section data.




Fig. 1. Polarization as function of momentum.




Fig. 2. NMR spectra.

Comparing nuclear moments of mirror pairs, information on the motion of nucleons inside nucleus can be extracted with less ambiguity, because of the mirror symmetry. The total orbital angular momentum $\langle l_z \rangle$ and the intrinsic spin $\langle S_z \rangle$ can be separated from the iso-scalar moment $\mu(i, s)$, as is shown in equation (1)^[5, 6], assuming free nucleon values for g -factors. On the other hand, $\langle \tau_3 l_z \rangle$ and $\langle \tau_3 S_z \rangle$ can be derived from the iso-vector moment $\mu(i, v)$ and the Gamow-Teller matrix element $\int \sigma$ as shown in Eq. (2).

$$\begin{aligned} \mu(i, s) &= \frac{1}{2} \{ \mu(^{28}\text{P}) + \mu(^{28}\text{Al}) \} = \\ &= \frac{J}{2} + \left(\mu_p + \mu_n - \frac{1}{2} \right) \langle S_z \rangle, \langle l_z \rangle = J - \langle S_z \rangle, \end{aligned} \quad (1)$$

$$\begin{aligned}\mu(i, \nu) &= \frac{1}{2} \{ \mu(^{28}\text{P}) - \mu(^{28}\text{Al}) = \\ &\frac{1}{2} \langle \tau_3 l_z + (\mu_p - \mu_n) \langle \tau_3 S_z \rangle, \langle \tau_3 S_z \rangle = \\ &\frac{1}{2} \sqrt{\frac{2TJ}{J+1}} \int \sigma, \end{aligned} \quad (2)$$

$\langle l_z \rangle$ and $\langle S_z \rangle$ are deduced to be 2.28(1) and 0.73(1), respectively, from the present magnetic moment of ^{28}P and the known magnetic moment of the mirror partner, $|\mu(^{28}\text{Al})| = 3.242(5)\mu_N^{[7]}$. The ft -value of the β transition from the ground state of ^{28}P to the IAS (isobaric analogue state) in the excited state of ^{28}Si at 9.314 MeV, is given as $ft = 2925(111)$ s, from the f and t values given from the end-point energy^[8] and the branching ratio $(0.113(4))^{[1]}$, respectively. Considering Fermi matrix element $|\int 1|^2=2$, the Gamow-Teller matrix element $\int \sigma$, hence $|\langle \tau_3 S_z \rangle| = 0.16(6)$ is given. Combined with this information, we obtain expectation values as,

$$\langle l_p \rangle = 0.43(29), \quad \langle l_n \rangle = 1.85(29),$$

$$\langle S_p \rangle = 0.28(4), \quad \langle S_n \rangle = 0.44(4),$$

while the shell model prediction gives,

$$\langle l_p \rangle_{\text{SM}} = 0.62, \quad \langle l_n \rangle_{\text{SM}} = 1.87,$$

$$\langle S_p \rangle_{\text{SM}} = 0.17, \quad \langle S_n \rangle_{\text{SM}} = 0.34,$$

where $\langle l_p \rangle$ denotes the z -component of the orbital angular momentum for proton group, etc. Although the magnetic moment is well reproduced by the shell model calculation, the individual angular momenta are not necessarily reproduced well (Fig. 3). The shell model prediction seems to develop configuration mixing too much. Especially, the total spin is too small.




Fig. 3. Decomposition of nuclear spin.

The present work was performed at NIRS (National Institute of Radiological Sciences)-HIMAC under the research project with heavy ions (program No.P026). The authors are grateful to the staffs of HIMAC. The work was partly supported by the Grant in aid for scientific research from the Japan Society for the Promotion of Science.

References

- 1 Endt P M. Nucl. Phys. A, 1990, **521**: 1
- 2 LIU Z H et al. P. R. C, 2004, **69**: 034326
- 3 Matsuta K et al. Hyperfine Interaction, 1988, **120/121**: 719; Mihara M. presented at this meeting
- 4 Brown B A et al. MSUCL report, 1988, 524
- 5 Sugimoto K. Proc. Int. Conf. on Nuclear Moments and Nuclear Structure. Edt. Horie H, Sugimoto K; Suppl. to J. Phys. Soc. Jpn., 1973, **34**: 197
- 6 Hanna S S, Hugg J W. Hyperfine Interactions 21, 59 (1985)
- 7 Minanisono T et al. Phys. Lett. B, 1981, **106**: 38
- 8 Wilkinson D H et al. Nucl. Phys. A, 1974, **232**: 58