UCSF UC San Francisco Previously Published Works

Title

Influential and Highest Cited Shoulder Instability Articles: A Bibliometric Analysis.

Permalink

https://escholarship.org/uc/item/5rd7q148

Journal

Orthopaedic journal of sports medicine, 9(3)

ISSN 2325-9671

Authors

Allahabadi, Sachin Eftekhari, Ava Feeley, Sonali E <u>et al.</u>

Publication Date

2021-03-01

DOI

10.1177/2325967121992577

Peer reviewed

Influential and Highest Cited Shoulder Instability Articles

A Bibliometric Analysis

Sachin Allahabadi,*[†] MD, Ava Eftekhari,[†], Sonali E. Feeley,[†], Brian T. Feeley,[†] MD, and Drew A. Lansdown,[†] MD

Investigation performed at the Department of Orthopaedic Surgery, University of California, San Francisco, San Francisco, California, USA

Background: In orthopaedic surgery and other fields, the effect of influential journal articles has been evaluated by their citation counts and their correlation with methodological quality.

Purpose: To identify and characterize the 50 most cited articles on shoulder instability, examine trends in publication characteristics, and evaluate the correlation of citations with quality of evidence.

Study Design: Cross-sectional study.

Methods: The Web of Science and Scopus online databases were searched to identify the top 50 most cited articles on shoulder instability, based on the average number of citations from each database. Article characteristics were recorded, and level of evidence and methodological quality were assessed using the modified Coleman Methodology Score (mCMS), Jadad scale, and Methodological Index for Non-Randomized Studies (MINORS). Spearman correlation was used to evaluate relationships between citations or citation density (citations/y) and level of evidence or methodological scoring. Top cited articles from recent years were also aggregated.

Results: The top 50 most cited papers had a mean \pm SD number of citations of 381.5 \pm 166.7, with a mean of 15.0 \pm 8.8 citations/y. Overall, 15 articles (30%) were biomechanical/cadaveric studies, and 15 (30%) were case series. Only 3 (6%) were considered to have level 1 evidence. The mean \pm SD mCMS was 54.4 \pm 12.7, mean \pm SD Jadad score was 3.1 \pm 1.4, and mean \pm SD MINORS score was 10.5 \pm 3.3. There were no significant correlations between citation rank or density and methodological assessments. There were weak correlations between citation rank and publication year ($r_s = 0.32$; P = .022) and between rank and level of evidence ($r_s = -0.38$; P = .047). The correlation between citation density and publication year was moderate ($r_s = 0.70$; P < .0001). There was no difference in citation density of the top 10 articles from 2010 to 2020 compared with the top 10 from the overall list (23.8 \pm 5.3 vs 28.8 \pm 9.5; P = .16).

Conclusion: Influential articles in shoulder instability included a high proportion of biomechanical/cadaveric studies. The majority of top cited articles had lower evidence levels and poorer methodological quality without strong correlation with citations or citation density. There was a moderate correlation between citation density and year of publication.

Keywords: shoulder; instability; dislocation; quality; Bankart; Latarjet; stabilization

The modern understanding of shoulder instability derives from the description of the essential lesion of the glenoid labrum. The presence of this lesion was initially hypothesized in 1890 by Broca and Hartmann and ultimately confirmed in 1906 by Georg Clemens von Perthes and in 1923 by A.S. Blundell Bankart.^{6,59} Bankart described his technique for fixing the anteroinferior labral detachment seen in cases of recurrent shoulder instability, and he would become the namesake of the Bankart lesion and Bankart repair.^{6,7} Since that time, the understanding of the pathoanatomy, biomechanics, natural history, and treatment modalities—including soft tissue and bony procedures to restore shoulder joint stability—of shoulder instability has evolved significantly. Arthroscopic shoulder surgery has become a mainstay of restoration of shoulder stability, and our understanding of its limitations—particularly relating to bone loss patterns has also allowed for advancement of open operations such as the procedure described by Latarjet in 1954.⁴⁶

As with other complex orthopaedic pathologies, it is critical to understand previous literature and the progression

The Orthopaedic Journal of Sports Medicine, 9(3), 2325967121992577 DOI: 10.1177/2325967121992577 © The Author(s) 2021

This open-access article is published and distributed under the Creative Commons Attribution - NonCommercial - No Derivatives License (https://creativecommons.org/ licenses/by-nc-nd/4.0/), which permits the noncommercial use, distribution, and reproduction of the article in any medium, provided the original author and source are credited. You may not alter, transform, or build upon this article without the permission of the Author(s). For article reuse guidelines, please visit SAGE's website at http://www.sagepub.com/journals-permissions.

of knowledge to continue to advance treatment strategies to improve patient outcomes. There has been a recent emphasis on evaluating influencial articles relating to various topics in orthopaedics and shoulder surgery.^{40,42,47,50,51,54,70} Evaluation of influential articles may be assisted by reviewing articles from high-impact journals. However, in orthopaedics, a journal's impact factor has been shown to result largely from several highly cited articles within the journal, skewing the mean number of citations upward, resulting in 85% of orthopaedic articles being cited fewer times than expected by the impact factor.¹² Therefore, studies have measured an article's influence by the mean number of citations.^{40,42,47,50,51,54,70} Aggregating the most commonly cited studies may provide clinicians and researchers with a foundational set of literature.

Although the most cited studies may be influential, authors have recently highlighted the lack of correlation of quality of evidence and number of citations.⁷⁰ Evaluation of study quality and additional study characteristics may elucidate why some studies are more cited than are others. Despite several recent studies on highly cited articles on various topics in orthopaedics, no similar study has been performed regarding shoulder instability.

The purpose of the present study was to identify and characterize the 50 most cited articles relating to shoulder instability, examine trends in publication characteristics, and evaluate correlations of study citations with quality of evidence. We hypothesized that the top 50 cited articles would have poor correlation with methodological quality and would have more historical and foundational value than value as literature that encompasses more modern concepts.

METHODS

The methods were adapted from previous recommendations and strategies for bibliometric analysis.⁷⁰ An online search was performed utilizing 2 databases: (1) Web of Science (all databases) and (2) Scopus. These databases were selected for their ability to search independent terms and present the number of citations per article. The following search terms and Boolean operators were utilized to identify articles relating to shoulder instability: (shoulder instability) OR (shoulder stabilization) OR (shoulder dislocation) OR (shoulder subluxation) OR (Bankart) OR (glenoid bone loss) OR (shoulder labrum). The search was performed on August 28, 2020, on both databases. All articles extracted from the search were included in initial screening. Articles were sorted from most to least number of citations on each database individually. Three authors (S.A., B.T.F., and D.A.L.) evaluated up to 100 articles from each database for inclusion and relation to shoulder instability. Articles relating to shoulder arthroplasty were excluded. The number of citations from each database was averaged for each study, and the articles were re-sorted using the new means from most to least citations to generate the top 50 list.

In addition to the number of citations on each database, the following data were collected from each article: digital object identifier or PubMed identifier, title, author list, year of publication, journal name, country of origin, study type (ie, case series, cohort study, case control, randomized controlled trial, biomechanical/cadaveric, imaging/ diagnostic, technique/review), study category (diagnostic, prognostic, therapeutic), and level of evidence. Each article was assigned a decade of publication based on the publication year. The level of evidence was agreed upon by the authors; the level of evidence was not scored for biomechanical/cadaveric studies or technique/review studies. The methodological quality of studies was evaluated using the modified Coleman Methodology Score (mCMS) (scoring range, 0-100), a modified version of the Jadad scale (scoring range, 0-8), and Methodological Index for Non-Randomized Studies (MINORS) (scoring range, 0-16 for noncomparative studies and 0-24 for comparative studies).^{19,37,69} mCMS was only applied to studies on surgical treatment, and MINORS was not applied to randomized trials. Biomechanical/cadaveric studies, imaging/diagnostic studies, and technique/reviews were not assessed for methodological quality. The number of citations per year (citation density) was calculated from the mean obtained divided by the number of years since publication.

The search was repeated with the years of inclusion limited to between 2010 and 2020 to generate a top 10 list of more recent publications, which were ranked by citation density.

Data were aggregated and presented utilizing descriptive statistics. The Spearman correlation coefficient (r_s) was used to evaluate correlations between citation values (mean number citations, citation density) and level of evidence and methodological quality, in addition to between year of publication and level of evidence. The strength of correlation was denoted by the absolute value of r_s : weak, <0.4; moderate, 0.4 to 0.7; and strong, >0.7.²¹ One-way analysis of variance was performed to evaluate for differences in articles between citations per year and level of evidence. Independent-samples t tests were performed to compare the citation density of the top 10 from the overall top 50 cited articles list and the top 10 between 2010 and 2020. Significance was set at P < .05 for 2-tailed testing. Statistical analysis was performed on STATA v16.1 (Stata-Corp, College Station, TX).

^{*}Address correspondence to Sachin Allahabadi, MD, UCSF Department of Orthopaedic Surgery, 500 Parnassus Avenue, MU 320-W, San Francisco, CA 94143, USA (email: sachin.allahabadi@gmail.com) (Twitter: Sachin Allahabadi: @sach_MD; Brian Feeley: @DrBrianFeeley).

[†]Department of Orthopaedic Surgery, University of California, San Francisco, San Francisco, California, USA.

Final revision submitted August 29, 2020; accepted November 3, 2020.

One or more of the authors has declared the following potential conflict of interest or source of funding: B.T.F. has received consulting fees from Kaliber and hospitality and education payments from Zimmer Biomet. D.A.L. has received research/education funding from Arthrex and Smith & Nephew; education payments from Medwest; and hospitality payments from Wright Medical. AOSSM checks author disclosures against the Open Payments Database (OPD). AOSSM has not conducted an independent investigation on the OPD and disclaims any liability or responsibility relating thereto.

Ethical approval was not sought for the present study.




Figure 1. Number of 50 most cited articles relating to shoulder instability by decade of publication.

RESULTS

The top 50 most cited articles relating to shoulder instability were published between 1923 and 2010 (Appendix Table A1). The mean difference in citation numbers between the Web of Science and Scopus databases was 59.3 ± 31.2 (range, 12-139). The Scopus database yielded a larger number of citations than did the Web of Science database for all articles except 1.⁷

The majority of publications (n = 33; 66%) were published in the 2 decades between 1990 and 2009 (Figure 1). Articles were published in 12 different journals. The majority (n = 27/50; 54%) were published in the affiliated journals of *The Journal of Bone and Joint Surgery* (Figure 2). The most common country of origin was the United States, representing 37 of 50 (74%) publications (Figure 3).

The mean \pm SD number of citations was 381.5 ± 166.7 (range, 240-1015) (Appendix Table A1). The mean \pm SD citation density was 15.0 ± 8.8 citations/y (range, 4.1-50.8 citations/y), with the same paper¹³ being the most cited for both mean number of citations and citation density.

The majority of studies evaluated for level of evidence were Level 4, with only 3 studies (6%) considered Level 1 (Figure 4). Of the studies, 30% were classified as case series, and 30% were classified as biomechanical/cadaveric (Figure 5). The majority of studies were categorized as diagnostic (n = 28/50; 56%). Two studies (4%) were on posterior instability.^{30,52}

A total of 24 studies were assessed for methodological quality in total (Appendix Table A1). The 26 studies not evaluated for methodological quality included 15 biomechanical/cadaveric, 7 technique/review, 3 imaging/diagnostic, and 1 descriptive epidemiology.

The mean \pm SD mCMS was 54.4 \pm 12.7 (range, 26-82). The mean \pm SD modified Jadad scale score was 3.1 \pm 1.4 (range, 1-6). The mean \pm SD MINORS score was 10.5 \pm 3.3 (range, 7-19). There was no significant correlation between year of publication and level of evidence ($r_{\rm S} = -0.13$; P = .53). There was a statistically significant, although weak, correlation between citation rank and year of publication ($r_{\rm S} = 0.32$; P = .022. There was a weak correlation




Figure 2. Number of 50 most cited articles relating to shoulder instability by journal of publication. AJR, American Journal of Roentgenology; AJSM, American Journal of Sports Medicine; Arthroscopy, The Journal of Arthroscopic and Related Surgery; BJS, British Journal of Surgery; BMJ, British Medical Journal; CORR, Clinical Orthopaedics and Related Research; JOR, Journal of Orthopaedic Research; JBJS, Journal of Bone and Joint Surgery (-Am, American Volume; -Br, British Volume); JSES, Journal of Shoulder and Elbow Surgery; Orthop Rev, Orthopedic Reviews.

between citation rank and level of evidence ($r_{\rm S} = -0.38$; P = .047). The correlation between citation density and year of publication was moderate and statistically significant ($r_{\rm S} = 0.70$; P < .0001). Tables 1 and 2 demonstrate the Spearman correlation coefficients for the mean number of citations and citation density.

There was no significant difference in citation density by level of evidence group (P = .77).

The top 10 most cited articles by citation density between 2010 and 2020^{\ddagger} are listed in Table 3. The mean \pm SD citation density of the top 10 between 2010 and 2020 was 23.8 ± 5.3 compared with 28.8 ± 9.5 for the top 10 citations by density from the overall top 50 list (P = .16). The top cited article from each year between 2015 and 2019^{3,23,35,68,80} is demonstrated in Table 4.

DISCUSSION

The present study identified the top 50 articles relating to native shoulder instability. The vast majority of the most cited articles on shoulder instability are of a lower level of evidence, with only 3 included studies that were level I evidence. Of the studies for which methodological quality was assessed, there were no significant correlations between methodology scores and number of citations.

The topics within the realm of shoulder instability represented in the presented list are vast, ranging from studies critical to the anatomic considerations to those on the natural history of shoulder instability and those on outcomes after surgery. In the present study of the top 50 articles,

[‡]References 22, 27, 28, 34, 45, 53, 58, 67, 68, 79.




Figure 3. Map of countries of origin for the most cited articles pertaining to shoulder instability.




Figure 4. Number of 50 most cited articles relating to shoulder instability by assigned level of evidence. Articles classified as biomechanical/cadaveric or technique/review were excluded from level of evidence analysis (n = 22 excluded; n = 28 included in level of evidence analysis).

15 (30%) were classified as biomechanical/cadaveric, whereas in a study on most cited articles on rotator cuff surgery, 8 articles (16%) were classified as animal or cadaveric.⁷⁰ Although the inclusion of articles was confounded by search terms used, author screening methods, and article classifications, the higher proportion of biomechanical/




Figure 5. Number of 50 most cited articles relating to shoulder instability by study design type. RCT, randomized controlled trial.

cadaveric studies on shoulder instability may reflect the relative effect of studies illuminating the mechanisms underlying the unstable shoulder.

Only 2 (4%) of the studies were specifically on posterior instability, which reflects the epidemiology of the pathology relative to anterior instability.^{30,52} McLaughlin⁵² initially described posterior instability as a "diagnostic trap for the unwary surgeon" in 1952, citing a rate of 3.8% of unstable shoulders with posterior instability at that time. Similarly low rates relative to those of anterior instability have been

TABLE 1 Spearman Correlations Between Number of Citations and Level of Evidence and Methodological Quality Assessments^a

Correlate with Rank of Mean No. of Citations	No. of Observations	Spearman Coefficient $(r_{\rm S})$	P Value
Year of publication	50	$\begin{array}{c} 0.32 \\ -0.38 \\ 0.35 \\ 0.21 \\ 0.35 \end{array}$	$.022^b$
Level of evidence	28		$.047^b$
mCMS	20		.13
Jadad	24		.33
MINORS	21		.12

^amCMS, modified Coleman Methodology Score; MINORS, Methodological Index for Non-Randomized Studies.

^bSignificant for 2-tailed alpha (P < .05).

TABLE 2 Spearman Correlations Between Citation Density and Level of Evidence and Methodological Quality Assessments^a

Correlate With Citation Density (Citations/y)	No. of Observations	Spearman Coefficient (r_S)	P Value
Year of publication	50	0.70	$<.0001^{b}$
Level of evidence	28	0.09	.63
mCMS	20	0.39	.088
Jadad	24	0.36	.080
MINORS	21	0.29	.21

^amCMS, modified Coleman Methodology Score; MINORS, Methodological Index for Non-Randomized Studies.

^bSignificant for 2-tailed alpha (P < .05).

cited since then, with a slightly higher incidence in young athletic populations.^{56,57} While the proportion of articles in the top 50 on posterior shoulder instability was small, there has been increasing attention on the pathology in recent years.

The majority of studies included were published between 1999 and 2009. Relative to the date of the present evaluation, 1999 to 2009 would allow for approximately 10 to 20 years for studies to be cited. This would allow for not only some recency but also adequate time for citations to accrue. In the same fashion, more recent studies would be less likely to be included in the top 50 mosts cited studies based on number of citations. Of note, the Multicenter Orthopaedic Outcomes Network (MOON) Shoulder Instability Cohort has published several recent studies with the largest cohorts of patients undergoing shoulder stabilization to date. but none of the MOON group papers were in the top 50, likely attributable to their recency.⁴³ Additional modern concepts that were not emphasized in this list include that of the glenoid track, with only the original article from 2007 focusing on this subject.⁷⁸ Correspondingly, descriptions of Hill-Sachs lesions were also not emphasized in this list, although the top cited article does reference these lesions.¹³

TABLE 3
10 Most Cited Articles by Approximate Citation Density,
2010-2020

2010-2020							
Rank	Article	Mean No. of Citations	Citation Density, Citations/y	General Topic			
1	Di Giacomo et al, 2014 ²²	195	32.5	Bone loss; on-track/off- track lesions			
2	Griesser et al, 2013 ²⁷	209	29.9	Bristow-Latarjet procedure			
3	Shaha et al, 2015 ⁶⁸	139	27.8	Bone loss			
4	Zacchilli and Owens, 2010 ^{79,a}	259.5	26.0	Epidemiology			
5	$\begin{array}{c} \text{Mizuno et al,} \\ 2014^{53} \end{array}$	152.5	25.4	Latarjet procedure			
6	Shah et al, 2012 ⁶⁷	165.5	20.7	Latarjet procedure			
7	Harris et al, 2013 ²⁸	141	20.1	Bankart stabilization			
8	Provencher et al, 2010 ⁵⁸	200.5	20.1	Bone loss			
9	Hovelius et al, 2012^{34}	142.5	17.8	Bristow-Latarjet procedure; bone block			
10	Lafosse and Boyle, 2010 ⁴⁵	171.5	17.2	Latarjet procedure			

 $^a{\rm A}$ publication by Zacchilli and Owens⁷⁹ that was included in the overall top 50 list was ranked fourth on citation density in this time frame.

TABLE 4
Top Cited Article From Each Year, 2015-2019

Year	Article	Mean No. of Citations	Citation Density, Citations/y	General Topic
2015	Shaha et al ⁶⁸	139	27.8	Bone loss
2016	Zimmermann et al ⁸⁰	76	19	Latarjet procedure; Bankart stabilization
2017	Anthony et al ³	51	17	Patient-reported outcomes
2018	Flinkkilä et al ²³	21.5	10.8	Bankart stabilization
2019	Hurley et al ³⁵	24	24	Latarjet procedure

When evaluating the top 10 most cited articles by citation density in more recent years (2010-2020), we found an increased emphasis on the topics of bone loss and Latarjet procedures relative to the overall top 50 most cited articles list. It is likely that future analysis of more recent years will highlight concepts of glenoid and bipolar bone loss, glenoid track, bone block procedures, and remplissage. The most common journals in which the studies were published were affiliations of *The Journal of Bone and Joint Surgery* and *The American Journal of Sports Medicine*. Each of these journals has high impact factors in orthopaedics and therefore would be expected to contain meaningful or influential papers.

For the papers describing shoulder instability, we did not identify a relationship between the number of citations and any measure of study quality, including level of evidence and various scoring systems for study quality. This finding is in contrast to a previous report on influential articles in rotator cuff surgery.⁷⁰ In that study, a weak correlation was identified between number of citations and Jadad score (r =-0.36).⁷⁰ However, no significant correlations were identified for mCMS or MINORS score.⁷⁰ Therefore, the methodological quality of a study may not appropriately capture a study's literature influence, at least that measured by citations. The effect of a study may derive from its contribution to foundational understanding or development of new concepts from which further higher-level research may arise. Longer-term follow-up studies are more likely to be retrospective in nature and therefore have a lower level of evidence that may not capture the benefit of elucidating long-term outcomes. The lack of correlation between methodological quality and citations also highlights the trend of lower quality studies in orthopaedics in general.^{47,54}

The studies identified through this search were also different from previous bibliometric publications. In a recent update article citing the top 100 most cited articles in orthopaedic surgery,⁵⁰ the only study on shoulder instability included was that by Rowe et al⁶³ on the Bankart procedure, which was ranked as the second-most cited article on shoulder instability in the present study. Interestingly, the most frequently cited article in the present study, which was by Burkhart et al,¹³ was not present in the most recently published or previously published top 100 most cited articles lists. 40,47,50 A study by Namdari et al⁵⁴ evaluating the top 50 most cited articles in shoulder surgery also did not include Burkhart et al.¹³ Differences in study technique, including databases searched and search terms, likely accounted for the discrepancies in articles included. Namdari et al⁵⁴ and Lum et al⁵⁰ both utilized the Web of Science database for expanded citations of articles published in journals classified as orthopaedics. On the other hand, the methods of Sochacki et al⁷⁰ and the present study included a search for specific terms among 2 databases with averaging of citation numbers to generate the highly cited article list. However, 11 of the listed top 50 (22%) cited studies in shoulder surgery were also aggregated in the present study.[§]

This study has several limitations that largely derive from the nature of the study. The decision to analyze the top 50 most cited articles was arbitrary, and numerous influential and groundbreaking studies exist outside of the top 50. There was inherent bias toward studies published earlier, with more recent studies unlikely to make the top 50 most cited articles list. For example, the most recently published study in the overall top 50 list was from 2010.79 Current topics of interest, such as those on subcritical glenoid bone loss, glenoid track, remplissage, and arthroscopic bone block procedures, were minimally represented. Earlier published studies may have been cited frequently for historical purposes, although there may have been less current clinical relevance of these papers. A "snowball" effect may have occurred where new articles cited review papers more frequently than previously.⁴⁴ On the contrary, it is also possible that over time, more historical studies began to be cited less frequently as their contents became a part of the accepted body of knowledge. The number of citations an article has is always in flux, and perhaps analyzing citation density within a shorter time frame (ie, first 5 years after publication) may be more indicative of the current or immediate effect and relevance of a study. In the present study, the correlations evaluated were limited by small numbers because of the higher proportion of studies that were not assessed for level of evidence or methodological quality.

CONCLUSION

Influential articles in shoulder instability included a high proportion of biomechanical/cadaveric studies. The majority of data in top cited articles in shoulder instability were lower-level evidence and poorer methodological quality without strong correlation with citations or citation density. There was a moderate correlation between citation density and year of publication.

REFERENCES

- Allain J, Goutallier D, Glorion C. Long-term results of the Latarjet procedure for the treatment of anterior instability of the shoulder. *J Bone Joint Surg Am.* 1998;80(6):841-852.
- Altchek DW, Warren RF, Skyhar MJ, Ortiz G. T-plasty modification of the Bankart procedure for multidirectional instability of the anterior and inferior types. J Bone Joint Surg Am. 1991;73(1):105-112.
- Anthony CA, Glass NA, Hancock K, Bollier M, Wolf BR, Hettrich CM. Performance of PROMIS instruments in patients with shoulder instability. *Am J Sports Med.* 2017;45(2):449-453.
- Arciero RA, Wheeler JH, Ryan JB, McBride JT. Arthroscopic Bankart repair versus nonoperative treatment for acute, initial anterior shoulder dislocations. *Am J Sports Med.* 1994;22(5):589-594.
- Balg F, Boileau P. The Instability Severity Index Score: a simple preoperative score to select patients for arthroscopic or open shoulder stabilisation. J Bone Joint Surg Br. 2007;89(11):1470-1477.
- 6. Bankart AS. Recurrent or habitual dislocation of the shoulder-joint. *Br Med J.* 1923;2(3285):1132-1133.
- Bankart ASB. The pathology and treatment of recurrent dislocation of the shoulder-joint. Br J Surg. 1938;26(101):23-29.
- Bigliani LU, Newton PM, Steinmann SP, Connor PM, McIlveen SJ. Glenoid rim lesions associated with recurrent anterior dislocation of the shoulder. *Am J Sports Med.* 1998;26(1):41-45.
- Bigliani LU, Pollock RG, Soslowsky LJ, Flatow EL, Pawluk RJ, Mow VC. Tensile properties of the inferior glenohumeral ligament. *J Orthop Res.* 1992;10(2):187-197.
- Boileau P, Villalba M, Héry J-Y, Balg F, Ahrens P, Neyton L. Risk factors for recurrence of shoulder instability after arthroscopic Bankart repair. J Bone Joint Surg Am. 2006;88(8):1755-1763.
- 11. Bottoni CR, Wilckens JH, DeBerardino TM, et al. A prospective, randomized evaluation of arthroscopic stabilization versus nonoperative

[§]References 26, 29, 32, 55, 62-64, 66, 74-76.

treatment in patients with acute, traumatic, first-time shoulder dislocations. Am J Sports Med. 2002;30(4):576-580.

- Bozzo A, Oitment C, Evaniew N, Ghert M. The journal impact factor of orthopaedic journals does not predict individual paper citation rate. *J Am Acad Orthop Surg Glob Res Rev.* 2017;1(2):e007.
- Burkhart SS, De Beer JF. Traumatic glenohumeral bone defects and their relationship to failure of arthroscopic Bankart repairs: significance of the inverted-pear glenoid and the humeral engaging Hill-Sachs lesion. *Arthroscopy*. 2000;16(7):677-694.
- 14. Burkhart SS, De Beer JF, Barth JRH, et al. Results of modified Latarjet reconstruction in patients with anteroinferior instability and significant bone loss. *Arthroscopy*. 2007;23(10):1033-1041.
- Burkhart SS, Debeer JF, Tehrany AM, Parten PM. Quantifying glenoid bone loss arthroscopically in shoulder instability. *Arthroscopy*. 2002; 18(5):488-491.
- Burkhead WZ, Rockwood CA. Treatment of instability of the shoulder with an exercise program. J Bone Joint Surg Am. 1992;74(6):890-896.
- Chandnani VP, Yeager TD, DeBerardino T, et al. Glenoid labral tears: prospective evaluation with MRI imaging, MR arthrography, and CT arthrography. *AJR Am J Roentgenol*. 1993;161(6):1229-1235.
- Cole BJ, L'Insalata J, Irrgang J, Warner JJ. Comparison of arthroscopic and open anterior shoulder stabilization: a two to six-year follow-up study. *J Bone Joint Surg Am*. 2000;82(8):1108-1114.
- Coleman BD, Khan KM, Maffulli N, Cook JL, Wark JD. Studies of surgical outcome after patellar tendinopathy: clinical significance of methodological deficiencies and guidelines for future studies. Victorian Institute of Sport Tendon Study Group. *Scand J Med Sci Sports*. 2000;10(1):2-11.
- Cooper DE, Arnoczky SP, O'Brien SJ, Warren RF, DiCarlo E, Allen AA. Anatomy, histology, and vascularity of the glenoid labrum: an anatomical study. *J Bone Joint Surg Am.* 1992;74(1):46-52.
- 21. Dancey C, Reidy J. Statistics Without Maths for Psychology. 7th ed. Pearson; 2017.
- Di Giacomo G, Itoi E, Burkhart SS. Evolving concept of bipolar bone loss and the Hill-Sachs lesion: from "engaging/non-engaging" lesion to "on-track/off-track" lesion. *Arthroscopy*. 2014;30(1):90-98.
- Flinkkilä T, Knape R, Sirniö K, Ohtonen P, Leppilahti J. Long-term results of arthroscopic Bankart repair: minimum 10 years of followup. *Knee Surg Sports Traumatol Arthrosc.* 2018;26(1):94-99.
- 24. Gartsman GM, Roddey TS, Hammerman SM. Arthroscopic treatment of anterior-inferior glenohumeral instability: two to five-year follow-up. *J Bone Joint Surg Am.* 2000;82(7):991-1003.
- 25. Gerber C, Nyffeler RW. Classification of glenohumeral joint instability. *Clin Orthop Relat Res.* 2002;400:65-76.
- Glousman R, Jobe F, Tibone J, Moynes D, Antonelli D, Perry J. Dynamic electromyographic analysis of the throwing shoulder with glenohumeral instability. J Bone Joint Surg Am. 1988;70(2):220-226.
- Griesser MJ, Harris JD, McCoy BW, et al. Complications and reoperations after Bristow-Latarjet shoulder stabilization: a systematic review. J Shoulder Elbow Surg. 2013;22(2):286-292.
- Harris JD, Gupta AK, Mall NA, et al. Long-term outcomes after Bankart shoulder stabilization. *Arthroscopy*. 2013;29(5):920-933.
- Harryman DT, Sidles JA, Clark JM, McQuade KJ, Gibb TD, Matsen FA. Translation of the humeral head on the glenoid with passive glenohumeral motion. *J Bone Joint Surg Am*. 1990;72(9): 1334-1343.
- Hawkins RJ, Neer CS, Pianta RM, Mendoza FX. Locked posterior dislocation of the shoulder. *J Bone Joint Surg Am.* 1987;69(1): 9-18.
- Helfet AJ. Coracoid transplantation for recurring dislocation of the shoulder. J Bone Joint Surg Br. 1958;40(2):198-202.
- Hovelius L, Augustini BG, Fredin H, Johansson O, Norlin R, Thorling J. Primary anterior dislocation of the shoulder in young patients: a tenyear prospective study. *J Bone Joint Surg Am.* 1996;78(11): 1677-1684.
- Hovelius L, Eriksson K, Fredin H, et al. Recurrences after initial dislocation of the shoulder: results of a prospective study of treatment. *J Bone Joint Surg Am.* 1983;65(3):343-349.

- 34. Hovelius L, Sandström B, Olofsson A, Svensson O, Rahme H. The effect of capsular repair, bone block healing, and position on the results of the Bristow-Latarjet procedure (study III): long-term follow-up in 319 shoulders. *J Shoulder Elbow Surg.* 2012;21(5): 647-660.
- Hurley ET, Lim Fat D, Farrington SK, Mullett H. Open versus arthroscopic Latarjet procedure for anterior shoulder instability: a systematic review and meta-analysis. *Am J Sports Med.* 2019;47(5): 1248-1253.
- Itoi E, Lee SB, Berglund LJ, Berge LL, An KN. The effect of a glenoid defect on anteroinferior stability of the shoulder after Bankart repair: a cadaveric study. J Bone Joint Surg Am. 2000;82(1):35-46.
- Jadad AR, Moore RA, Carroll D, et al. Assessing the quality of reports of randomized clinical trials: is blinding necessary? *Control Clin Trials*. 1996;17(1):1-12.
- Jobe FW, Giangarra CE, Kvitne RS, Glousman RE. Anterior capsulolabral reconstruction of the shoulder in athletes in overhand sports. *Am J Sports Med.* 1991;19(5):428-434.
- Jobe FW, Kvitne RS, Giangarra CE. Shoulder pain in the overhand or throwing athlete: the relationship of anterior instability and rotator cuff impingement. *Orthop Rev.* 1989;18(9):963-975.
- Kelly JC, Glynn RW, O'Briain DE, Felle P, McCabe JP. The 100 classic papers of orthopaedic surgery: a bibliometric analysis. *J Bone Joint Surg Br.* 2010;92(10):1338-1343.
- Kirkley A, Griffin S, McLintock H, Ng L. The development and evaluation of a disease-specific quality of life measurement tool for shoulder instability: the Western Ontario Shoulder Instability Index (WOSI). *Am J Sports Med.* 1998;26(6):764-772.
- Kraeutler MJ, Freedman KB, MacLeod RA, Schrock JB, Tjoumakaris FP, McCarty EC. The 50 most cited articles in rotator cuff repair research. *Orthopedics*. 2016;39(6):e1045-e1051.
- Kraeutler MJ, McCarty EC, Belk JW, et al. Descriptive epidemiology of the MOON Shoulder Instability Cohort. *Am J Sports Med.* 2018; 46(5):1064-1069.
- Kuhn TS. Historical structure of scientific discovery. Science. 1962; 136(3518):760-764.
- Lafosse L, Boyle S. Arthroscopic Latarjet procedure. J Shoulder Elbow Surg. 2010;19(2):2-12.
- Latarjet M. Treatment of recurrent dislocation of the shoulder. Lyon Chir. 1954;49(8):994-997.
- 47. Lefaivre KA, Shadgan B, O'Brien PJ. 100 most cited articles in orthopaedic surgery. *Clin Orthop Relat Res.* 2011;469(5):1487-1497.
- Lippitt S, Matsen F. Mechanisms of glenohumeral joint stability. *Clin* Orthop Relat Res. 1993;291:20-28.
- Lo IKY, Parten PM, Burkhart SS. The inverted pear glenoid: an indicator of significant glenoid bone loss. *Arthroscopy*. 2004;20(2): 169-174.
- Lum ZC, Pereira GC, Giordani M, Meehan JP. Top 100 most cited articles in orthopaedic surgery: an update. *J Orthop*. 2020;19: 132-137.
- Marquez-Lara A, Stone AV, Luo TD, Parker BR, Sharma A, Freehill MT. Top 50 cited journal articles on overhead throwing athletes: a bibliographic analysis. *JSES Open Access*. 2017;1(2):55-62.
- McLaughlin HL. Posterior dislocation of the shoulder. J Bone Joint Surg Am. 1952;24(3):584-590.
- Mizuno N, Denard PJ, Raiss P, Melis B, Walch G. Long-term results of the Latarjet procedure for anterior instability of the shoulder. *J Shoulder Elbow Surg.* 2014;23(11):1691-1699.
- Namdari S, Baldwin K, Kovatch K, Huffman GR, Glaser D. Fifty most cited articles in orthopedic shoulder surgery. *J Shoulder Elbow Surg*. 2012;21(12):1796-1802.
- 55. Neer CS, Foster CR. Inferior capsular shift for involuntary inferior and multidirectional instability of the shoulder: a preliminary report. *J Bone Joint Surg Am*. 1980;62(6):897-908.
- Owens BD, Campbell SE, Cameron KL. Risk factors for posterior shoulder instability in young athletes. *Am J Sports Med.* 2013; 41(11):2645-2649.
- Pollock RG, Bigliani LU. Recurrent posterior shoulder instability: diagnosis and treatment. *Clin Orthop Relat Res.* 1993;291:85-96.

- Provencher CMT, Bhatia S, Ghodadra NS, et al. Recurrent shoulder instability: current concepts for evaluation and management of glenoid bone loss. J Bone Joint Surg Am. 2010;92(suppl 2):133-151.
- Randelli P, Cucchi D, Butt U. History of shoulder instability surgery. Knee Surg Sports Traumatol Arthrosc. 2016;24(2):305-329.
- Robinson CM, Howes J, Murdoch H, Will E, Graham C. Functional outcome and risk of recurrent instability after primary traumatic anterior shoulder dislocation in young patients. *J Bone Joint Surg Am*. 2006;88(11):2326-2336.
- Rodosky MW, Harner CD, Fu FH. The role of the long head of the biceps muscle and superior glenoid labrum in anterior stability of the shoulder. *Am J Sports Med.* 1994;22(1):121-130.
- Rowe CR. Prognosis in dislocations of the shoulder. J Bone Joint Surg Am. 1956;38(5):957-977.
- 63. Rowe CR, Patel D, Southmayd WW. The Bankart procedure: a long-term end-result study. *J Bone Joint Surg Am*. 1978;60(1):1-16.
- 64. Rowe CR, Zarins B. Recurrent transient subluxation of the shoulder. *J Bone Joint Surg Am.* 1981;63(6):863-872.
- Rowe CR, Zarins B, Ciullo JV. Recurrent anterior dislocation of the shoulder after surgical repair: apparent causes of failure and treatment. J Bone Joint Surg Am. 1984;66(2):159-168.
- 66. Samilson RL, Prieto V. Dislocation arthropathy of the shoulder. *J Bone Joint Surg Am.* 1983;65(4):456-460.
- Shah AA, Butler RB, Romanowski J, Goel D, Karadagli D, Warner JJ. Short-term complications of the Latarjet procedure. *J Bone Joint Surg Am.* 2012;94(6):495-501.
- Shaha JS, Cook JB, Song DJ, et al. Redefining "critical" bone loss in shoulder instability: functional outcomes worsen with "subcritical" bone loss. *Am J Sports Med.* 2015;43(7):1719-1725.
- Slim K, Nini E, Forestier D, Kwiatkowski F, Panis Y, Chipponi J. Methodological Index for Non-Randomized Studies (MINORS): development and validation of a new instrument. ANZ J Surg. 2003;73(9):712-716.
- Sochacki KR, Jack RA, Nauert R, Harris JD. Correlation between quality of evidence and number of citations in top 50 cited articles in rotator cuff repair surgery. *Orthop J Sports Med.* 2018;6(6): 2325967118776635.

- Speer KP, Deng X, Borrero S, Torzilli PA, Altchek DA, Warren RF. Biomechanical evaluation of a simulated Bankart lesion. *J Bone Joint Surg Am.* 1994;76(12):1819-1826.
- Sugaya H, Moriishi J, Dohi M, Kon Y, Tsuchiya A. Glenoid rim morphology in recurrent anterior glenohumeral instability. *J Bone Joint Surg Am.* 2003;85(5):878-884.
- Taylor DC, Arciero RA. Pathologic changes associated with shoulder dislocations: arthroscopic and physical examination findings in firsttime, traumatic anterior dislocations. *Am J Sports Med.* 1997;25(3): 306-311.
- Turkel SJ, Panio MW, Marshall JL, Girgis FG. Stabilizing mechanisms preventing anterior dislocation of the glenohumeral joint. *J Bone Joint Surg Am.* 1981;63(8):1208-1217.
- Warner JJ, Deng XH, Warren RF, Torzilli PA. Static capsuloligamentous restraints to superior-inferior translation of the glenohumeral joint. *Am J Sports Med.* 1992;20(6):675-685.
- Warner JJ, Micheli LJ, Arslanian LE, Kennedy J, Kennedy R. Patterns of flexibility, laxity, and strength in normal shoulders and shoulders with instability and impingement. *Am J Sports Med.* 1990;18(4): 366-375.
- 77. Warner JJ, Micheli LJ, Arslanian LE, Kennedy J, Kennedy R. Scapulothoracic motion in normal shoulders and shoulders with glenohumeral instability and impingement syndrome: a study using Moiré topographic analysis. *Clin Orthop Relat Res.* 1992; 285:191-199.
- Yamamoto N, Itoi E, Abe H, et al. Contact between the glenoid and the humeral head in abduction, external rotation, and horizontal extension: a new concept of glenoid track. *J Shoulder Elbow Surg*. 2007;16(5):649-656.
- Zacchilli MA, Owens BD. Epidemiology of shoulder dislocations presenting to emergency departments in the United States. *J Bone Joint Surg Am*. 2010;92(3):542-549.
- Zimmermann SM, Scheyerer MJ, Farshad M, Catanzaro S, Rahm S, Gerber C. Long-term restoration of anterior shoulder stability: a retrospective analysis of arthroscopic Bankart repair versus open Latarjet procedure. *J Bone Joint Surg Am.* 2016;98(23):1954-1961.

APPENDIX

TABLE A1
Top 50 Most Cited Articles Pertaining to Shoulder Instability

				Citations					
Rank	Lead Author (Year)	Country	Study Design	Mean No.	Density, Citations/y	LOE	mCMS	Jadad	MINORS
1	Burkhart SS (2000) ¹³	USA	Case series	1015	50.8	4	54	3	10
2	Rowe CR (1978) ⁶³	USA	Case series	969.5	23.1	4	57	3	8
3	Neer CS (1980) ⁵⁵	USA	Case series	690	17.3	4	47	2	9
4	Turkel SJ (1981) ⁷⁴	USA	Biomechanical/cadaveric	636.5	16.3	N/A	N/A	N/A	N/A
5	Bankart ASB (1938) ⁷	UK	Technique/review	588	7.2	N/A	N/A	N/A	N/A
6	Boileau P (2006) ¹⁰	France	Case series	556	39.7	4	65	4	10
7	Harryman DT (1990) ²⁹	USA	Biomechanical/cadaveric	552.5	18.4	N/A	N/A	N/A	N/A
8	Samilson RL (1983) ⁶⁶	USA	Imaging/diagnostic	512	13.8	4	N/A	N/A	N/A
9	Itoi E (2000) ³⁶	USA	Biomechanical/cadaveric	511.5	25.6	N/A	N/A	N/A	N/A
10	Allain J (1998) ¹	France	Case series	440	20	4	59	4	8
11	Hovelius L (1996) ³²	Sweden	RCT	439.5	18.3	1	N/A	6	N/A
12	Sugaya H (2003) ⁷²	Japan	Biomechanical/cadaveric	416.5	24.5	N/A	N/A	N/A	N/A
13	Rowe CR (1956) ⁶²	USA	Case series	409.5	6.4	4	33	1	8
14	Balg F (2007) ⁵	France	Case-control	402.5	31	3	56	3	16
15	Rowe CR (1981) ⁶⁴	USA	Cohort study	398.5	10.2	2	53	2	8
16	Glousman R (1988) ²⁶	USA	Imaging/diagnostic	396.5	12.4	2	N/A	N/A	N/A
17	Bankart AS (1923) ⁶	UK	Technique/review	396	4.1	N/A	N/A	N/A	N/A

(continued)

Rank	Lead Author (Year)	Country	Study Design	Citations					
				Mean No.	Density, Citations/y	LOE	mCMS	Jadad	MINORS
18	Bigliani LU (1992) ⁹	USA	Biomechanical/cadaveric	387.5	13.8	N/A	N/A	N/A	N/A
19	Kirkley A (1998) ⁴¹	Canada	Technique/review	378	17.2	N/A	N/A	N/A	N/A
20	Latarjet M (1954) ⁴⁶	France	Technique/review	370	5.6	N/A	N/A	N/A	N/A
21	Jobe FW (1989) ³⁹	USA	Technique/review	365.5	11.8	N/A	N/A	N/A	N/A
22	Burkhead WZ (1992) ¹⁶	USA	Case series	345	12.3	4	N/A	2	10
23	Rowe CR (1984) ⁶⁵	USA	Case series	343	9.5	4	46	3	8
24	Warner JJ (1990) ⁷⁶	USA	Biomechanical/cadaveric	342.5	11.4	N/A	N/A	N/A	N/A
25	Rodosky MW (1994) ⁶¹	USA	Biomechanical/cadaveric	337	13	N/A	N/A	N/A	N/A
26	Burkhart SS (2007) ¹⁴	USA	Case series	331.5	25.5	4	56	3	7
27	Altchek DW $(1991)^2$	USA	Case series	314.5	10.8	4	50	3	8
28	Bigliani LU (1998) ⁸	USA	Case series	312.5	14.2	4	46	2	10
29	McLaughlin HL $(1952)^{52}$	USA	Case series	305.5	4.5	4	26	2	7
30	Taylor DC (1997) ⁷³	USA	Cohort study	302	13.1	3	62	1	14
31	Arciero RA (1994) ⁴	USA	Cohort study	297	11.4	2	73	3	19
32	Cooper DE (1992) ²⁰	USA	Biomechanical/cadaveric	296.5	10.6	N/A	N/A	N/A	N/A
33	Yamamoto N (2007) ⁷⁸	Japan	Biomechanical/cadaveric	286.5	22	N/A	N/A	N/A	N/A
34	Jobe FW (1991) ³⁸	USA	Case series	283.5	9.8	4	49	2	8
35	Lippitt S (1993) ⁴⁸	USA	Biomechanical/cadaveric	283	10.5	N/A	N/A	N/A	N/A
36	Helfet AJ $(1958)^{31}$	UK	Technique/review	282	4.5	N/A	N/A	N/A	N/A
37	Warner JJ $(1992)^{77}$	USA	Biomechanical/cadaveric	281	10	N/A	N/A	N/A	N/A
38	Warner JJ $(1992)^{75}$	USA	Biomechanical/cadaveric	280	10	N/A	N/A	N/A	N/A
39	Chandnani VP $(1993)^{17}$	USA	Imaging/diagnostic	277.5	10.3	3	N/A	N/A	N/A
40	Hawkins RJ (1987) ³⁰	USA	Case series	268.5	8.1	4	56	2	10
41	Zacchilli MA (2010) ⁷⁹	USA	Descriptive epidemiology	259.5	26	4	N/A	N/A	N/A
42	Burkhart SS $(2002)^{15}$	USA	Biomechanical/cadaveric	253.5	14.1	N/A	N/A	N/A	N/A
43	Gartsman GM $(2000)^{24}$	USA	Case series	250.0 251	12.6	4	49	4	14
44	Cole BJ $(2000)^{18}$	USA	Cohort study	249.5	12.5	3	82	4	14
45	Lo IKY $(2004)^{49}$	USA	Biomechanical/cadaveric	247	15.4	N/A	N/A	N/A	N/A
46	Gerber C $(2002)^{25}$	Switzerland	Technique/review	244.5	13.6	N/A	N/A	N/A	N/A
40 47	Robinson CM (2006) ⁶⁰	UK	Cohort study	244.5 244.5	17.5	2	N/A	4	14
48	Speer KP $(1994)^{71}$	USA	Biomechanical/cadaveric	244.5 243	9.3	N/A	N/A	N/A	N/A
40 49	Hovelius L $(1983)^{33}$	USA	RCT	240.5	5.5 6.5	1	N/A	6	N/A
49 50	Bottoni CR $(2002)^{11}$	USA	RCT	$\frac{240.5}{240}$	13.3	1	69	6	N/A
90	DOLIONI UK (2002)	USA	NU1	240	13.3	T	69	Ø	IN/A

Table A1 (continued)

 a LOE, level of evidence; mCMS, modified Coleman Methodology Score; MINORS, Methodological Index for Non-Randomized Studies; N/A, not available; RCT, randomized controlled trial.