Lawrence Berkeley National Laboratory

Recent Work

Title

INHIBITION OF PHOTOSYNTHESIS IN ISOLATED SPINACH CHLOROPLASTS BY INORGANIC PHOSPHATE OR INORGANIC PYROPHOSPHATASE IN THE PRESENCE OF PYROPHOSPHATE AND MAGNESIUM IONS

Permalink

<https://escholarship.org/uc/item/5rk7q8gr>

Authors

Levine, Gerri
Bassham, J.A.

Publication Date

1973-07-01

INHIBITION OF PHOTOSYNTHESIS IN ISOLATED SPINACH
CHLOROPLASTS BY INORGANIC PHOSPHATE OR
INORGANIC PYROPHOSPHATASE IN THE PRESENCE
OF PYROPHOSPHATE AND MAGNESIUM IONS

Gerri Levine and J. A. Bassham

July 1973

Prepared for the U. S. Atomic Energy Commission
under Contract W-7405-ENG-48

For Reference

Not to be taken from this room

DISCLAIMER

This document was prepared as an account of work sponsored by the United States Government. While this document is believed to contain correct information, neither the United States Government nor any agency thereof, nor the Regents of the University of California, nor any of their employees, makes any warranty, express or implied, or assumes any legal responsibility for the accuracy, completeness, or usefulness of any information, apparatus, product, or process disclosed, or represents that its use would not infringe privately owned rights. Reference herein to any specific commercial product, process, or service by its trade name, trademark, manufacturer, or otherwise, does not necessarily constitute or imply its endorsement, recommendation, or favoring by the United States Government or any agency thereof, or the Regents of the University of California. The views and opinions of authors expressed herein do not necessarily state or reflect those of the United States Government or any agency thereof or the Regents of the University of California.

INHIBITION OF PHOTOSYNTHESIS IN ISOLATED SPINACH CHLOROPLASTS BY
INORGANIC PHOSPHATE OR INORGANIC PYROPHOSPHATASE IN THE PRESENCE
OF PYROPHOSPHATE AND MAGNESIUM IONS

GERRI LEVINE and J. A. BASSHAM

Laboratory of Chemical Biodynamics, Lawrence Berkeley Laboratory,
University of California, Berkeley, California 94720 (U.S.A.)

SUMMARY

Inhibition by inorganic phosphate of photosynthesis by isolated spinach chloroplasts is decreased by the presence of pyrophosphate and increased with increasing Mg^{++} ions. Previously reported regulation of this photosynthesis by protein factors from spinach leaves appears to be due mostly to pyrophosphate phosphohydrolase (E.C. 3.6.1.1) activity which converts pyrophosphate to phosphate and to the effects of pyrophosphate and Mg^{++} on this pyrophosphatase activity.

Spinach leaf juice exerts a strong inhibitory effect on the photosynthetic fixation of CO_2 by isolated spinach chloroplasts in a medium containing inorganic pyrophosphate (PP_i)¹. One of the inhibitory factors was found to be magnesium ions, while another factor was a

Abbreviations: FDPase, D-fructose-1,6-diphosphate 1-phosphohydrolase; PP_i , inorganic pyrophosphate; P_i , orthophosphate.

protein^{2,3}. In the presence of an appropriate amount of inorganic pyrophosphate, a concentration of Mg^{++} about equal to or slightly less than the pyrophosphate concentration could stimulate the rate of CO_2 fixation, while addition of a small amount of the protein factor, in the absence of Mg^{++} ions, could also stimulate the rate of CO_2 fixation. The presence of these amounts of Mg^{++} and protein factor together could cause a very large (90% or more) inhibition of the rate of CO_2 inhibition ("synergistic inhibition").

Other characteristics of this inhibitory effect are: (1) a requirement for a low level of PP_i , (2) the inhibition can be overcome by higher levels of PP_i , (3) either the protein factor or Mg^{++} alone can inhibit when added in higher concentrations, and (4) inhibition by Mg^{++} alone, protein factor alone, or a combination of the two is accompanied by increased movement of intermediate compounds of the reductive pentose phosphate cycle, especially dihydroxyacetone phosphate, out of the chloroplasts into the suspending medium³.

A partially purified preparation of D-fructose-1,6-diphosphate 1-phosphohydrolase, E.C.3.1.3.11 (FDPase), was found to reproduce the activity of the protein factor, leading at first to a conclusion that the protein factor might be FDPase². However, heat treatment which denatured the protein and completely destroyed the FDPase activity released a factor which still exhibited inhibitory activity³. Moreover, the protein factor could be separated from FDPase activity by gel filtration⁴, and has a lower molecular weight than FDPase.

A Mg^{++} dependent pyrophosphate phosphohydrolase, E.C. 3.6.1.1 (inorganic pyrophosphatase), was previously isolated from spinach chloro-

plasts by procedures which follow some of the same steps that are used in the purification of the protein factor⁵. The protein factor contains considerable pyrophosphatase activity, and significant amounts of inorganic phosphate are released by the hydrolysis of PP_i under the conditions used by us in our studies. Photosynthesis by isolated chloroplasts is inhibited by such levels of inorganic phosphate. The synergistic inhibitory activity we reported appears to be due to the stimulation of pyrophosphatase activity by Mg^{++} , resulting in the release of inorganic phosphate into the medium. Pyrophosphate reduces the inhibitory effect of inorganic phosphate on chloroplast CO_2 fixation.

EXPERIMENTAL

Isolation of inhibition protein factor

A partially purified preparation of FDPase was prepared from spinach leaves as previously reported², except that the spinach leaves were ground and used directly (no chloroplast isolation) and the $(NH_4)_2SO_4$ precipitation step was eliminated. Gel filtration on Sephadex G-150 (instead of G-200) resulted in separation of an FDPase-active fraction from a smaller molecular weight protein fraction, which was inhibitory to CO_2 fixation in isolated chloroplasts. The further purified FDPase fraction did not cause such inhibition. The inhibitory protein factor has pyrophosphatase activity.

Chloroplast photosynthesis

Chloroplasts were prepared from fresh spinach as described earlier⁶. These chloroplasts were suspended in Solution C and placed in small serum-stoppered flasks (total suspension volume, 0.5 ml) on a shaker

over an illuminated light table⁶. Varying amounts of protein factors, inorganic pyrophosphatase, or of inorganic phosphate were added to flasks as indicated in the tables. After 5 min preillumination, 25 μ l of solution of $\text{NaH}^{14}\text{CO}_3$ (0.15 mM, 20 $\mu\text{C}/\mu\text{mole}$) was added and the chloroplasts were allowed to photosynthesize for 5 min before killing with 4.5 ml methanol.

For determination of acid-stable radioactive compounds formed, an aliquot portion of the killed chloroplast suspension was run onto filter paper, two drops of glacial acetic acid were added, and the wet area of filter paper was dried with a stream of N_2 . After standing several hours, the radioactivity on the paper was measured with a pair of thin-window GM tubes. Chlorophyll content was determined, and $^{14}\text{CO}_2$ fixation was expressed as $\mu\text{moles } ^{14}\text{CO}_2$ fixed per mg of chlorophyll per hour⁶.

Assay for pyrophosphatase activity

Inorganic pyrophosphatase activity was assayed by measuring inorganic phosphate released into the reaction mixture⁷. Yeast inorganic pyrophosphatase was purchased from Worthington Biochemical Corp.

Comparison of effect of protein factor and of inorganic phosphate on chloroplast photosynthesis

In our usual assay of protein factor activity, the chloroplast preparation is in contact with the protein for a total of 10 min (5 min preillumination and 5 min with $\text{H}^{14}\text{CO}_3^-$) for measurement of fixation. The amounts of protein used in our experiments would release a total amount of inorganic phosphate sufficient to bring the concentration to 5.6 mM in the flask by the end of the 10-min period. We added amounts of P_i

sufficient to give concentrations ranging from 0.5 mM to 8 mM to the flasks at the beginning of the preillumination period.

RESULTS

Up to 93% decrease in the rate of $^{14}\text{CO}_2$ uptake is seen with 10.5 mM inorganic phosphate and no PP_i (Table I). These are the concentrations which would result from the complete conversion of 5 mM PP_i to P_i (the solution C contains 0.5 mM P_i). Thus the severe inhibition reported to result from addition of protein factor plus Mg^{++} can be accounted for if the additions result in substantial hydrolysis of PP_i . It is noteworthy that, at all levels of P_i , inhibition is greater with 4 mM Mg^{++} than with 1 mM Mg^{++} , and is less in the presence of PP_i than in the absence of PP_i . Thus there are some effects of Mg^{++} and PP_i on the inhibition by P_i which do not require any exogenous pyrophosphatase activity.

The "synergistic inhibitory activity" seen upon addition of protein factor (now known to contain inorganic pyrophosphatase activity) plus 4 mM Mg^{++} can be exactly duplicated by the addition of purified yeast pyrophosphatase (Table II). Little inhibition is seen with 1 mM Mg^{++} plus either protein factor or yeast pyrophosphatase.

DISCUSSION

The "synergistic" inhibition of CO_2 fixation in isolated spinach chloroplasts caused by addition of "protein factor" plus 4 mM Mg^{++} in the presence of 5 mM PP_i is now seen to be the result of pyrophosphatase activity resulting in conversion of 5 mM PP_i to 10 mM P_i , which severely inhibits CO_2 fixation. The "synergistic" requirement for Mg^{++}

is not required for the phosphate inhibition, since one of the roles of Mg^{++} is to activate the pyrophosphatase⁵. There remains the observation that Mg^{++} alone (with low levels of PP_i) can stimulate (at low levels) or inhibit (at higher levels) CO_2 fixation by isolated chloroplasts. Also, Mg^{++} alone increases the rate of movement of metabolites out of the chloroplasts, while increased PP_i (with low levels of Mg^{++}) has the opposite effects. These phenomena are perhaps due, in part, to endogenous pyrophosphatase and the resulting inorganic phosphate released, and to the effects of Mg^{++} in activating this enzyme. Excess PP_i could inactivate the enzyme by complexing Mg^{++} .

That we failed to accept this rather simple explanation of the effects earlier was due in part to an early observed severe "synergistic" inhibition with only 1 mM PP_i plus 1.5 mM Mg^{++} ³, which could only give a maximum of 2 mM inorganic phosphate. This level of inorganic phosphate does not cause severe inhibition. However, with spinach chloroplasts currently being isolated in this laboratory, severe synergistic inhibition with protein factor or yeast inorganic pyrophosphatase plus 1 mM PP_i and 1.5 mM Mg^{++} is no longer observed, despite many experiments. Presumably this is yet another example of the variability of isolated spinach chloroplast preparations over the seasons and with differently grown spinach. It is possible that the impure factor used in earlier experiments contained factors other than pyrophosphatase which influence the rate of photosynthesis during the first 5 min. However, the severe inhibitory effect, dependent on Mg^{++} and PP_i concentrations, appears to be due for the most part to the inorganic pyrophosphatase as activated by Mg^{++} in the presence of low levels of PP_i .

REFERENCES

- 1 Moore, R. E., Springer-Lederer, H., Ottenheim, H.C.J. and Bassham, J. A. (1969) Biochim. Biophys. Acta 180, 368-376
- 2 Springer-Lederer, H., El Badry, A. M., Ottenheim, H.C.J. and Bassham, J. A. (1969) Biochim. Biophys. Acta 189, 464-467
- 3 Bassham, J. A., El Badry, A. M., Kirk, M. R., Ottenheim, H.C.J. and Springer-Lederer, H. (1970) Biochim. Biophys. Acta 223, 261-274
- 4 Bassham, J. A. (1971) in Energy Transduction in Respiration and Photosynthesis (Quagliariello, E., Papa, S. and Rossi, C. S., eds.), pp. 869-882, Adriatica Editrice, Bari
- 5 El Badry, A. M. and Bassham, J. A. (1970) Biochim. Biophys. Acta 197, 308-316
- 6 Jensen, R. G. and Bassham, J. A. (1966) Proc. Nat. Acad. Sci. U.S. 56, 1095-1101
- 7 Chen, P. S., Jr., Torebara, T. Y. and Warner, H. (1956) Anal. Chem. 28, 1756-1758

*TABLE I

EFFECTS OF P_i , PP_i AND Mg^{++} IONS ON RATE OF PHOTOSYNTHESIS BY ISOLATED SPINACH CHLOROPLASTS

Relative rates*			
Mg^{++}	Phosphate	$-PP_i$	+5 mM PP_i
1.0	0.5	.82	1.00
1.0	1.5	.73	.95
1.0	2.5	.63	.87
1.0	5.5	.35	.63
1.0	10.5	.065	.31
4.0	0.5	.47	.96
4.0	1.5	.32	.89
4.0	2.5	.21	.73
4.0	5.5	.079	.49
4.0	10.5	.027	.18

*Control rate (1.00 in table) was 170 μ moles of $^{14}CO_2/hr \cdot m$ g Chl.

TABLE II

EFFECTS OF "PROTEIN FACTOR" AND OF YEAST PYROPHOSPHATASE ON RATE OF PHOTOSYNTHESIS BY ISOLATED SPINACH CHLOROPLASTS

	Relative rates	
	1 mM Mg ⁺⁺	4 mM Mg ⁺⁺
No addition	1.00*	0.76
Protein factor (240 µg)	0.83	0.14
Yeast pyrophosphatase (0.5 µg)	0.93	0.14

*Control rate (1.00 in table) was 125 µmoles ¹⁴CO₂/hr·mg Chl.

Concentration of PP_i was 5.0 mM. The pyrophosphatase activity of the protein factor was 2.8 units/mg, while that of yeast pyrophosphatase was 600 units/mg (one unit = 1 µmole P_i released/min).

LEGAL NOTICE

This report was prepared as an account of work sponsored by the United States Government. Neither the United States nor the United States Atomic Energy Commission, nor any of their employees, nor any of their contractors, subcontractors, or their employees, makes any warranty, express or implied, or assumes any legal liability or responsibility for the accuracy, completeness or usefulness of any information, apparatus, product or process disclosed, or represents that its use would not infringe privately owned rights.

TECHNICAL INFORMATION DIVISION
LAWRENCE BERKELEY LABORATORY
UNIVERSITY OF CALIFORNIA
BERKELEY, CALIFORNIA 94720