

UC Berkeley

UC Berkeley Previously Published Works

Title

Study of CP-violating asymmetries in $B^0 \rightarrow \pi^+ \pi^-$, $K^+ \pi^-$ decays

Permalink

<https://escholarship.org/uc/item/5s21h7ww>

Journal

Physical Review D, 65(5)

ISSN

2470-0010

Authors

Aubert, B
Boutigny, D
Gaillard, J-M
[et al.](#)

Publication Date

2002-03-01

DOI

10.1103/physrevd.65.051502

Copyright Information

This work is made available under the terms of a Creative Commons Attribution License, available at <https://creativecommons.org/licenses/by/4.0/>

Peer reviewed

Study of CP -violating asymmetries in $B^0 \rightarrow \pi^+ \pi^-$, $K^+ \pi^-$ decays

B. Aubert,¹ D. Boutigny,¹ J.-M. Gaillard,¹ A. Hicheur,¹ Y. Karyotakis,¹ J. P. Lees,¹ P. Robbe,¹ V. Tisserand,¹ A. Palano,² A. Pompili,² G. P. Chen,³ J. C. Chen,³ N. D. Qi,³ G. Rong,³ P. Wang,³ Y. S. Zhu,³ G. Eigen,⁴ B. Stugu,⁴ G. S. Abrams,⁵ A. W. Borgland,⁵ A. B. Breon,⁵ D. N. Brown,⁵ J. Button-Shafer,⁵ R. N. Cahn,⁵ A. R. Clark,⁵ M. S. Gill,⁵ A. V. Gritsan,⁵ Y. Groysman,⁵ R. G. Jacobsen,⁵ R. W. Kadel,⁵ J. Kadyk,⁵ L. T. Kerth,⁵ Yu. G. Kolomensky,⁵ J. F. Kral,⁵ C. LeClerc,⁵ M. E. Levi,⁵ G. Lynch,⁵ P. J. Oddone,⁵ A. Perazzo,⁵ M. Pripstein,⁵ N. A. Roe,⁵ A. Romosan,⁵ M. T. Ronan,⁵ V. G. Shelkov,⁵ A. V. Telnov,⁵ W. A. Wenzel,⁵ P. G. Bright-Thomas,⁶ T. J. Harrison,⁶ C. M. Hawkes,⁶ D. J. Knowles,⁶ S. W. O'Neale,⁶ R. C. Penny,⁶ A. T. Watson,⁶ N. K. Watson,⁶ T. Deppermann,⁷ K. Goetzen,⁷ H. Koch,⁷ M. Kunze,⁷ B. Lewandowski,⁷ K. Peters,⁷ H. Schmucker,⁷ M. Steinke,⁷ J. C. Andress,⁸ N. R. Barlow,⁸ W. Bhimji,⁸ N. Chevalier,⁸ P. J. Clark,⁸ W. N. Cottingham,⁸ N. Dyce,⁸ B. Foster,⁸ C. Mackay,⁸ D. Wallom,⁸ F. F. Wilson,⁸ K. Abe,⁹ C. Hearty,⁹ T. S. Mattison,⁹ J. A. McKenna,⁹ D. Thiessen,⁹ S. Jolly,¹⁰ A. K. McKemey,¹⁰ V. E. Blinov,¹¹ A. D. Bukin,¹¹ D. A. Bukin,¹¹ A. R. Buzykaev,¹¹ V. B. Golubev,¹¹ V. N. Ivanchenko,¹¹ A. A. Korol,¹¹ E. A. Kravchenko,¹¹ A. P. Onuchin,¹¹ A. A. Salnikov,¹¹ S. I. Serednyakov,¹¹ Yu. I. Skovpen,¹¹ V. I. Telnov,¹¹ A. N. Yushkov,¹¹ D. Best,¹² M. Chao,¹² A. J. Lankford,¹² M. Mandelkern,¹² S. McMahon,¹² D. P. Stoker,¹² K. Arisaka,¹³ C. Buchanan,¹³ S. Chun,¹³ D. B. MacFarlane,¹⁴ S. Prell,¹⁴ Sh. Rahatlou,¹⁴ G. Raven,¹⁴ V. Sharma,¹⁴ C. Campagnari,¹⁵ B. Dahmes,¹⁵ P. A. Hart,¹⁵ N. Kuznetsova,¹⁵ S. L. Levy,¹⁵ O. Long,¹⁵ A. Lu,¹⁵ J. D. Richman,¹⁵ W. Verkerke,¹⁵ M. Witherell,¹⁵ S. Yellin,¹⁵ J. Beringer,¹⁶ D. E. Dorfan,¹⁶ A. M. Eisner,¹⁶ A. A. Grillo,¹⁶ M. Grothe,¹⁶ C. A. Heusch,¹⁶ R. P. Johnson,¹⁶ W. S. Lockman,¹⁶ T. Pulliam,¹⁶ H. Sadrozinski,¹⁶ T. Schalk,¹⁶ R. E. Schmitz,¹⁶ B. A. Schumm,¹⁶ A. Seiden,¹⁶ M. Turri,¹⁶ W. Walkowiak,¹⁶ D. C. Williams,¹⁶ M. G. Wilson,¹⁶ E. Chen,¹⁷ G. P. Dubois-Felsmann,¹⁷ A. Dvoretzki,¹⁷ D. G. Hitlin,¹⁷ S. Metzler,¹⁷ J. Oyang,¹⁷ F. C. Porter,¹⁷ A. Ryd,¹⁷ A. Samuel,¹⁷ M. Weaver,¹⁷ S. Yang,¹⁷ R. Y. Zhu,¹⁷ S. Devmal,¹⁸ T. L. Geld,¹⁸ S. Jayatilake,¹⁸ G. Mancinelli,¹⁸ B. T. Meadows,¹⁸ M. D. Sokoloff,¹⁸ T. Barillari,¹⁹ P. Bloom,¹⁹ M. O. Dima,¹⁹ S. Fahey,¹⁹ W. T. Ford,¹⁹ D. R. Johnson,¹⁹ U. Nauenberg,¹⁹ A. Olivas,¹⁹ P. Rankin,¹⁹ J. Roy,¹⁹ S. Sen,¹⁹ J. G. Smith,¹⁹ W. C. van Hoek,¹⁹ D. L. Wagner,¹⁹ J. Blouw,²⁰ J. L. Harton,²⁰ M. Krishnamurthy,²⁰ A. Soffer,²⁰ W. H. Toki,²⁰ R. J. Wilson,²⁰ J. Zhang,²⁰ R. Aleksan,²¹ A. de Lesquen,²¹ S. Emery,²¹ A. Gaidot,²¹ S. F. Ganzhur,²¹ P.-F. Giraud,²¹ G. Hamel de Monchenault,²¹ W. Kozanecki,²¹ M. Langer,²¹ G. W. London,²¹ B. Mayer,²¹ B. Serfass,²¹ G. Vasseur,²¹ Ch. Yèche,²¹ M. Zito,²¹ T. Brandt,²² J. Brose,²² T. Colberg,²² M. Dickopp,²² R. S. Dubitzky,²² A. Hauke,²² E. Maly,²² R. Müller-Pfefferkorn,²² S. Otto,²² K. R. Schubert,²² R. Schwierz,²² B. Spaan,²² L. Wilden,²² D. Bernard,²³ G. R. Bonneaud,²³ F. Brochard,²³ J. Cohen-Tanugi,²³ S. Ferrag,²³ E. Roussot,²³ S. T'Jampens,²³ Ch. Thiebaux,²³ G. Vasileiadis,²³ M. Verderi,²³ A. Anjomshoa,²⁴ R. Bernet,²⁴ A. Khan,²⁴ D. Lavin,²⁴ F. Muheim,²⁴ S. Playfer,²⁴ J. E. Swain,²⁴ J. Tinslay,²⁴ M. Falbo,²⁵ C. Borean,²⁶ C. Bozzi,²⁶ S. Dittongo,²⁶ L. Piemontese,²⁶ E. Treadwell,²⁷ F. Anulli,^{28,*} R. Baldini-Ferrolì,²⁸ A. Calcaterra,²⁸ R. de Sangro,²⁸ D. Falciari,²⁸ G. Finocchiaro,²⁸ P. Patteri,²⁸ I. M. Peruzzi,^{28,*} M. Piccolo,²⁸ Y. Xie,²⁸ A. Zallo,²⁸ S. Bagnasco,²⁹ A. Buzzo,²⁹ R. Contri,²⁹ G. Crosetti,²⁹ M. Lo Vetere,²⁹ M. Macri,²⁹ M. R. Monge,²⁹ S. Passaggio,²⁹ F. C. Pastore,²⁹ C. Patrignani,²⁹ M. G. Pia,²⁹ E. Robutti,²⁹ A. Santroni,²⁹ S. Tosi,²⁹ M. Morii,³⁰ R. Bartoldus,³¹ R. Hamilton,³¹ U. Mallik,³¹ J. Cochran,³² H. B. Crawley,³² P.-A. Fischer,³² J. Lamsa,³² W. T. Meyer,³² E. I. Rosenberg,³² G. Grosdidier,³³ C. Hast,³³ A. Höcker,³³ H. M. Lacker,³³ S. Laplace,³³ V. Lepeltier,³³ A. M. Lutz,³³ S. Plaszczynski,³³ M. H. Schune,³³ S. Trincaz-Duviois,³³ G. Wormser,³³ R. M. Bionta,³⁴ V. Brigljević,³⁴ D. J. Lange,³⁴ M. Mugge,³⁴ K. van Bibber,³⁴ D. M. Wright,³⁴ M. Carroll,³⁵ J. R. Fry,³⁵ E. Gabathuler,³⁵ R. Gamet,³⁵ M. George,³⁵ M. Kay,³⁵ D. J. Payne,³⁵ R. J. Sloane,³⁵ C. Touramanis,³⁵ M. L. Aspinwall,³⁶ D. A. Bowerman,³⁶ P. D. Dauncey,³⁶ U. Egede,³⁶ I. Eschrich,³⁶ N. J. W. Gunawardane,³⁶ J. A. Nash,³⁶ P. Sanders,³⁶ D. Smith,³⁶ D. E. Azzopardi,³⁷ J. J. Back,³⁷ P. Dixon,³⁷ P. F. Harrison,³⁷ R. J. L. Potter,³⁷ H. W. Shorthouse,³⁷ P. Strother,³⁷ P. B. Vidal,³⁷ M. I. Williams,³⁷ G. Cowan,³⁸ S. George,³⁸ M. G. Green,³⁸ A. Kurup,³⁸ C. E. Marker,³⁸ P. McGrath,³⁸ T. R. McMahon,³⁸ S. Ricciardi,³⁸ F. Salvatore,³⁸ I. Scott,³⁸ G. Vaitsas,³⁸ D. Brown,³⁹ C. L. Davis,³⁹ J. Allison,⁴⁰ R. J. Barlow,⁴⁰ J. T. Boyd,⁴⁰ A. C. Forti,⁴⁰ J. Fullwood,⁴⁰ F. Jackson,⁴⁰ G. D. Lafferty,⁴⁰ N. Savvas,⁴⁰ E. T. Simopoulos,⁴⁰ J. H. Weatherall,⁴⁰ A. Farbin,⁴¹ A. Jawahery,⁴¹ V. Lillard,⁴¹ J. Olsen,⁴¹ D. A. Roberts,⁴¹ J. R. Schieck,⁴¹ G. Blaylock,⁴² C. Dallapiccola,⁴² K. T. Flood,⁴² S. S. Hertzbach,⁴² R. Kofler,⁴² V. G. Koptchev,⁴² T. B. Moore,⁴² H. Staengle,⁴² S. Willocq,⁴² B. Brau,⁴³ R. Cowan,⁴³ G. Sciolla,⁴³ F. Taylor,⁴³ R. K. Yamamoto,⁴³ M. Milek,⁴⁴ P. M. Patel,⁴⁴ F. Palombo,⁴⁵ J. M. Bauer,⁴⁶ L. Cremaldi,⁴⁶ V. Eschenburg,⁴⁶ R. Kroeger,⁴⁶ J. Reidy,⁴⁶ D. A. Sanders,⁴⁶ D. J. Summers,⁴⁶ J. P. Martin,⁴⁷ J. Y. Nief,⁴⁷ R. Seitz,⁴⁷ P. Taras,⁴⁷ V. Zacek,⁴⁷ H. Nicholson,⁴⁸ C. S. Sutton,⁴⁸ C. Cartaro,⁴⁹ N. Cavallo,^{49,†} G. De Nardo,⁴⁹ F. Fabozzi,⁴⁹ C. Gatto,⁴⁹ L. Lista,⁴⁹ P. Paolucci,⁴⁹ D. Piccolo,⁴⁹ C. Sciacca,⁴⁹ J. M. LoSecco,⁵⁰ J. R. G. Alsmiller,⁵¹ T. A. Gabriel,⁵¹ T. Handler,⁵¹ J. Brau,⁵² R. Frey,⁵² M. Iwasaki,⁵² N. B. Sinev,⁵² D. Strom,⁵² F. Colecchia,⁵³ F. Dal Corso,⁵³ A. Dorigo,⁵³ F. Galeazzi,⁵³ M. Margoni,⁵³ G. Michelon,⁵³ M. Morandin,⁵³ M. Posocco,⁵³ M. Rotondo,⁵³ F. Simonetto,⁵³ R. Stroili,⁵³ E. Torassa,⁵³ C. Voci,⁵³ M. Benayoun,⁵⁴ H. Briand,⁵⁴ J. Chauveau,⁵⁴ P. David,⁵⁴ Ch. de la Vaissière,⁵⁴ L. Del Buono,⁵⁴ O. Hamon,⁵⁴ F. Le Diberder,⁵⁴ Ph. Leruste,⁵⁴ J. Ocariz,⁵⁴ L. Roos,⁵⁴ J. Stark,⁵⁴ S. Versillé,⁵⁴ P. F. Manfredi,⁵⁵ V. Re,⁵⁵ V. Speziali,⁵⁵ E. D. Frank,⁵⁶ L. Gladney,⁵⁶ Q. H. Guo,⁵⁶ J. Panetta,⁵⁶ C. Angelini,⁵⁷ G. Batignani,⁵⁷ S. Bettarini,⁵⁷ M. Bondioli,⁵⁷ M. Carpinelli,⁵⁷ F. Forti,⁵⁷ M. A. Giorgi,⁵⁷ A. Lusiani,⁵⁷ F. Martinez-Vidal,⁵⁷ M. Morganti,⁵⁷ N. Neri,⁵⁷ E. Paoloni,⁵⁷ M. Rama,⁵⁷ G. Rizzo,⁵⁷ F. Sandrelli,⁵⁷ G. Simi,⁵⁷ G. Triggiani,⁵⁷ J. Walsh,⁵⁷ M. Haire,⁵⁸ D. Judd,⁵⁸ K. Paick,⁵⁸ L. Turnbull,⁵⁸ D. E. Wagoner,⁵⁹ J. Albert,⁵⁹ P. Elmer,⁵⁹ C. Lu,⁵⁹ K. T. McDonald,⁵⁹ V. Miftakov,⁵⁹ S. F. Schaffner,⁵⁹ A. J. S. Smith,⁵⁹ A. Tumanov,⁵⁹ E. W. Varnes,⁵⁹ G. Cavoto,⁶⁰ D. del Re,⁶⁰ R. Faccini,⁶⁰ F. Ferrarotto,⁶⁰ F. Ferroni,⁶⁰

E. Lamanna,⁶⁰ E. Leonardi,⁶⁰ M. A. Mazzoni,⁶⁰ S. Morganti,⁶⁰ G. Piredda,⁶⁰ F. Safai Tehrani,⁶⁰ M. Serra,⁶⁰ C. Voena,⁶⁰ S. Christ,⁶¹ R. Waldi,⁶¹ T. Adye,⁶² N. De Groot,^{8,62} B. Franek,⁶² N. I. Geddes,⁶² G. P. Gopal,⁶² S. M. Xella,⁶² N. Coptý,⁶³ M. V. Purohit,⁶³ H. Singh,⁶³ F. X. Yumiceva,⁶³ I. Adam,⁶⁴ P. L. Anthony,⁶⁴ D. Aston,⁶⁴ K. Baird,⁶⁴ N. Berger,⁶⁴ E. Bloom,⁶⁴ A. M. Boyarski,⁶⁴ F. Bulos,⁶⁴ G. Calderini,⁶⁴ M. R. Convery,⁶⁴ D. P. Coupal,⁶⁴ D. H. Coward,⁶⁴ J. Dorfan,⁶⁴ W. Dunwoodie,⁶⁴ R. C. Field,⁶⁴ T. Glanzman,⁶⁴ G. L. Godfrey,⁶⁴ S. J. Gowdy,⁶⁴ P. Grosso,⁶⁴ T. Haas,⁶⁴ T. Himel,⁶⁴ T. Hryn'ova,⁶⁴ M. E. Huffer,⁶⁴ W. R. Innes,⁶⁴ C. P. Jessop,⁶⁴ M. H. Kelsey,⁶⁴ P. Kim,⁶⁴ M. L. Kocian,⁶⁴ U. Langenegger,⁶⁴ D. W. G. S. Leith,⁶⁴ S. Luitz,⁶⁴ V. Luth,⁶⁴ H. L. Lynch,⁶⁴ H. Marsiske,⁶⁴ S. Menke,⁶⁴ R. Messner,⁶⁴ K. C. Moffeit,⁶⁴ R. Mount,⁶⁴ D. R. Muller,⁶⁴ C. P. O'Grady,⁶⁴ V. E. Ozcan,⁶⁴ M. Perl,⁶⁴ S. Petrak,⁶⁴ H. Quinn,⁶⁴ B. N. Ratcliff,⁶⁴ S. H. Robertson,⁶⁴ L. S. Rochester,⁶⁴ A. Roodman,⁶⁴ T. Schietinger,⁶⁴ R. H. Schindler,⁶⁴ J. Schwiening,⁶⁴ V. V. Serbo,⁶⁴ A. Snyder,⁶⁴ A. Soha,⁶⁴ S. M. Spanier,⁶⁴ J. Stelzer,⁶⁴ D. Su,⁶⁴ M. K. Sullivan,⁶⁴ H. A. Tanaka,⁶⁴ J. Va'vra,⁶⁴ S. R. Wagner,⁶⁴ A. J. R. Weinstein,⁶⁴ W. J. Wisniewski,⁶⁴ D. H. Wright,⁶⁴ C. C. Young,⁶⁴ P. R. Burchat,⁶⁵ C. H. Cheng,⁶⁵ D. Kirkby,⁶⁵ T. I. Meyer,⁶⁵ C. Roat,⁶⁵ R. Henderson,⁶⁶ W. Bugg,⁶⁷ H. Cohn,⁶⁷ A. W. Weidemann,⁶⁷ J. M. Izen,⁶⁸ I. Kitayama,⁶⁸ X. C. Lou,⁶⁸ F. Bianchi,⁶⁹ M. Bona,⁶⁹ D. Gamba,⁶⁹ A. Smol,⁶⁹ L. Bosisio,⁷⁰ G. Della Ricca,⁷⁰ L. Lanceri,⁷⁰ P. Poropat,⁷⁰ G. Vuagnin,⁷⁰ R. S. Panvini,⁷¹ C. M. Brown,⁷² P. D. Jackson,⁷² R. Kowalewski,⁷² J. M. Roney,⁷² H. R. Band,⁷³ E. Charles,⁷³ S. Dasu,⁷³ F. Di Lodovico,⁷³ A. M. Eichenbaum,⁷³ H. Hu,⁷³ J. R. Johnson,⁷³ R. Liu,⁷³ Y. Pan,⁷³ R. Prepost,⁷³ I. J. Scott,⁷³ S. J. Sekula,⁷³ J. H. von Wimmersperg-Toeller,⁷³ S. L. Wu,⁷³ Z. Yu,⁷³ T. M. B. Kordich,⁷⁴ and H. Neal⁷⁴

(BABAR Collaboration)

¹Laboratoire de Physique des Particules, F-74941 Annecy-le-Vieux, France²Università di Bari, Dipartimento di Fisica and INFN, I-70126 Bari, Italy³Institute of High Energy Physics, Beijing 100039, China⁴University of Bergen, Institute of Physics, N-5007 Bergen, Norway⁵Lawrence Berkeley National Laboratory and University of California, Berkeley, California 94720⁶University of Birmingham, Birmingham B15 2TT, United Kingdom⁷Ruhr Universität Bochum, Institut für Experimentalphysik 1, D-44780 Bochum, Germany⁸University of Bristol, Bristol BS8 1TL, United Kingdom⁹University of British Columbia, Vancouver, British Columbia, Canada V6T 1Z1¹⁰Brunel University, Uxbridge, Middlesex UB8 3PH, United Kingdom¹¹Budker Institute of Nuclear Physics, Novosibirsk 630090, Russia¹²University of California at Irvine, Irvine, California 92697¹³University of California at Los Angeles, Los Angeles, California 90024¹⁴University of California at San Diego, La Jolla, California 92093¹⁵University of California at Santa Barbara, Santa Barbara, California 93106¹⁶Institute for Particle Physics, University of California at Santa Cruz, Santa Cruz, California 95064¹⁷California Institute of Technology, Pasadena, California 91125¹⁸University of Cincinnati, Cincinnati, Ohio 45221¹⁹University of Colorado, Boulder, Colorado 80309²⁰Colorado State University, Fort Collins, Colorado 80523²¹DAPNIA, Commissariat à l'Energie Atomique/Saclay, F-91191 Gif-sur-Yvette, France²²Technische Universität Dresden, Institut für Kern- und Teilchenphysik, D-01062 Dresden, Germany²³Ecole Polytechnique, F-91128 Palaiseau, France²⁴University of Edinburgh, Edinburgh EH9 3JZ, United Kingdom²⁵Elon University, Elon University, North Carolina 27244-2010²⁶Università di Ferrara, Dipartimento di Fisica and INFN, I-44100 Ferrara, Italy²⁷Florida A&M University, Tallahassee, Florida 32307²⁸Laboratori Nazionali di Frascati dell'INFN, I-00044 Frascati, Italy²⁹Università di Genova, Dipartimento di Fisica and INFN, I-16146 Genova, Italy³⁰Harvard University, Cambridge, Massachusetts 02138³¹University of Iowa, Iowa City, Iowa 52242³²Iowa State University, Ames, Iowa 50011-3160³³Laboratoire de l'Accélérateur Linéaire, F-91898 Orsay, France³⁴Lawrence Livermore National Laboratory, Livermore, California 94550³⁵University of Liverpool, Liverpool L69 3BX, United Kingdom³⁶University of London, Imperial College, London SW7 2BW, United Kingdom³⁷Queen Mary, University of London, E1 4NS, United Kingdom³⁸University of London, Royal Holloway and Bedford New College, Egham, Surrey TW20 0EX, United Kingdom³⁹University of Louisville, Louisville, Kentucky 40292⁴⁰University of Manchester, Manchester M13 9PL, United Kingdom

- ⁴¹University of Maryland, College Park, Maryland 20742
⁴²University of Massachusetts, Amherst, Massachusetts 01003
⁴³Massachusetts Institute of Technology, Laboratory for Nuclear Science, Cambridge, Massachusetts 02139
⁴⁴McGill University, Montréal, Québec, Canada H3A 2T8
⁴⁵Università di Milano, Dipartimento di Fisica and INFN, I-20133 Milano, Italy
⁴⁶University of Mississippi, University, Mississippi 38677
⁴⁷Université de Montréal, Laboratoire René J. A. Lévesque, Montréal, Québec, Canada H3C 3J7
⁴⁸Mount Holyoke College, South Hadley, Massachusetts 01075
⁴⁹Università di Napoli Federico II, Dipartimento di Scienze Fisiche and INFN, I-80126 Napoli, Italy
⁵⁰University of Notre Dame, Notre Dame, Indiana 46556
⁵¹Oak Ridge National Laboratory, Oak Ridge, Tennessee 37831
⁵²University of Oregon, Eugene, Oregon 97403
⁵³Università di Padova, Dipartimento di Fisica and INFN, I-35131 Padova, Italy
⁵⁴Universités Paris VI et VII, Lab de Physique Nucléaire H. E., F-75252 Paris, France
⁵⁵Università di Pavia, Dipartimento di Elettronica and INFN, I-27100 Pavia, Italy
⁵⁶University of Pennsylvania, Philadelphia, Pennsylvania 19104
⁵⁷Università di Pisa, Scuola Normale Superiore and INFN, I-56010 Pisa, Italy
⁵⁸Prairie View A&M University, Prairie View, Texas 77446
⁵⁹Princeton University, Princeton, New Jersey 08544
⁶⁰Università di Roma La Sapienza, Dipartimento di Fisica and INFN, I-00185 Roma, Italy
⁶¹Universität Rostock, D-18051 Rostock, Germany
⁶²Rutherford Appleton Laboratory, Chilton, Didcot, Oxon OX11 0QX, United Kingdom
⁶³University of South Carolina, Columbia, South Carolina 29208
⁶⁴Stanford Linear Accelerator Center, Stanford, California 94309
⁶⁵Stanford University, Stanford, California 94305-4060
⁶⁶TRIUMF, Vancouver, British Columbia, Canada V6T 2A3
⁶⁷University of Tennessee, Knoxville, Tennessee 37996
⁶⁸University of Texas at Dallas, Richardson, Texas 75083
⁶⁹Università di Torino, Dipartimento di Fisica Sperimentale and INFN, I-10125 Torino, Italy
⁷⁰Università di Trieste, Dipartimento di Fisica and INFN, I-34127 Trieste, Italy
⁷¹Vanderbilt University, Nashville, Tennessee 37235
⁷²University of Victoria, Victoria, British Columbia, Canada V8W 3P6
⁷³University of Wisconsin, Madison, Wisconsin 53706
⁷⁴Yale University, New Haven, Connecticut 06511

(Received 25 October 2001; published 13 February 2002)

We present a measurement of the time-dependent CP -violating asymmetries in neutral B decays to the $\pi^+\pi^-$ CP eigenstate, and an updated measurement of the charge asymmetry in $B^0 \rightarrow K^+\pi^-$ decays. In a sample of 33 million $Y(4S) \rightarrow B\bar{B}$ decays collected with the BABAR detector at the SLAC PEP-II asymmetric B factory, we find $65_{-11}^{+12} \pi^+\pi^-$ and $217 \pm 18 K^+\pi^-$ candidates and measure the asymmetry parameters $S_{\pi\pi} = 0.03_{-0.56}^{+0.53} \pm 0.11$, $C_{\pi\pi} = -0.25_{-0.47}^{+0.45} \pm 0.14$, and $A_{K\pi} = -0.07 \pm 0.08 \pm 0.02$, where the first error is statistical and the second is systematic.

DOI: 10.1103/PhysRevD.65.051502

PACS number(s): 13.25.Hw, 11.30.Er, 12.15.Hh

In the Standard Model, all CP -violating effects arise from a single complex phase in the three-generation Cabibbo-Kobayashi-Maskawa (CKM) quark-mixing matrix [1]. One of the central questions in particle physics is whether this mechanism is sufficient to explain the pattern of CP violation observed in nature. Recent measurements of the parameter $\sin 2\beta$ by the BABAR [2] and BELLE [3] Collaborations establish that CP symmetry is violated in the neutral B -meson system. In addition to measuring $\sin 2\beta$ more precisely, one of the primary goals of the B -factory experiments

in the future will be to measure the remaining angles (α and γ) and sides of the Unitarity Triangle in order to further test whether the Standard Model description of CP violation is correct.

The study of B decays to charmless hadronic two-body final states will play an increasingly important role in our understanding of CP violation. In the Standard Model, the time-dependent CP -violating asymmetry in the reaction $B^0 \rightarrow \pi^+\pi^-$ is related to the angle α . In addition, observation of a significant rate asymmetry between $B^0 \rightarrow K^+\pi^-$ and $\bar{B}^0 \rightarrow K^-\pi^+$ decays would be evidence for direct CP violation, and ratios of branching fractions for various $\pi\pi$ and $K\pi$ decay modes are sensitive to the angle γ . Finally, branching fraction measurements provide critical tests of the-

*Also with Università di Perugia, Perugia, Italy.

†Also with Università della Basilicata, Potenza, Italy.

oretical models that are needed to extract reliable information on CP violation from the experimental observables.

The BABAR Collaboration recently reported measurements of branching fractions and charge asymmetries for several charmless two-body B decays using a data set of 23 million $B\bar{B}$ pairs [4]. In this paper, using a data sample of approximately 33 million $B\bar{B}$ pairs, we report a measurement of time-dependent CP -violating asymmetries in neutral B decays to the $\pi^+\pi^-$ CP eigenstate and an updated measurement of the charge asymmetry in $B^0 \rightarrow K^+\pi^-$ decays.

The time-dependent CP -violating asymmetry in the decay $B^0 \rightarrow \pi^+\pi^-$ arises from interference between mixing and decay amplitudes, and interference between the $b \rightarrow uW^-$ (tree) and $b \rightarrow dg$ (penguin) decay amplitudes. A $B^0\bar{B}^0$ pair produced in $Y(4S)$ decay evolves in time in a coherent P -wave state until one of the two mesons decays. We reconstruct a sample of B mesons ($B_{hh'}$) decaying to the $h^+h'^-$ final state, where h and h' refer to π or K , and examine the remaining charged particles in each event to “tag” the flavor of the other B meson (B_{tag}). The decay rate distribution f_+ (f_-) when $h^+h'^- = \pi^+\pi^-$ and $B_{\text{tag}} = B^0$ (\bar{B}^0) is given by [5]

$$f_{\pm}(\Delta t) = \frac{e^{-|\Delta t|/\tau}}{4\tau} [1 \pm S_{\pi\pi} \sin(\Delta m_d \Delta t) \mp C_{\pi\pi} \cos(\Delta m_d \Delta t)], \quad (1)$$

where τ is the B^0 lifetime, Δm_d is the $B^0\bar{B}^0$ mixing frequency, and $\Delta t = t_{hh'} - t_{\text{tag}}$ is the time between the $B_{hh'}$ and B_{tag} decays. The CP -violating parameters $S_{\pi\pi}$ and $C_{\pi\pi}$ are defined as

$$S_{\pi\pi} = \frac{2 \text{Im} \lambda}{1 + |\lambda|^2} \quad \text{and} \quad C_{\pi\pi} = \frac{1 - |\lambda|^2}{1 + |\lambda|^2}. \quad (2)$$

If the decay proceeds purely through the tree process, the complex parameter λ is directly related to CKM matrix elements,

$$\lambda(B \rightarrow \pi^+\pi^-) = \left(\frac{V_{tb}^* V_{td}}{V_{tb} V_{td}^*} \right) \left(\frac{V_{ud}^* V_{ub}}{V_{ud} V_{ub}^*} \right), \quad (3)$$

where we are assuming equal widths ($\Delta\Gamma_B = 0$) for the heavy and light mass eigenstates. Thus, at tree level in the standard model, $|\lambda| = 1$ and $\text{Im} \lambda = \sin 2\alpha$, where $\alpha \equiv \arg[-V_{td} V_{tb}^*/V_{ud} V_{ub}^*]$.

Recent theoretical estimates indicate that the contribution from the gluonic penguin amplitude can be significant [6–8]. The process $b \rightarrow dg$ carries the weak phase $\arg(V_{td}^* V_{tb})$, which can modify both the magnitude and phase of λ . Thus, in general, $|\lambda| \neq 1$ and $\text{Im} \lambda = |\lambda| \sin 2\alpha_{\text{eff}}$, where α_{eff} depends on the magnitudes and strong phases of the tree and penguin amplitudes. Several approaches have been proposed to obtain information on α in the presence of penguins [6,9].

In this analysis, we extract signal and background yields for $\pi^+\pi^-$, $K^+\pi^-$, and K^+K^- decays [10], and the amplitudes of the $\pi\pi$ sine ($S_{\pi\pi}$) and cosine ($C_{\pi\pi}$) oscillation

terms simultaneously from an unbinned maximum likelihood fit. We parametrize the $K\pi$ component in terms of the total yield and the CP -violating charge asymmetry

$$\mathcal{A}_{K\pi} \equiv \frac{N_{K^-\pi^+} - N_{K^+\pi^-}}{N_{K^-\pi^+} + N_{K^+\pi^-}}. \quad (4)$$

The data sample used in this analysis consists of 33.7 fb⁻¹ collected with the BABAR detector at the SLAC e^+e^- storage ring PEP-II between October 1999 and June 2001. The PEP-II facility operates nominally at the $Y(4S)$ resonance, providing collisions of 9.0 GeV electrons on 3.1 GeV positrons. The data set includes 30.4 fb⁻¹ collected in this configuration (on-resonance) and 3.3 fb⁻¹ collected below the $B\bar{B}$ threshold (off-resonance) that are used for continuum background studies.

A detailed description of the BABAR detector is presented in Ref. [11]. Charged particle (track) momenta are measured in a tracking system consisting of a 5-layer double-sided silicon vertex tracker (SVT) and a 40-layer drift chamber (DCH) filled with a gas mixture of helium and isobutane. The SVT and DCH operate within a 1.5 T superconducting solenoidal magnet. The typical decay vertex resolution for fully reconstructed B decays is approximately 65 μm along the center-of-mass (c.m.) boost direction. Photons are detected in an electromagnetic calorimeter (EMC) consisting of 6580 CsI(Tl) crystals arranged in barrel and forward endcap subdetectors. The flux return for the solenoid is composed of multiple layers of iron and resistive plate chambers for the identification of muons and long-lived neutral hadrons.

Tracks from the $B_{hh'}$ decay are identified as pions or kaons by the Cherenkov angle θ_c measured with a detector of internally reflected Cherenkov light (DIRC). The typical separation between pions and kaons varies from 8σ at 2 GeV/ c to 2.5σ at 4 GeV/ c , where σ is the average resolution on θ_c . Lower momentum kaons used in B flavor tagging are identified with a selection algorithm that combines θ_c (for momenta down to 0.6 GeV/ c) with measurements of ionization energy loss dE/dx in the DCH and SVT. The selection efficiency is approximately 85% for a pion misidentification probability of 2.5%.

Hadronic events are selected based on track multiplicity and event topology. We require at least three tracks in the laboratory polar angle region $0.41 < \theta_{\text{lab}} < 2.54$ satisfying the following requirements: transverse momentum greater than 100 MeV/ c , at least 12 DCH hits, and originating from the interaction point within 10 cm in z and 1.5 cm in $r - \phi$ [12]. Residual two-prong events from the reaction $e^+e^- \rightarrow l^+l^-$ ($l = e, \mu, \tau$) are suppressed by requiring the ratio of Fox-Wolfram moments H_2/H_0 [13] to be less than 0.95 and the sphericity [14] of the event to be greater than 0.01.

Candidate $B_{hh'}$ decays are reconstructed from pairs of oppositely charged tracks forming a good quality vertex, where the $B_{hh'}$ four-vector is calculated assuming the pion mass for both tracks. We require each track to have an associated θ_c measurement with a minimum of six Cherenkov photons above background, where the average is approximately 30 for both pions and kaons. Protons are rejected based on θ_c and electrons are rejected based on dE/dx ,

shower shape in the EMC, and the ratio of shower energy and track momentum. Background from the reaction $e^+e^- \rightarrow q\bar{q}$ ($q=u,d,s,c$) is suppressed by removing jet-like events from the sample: we define the c.m. angle θ_s between the sphericity axes of the B candidate and the remaining tracks and photons in the event, and require $|\cos\theta_s| < 0.8$, which removes 83% of the background. The total efficiency on signal events for all of the above selection is approximately 38%.

We define a beam-energy substituted mass $m_{ES} = \sqrt{E_b^2 - \mathbf{p}_B^2}$. The candidate energy is defined as $E_b = (s/2 + \mathbf{p}_i \cdot \mathbf{p}_B)/E_i$, where \sqrt{s} and E_i are the total energies of the e^+e^- system in the c.m. and laboratory frames, respectively, and \mathbf{p}_i and \mathbf{p}_B are the momentum vectors in the laboratory frame of the e^+e^- system and the $B_{hh'}$ candidate, respectively. Signal events are Gaussian distributed in m_{ES} with a mean near the B mass and a resolution of $2.6 \text{ MeV}/c^2$, dominated by the beam energy spread. The background shape is parametrized by a threshold function [15] with a fixed end point given by the average beam energy.

We define a second kinematic variable ΔE as the difference between the energy of the $B_{hh'}$ candidate in the c.m. frame and $\sqrt{s}/2$. The ΔE distribution is peaked near zero for $\pi^+\pi^-$ decays. For decays with one (two) kaons, the distribution is shifted relative to $\pi\pi$ on average by -45 MeV (-91 MeV), respectively, where the exact separation depends on the laboratory momentum of the kaon(s). The resolution on ΔE for signal decays is approximately 26 MeV . The background is parametrized by a quadratic function.

Candidate $h^+h'^-$ pairs selected in the region $5.2 < m_{ES} < 5.3 \text{ GeV}/c^2$ and $|\Delta E| < 0.15 \text{ GeV}$ are used to extract yields and CP -violating asymmetries with an unbinned maximum likelihood fit. The total number of events in the fit region satisfying all of the above criteria is 9741. A sideband region, defined as $5.20 < m_{ES} < 5.26 \text{ GeV}/c^2$ and $|\Delta E| < 0.42 \text{ GeV}$, is used to extract various background parameters.

The analysis method combines the techniques used to measure charmless two-body branching fractions [4] and $\sin 2\beta$ [2]. The primary issues in this analysis are determination of the B_{tag} flavor, measurement of the distance Δz between the $B_{hh'}$ and B_{tag} decay vertices, discrimination of signal from background, identification of pions and kaons, and extraction of yields and CP asymmetries.

To determine the flavor of the B_{tag} meson we use the same B -tagging algorithm used in the $\sin 2\beta$ and $B^0-\bar{B}^0$ mixing [16] analyses. The algorithm relies on the correlation between the flavor of the b quark and the charge of the remaining tracks in the event after removal of the $B_{hh'}$ candidate. We define five mutually exclusive tagging categories: Lepton, Kaon, NT1, NT2, and Untagged. Lepton tags rely on primary electrons and muons from semileptonic B decays, while Kaon tags exploit the correlation in the process $b \rightarrow c \rightarrow s$ between the net kaon charge and the charge of the b quark. The NT1 (more certain tags) and NT2 (less certain tags) categories are derived from a neural network that is sensitive to charge correlations between the parent B and unidentified leptons and kaons, soft pions, or the charge and momentum of the track with the highest c.m. momentum.

TABLE I. Tagging efficiency ϵ , average dilution $D=1/2$ ($D_{B^0} + D_{\bar{B}^0}$), dilution difference $\Delta D = D_{B^0} - D_{\bar{B}^0}$, and effective tagging efficiency Q for signal events in each tagging category.

Category	ϵ (%)	D (%)	ΔD (%)	Q (%)
Lepton	11.0 ± 0.3	82.3 ± 2.7	-2.1 ± 4.5	7.5 ± 0.5
Kaon	35.8 ± 0.5	64.8 ± 2.0	3.5 ± 3.1	15.0 ± 1.0
NT1	8.0 ± 0.3	55.6 ± 4.2	-12.1 ± 6.7	2.5 ± 0.4
NT2	13.9 ± 0.4	30.2 ± 3.8	9.0 ± 5.7	1.3 ± 0.3
Untagged	31.3 ± 0.5			
Total Q				26.3 ± 1.2

The addition of Untagged events improves the signal yield estimates and provides a larger sample for determining background shape parameters directly in the maximum likelihood fit.

The quality of tagging is expressed in terms of the effective efficiency $Q = \sum_i \epsilon_i D_i^2$, where ϵ_i is the fraction of events tagged in category i and the dilution $D_i = 1 - 2w_i$ is related to the mistag fraction w_i . The statistical errors on $S_{\pi\pi}$ and $C_{\pi\pi}$ are proportional to $1/\sqrt{Q}$. Table I summarizes the tagging performance in a data sample B_{flav} of fully reconstructed neutral B decays into $D^{(*)-}h^+$ ($h^+ = \pi^+, \rho^+, a_1^+$) and $J/\psi K^{*0}$ ($K^{*0} \rightarrow K^+\pi^-$) flavor eigenstates. We use the same tagging efficiencies and dilutions for signal $\pi\pi$, $K\pi$, and KK decays. Separate background tagging efficiencies for each species are obtained from a fit to the $h^+h'^-$ on-resonance sideband data and reported in Table II.

The time difference Δt is obtained from the measured distance between the z position of the $B_{hh'}$ and B_{tag} decay vertices and the known boost of the e^+e^- system. The z position of the B_{tag} vertex is determined with an iterative procedure that removes tracks with a large contribution to the total χ^2 [2,16]. An additional constraint is constructed from the three-momentum and vertex position of the $B_{hh'}$ candidate, and the average e^+e^- interaction point and boost. The typical Δz resolution is $180 \mu\text{m}$. We require $|\Delta t| < 17 \text{ ps}$ and $0.3 < \sigma_{\Delta t} < 3.0 \text{ ps}$, where $\sigma_{\Delta t}$ is the error from the vertex fit. The resolution function for signal candidates is a sum of three Gaussians, identical to the one described in Ref. [2], with parameters determined from a fit to the B_{flav} sample (including events in all five tagging categories). The background resolution function is parametrized as the sum of three Gaussians, with the parameters determined from a fit to the $h^+h'^-$ on-resonance sideband data.

TABLE II. Tagging efficiencies (%) for background events in each species.

Category	$\epsilon(\pi\pi)$	$\epsilon(K\pi)$	$\epsilon(KK)$
Lepton	1.0 ± 0.1	1.0 ± 0.1	1.5 ± 0.2
Kaon	26.0 ± 0.4	33.1 ± 0.6	23.5 ± 0.7
NT1	6.6 ± 0.2	5.4 ± 0.3	6.9 ± 0.4
NT2	17.6 ± 0.4	15.3 ± 0.5	19.7 ± 0.6
Untagged	48.9 ± 0.7	45.2 ± 0.6	48.3 ± 0.8

The data sample used in the fit contains 97% background, mostly due to random combinations of tracks produced in $e^+e^- \rightarrow q\bar{q}$ events. Discrimination of signal from background in the maximum likelihood fit is enhanced by the use of a Fisher discriminant \mathcal{F} [4]. The discriminating variables are constructed from the scalar sum of the c.m. momenta of all tracks and photons (excluding tracks from the $B_{hh'}$ candidate) entering nine two-sided 10-degree concentric cones centered on the thrust axis of the $B_{hh'}$ candidate. The distribution of \mathcal{F} for signal events is parametrized as a single Gaussian, with parameters determined from Monte Carlo simulated decays and validated with $B^- \rightarrow D^0 \pi^-$ decays reconstructed in data. The background shape is parametrized as the sum of two Gaussians, with parameters determined directly in the maximum likelihood fit.

Identification of $h^+h'^-$ tracks as pions or kaons is accomplished with the Cherenkov angle measurement from the DIRC. We construct Gaussian probability density functions (PDFs) from the difference between measured and expected values of θ_c for the pion or kaon hypothesis, normalized by the resolution. The DIRC performance is parametrized using a sample of $D^{*+} \rightarrow D^0 \pi^+$, $D^0 \rightarrow K^- \pi^+$ decays reconstructed in data. Within the statistical precision of the control sample (approximately 10^5 events), we find similar response for positively and negatively charged tracks and use a single parametrization for both.

We use an unbinned extended maximum likelihood fit to extract yields and CP parameters from the $B_{hh'}$ sample. The likelihood for candidate j tagged in category c is obtained by summing the product of event yield n_i , tagging efficiency $\epsilon_{i,c}$, and probability $\mathcal{P}_{i,c}$ over the eight possible signal and background hypotheses i (referring to $\pi\pi$, $K^+ \pi^-$, $K^- \pi^+$, and KK decays),

$$\mathcal{L}_c = \exp\left(-\sum_i n_i \epsilon_{i,c}\right) \prod_j \left[\sum_i n_i \epsilon_{i,c} \mathcal{P}_{i,c}(\vec{x}_j; \vec{\alpha}_i)\right]. \quad (5)$$

For the $K^\mp \pi^\pm$ hypotheses, the yield is parametrized as $n_i = N_{K\pi}(1 \pm \mathcal{A}_{K\pi})/2$, where $N_{K\pi} = N_{K^- \pi^+} + N_{K^+ \pi^-}$. We fix the tagging efficiencies ϵ_i to the values in Tables I and II. The probabilities $\mathcal{P}_{i,c}$ are evaluated as the product of PDFs for each of the independent variables $\vec{x}_j = \{m_{ES}, \Delta E, \mathcal{F}, \theta_c^+, \theta_c^-, \Delta t\}$, where θ_c^+ and θ_c^- are the Cherenkov angles for the positively and negatively charged tracks. The total likelihood \mathcal{L} is the product of likelihoods for each tagging category and the free parameters are determined by minimizing the quantity $-2 \ln \mathcal{L}$.

The Δt PDF for signal $\pi^+ \pi^-$ decays is given by Eq. (1), modified to include the dilution and dilution difference for each tagging category, and convolved with the signal resolution function. The Δt PDF for signal $K\pi$ events takes into account $B^0 - \bar{B}^0$ mixing, depending on the charge of the kaon and the flavor of B_{tag} . We parametrize $B^0 \rightarrow K^+ K^-$ decays as an exponential convolved with the resolution function.

There are 18 free parameters in the fit. In addition to the CP -violating parameters $S_{\pi\pi}$, $C_{\pi\pi}$, and $\mathcal{A}_{K\pi}$, the fit determines signal and background yields (six parameters), the background $K\pi$ charge asymmetry, and eight parameters de-

TABLE III. Central values and 90% C.L. intervals for $S_{\pi\pi}$, $C_{\pi\pi}$, and $\mathcal{A}_{K\pi}$ from the maximum likelihood fit.

Parameter	Central value	90% C.L. interval
$S_{\pi\pi}$	$0.03^{+0.53}_{-0.56} \pm 0.11$	$[-0.89, +0.85]$
$C_{\pi\pi}$	$-0.25^{+0.45}_{-0.47} \pm 0.14$	$[-1.0, +0.47]$
$\mathcal{A}_{K\pi}$	$-0.07 \pm 0.08 \pm 0.02$	$[-0.21, +0.07]$

scribing the background shapes in m_{ES} , ΔE , and \mathcal{F} . We fix τ and Δm_d to the world-average values [17].

In a sample of 33 million $B\bar{B}$ pairs, we find $65^{+12}_{-11} \pi\pi$, $217 \pm 18 K\pi$, and $4.3^{+6.3}_{-4.3} KK$ events. These yields are consistent with the branching fractions reported in Ref. [4], as well as measurements from other experiments [18,19]. The results for CP -violating asymmetries are summarized in Table III. Statistical errors correspond to unit change in $\chi^2 \equiv -2 \ln \mathcal{L}$. For each parameter, we also calculate the 90% confidence level (C.L.) interval corresponding to a change in χ^2 of 2.69, and taking into account the systematic error. The correlation between $S_{\pi\pi}$ and $C_{\pi\pi}$ is -21% , while $\mathcal{A}_{K\pi}$ is uncorrelated with either $S_{\pi\pi}$ or $C_{\pi\pi}$.

Figure 1 shows distributions of m_{ES} and ΔE for events enhanced in signal decays based on likelihood ratios. We define $\mathcal{R}_{\text{sig}} = \sum_s n_s \mathcal{P}_s / \sum_i n_i \mathcal{P}_i$ and $\mathcal{R}_k = n_k \mathcal{P}_k / \sum_s n_s \mathcal{P}_s$, where \sum_s (\sum_i) indicates a sum over signal (all) hypotheses, and \mathcal{P}_k indicates the probability for signal hypothesis k . The probabilities include the PDFs for θ_c , \mathcal{F} , and m_{ES} (ΔE) when plotting ΔE (m_{ES}). The selection is defined by optimizing the signal significance with respect to \mathcal{R}_{sig} and \mathcal{R}_k . The solid curve in each plot represents the fit projection after correcting for the efficiency of the additional selection (approximately 55% for $\pi\pi$ and 85% for $K\pi$).

Figure 2 shows the Δt distributions and the asymmetry




FIG. 1. Distributions of m_{ES} and ΔE (unshaded histograms) for events enhanced in signal (a),(b) $\pi\pi$ and (c),(d) $K\pi$ decays based on the likelihood ratio selection described in the text. Solid curves represent projections of the maximum likelihood fit result after accounting for the efficiency of the additional selection, while dashed curves represent $q\bar{q}$ and $\pi\pi \leftrightarrow K\pi$ cross-feed background. Shaded histograms show the subset of events that are tagged.




FIG. 2. Distribution of Δt for events enhanced in signal $\pi\pi$ decays based on the likelihood ratio selection described in the text. (a) and (b) show events (points with errors) with $B_{\text{tag}}=B^0$ or \bar{B}^0 . Solid curves represent projections of the maximum likelihood fit, dashed curves represent the sum of $q\bar{q}$ and $K\pi$ background events, and the shaded region represents the contribution from signal $\pi\pi$ events. (c) shows $\mathcal{A}_{\pi\pi}(\Delta t)$ for data (points with errors), as well as fit projections for signal and background events (solid curve), and signal events only (dashed curve).

$\mathcal{A}_{\pi\pi}(\Delta t) = [N_{B^0}(\Delta t) - N_{\bar{B}^0}(\Delta t)] / [N_{B^0}(\Delta t) + N_{\bar{B}^0}(\Delta t)]$ for tagged events enhanced in signal $\pi\pi$ decays. The selection procedure is the same as Fig. 1, with the likelihoods defined including the PDFs for θ_c , \mathcal{F} , m_{ES} , and ΔE . Approximately 24 $\pi\pi$, 22 $q\bar{q}$, and 5 $K\pi$ events satisfy the selection.

Systematic uncertainties on $S_{\pi\pi}$, $C_{\pi\pi}$, and $\mathcal{A}_{K\pi}$ arise primarily from imperfect knowledge of the PDF shapes and uncertainties on tagging efficiencies, dilutions, τ , and Δm_d . The total systematic error is calculated as the sum in quadrature of the individual uncertainties. The error on $\mathcal{A}_{K\pi}$ is dominated by uncertainty in the mean of the ΔE PDF (0.01) and possible charge bias in track and θ_c reconstruction (0.01)

[20]. Errors on $S_{\pi\pi}$ and $C_{\pi\pi}$ are dominated by the parametrization of Δt resolution for signal and background (≈ 0.07 for $S_{\pi\pi}$, ≈ 0.03 for $C_{\pi\pi}$), tagging (0.05), and, for $C_{\pi\pi}$ only, the mean of the ΔE PDF (0.1).

Extensive studies were performed to validate the fit technique. A large ensemble of Monte Carlo pseudo-experiments was generated from the nominal PDFs with the statistics observed in the full data set. Parameter errors and the maximum value of the likelihood obtained in the data fit are all consistent with expectations based on these pseudo-experiments, and all free parameters are unbiased. We have checked that consistent results are obtained when separating events by B_{tag} flavor. As a validation of the Δt parametrization in data, we fit the full data set to simultaneously extract yields, background parameters, τ , Δm_d , $S_{\pi\pi}$, and $C_{\pi\pi}$. We find $\tau = (1.52 \pm 0.12)$ ps and $\Delta m_d = (0.54 \pm 0.09) \hbar \text{ ps}^{-1}$, and all other parameters are consistent with the nominal fit.

In summary, we have presented a measurement of time-dependent CP -violating asymmetries in $B^0 \rightarrow \pi^+ \pi^-$ decays and an updated measurement of the charge asymmetry $\mathcal{A}_{K\pi}$. The latter is consistent with our previous result reported in Ref. [4], as well as results from other experiments [21,22]. We observe no evidence for direct CP violation in the $K\pi$ mode and determine a 90% C.L. interval excluding a significant part of the allowed region. Although the current measurements of $S_{\pi\pi}$ and $C_{\pi\pi}$ do not significantly constrain the Unitarity Triangle, with the addition of more data and further improvements in detector performance and analysis techniques, future results will yield important information about CP violation in the B -meson system.

We are grateful for the excellent luminosity and machine conditions provided by our PEP-II colleagues. The collaborating institutions wish to thank SLAC for its support and kind hospitality. This work is supported by DOE and NSF (USA), NSERC (Canada), IHEP (China), CEA and CNRS-IN2P3 (France), BMBF (Germany), INFN (Italy), NFR (Norway), MIST (Russia), and PPARC (United Kingdom). Individuals have received support from the Swiss NSF, A. P. Sloan Foundation, Research Corporation, and Alexander von Humboldt Foundation.

[1] N. Cabibbo, Phys. Rev. Lett. **10**, 531 (1963); M. Kobayashi and T. Maskawa, Prog. Theor. Phys. **49**, 652 (1973).
 [2] BABAR Collaboration, B. Aubert *et al.*, Phys. Rev. Lett. **87**, 091801 (2001).
 [3] BELLE Collaboration, K. Abe *et al.*, Phys. Rev. Lett. **87**, 091802 (2001).
 [4] BABAR Collaboration, B. Aubert *et al.*, Phys. Rev. Lett. **87**, 151802 (2001).
 [5] For a general review of the formalism, see Y. Nir and H. Quinn, Annu. Rev. Nucl. Part. Sci. **42**, 211 (1992).
 [6] M. Beneke, G. Buchalla, M. Neubert, and C. T. Sachrajda, Nucl. Phys. **B606**, 245 (2001).
 [7] Y. Y. Keum, H.-n. Li, and A. I. Sanda, Phys. Rev. D **63**, 054008 (2001).

[8] M. Ciuchini *et al.*, Phys. Lett. B **515**, 33 (2001).
 [9] M. Gronau and D. London, Phys. Rev. Lett. **65**, 3381 (1990); Y. Grossman and H. R. Quinn, Phys. Rev. D **58**, 017504 (1998); J. Charles, *ibid.* **59**, 054007 (1999); M. Gronau, D. London, N. Sinha, and R. Sinha, Phys. Lett. B **514**, 315 (2001).
 [10] When only one charge mode is given, conjugate decay modes are implied.
 [11] BABAR Collaboration, B. Aubert *et al.*, BABAR-PUB-01/08, hep-ex/0105044.
 [12] The unit vector \hat{z} is aligned along the detector axis in the electron beam direction.
 [13] G. C. Fox and S. Wolfram, Phys. Rev. Lett. **41**, 1581 (1978).
 [14] S. L. Wu, Phys. Rep. **107**, 59 (1984).

- [15] ARGUS Collaboration, H. Albrecht *et al.*, *Z. Phys. C* **48**, 543 (1990).
- [16] BABAR Collaboration, B. Aubert *et al.*, BABAR-PUB-01/02, hep-ex/0112044.
- [17] Particle Data Group, D. E. Groom *et al.*, *Eur. Phys. J. C* **15**, 1 (2000).
- [18] CLEO Collaboration, D. Cronin-Hennessey *et al.*, *Phys. Rev. Lett.* **85**, 515 (2000).
- [19] BELLE Collaboration, K. Abe *et al.*, *Phys. Rev. Lett.* **87**, 101801 (2001).
- [20] BABAR Collaboration, B. Aubert *et al.*, *Phys. Rev. D* (to be published), hep-ex/0111087.
- [21] CLEO Collaboration, S. Chen *et al.*, *Phys. Rev. Lett.* **85**, 525 (2000).
- [22] BELLE Collaboration, K. Abe *et al.*, *Phys. Rev. D* **64**, 071101(R) (2001).