Lawrence Berkeley National Laboratory

Recent Work

Title

Rapid one-pot synthesis of pyrrole-appended isocorroles.

Permalink

<https://escholarship.org/uc/item/5sb7p3r5>

Journal

Organic & biomolecular chemistry, 17(12)

ISSN

1477-0520

Authors

Larsen, Simon
McCormick, Laura J
Ghosh, Abhik

Publication Date

2019-03-01

DOI

10.1039/c9ob00168a

Peer reviewed

Rapid one-pot synthesis of pyrrole-appended isocorroles

Simon Larsen, ^a Laura J. M^cCormick ^b and Abhik Ghosh ^{*a}

Free-base *meso*-triarylcorroles have been found to undergo oxidative coupling with an excess of pyrrole in dichloromethane in the presence of 2,3-dichloro-5,6-dicyano-1,4-benzoquinone (DDQ) affording 5/10-pyrrole-appended isocorroles in reasonable yields (35–60%) and in a matter of seconds. The free-base isocorrole ligands could all be complexed to copper with Cu(OAc)₂·H₂O in chloroform/methanol in 55–80% yields. Single-crystal X-ray structures of two of the new compounds (H₂[5-pyr-TpOMePiC] and Cu[10-pyr-TpOMePiC]) revealed planar macrocycles with rms atomic displacements of only 0.02 and 0.06 Å relative to their respective best-fit C₁₉N₄ planes. Both free-base and Cu(II)-complexed isocorroles exhibit richly featured UV-vis-NIR spectra with red/NIR absorption maxima at ~650 nm and ~725 nm for the free-bases and ~800–850 nm for the copper complexes, suggesting potential applications in photodynamic therapy. Cyclic voltammetric analyses of five of the Cu complexes revealed fully reversible redox cycles with multiple oxidation and reduction features.

Introduction

Over the last quarter-century, during which the chemistry of corroles has grown by leaps and bounds,^{1–3} isocorroles have to some extent languished in the shadow of their better-known isomers. Nevertheless, a string of papers over the last decade have highlighted their potential importance.^{4–7} They are relatively stable and practical to work with. They are also increasingly accessible, albeit in rather variable yields, *via* simple derivatization of corroles.^{4,8,9} Moreover, by combining a porphyrin-like 2– charge with a corrole-like, sterically constrained N₄ core, isocorroles and heteroisocorroles^{10,11} provide novel platforms for coordination chemistry.^{12,13} Finally, as homoconjugated and potentially homoaromatic¹⁴ compounds, they exhibit surprisingly porphyrin-like electronic absorption spectra that extend well into the near-infrared (NIR), potentially heralding applications in photodynamic therapy.^{15,16} Here we report that *meso*-pyrrole-appended isocorroles may be obtained in reasonable yields *via* brief, room-temperature oxidative coupling of a free-base *meso*-triarylcorrole and pyrrole (Scheme 1).^{17–19} Both 5- and 10-isocorroles were obtained, with the former predominating. Unambiguous proof of struc-

ture came from two single-crystal X-ray diffraction analyses, one for a free-base 5-isocorrole and the other for a Cu(II) 10-isocorrole.

Results and discussion

The free-base compounds described here were first observed as products of pyrrole-aldehyde condensations,^{1,20} particularly reaction conditions with a large excess of pyrrole. Mass spectrometric analyses indicated pentapyrrolic products with a

Scheme 1 Synthesis of pyrrole-appended *meso*-tris(*para*-X-phenyl) isocorroles, abbreviated hereafter as H₂[5/10-pyr-TpXPiC], where X = {CF₃, H, Me, OMe}.

^aDepartment of Chemistry, UiT – The Arctic University of Norway, 9037 Tromsø, Norway. E-mail: abhik.ghosh@uit.no

^bAdvanced Light Source, Lawrence Berkeley National Laboratory, Berkeley, CA 94720-8229, USA

molecular formula equivalent to ‘pyrrole + corrole-2H’, apparently consistent with an isosmaragdyrin. One of these products fortunately proved amenable to single-crystal X-ray structure determination, clearly indicating a 5-(2-pyrrolyl)isocorrole (as opposed to an isosmaragdyrin). With the nature of the products established, we devised an alternative, more convenient route based on the oxidative coupling of a corrole and pyrrole. According to the final protocol, a *meso*-tris(*para*-X-phenyl) corrole, H₃[TpXPC], where X ∈ {CF₃, H, Me, OMe}, and pyrrole underwent immediate coupling in dichloromethane in the presence of 2,3-dichloro-5,6-dicyano-1,4-benzoquinone (DDQ). Both the 5- and 10-(2-pyrrolyl)isocorrole isomers were obtained, denoted hereafter as H₂[5/10-pyr-TpXPiC], with the 5-isomer accounting for ≥90% of the combined yield, which ranged from about 35–40% for X = CF₃ and H to about 60% for X = Me and OMe.²¹ The free ligands were also complexed to Cu(II) using Cu(OAc)₂·H₂O (Ac = acetyl) in 4 : 1 v/v mixture of chloroform/methanol as solvent. The 5- and 10-regioisomers of both the free bases and Cu(II) complexes could be separated *via* preparative thin layer chromatography.

The ¹H NMR spectra of the eight free-base isocorroles could each be fully assigned, as depicted in Fig. 1 for H₂[5-pyr-TpOMePiC]. The chemical shifts of all the β- and *meso*-aryl protons, including the *meso*-pyrrolyl CH protons, were found to range from about 5.9 to 7.5 ppm, while the *meso*-pyrrolyl NH proton was found at ~8.1 ppm. Using the latter peak as a starting point, COSY analysis led to the assignment of the other *meso*-pyrrolyl protons. NOESY analysis then identified the nearby *meso*-aryl and β protons, with the remainder of the peaks identified *via* a combination of COSY and NOESY analysis. The two inner NH protons were located at 14.83 ± 0.3 ppm, suggesting a net global paratropic current. Such a proposition is in line with recent DFT calculations of magnetically induced current densities indicating that, while unsubstituted isocorrole is homoaromatic,^{15,16} substituents at the satu-

Fig. 2 Thermal ellipsoid plot for H₂[5-pyr-TpOMePiC], including key hydrogen bond distances.

Fig. 3 Thermal ellipsoid plot for Cu[10-pyr-TpOMePiC]. Selected distances (Å): Cu1–N1 1.915(3), Cu1–N2 1.925(3), Cu1–N3 1.925(3), and Cu1–N4 1.921(3).

Fig. 1 The ¹H NMR spectrum of H₂[5-pyr-TpOMePiC].

Table 1 Crystallographic data for H₂[5-pyr-TpOMePiC] and Cu[10-pyr-TpOMePiC]

Sample	H ₂ [5-pyr-TpOMePiC]·dichloromethane	Cu[10-pyr-TpOMePiC]·chloroform
Chemical formula	C ₄₅ H ₃₇ Cl ₂ N ₅ O ₃	C ₄₅ H ₃₄ Cl ₃ CuN ₅ O ₃
Formula mass	766.69	862.66
Crystal system	Triclinic	Triclinic
Crystal size (mm ³)	0.230 × 0.030 × 0.010	0.150 × 0.080 × 0.070
Space group	<i>P</i> $\bar{1}$	<i>P</i> $\bar{1}$
λ (Å)	0.7749	0.7749
<i>a</i> (Å)	9.7321(6)	11.1767(16)
<i>b</i> (Å)	12.1142(7)	11.6310(16)
<i>c</i> (Å)	16.7703(10)	15.297(2)
α (°)	85.618(2)	104.788(3)
β (°)	85.655(2)	91.215(3)
γ (°)	70.199(2)	96.469(3)
<i>Z</i>	2	2
<i>V</i> (Å ³)	1852.20(19)	1907.9(5)
Temperature (K)	100(2)	100(2)
Density (g cm ⁻³)	1.375	1.502
Measured reflections	24 592	94 061
Unique reflections	4811	10 084
Parameters	554	570
Restraints	46	56
<i>R</i> _{int}	0.0583	0.0568
θ range (°)	1.330 to 24.614	1.503 to 31.849
<i>R</i> ₁ , <i>wR</i> ₂ all data	0.0970, 0.2092	0.0842, 0.2105
<i>S</i> (GoF) all data	1.133	1.026
Max/min res. dens. (e Å ⁻³)	0.474/−0.538	1.383/−1.674

Fig. 4 UV-vis spectra of H₂[5-pyr-TpXPiC], where X ∈ {CF₃, H, Me, OMe}. **Fig. 5** UV-vis spectra of H₂[10-pyr-TpXPiC], where X ∈ {CF₃, H, Me, OMe}.

rated *meso* carbon can dramatically affect the global ring current. Substituted isocorroles accordingly may range from homoaromatic to antihomoaromatic.

Single-crystal X-ray structures could be obtained for two of the compounds prepared, H₂[5-pyr-TpOMePiC] (Fig. 2) and Cu[10-pyr-TpOMePiC] (Fig. 3 and Table 1). Like other isocorrole structures reported to date,^{4–9} both compounds exhibit remarkably planar macrocycles with rms atomic displacements of only 0.02 and 0.06 Å relative to their respective best-fit C₁₉N₄ planes. The relative planarity of the Cu complex may be contrasted with the saddled geometries of Cu corroles, in

which the saddling is thought to reflect a noninnocent Cu^{II}-corrole^{•2–} description.^{22–30} Not surprisingly, the observed Cu–N distances (1.915–1.925 Å) are intermediate between those observed for typical Cu triarylcorroles and Cu porphyrins.^{2,31} Aside from these, the other bond distances do not warrant much comment, the skeletal C–C and C–N bond distances being typical of those observed for dipyrins.

Both the free-base and Cu(II)-complexed isocorroles exhibit richly featured UV-vis-NIR spectra (Fig. 4–7 and Table 2). All the compounds exhibit a strong “Soret” band in the

Fig. 6 UV-visible spectra of Cu[5-pyr-TpXPiC], where X $\in$ {CF₃, H, Me, OMe}.

Fig. 7 UV-visible spectra of Cu[10-pyr-TpXPiC], where X $\in$ {CF₃, H, Me, OMe}.

400–500 nm region as well as one or more sharp post-Soret features in the near-UV. More notably, compared with their free-base precursors, the Cu isocorrole derivatives exhibit significantly more redshifted Soret maxima, by a margin of a few tens of nm. The compounds also exhibit strong “Q” bands. For the free base isocorroles, these are broad, double-humped features plateauing within approximately 650–725 nm. The Q bands of the Cu complexes are also double humped, but sharper and redshifted into the near-IR, with the most intense peaks occurring at $\sim$ 855 nm for the 5-regioisomers and $\sim$ 800 nm for the 10-regioisomers. We tentatively suggest that the considerably redshifted spectra of the Cu complexes relative to the free bases might reflect ligand-to-metal charge transfer transitions. That said, the optical properties of isocorrole derivatives clearly deserve more in-depth study, both *via* the synthesis of new complexes and *via* quantum chemical means.¹⁵ The potential for new isocorrole-based sensitizers for photodynamic therapy is indeed considerable.

Fig. 8 Cyclic voltammograms of selected Cu isocorrole complexes in CH₂Cl₂ containing 0.1 M TBAP. Scan rate = 100 mV s^{−1}.

Table 2 UV-vis absorption maxima (nm) and molar absorptivities ($\epsilon \times 10^{-4}$, M^{−1} cm^{−1}) in dichloromethane solution

	<450 nm	450–600 nm	>600 nm
H ₂ [5-pyr-TpOCH ₃ PiC]	345 (2.05), 415 (6.00)	—	660 (0.76), 708 (0.72).
H ₂ [5-pyr-TpMePiC]	345 (2.97), 406 (5.61)	—	661 (0.82), 705 (0.79)
H ₂ [5-pyr-TpPiC]	339 (2.94), 407 (4.72)	—	661 (0.71), 705 (0.69)
H ₂ [5-pyr-TpCF ₃ PiC]	338 (2.71), 407 (4.06)	—	661 (0.73), 706 (0.74)
H ₂ [10-pyr-TpOCH ₃ PiC]	375 (2.61), 434 (4.71)	—	656 (0.69), 698 (0.69)
H ₂ [10-pyr-TpMePiC]	359 (1.86), 432 (3.16)	—	658 (0.44), 699 (0.45)
H ₂ [10-pyr-TpPiC]	351 (1.41), 430 (2.58)	—	656 (0.37), 691 (0.36)
H ₂ [10-pyr-TpCF ₃ PiC]	347 (0.89), 429 (1.94)	—	648 (0.29), 700 (0.26)
Cu[5-pyr-TpOCH ₃ PiC]	368 (2.42)	459 (3.80), 556 (0.69)	786 (0.59), 862 (0.77)
Cu[5-pyr-TpMePiC]	285 (1.60), 361 (3.00)	459 (3.61), 557 (0.65)	793 (0.65), 860 (0.85)
Cu[5-pyr-TpPiC]	296 (1.41), 357 (2.68)	459 (3.13), 557 (0.57)	792 (0.58), 856 (0.80)
Cu[5-pyr-TpCF ₃ PiC]	308 (1.83), 357 (2.67)	465 (3.44), 560 (0.63)	792 (0.67), 858 (0.98)
Cu[10-pyr-TpOCH ₃ PiC]	271 (1.38), 387 (2.76)	470 (3.50), 544 (0.34)	726 (0.34), 800 (0.78)
Cu[10-pyr-TpMePiC]	315 (1.23), 370 (2.81)	468 (3.49), 539 (0.38)	728 (0.33), 800 (0.80)
Cu[10-pyr-TpPiC]	305 (0.87), 361 (1.28)	470 (1.96), 540 (0.21)	729 (0.18), 800 (0.44)
Cu[10-pyr-TpCF ₃ PiC]	302 (1.42), 356 (1.39), 388 (1.31)	474 (2.62), 545 (0.24)	734 (0.27), 806 (0.62)

Table 3 Redox potentials (V vs. SCE) of selected Cu isocorrole derivatives in CH₂Cl₂ containing 0.1 M TBAP. Scan rate = 100 mV s⁻¹

Compound	$E_{1/2ox-c}$	$E_{1/2ox-b}$	E_{ox-a}	E_{red-a}	$E_{1/2red-b}$	$E_{1/2red-c}$
Cu[10-pyr-TpCF ₃ PiC]	1.05	0.79	—	—	-0.75	-1.34
Cu[5-pyr-TpOCH ₃ PiC]	—	—	0.68 ^a	-0.29 ^a	-0.75	-1.23
Cu[5-pyr-TpCH ₃ PiC]	—	—	0.70 ^a	-0.27 ^a	-0.75	-1.22
Cu[5-pyr-TPiC]	—	—	0.74 ^a	-0.26 ^a	-0.74	-1.20
Cu[5-pyr-TpCF ₃ PiC]	—	—	0.79 ^a	-0.13 ^a	-0.63	-1.06

^a Peak potential.

Cyclic voltammetry on the four Cu(II) 5-isocorroles and on Cu[10-pyr-TpCF₃PiC] revealed fully reversible redox cycles with multiple oxidation and reduction features (Fig. 8 and Table 3). Interestingly, the 5-isocorrole series was found to exhibit an isolated oxidation peak 0.74 ± 0.05 V vs. SCE with no nearby, corresponding reduction peak and a reduction peak at -0.21 ± 0.9 V vs. SCE with no nearby, corresponding oxidation peak. It is reasonable to suppose that oxidation leads to a persistent π -cation radical, potentially involving the pendant pyrrole, which is reduced at a much more negative potential relative to the oxidation peak potential. In contrast, the 10-isocorrole complex Cu[10-pyr-TpCF₃PiC] was found to exhibit normal, reversible oxidation and reduction features. Interestingly, the second reduction peak of the 5-isocorrole series (labeled $E_{1/2red-b}$ in Table 3) occurs at the same potential as the first reduction of *meso*-methoxy isocorrole complexes Cu[5/10-MeO-TpCH₃PiC] studied by Pomarico *et al.* and therefore by analogy may be tentatively assigned to a reduction leading to the molecular anion.⁴ Detailed spectroelectrochemical and density functional theory-based computational studies are currently in progress, which should result in full assignment of the electrochemical features observed here.⁴

Conclusion

Interaction of free-base *meso*-triarylcorroles with an excess of pyrrole in the presence of DDQ in dichloromethane at room temperature results in moderate yields (35–60%) of 5/10-(2-pyrrolyl)isocorroles in a matter of seconds. Metal coordination to each isocorrole ligand was demonstrated by complexation with Cu(II). Both the free ligands and their metal complexes were found to exhibit strong absorption in the near-infrared, foreshadowing potential applications as sensitizers in photodynamic therapy.

Experimental section

Materials

All free-base corroles were synthesized *via* the now-standard water-methanol method.³² All reagents, except pyrrole, were purchased from Sigma-Aldrich and used as received. Pyrrole was passed through basic alumina (aluminium oxide 60, active basic activity I, 0.063–0.200 mm particle size, 70–230 mesh, Merck) and stored in the freezer. Silica gel 60 (0.04–0.063 mm

particle size, 230–400 mesh, Merck) was employed for flash chromatography. Silica gel 60 preparative thin-layer chromatographic plates (20 cm × 20 cm × 0.5 mm, Merck) were used for final purification of all compounds.

General instrumental methods

UV-visible spectra were recorded on an HP 8453 spectrophotometer. ¹H NMR spectra were recorded on a 400 MHz Bruker Avance III HD spectrometer equipped with a 5 mm BB/1H SmartProbe and referenced to either residual CH₂Cl₂ at 5.32 ppm or residual CHCl₃ at 7.26 ppm. High-resolution electrospray-ionization (HR-ESI) mass spectra were recorded on an LTQ Orbitrap XL spectrometer, using methanolic solutions and typically in positive ion mode. Elemental analyses were performed by Atlantic Microlab Inc., USA³³

Cyclic voltammetry was carried out at 298 K with an EG&G Model 263A potentiostat equipped with a three-electrode system: a glassy carbon working electrode, a platinum wire counterelectrode, and a saturated calomel reference electrode (SCE). Tetra(*n*-butyl)ammonium perchlorate (**CAUTION!**), recrystallized twice from absolute ethanol and dried in a desiccator for at least 2 weeks, was used as the supporting electrolyte. Anhydrous CH₂Cl₂ (Aldrich) was used as solvent. The reference electrode was separated from the bulk solution by a fritted-glass bridge filled with the solvent/supporting electrolyte mixture. The electrolyte solution was purged with argon for at least 2 min prior to all measurements, which were carried out under an argon blanket. All potentials were referenced to the SCE.

General procedure for preparation of free-base 5/10-(2-pyrrolyl) isocorroles

A 100 mL round-bottom flask equipped with a magnetic stirring bar was charged with a free-base *meso*-triarylcorrole (20–25 mg) and dichloromethane (10 mL). To the stirred, degassed solution under argon were added DDQ (1 eq.) and pyrrole (10 eq.) in succession, whereupon the color of the solution turned olive-green within seconds. The reaction mixture was then passed through a silica plug with dichloromethane to yield the crude product as a mixture of the 5- and 10-isomer. Separation of the two regioisomers was accomplished with preparative thin-layer chromatography employing the following solvents: dichloromethane/*n*-hexane (8 : 1) for X = *p*-OCH₃, dichloromethane/*n*-hexane (2 : 1) for *p*-CH₃, dichloromethane/pentane (1 : 1) for *p*-H, and dichloromethane/

pentane (1 : 2) for *p*-CF₃. In the event the isomers did not separate, the isocorrole band was divided in fractions and prep-TLC repeated on each individual fraction. The process continued until full separation was achieved. Yields and analytical details for each compound are as follows. Because of the low yields of the 10-isomers, elemental analyses were only carried out for the 5-isomers.

5-(2-Pyrrolyl)-5,10,15-tris(4-methoxyphenyl)isocorrole. Yield 14.8 mg (56%). UV-Vis (CH₂Cl₂) $\lambda_{\max}$ (nm) [$\epsilon \times 10^{-4}$ (M⁻¹ cm⁻¹)]: 345 (2.05), 415 (6.00), 660 (0.76), 708 (0.72). ¹H NMR (400 MHz, CDCl₃, δ): 14.86 (s, 1H, NH), 14.80 (s, 1H, NH), 8.12 (s, 1H, 1-pyrrolyl), 7.49 (d, *J* = 8.8 Hz, 2H, 15-*o*-Ph), 7.40 (d, *J* = 8.8 Hz, 2H, 10-*o*-Ph), 7.16 (d, *J* = 8.9 Hz, 2H, 5-*o*-Ph), 7.04 (d, *J* = 4.5 Hz, 1H, β -H), 6.99 (d, *J* = 8.8 Hz, 2H, 15-*m*-Ph), 6.96–6.91 (m, 3H, overlapping 10-*m*-Ph and β -H), 6.79 (d, *J* = 8.8 Hz, 2H, 5-*m*-Ph), 6.67–6.63 (m, 2H, overlapping β -H and 5-pyrrolyl), 6.58 (dd, *J* = 3.6, 2.5 Hz, 1H, β -H), 6.45 (dd, *J* = 4.2, 2.0 Hz, 1H, β -H), 6.35 (d, *J* = 4.5 Hz, 1H, β -H), 6.28 (dd, *J* = 4.3, 2.5 Hz, 1H, β -H), 6.15–6.12 (m, 1H, 4-pyrrolyl), 6.10–6.07 (m, 1H, 3-pyrrolyl), 5.91 (dd, *J* = 3.6, 2.4 Hz, 1H, β -H), 3.89 (s, 3H, 15-*p*-OCH₃), 3.88 (s, 3H, 10-*p*-OCH₃), 3.78 (s, 3H, 5-*p*-OCH₃). MS (ESI) *m/z*: 682.2815; calcd for C₄₄H₃₅N₅O₃H: 682.2813 [M + H⁺]. Elemental analysis: C 77.45, H 5.25, N 10.09; calcd for C₄₄H₃₅N₅O₃: C 77.51, H 5.17, N 10.27.

10-(2-Pyrrolyl)-5,10,15-tris(4-methoxyphenyl)isocorrole. Yield 1.2 mg (4.5%). UV-Vis (CH₂Cl₂) $\lambda_{\max}$ (nm) [$\epsilon \times 10^{-4}$ (M⁻¹ cm⁻¹)]: 375 (2.61), 434 (4.71), 656 (0.69), 698 (0.69). ¹H NMR (400 MHz, CDCl₃, δ): 14.72 (s, 2H, NH), 7.93 (s, 1H, 1-pyrrolyl), 7.56 (d, *J* = 8.7 Hz, 4H, 5,15-*o*-Ph), 7.04–6.97 (m, 6H, overlapping 5,15-*m*-Ph and 10-*o*-Ph), 6.82 (d, *J* = 8.9 Hz, 2H, 10-*m*-Ph), 6.77 (d, *J* = 4.1 Hz, 2H, β -H), 6.73 (d, *J* = 4.1 Hz, 2H, β -H), 6.68–6.65 (m, 1H, 5-pyrrolyl), 6.64 (d, *J* = 4.2 Hz, 2H, β -H), 6.11–6.07 (m, 1H, 4-pyrrolyl), 5.99 (d, *J* = 4.2 Hz, 2H, β -H), 5.85–5.81 (m, 1H, 3-pyrrolyl), 3.90 (s, 6H, 5,15-*p*-OCH₃), 3.80 (s, 3H, 10-*p*-OCH₃). MS (ESI) *m/z*: 682.2814; calcd for C₄₄H₃₅N₅O₃H: 682.2813 [M + H⁺].

5-(2-Pyrrolyl)-5,10,15-tris(4-methylphenyl)isocorrole. Yield 17.1 mg (58%). UV-Vis (CH₂Cl₂) $\lambda_{\max}$ (nm) [$\epsilon \times 10^{-4}$ (M⁻¹ cm⁻¹)]: 345 (2.97), 406 (5.61), 661 (0.82), 705 (0.79); ¹H NMR (400 MHz, CDCl₃, δ): 14.85 (s, 1H, NH), 14.79 (s, 1H, NH), 8.12 (s, 1H, 1-pyrrolyl), 7.44 (d, *J* = 8.0 Hz, 2H, 10-*o*-Ph), 7.35 (d, *J* = 7.9 Hz, 2H, 10-*o*-Ph), 7.28 (d, *J* = 7.0 Hz, 2H, 15-*m*-Ph), 7.22 (d, *J* = 7.2 Hz, 2H, 10-*m*-Ph), 7.14 (d, *J* = 8.2 Hz, 2H, 5-*o*-Ph), 7.09–7.02 (m, 3H, overlapping 5-*m*-Ph and β -H), 6.94 (d, *J* = 4.5 Hz, 1H, β -H), 6.67–6.62 (m, 2H, overlapping 5-pyrrolyl and β -H), 6.60 (dd, *J* = 2.6 Hz, 1H, β -H), 6.43 (dd, *J* = 4.3, 1.5 Hz, 1H, β -H), 6.37 (d, *J* = 4.6 Hz, 1H, β -H), 6.25 (dd, *J* = 4.5, 2.0 Hz, 1H, β -H), 6.15–6.09 (m, 2H, overlapping 3-pyrrolyl and 4-pyrrolyl), 5.92 (dd, *J* = 3.8, 1.7 Hz, 1H, β -H), 2.45 (s, 3H, 15-*p*-CH₃), 2.44 (s, 3H, 10-*p*-CH₃), 2.32 (s, 3H, 5-*p*-CH₃). MS (ESI) *m/z*: 634.2958; calcd for C₄₄H₃₅N₅H: 634.2965 [M + H⁺]. Elemental analysis: C 82.68, H 5.79, N 10.59; calcd for C₄₄H₃₅N₅: C 83.38, H 5.57, N 11.05.

10-(2-Pyrrolyl)-5,10,15-tris(4-methylphenyl)isocorrole. Yield 0.8 mg (2.7%). UV-Vis (CH₂Cl₂) $\lambda_{\max}$ (nm) [$\epsilon \times 10^{-4}$ (M⁻¹ cm⁻¹)]: 359 (1.86), 432 (3.16), 658 (0.44), 699 (0.45); ¹H NMR

(400 MHz, CDCl₃, δ): 14.73 (s, 2H, NH), 7.94 (s, 1H, 1-pyrrolyl), 7.49 (d, *J* = 7.8 Hz, 4H, 5,15-*o*-Ph), 7.28 (d, 4H, 5,15-*m*-Ph), 7.09 (d, *J* = 8.1 Hz, 2H, 10-*m*-Ph), 6.99 (d, *J* = 7.9 Hz, 2H, 10-*o*-Ph), 6.76 (d, *J* = 4.1 Hz, 2H, β -H), 6.72 (d, *J* = 4.1 Hz, 2H, β -H), 6.67–6.64 (m, 1H, 2-pyrrolyl), 6.62 (d, *J* = 4.2 Hz, 2H, β -H), 6.11–6.08 (m, 1H, 3-pyrrolyl), 5.98 (d, *J* = 4.3 Hz, 2H, β -H), 5.87–5.83 (m, 1H, 4-pyrrolyl), 2.46 (s, 6H, 5,15-*p*-CH₃), 2.33 (s, 3H, 10-*p*-CH₃). MS (ESI) *m/z*: 634.2968; calcd for C₄₄H₃₅N₅H: 634.2965 [M + H⁺].

5-(2-Pyrrolyl)-5,10,15-triphenylisocorrole. Yield 11.6 mg (38%). UV-Vis (CH₂Cl₂) $\lambda_{\max}$ (nm) [$\epsilon \times 10^{-4}$ (M⁻¹ cm⁻¹)]: 339 (2.94), 407 (4.72), 661 (0.71), 705 (0.69). ¹H NMR (400 MHz, CD₂Cl₂, δ): 14.85 (s, 1H, NH), 14.81 (s, 1H, NH), 8.21 (s, 1H, 1-pyrrolyl), 7.59–7.55 (m, 2H, Ph), 7.52–7.43 (m, 8H, overlapping 5-*o*-Ph, 10-*o*-Ph and Ph), 7.33–7.23 (m, 5H, overlapping 15-*o*-Ph and Ph), 7.05 (d, *J* = 4.5 Hz, 1H, β -H), 7.00 (d, *J* = 4.5 Hz, 1H, β -H), 6.70–6.62 (m, 3H, overlapping 5-pyrrolyl and β -H), 6.46 (dd, *J* = 4.3, 1.8 Hz, 1H, β -H), 6.38 (d, *J* = 4.5 Hz, 1H, β -H), 6.27 (dd, *J* = 4.3, 2.3 Hz, 1H, β -H), 6.14–6.11 (m, 1H, 4-pyrrolyl), 6.11–6.08 (m, 1H, 3-pyrrolyl), 5.94 (dd, *J* = 2.9 Hz, 1H, β -H). MS (ESI) *m/z*: 592.2495; calcd for C₄₁H₂₉N₅H: 592.2496 [M + H⁺]. Elemental analysis (%): C 82.60, H 5.41, N 11.15; calcd for C₄₁H₂₉N₅: C 83.22, H 4.94, N 11.84.

10-(2-Pyrrolyl)-5,10,15-triphenylisocorrole. Yield 1.2 mg (3.9%). UV-Vis (CH₂Cl₂) $\lambda_{\max}$ (nm) [$\epsilon \times 10^{-4}$ (M⁻¹ cm⁻¹)]: 351 (1.41), 430 (2.58), 656 (0.37), 691 (0.36). ¹H NMR (400 MHz, CDCl₃, δ): 14.73 (s, 2H, NH), 7.94 (s, 1H, 1-pyrrolyl), 7.63–7.58 (m, 4H, 5,15-*o*-Ph), 7.50–7.44 (m, 6H, overlapping 5,15-*m*-Ph and 5,15-*p*-Ph), 7.34–7.28 (m, 3H, overlapping 10-*m*-Ph and 10-*p*-Ph), 7.13–7.08 (m, 2H, 10-*o*-Ph), 6.77–6.73 (m, 4H, overlapping β -H), 6.69–6.65 (m, 1H, 2-pyrrolyl), 6.61 (d, *J* = 4.3 Hz, 2H, β -H), 6.10 (q, *J* = 3.0 Hz, 1H, 3-pyrrolyl), 5.97 (d, *J* = 4.3 Hz, 2H, β -H), 5.87–5.84 (m, 1H, 4-pyrrolyl). MS (ESI) *m/z*: 592.2499; calcd for C₄₁H₂₉N₅H: 592.2496 [M + H⁺].

5-(2-Pyrrolyl)-5,10,15-tris[(4-trifluoromethyl)phenyl]isocorrole. Yield 6.1 mg (30%). UV-Vis (CH₂Cl₂) $\lambda_{\max}$ (nm) [$\epsilon \times 10^{-4}$ (M⁻¹ cm⁻¹)]: 338 (2.71), 407 (4.06), 661 (0.73), 706 (0.74). ¹H NMR (400 MHz, CDCl₃, δ): 14.75 (s, 2H, NH), 8.11 (s, 1H, 1-pyrrolyl), 7.77–7.69 (m, 4H, overlapping 10-*m*-Ph and 15-*m*-Ph), 7.67 (d, *J* = 8.0 Hz, 2H, 15-*o*-Ph), 7.61 (d, *J* = 7.9 Hz, 2H, 10-*o*-Ph), 7.53 (d, *J* = 8.2 Hz, 2H, 5-*m*-Ph), 7.36 (d, *J* = 8.2 Hz, 2H, 5-*o*-Ph), 7.01 (d, 2H, 2 overlapping β -H), 6.70–6.67 (m, 2H, overlapping 5-pyrrolyl and β -H), 6.58 (d, *J* = 4.6 Hz, 1H, β -H), 6.40–6.36 (m, 2H, 2 overlapping β -H), 6.20 (dd, *J* = 4.3, 2.4 Hz, 1H, β -H), 6.18–6.15 (m, 1H, 4-pyrrolyl), 6.15–6.11 (m, 1H, 3-pyrrolyl), 5.95 (dd, *J* = 3.6, 2.5 Hz, 1H, β -H). MS (ESI) *m/z*: 796.2114; calcd for C₄₄H₂₆N₅F₉H: 796.2117 [M + H⁺]. Elemental analysis (%): C 66.62, H 3.42, N 8.69; calcd for C₄₄H₂₆N₅F₉: C 66.42, H 3.29, N 8.80.

10-(2-Pyrrolyl)-5,10,15-tris[(4-trifluoromethyl)phenyl]isocorrole. Yield 0.4 mg (2.0%). UV-Vis (CH₂Cl₂) $\lambda_{\max}$ (nm) [$\epsilon \times 10^{-4}$ (M⁻¹ cm⁻¹)]: 347 (0.89), 429 (1.94), 648 (0.29), 700 (0.26). ¹H NMR (400 MHz, CDCl₃, δ): 14.58 (s, 2H, NH), 7.86 (s, 1H, 1-pyrrolyl), 7.71–7.62 (m, 8H, overlapping 5,15-*o*-Ph and 5,15-*m*-Ph), 7.50 (d, *J* = 8.2 Hz, 2H, 10-*m*-Ph), 7.15 (d, *J* = 8.2 Hz, 2H, 10-*o*-Ph), 6.72 (d, *J* = 4.2 Hz, 2H, β -H), 6.67–6.61 (m, 3H, overlapping

β -H and 2-pyrrolyl), 6.50 (d, $J = 4.3$ Hz, 2H, β -H), 6.06 (q, $J = 3.0$ Hz, 1H, 3-pyrrolyl), 5.90 (d, $J = 4.3$ Hz, 2H, β -H), 5.79 (d, $J = 3.9$ Hz, 1H, 4-pyrrolyl). MS (ESI) m/z : 796.2112; calcd for $C_{44}H_{26}N_5F_9H$: 796.2117 $[M + H]^+$.

General procedure for preparation of copper 5/10-(2-pyrrolyl) triarylisocorroles

To a 100 ml round-bottom flask equipped with a magnetic stirring bar was added free-base isocorrole (15–25 mg, mixture of isomers), $Cu(OAc)_2 \cdot H_2O$ (1.5 eq.), $CHCl_3$ (8 mL) and MeOH (2 mL). The mixture was refluxed for 30 minutes (room temperature for p -CF₃). The solvents were removed under vacuum and the solids passed through silica with dichloromethane. Separation of 5- and 10-isomers required preparative thin-layer chromatography employing DCM for p -OCH₃, DCM/pentane (1 : 1) for p -CH₃, and DCM/pentane (1 : 2) for p -H and p -CF₃. In the event the isomers did not separate, the copper isocorrole band was divided in fractions and prep-TLC repeated on each individual fraction. The process continued until full separation was achieved. Yields and analytical details for each compound are as follows. Because of the low yields of the 10-isomers, elemental analyses were only carried out for the 5-isomers.

Copper 5-(2-pyrrolyl)-5,10,15-tris(4-methoxyphenyl)isocorrole. Yield 9.2 mg (54%). UV-Vis (CH_2Cl_2) λ_{max} (nm) [$\epsilon \times 10^{-4}$ ($M^{-1} cm^{-1}$)]: 368 (2.42), 459 (3.80), 556 (0.69), 786 (0.59), 862 (0.77). MS (ESI) m/z : 742.1869; calcd for $C_{44}H_{33}N_5O_3Cu$: 742.1874 $[M]^+$. Elemental analysis (%): C 71.19, H 4.70, N 9.03; calcd for $C_{44}H_{33}N_5O_3Cu$: C 71.10, H 4.47, N 9.42.

Copper 10-(2-pyrrolyl)-5,10,15-tris(4-methoxyphenyl)isocorrole. Yield 0.9 mg (5.3%). UV-Vis (CH_2Cl_2) λ_{max} (nm) [$\epsilon \times 10^{-4}$ ($M^{-1} cm^{-1}$)]: 271 (1.38), 387 (2.76), 470 (3.50), 544 (0.34), 726 (0.34), 800 (0.78). MS (ESI) m/z : 742.1876; calcd for $C_{44}H_{33}N_5O_3Cu$: 742.1874 $[M]^+$.

Copper 5-(2-pyrrolyl)-5,10,15-tris(4-methylphenyl)isocorrole. Yield 21.7 mg (77%). UV-Vis (CH_2Cl_2) λ_{max} (nm) [$\epsilon \times 10^{-4}$ ($M^{-1} cm^{-1}$)]: 285 (1.60), 361 (3.00), 459 (3.61), 557 (0.65), 793 (0.65), 860 (0.85). MS (ESI) m/z : 694.2032; calcd for $C_{44}H_{33}N_5Cu$: 694.2026 $[M]^+$. Elemental analysis (%): C 75.41, H 5.06, N 9.61; calcd for $C_{44}H_{33}N_5Cu$: C 76.00, H 4.78, N 10.07.

Copper 10-(2-pyrrolyl)-5,10,15-tris(4-methylphenyl)isocorrole. Yield 1.5 mg (5.3%). UV-Vis (CH_2Cl_2) λ_{max} (nm) [$\epsilon \times 10^{-4}$ ($M^{-1} cm^{-1}$)]: 315 (1.23), 370 (2.81), 468 (3.49), 539 (0.38), 728 (0.33), 800 (0.80). MS (ESI) m/z : 694.2031; calcd for $C_{44}H_{33}N_5Cu$: 694.2026 $[M]^+$.

Copper 5-(2-pyrrolyl)-5,10,15-triphenylisocorrole. Yield 14.7 mg (73%). UV-Vis (CH_2Cl_2) λ_{max} (nm) [$\epsilon \times 10^{-4}$ ($M^{-1} cm^{-1}$)]: 296 (1.41), 357 (2.68), 459 (3.13), 557 (0.57), 792 (0.58), 856 (0.80). MS (ESI) m/z : 652.1558; calcd for $C_{41}H_{27}N_5Cu$: 652.1557 $[M]^+$. Elemental analysis (%): C 75.44, H 4.57, N 10.25; calcd for $C_{41}H_{27}N_5Cu$: C 75.38, H 4.17, N 10.72.

Copper 10-(2-pyrrolyl)-5,10,15-triphenylisocorrole. Yield 1 mg (5.0%). UV-Vis (CH_2Cl_2) λ_{max} (nm) [$\epsilon \times 10^{-4}$ ($M^{-1} cm^{-1}$)]: 305 (0.87), 361 (1.28), 470 (1.96), 540 (0.21), 729 (0.18), 800 (0.44); MS (ESI) m/z : 652.1556; calcd for $C_{41}H_{27}N_5Cu$: 652.1557 $[M]^+$.

Copper 5-(2-pyrrolyl)-5,10,15-tris[(4-trifluoromethyl)phenyl]isocorrole. Yield 11.6 mg (48%). UV-Vis (CH_2Cl_2) λ_{max} (nm) [$\epsilon \times 10^{-4}$ ($M^{-1} cm^{-1}$)]: 308 (1.83), 357 (2.67), 465 (3.44), 560 (0.63), 792 (0.67), 858 (0.98). MS (ESI) m/z : 856.1171; calcd for $C_{44}H_{24}N_5F_9Cu$: 856.1179 $[M]^+$. Elemental analysis (%): C 61.17, H 3.06, N 7.70; calcd for $C_{44}H_{24}N_5F_9Cu$: C 61.65, H 2.82, N 8.17.

Copper 10-(2-pyrrolyl)-5,10,15-tris[(4-trifluoromethyl)phenyl]isocorrole. Yield 1.2 mg (5.0%). UV-Vis (CH_2Cl_2) λ_{max} (nm) [$\epsilon \times 10^{-4}$ ($M^{-1} cm^{-1}$)]: 302 (1.42), 356 (1.39), 388 (1.31), 474 (2.62), 545 (0.24), 734 (0.27), 806 (0.62). MS (ESI) m/z : 856.1173; calcd for $C_{44}H_{24}N_5F_9Cu$: 856.1179 $[M]^+$.

Single-crystal X-ray structure determination. X-ray structure determination. Suitable crystals were obtained by diffusion of methanol vapor into concentrated solutions of $H_2[5-pyr-TpOMePiC]$ in dichloromethane and $Cu[10-pyr-TpOMePiC]$ in chloroform. X-ray data were collected on beamline 11.3.1 at the Advanced Light Source of Lawrence Berkeley National Laboratory, Berkeley, California. The samples were mounted on MiTeGen® kapton loops and placed in a 100(2) K nitrogen cold stream provided by an Oxford Cryostream 700 Plus low temperature apparatus on the goniometer head of a Bruker D8 diffractometer equipped with PHOTONII CPAD detector. Diffraction data were collected using synchrotron radiation monochromated with silicon(111) to a wavelength of 0.7749(1) Å. In each case, an approximate full-sphere of data was collected using $1^\circ \omega$ scans. Absorption corrections were applied using SADABS.³⁴ The structure was solved by intrinsic phasing (SHELXT)³⁵ and refined by full-matrix least squares on F^2 (SHELXL-2014)³⁶ using the ShelXle GUI.³⁷ All non-hydrogen atoms were refined anisotropically. Hydrogen atoms were geometrically calculated and refined as riding atoms.

Conflicts of interest

There are no conflicts of interest to declare.

Acknowledgements

This work was supported by the Research Council of Norway (grant no. 262229 to AG) and by the Advanced Light Source, Berkeley, California. The Advanced Light Source is supported by the Director, Office of Science, Office of Basic Energy Sciences, of the U.S. Department of Energy under Contract No. DE-AC02-05CH11231.

References

- 1 R. Orlowski, D. Gryko and D. T. Gryko, *Chem. Rev.*, 2017, **117**, 3102–3137.
- 2 A. Ghosh, *Chem. Rev.*, 2017, **117**, 3798–3881.
- 3 R. D. Teo, J. Y. Hwang, J. Termini, Z. Gross and H. B. Gray, *Chem. Rev.*, 2017, **117**, 2711–2729.

- 4 G. Pomarico, X. Xiao, S. Nardis, R. Paolesse, F. R. Fronczek, K. M. Smith, Y. Fang, Z. Ou and K. M. Kadish, *Inorg. Chem.*, 2010, **49**, 5766–5774.
- 5 R. Costa, G. R. Geir III and C. J. Ziegler, *Dalton Trans.*, 2011, **40**, 4384–4386.
- 6 M. Hoffmann, B. Cordes, C. Kleeberg, P. Schweyen, B. Wolfram and M. Bröring, *Eur. J. Inorg. Chem.*, 2016, 3076–3085.
- 7 K. E. Thomas, C. M. Beavers, K. J. Gagnon and A. Ghosh, *ChemistryOpen*, 2017, **6**, 402–409.
- 8 S. Nardis, G. Pomarico, F. Mandoj, F. R. Fronczek, K. M. Smith and R. Paolesse, *J. Porphyrins Phthalocyanines*, 2010, **14**, 752–757.
- 9 S. Nardis, G. Pomarico, F. R. Fronczek, M. G. H. Vicente and R. Paolesse, *Tetrahedron Lett.*, 2007, **48**, 8643–8646.
- 10 D. Sakow, B. Böker, K. Brandhorst, O. Burghaus and M. Bröring, *Angew. Chem., Int. Ed.*, 2013, **52**, 4912–4915.
- 11 B. Umasekhar, V. S. Shetti and M. Ravikanth, *RSC Adv.*, 2018, **8**, 21100–21132.
- 12 M. Bröring, S. Köhler and C. Pietzonka, *J. Porphyrins Phthalocyanines*, 2012, **16**, 641–650.
- 13 D. Sakow, D. Baabe, B. Böker, O. Burghaus, M. Funk, C. Kleeberg, D. Menzel, C. Pietzonka and M. Bröring, *Chem. – Eur. J.*, 2014, **20**, 2913–2924.
- 14 S. Winstein, *J. Am. Chem. Soc.*, 1959, **81**, 6524–6525.
- 15 C. Foroutan-Nejad, S. Larsen, J. Conradie and A. Ghosh, *Sci. Rep.*, 2018, **8**, 11952.
- 16 C. Foroutan-Nejad and A. Ghosh, *ACS Omega*, 2018, **3**, 15865–15869.
- 17 Several pyrrole-appended porphyrinoid macrocycles are reported in the literature.^{18,19}
- 18 M. Pawlicki, I. Kańska and L. Latos-Grażyński, *Inorg. Chem.*, 2007, **46**, 6575–6584, and references therein.
- 19 M. Li, P. Wei, M. Ishida, X. Li, M. Savage, R. Guo, Z. Ou, S. Yang, H. Furuta and Y. Xie, *Angew. Chem., Int. Ed.*, 2016, **55**, 3063–3067.
- 20 A. Ghosh, *Angew. Chem., Int. Ed.*, 2004, **43**, 1918–1931.
- 21 The overwhelming preference for the 5-isocorrole regioisomer observed here differs from the much more even 5/10 regioisomer ratio reported in ref. 9. In our hands, however, the conditions used in ref. 9 (*i.e.*, DDQ-promoted addition of methanol) applied to free-base triphenylcorrole also led to a strong preponderance of the 5-isocorrole.¹⁵
- 22 I. H. Wasbotten, T. Wondimagegn and A. Ghosh, *J. Am. Chem. Soc.*, 2002, **124**, 8104–8116.
- 23 M. Bröring, F. Bregier, E. C. Tejero, C. Hell and M. C. Holthausen, *Angew. Chem., Int. Ed.*, 2007, **46**, 445–448.
- 24 A. B. Alemayehu, E. Gonzalez, L. K. Hansen and A. Ghosh, *Inorg. Chem.*, 2009, **48**, 7794–7799.
- 25 K. E. Thomas, J. Conradie, L.-K. Hansen and A. Ghosh, *Eur. J. Inorg. Chem.*, 2011, 1865–1870.
- 26 A. B. Alemayehu, L.-K. Hansen and A. Ghosh, Nonplanar, Noninnocent, and Chiral: A Strongly Saddled Metalloporrole, *Inorg. Chem.*, 2010, **49**, 7608–7610.
- 27 K. E. Thomas, H. Vazquez-Lima, Y. Fang, Y. Song, K. J. Gagnon, C. M. Beavers, K. M. Kadish and A. Ghosh, *Chem. – Eur. J.*, 2015, **21**, 16839–16847.
- 28 K. E. Thomas, L. J. McCormick, D. Carrié, H. Vazquez-Lima, G. Simonneaux and A. Ghosh, *Inorg. Chem.*, 2018, **57**, 4270–4276.
- 29 S. Berg, K. E. Thomas, C. M. Beavers and A. Ghosh, *Inorg. Chem.*, 2012, **51**, 9911–9916.
- 30 I. K. Thomassen, L. J. McCormick and A. Ghosh, *ACS Omega*, 2018, **3**, 5106–5110.
- 31 K. E. Thomas, A. B. Alemayehu, J. Conradie, C. M. Beavers and A. Ghosh, *Acc. Chem. Res.*, 2012, **45**, 1203–1214.
- 32 B. Koszarna and D. T. Gryko, *J. Org. Chem.*, 2006, **71**, 3707–3717.
- 33 Many metalloporphyrins and metalloporroles, in particular hemes and Mn and Fe corroles do not yield satisfactory elemental analyses. Others, such as coinage metal corroles, do so readily. The present complexes turned out to be intermediate. Although the majority of analyses proved to be within 0.5% of their theoretical values, a few exhibited errors of 1–2%. Although the latter can be accounted for by assuming tightly held solvent molecules, the theoretical analyses quoted here refer to simply the solvent-free molecular formulas.
- 34 L. Krause, R. Herbst-Irmer, G. M. Sheldrick and D. Stalke, Comparison of silver and molybdenum microfocus X-ray sources for single-crystal structure determination, *J. Appl. Crystallogr.*, 2015, **48**, 3–10.
- 35 G. M. Sheldrick, SHELXT - Integrated Space-Group and Crystal-Structure Determination, *Acta Crystallogr., Sect. A: Found. Adv.*, 2015, **A71**, 3–8.
- 36 G. M. Sheldrick, Crystal Structure Refinement with SHELXL, *Acta Crystallogr., Sect. C: Struct. Chem.*, 2015, **C71**, 3–8.
- 37 C. B. Hübschle, G. M. Sheldrick and B. Dittrich, ShelXle: a Qt graphical user interface for SHELXL, *J. Appl. Crystallogr.*, 2011, **44**, 1281–1284.