

UCSF

UC San Francisco Previously Published Works

Title

Comparison of Best Versus Worst Clinical Outcomes for Adult Cervical Deformity Surgery.

Permalink

<https://escholarship.org/uc/item/5sk761qp>

Journal

Global spine journal, 9(3)

ISSN

2192-5682

Authors

Smith, Justin S
Shaffrey, Christopher I
Kim, Han Jo
[et al.](#)

Publication Date

2019-05-01

DOI


10.1177/2192568218794164

Peer reviewed

Comparison of Best Versus Worst Clinical Outcomes for Adult Cervical Deformity Surgery

Global Spine Journal
2019, Vol. 9(3) 303-314
© The Author(s) 2018
Article reuse guidelines:
sagepub.com/journals-permissions
DOI: 10.1177/2192568218794164
journals.sagepub.com/home/gsj



Justin S. Smith, MD, PhD¹, Christopher I. Shaffrey, MD¹, Han Jo Kim, MD², Peter Passias, MD³, Themistocles Protopsaltis, MD³, Renaud Lafage, MS², Gregory M. Mundis Jr, MD⁴, Eric Klineberg, MD⁵, Virginie Lafage, PhD², Frank J. Schwab, MD², Justin K. Scheer, MD⁶, Michael Kelly, MD⁷ , D. Kojo Hamilton, MD⁸, Munish Gupta, MD⁷, Vedat Deviren, MD⁹, Richard Hostin, MD¹⁰, Todd Albert, MD², K. Daniel Riew, MD¹¹, Robert Hart, MD¹², Doug Burton, MD¹³, Shay Bess, MD¹⁴, Christopher P. Ames, MD⁹; on behalf of the International Spine Study Group

Abstract

Study Design: Retrospective cohort study.

Objective: Factors that predict outcomes for adult cervical spine deformity (ACSD) have not been well defined. To compare ACSD patients with best versus worst outcomes.

Methods: This study was based on a prospective, multicenter observational ACSD cohort. Best versus worst outcomes were compared based on Neck Disability Index (NDI), Neck Pain Numeric Rating Scale (NP-NRS), and modified Japanese Orthopaedic Association (mJOA) scores.

Results: Of 111 patients, 80 (72%) had minimum 1-year follow-up. For NDI, compared with best outcome patients ($n = 28$), worst outcome patients ($n = 32$) were more likely to have had a major complication ($P = .004$) and to have undergone a posterior-only procedure ($P = .039$), had greater Charlson Comorbidity Index ($P = .009$), and had worse postoperative C7-S1 sagittal vertical axis (SVA; $P = .027$). For NP-NRS, compared with best outcome patients ($n = 26$), worst outcome patients ($n = 18$) were younger ($P = .045$), had worse baseline NP-NRS ($P = .034$), and were more likely to have had a minor complication ($P = .030$). For the mJOA, compared with best outcome patients ($n = 16$), worst outcome patients ($n = 18$) were more likely to have had a major complication ($P = .007$) and to have a better baseline mJOA ($P = .030$). Multivariate models for NDI included posterior-only surgery ($P = .006$), major complication ($P = .002$), and postoperative C7-S1 SVA ($P = .012$); models for NP-NRS included baseline NP-NRS ($P = .009$), age ($P = .017$), and posterior-only surgery ($P = .038$); and models for mJOA included major complication ($P = .008$).

¹ University of Virginia, Charlottesville, VA, USA

² Hospital for Special Surgery, New York, NY, USA

³ NYU Hospital for Joint Diseases, New York, NY, USA

⁴ Scripps Clinic, San Diego, CA, USA

⁵ University of California, Davis, Sacramento, CA, USA

⁶ University of Illinois at Chicago, Chicago, IL, USA

⁷ Washington University, St Louis, MO, USA

⁸ University of Pittsburgh Medical Center, Pittsburgh, PA, USA

⁹ University of California, San Francisco, San Francisco, CA, USA

¹⁰ Baylor Scoliosis Center, Plano, TX, USA

¹¹ Columbia University, New York, NY, USA

¹² Swedish Medical Center, Seattle, WA, USA

¹³ University of Kansas Medical Center, Kansas City, KA, USA

¹⁴ Presbyterian St Lukes Medical Center, Denver, CO, USA

Corresponding Author:

Justin S. Smith, Department of Neurosurgery, University of Virginia Health Sciences Center, PO Box 800212, Charlottesville, VA 22908, USA.

Email: jss7f@virginia.edu



Conclusions: Factors distinguishing best and worst ACSD surgery outcomes included patient, surgical, and radiographic factors. These findings suggest areas that may warrant greater awareness to optimize patient counseling and outcomes.

Keywords

adult, cervical deformity, outcomes, surgery

Introduction

Early attempts to surgically treat adult cervical spine deformity (ACSD) focused on the most severe forms, and the treatments were considered high risk and had high complication rates.¹⁻⁴ As advances have been made in anesthesia and critical care, as well as surgical techniques and instrumentation, there has been an increased interest in surgically treating these deformities.⁵⁻¹³ In contrast to the progress that has been made in treating adult thoracolumbar deformities, only recently have similar advances begun for ACSD.¹⁴ Recent advances have included proposal of a cervical deformity classification¹⁵ and development of a standardized nomenclature to describe the osteotomies used to treat these deformities.¹⁶ Other recent studies have highlighted the remarkable lack of consensus for ACSD treatment strategies¹⁷ and have demonstrated that, similar to thoracolumbar deformity surgery, ACSD surgery is associated with high complication rates.^{18,19}

Although surgery for ACSD has the potential to significantly improve pain, disability, and neurological function, reported outcomes are often presented as averages.⁸ While averages may be useful for summarizing data, they do not fully reflect the spectrum of outcomes. As has been shown for thoracolumbar deformity surgeries,^{20,21} there is a range of outcomes for ACSD. Not every patient can expect to achieve the average outcome, and at the 2 extreme ends of the spectrum are those who have been left with little or no residual pain and disability (“best outcome”) and those who have improved minimally, not at all, or even worsened (“worst outcome”). Comparing patients with the best and worst outcomes following surgery for ACSD may prove useful for better optimization of patient selection, treatment strategies, and patient counseling.

In the present study, we provide assessment of a prospective, multicenter observational cohort of surgically treated ACSD patients. Our objectives were to assess the range of patient outcomes following surgical treatment for ACSD based on multiple outcomes measures. In addition, we sought to compare the patients with the best and worst outcomes based on each standardized outcome measure in order to identify factors that may distinguish these 2 groups.

Methods

Patient Population

As part of a prospective, multicenter observational cohort study, consecutive adult patients presenting for surgical

evaluation of cervical and cervicothoracic deformities were enrolled at 13 centers across the United States. Each participating site obtained institutional review board approval, and each patient provided written consent. Inclusion criteria were the following: age ≥ 18 years, cervical deformity, and plan for surgical deformity correction. Cervical deformity was defined as the presence of at least one of the following: cervical kyphosis (C2-7 sagittal Cobb angle $>10^\circ$), cervical scoliosis (C2-7 coronal Cobb angle $>10^\circ$), C2-7 sagittal vertical axis (C2-7 SVA) >4 cm, or chin-brown vertical angle $>25^\circ$. Patients with active tumor or infection were excluded. For the present study, only patients with complete baseline data and minimum 1-year follow-up were included.

Data Collection and Radiographic Assessment

Patient demographics, clinical data, surgical procedure details, and complications were collected using standardized forms. Standardized measures of health status included Neck Disability Index (NDI), modified Japanese Orthopaedic Association (mJOA), and the Neck Pain Numerical Rating Scale (NP-NRS). Osteotomies were classified as low-grade or high-grade based on a recently published classification system.¹⁶ Complications were classified as major or minor. A complication was termed major if it involved invasive intervention, had prolonged or permanent morbidity, or resulted in death.

Full-length, free-standing spine radiographs, in combination with focused cervical lateral radiographs, were obtained at baseline and at regular follow-up intervals. Radiographs were analyzed using validated software at a central core facility based on previously reported techniques.^{5,22-25} Measures included C2-C7 lordosis, C2-C7 sagittal vertical axis (C2-C7 SVA), C7-S1 SVA, cervical-thoracic pelvic angle (CTPA), pelvic tilt (PT), and mismatch between pelvic incidence and lumbar lordosis (PI-LL).

Data and Statistical Analysis

Patients with the best and worst clinical outcomes were defined separately based on 3 measures, NDI, mJOA, and NP-NRS. The changes from preoperative baseline to last follow-up were calculated for each measure for each patient and were used to create distribution plots. Thresholds for best and worst outcomes were established based on the tails at the extremes of the distribution plots and previously reported score correlates to health states.²⁶⁻³⁰ The resulting best and worst outcomes groups within each measure were then compared. To minimize

floor effects of the NDI, comparisons of best versus worst outcomes were based on patients with a preoperative baseline NDI of at least 20. To minimize the floor effects of the NP-NRS, comparisons between best versus worst outcomes were made with patients having a baseline NP-NRS score of at least 3. Based on assessment of distribution plots and established threshold cutoffs for the best and worst groups, similar cutoffs were not necessary for the mJOA. These baseline threshold cutoffs for NDI and NP-NRS were selected to reflect levels of at least moderate disability and pain based on previous reports.²⁶⁻³⁰

Demographic, clinical, and surgical variables were summarized using means and standard deviations for continuous variables and frequencies and percentages for categorical variables. For continuous variables, normality of distribution was assessed using the Kolmogorov-Smirnov test. *T* tests were used to assess for differences between normally distributed continuous variables, and the independent-samples Mann-Whitney *U* test was used for comparisons between continuous variables without normal distribution. Binary logistic regression analysis was used to adjust for the effects of multiple covariates predictive of best versus worst outcomes. Forward stepwise regression analyses were performed using variables with $P < .1$ on univariate assessment to distinguish between patients with best versus worst outcomes. Statistical analyses were performed using IBM SPSS Statistics for Windows, Version 24 (IBM Corp, Armonk, NY). Statistical tests were 2-tailed, and $P < .05$ was considered statistically significant.

Results

Patient Population

Of 111 patients who met inclusion criteria, 80 (72%) had minimum 1-year follow-up and were included in the present study. Baseline demographic, clinical, and operative parameters are summarized in Table 1. The mean available postoperative follow-up was 1.4 years and ranged from 1 to 2 years.

Mean preoperative and follow-up clinical outcomes scores and radiographic measures are summarized in Table 2. Compared with baseline, the mean NDI and NP-NRS scores improved significantly at last follow-up. The overall mJOA modestly improved, but this change did not reach statistical significance. The mean C2-C7 lordosis and overall global sagittal alignment (C7-S1 SVA) improved significantly, while significant changes were not appreciated for C2-C7 SVA, CTPA, PT, or PI-LL mismatch.

Best and Worst Outcomes Based on NDI

Of the 80 patients in the study cohort, 77 (96.3%) had a baseline NDI of ≥ 20 . For these 77 patients, the mean baseline NDI was 48.8 (SD = 15.7, range 22.2 to 92.0) and the mean follow-up NDI was 36.0 (SD = 20.3, range 0.0 to 80.0; Figure 1). Based on the NDI, the best outcome group (NDI improved by ≥ 20 points) consisted of 28 patients, and the worst outcome

Table 1. Demographic, Clinical, and Operative Parameters for 80 Surgically Treated Adults With Cervical Spinal Deformity.

Parameter	Value
Mean follow-up, years (SD, range)	1.4 (0.5, 1-2)
Mean age, years (SD, range)	62.2 (10.7, 36.5-82.6)
Male/female	33:47
Mean BMI (SD, range)	29.1 (7.6, 16.8-58.3)
Smoker (%)	5.1
Depression/anxiety (%)	26.3
Mean CCI (SD, range)	0.7 (1.0, 0-6)
Diagnosis	
Degenerative kyphosis (%)	47.5
Iatrogenic kyphosis (%)	20.0
Degenerative kyphoscoliosis (%)	5.0
Degenerative scoliosis (%)	5.0
Traumatic kyphosis (%)	2.5
Congenital kyphosis (%)	2.5
Other (%)	17.5
Previous cervical spine surgery (%)	42.3
Mean estimated blood loss, L (SD, range)	0.88 (0.89, 0.005-4.5)
Mean operative time, hours (SD, range)	6.2 (3.8, 1.9-22.7)
Mean number of fusion levels (SD, range)	7.8 (3.8, 3-19)
Surgical approach	
Anterior-only (%)	15.0
Posterior-only (%)	47.5
Combined (%)	37.5
Surgical procedures	
ACDF (%)	47.5
Corpectomy (%)	18.8
Low-grade posterior osteotomy (%) ^a	30.0
High-grade posterior osteotomy (%) ^a	20.0
Posterior neural decompression (%) ^b	56.3
Complications	
Any major complication (%)	43.8
Any minor complication (%)	36.3
Any complication (%)	66.3

Abbreviations: SD, standard deviation; BMI, body mass index; CCI, Charlson Comorbidity Index; mJOA, modified Japanese Orthopaedic Association; NDI, Neck Disability Index; ACDF, anterior cervical discectomy and fusion.

^aLow-grade osteotomies included partial or complete facetectomy and Smith-Petersen osteotomy. High-grade osteotomies included pedicle subtraction osteotomy, vertebral column resection, and opening wedge 3-column osteotomy.

^bIncludes laminectomy and/or foraminotomy for the purposes of decompression.

group (NDI improved by < 10 points or worsened) consisted of 32 patients (Figure 1). The best outcome group had a mean baseline NDI of 49.7 (SD = 14.9) that improved to a mean of 21.4 (SD = 14.1) at follow-up. In contrast, the worst outcome group had a mean baseline NDI of 48.5 (SD = 14.4) that increased to a mean follow-up NDI of 50.3 (SD = 14.7).

Univariate assessments between the best and worst outcomes groups based on NDI are summarized in Table 3. Compared with best outcome patients, at baseline those with the worst outcomes had a higher comorbidity score ($P = .009$), lower PT ($P = .003$), and more negative PI-LL mismatch ($P = .012$). A significantly higher proportion of patients in the worst outcome group was treated with a posterior-only approach ($P = .039$), had a high-grade osteotomy ($P = .004$),

and had a major complication ($P = .004$). At follow-up the worst outcome group based on NDI also had worse mJOA ($P = .042$) and NP-NRS ($P < .001$), had worse global sagittal alignment based on the C7-S1 SVA ($P = .027$), and had less increase in cervical lordosis ($P = .041$; Table 3).

The best-fit model based on logistic regression analysis included 3 parameters: posterior-only approach ($P = .006$), occurrence of a major complication ($P = .012$), and C7-S1 SVA at last follow-up ($P = .012$; Table 4).

Best and Worst Outcomes Based on NP-NRS

Of the 80 patients in the study cohort, 74 (92.5%) had NP-NRS ≥ 3 . For these 74 patients, the mean baseline NP-NRS was

Table 2. Radiographic and Clinical Outcomes at Baseline and Minimum 1-Year Follow-up for 80 Surgically Treated Adults With Cervical Spinal Deformity.¹

Parameter	Preoperative	Minimum 1-Year Follow-up	P Value
Mean NDI (SD)	47.4 (16.9)	35.4 (20.3)	<.001
Mean mJOA score (SD)	13.6 (2.6)	14.3 (2.7)	.050
Mean NP-NRS score (SD)	6.6 (2.4)	4.0 (3.0)	<.001
Mean C2-C7 lordosis, ° (SD)	4.9 (20.2)	-8.2 (16.3)	<.001
Mean C2-C7 SVA, mm (SD)	45.3 (25.4)	43.0 (18.5)	.38
Mean C7-S1 SVA, mm (SD)	7.7 (68.2)	35.3 (59.4)	<.001
Mean CTPA, ° (SD)	4.9 (2.9)	4.6 (2.1)	.32
Mean PT, ° (SD)	18.6 (12.1)	17.9 (12.1)	.49
Mean PI-LL mismatch, ° (SD)	-0.5 (18.7)	0.5 (17.7)	.48

Abbreviations: SD, standard deviation; NDI, Neck Disability Index; mJOA, modified Japanese Orthopaedic Association; NP-NRS, Neck Pain Numeric Rating Scale; SVA, sagittal vertical axis; CTPA, cervical-thoracic pelvic angle; PT, pelvic tilt; PI, pelvic incidence; LL, lumbar lordosis.

7.1 (SD = 1.8, range 3 to 10) and the mean follow-up NP-NRS was 4.2 (SD = 3.0, range 0 to 10.0; Figure 2). Based on the NP-NRS, the best outcome group (NP-NRS improved by ≥ 5 points) consisted of 26 patients, and the worst outcome group (NP-NRS without improvement or worsened) consisted of 18 patients (Figure 2). The best outcome group had a mean baseline NP-NRS of 7.6 (SD = 1.4) that improved to a mean of 1.3 (SD = 1.4) at follow-up. In contrast, the worst outcome group had a mean baseline NP-NRS of 6.3 (SD = 1.8) that increased to a mean follow-up NP-NRS of 7.6 (SD = 1.7).

Univariate assessments between the best and worst outcomes groups based on NP-NRS are summarized in Table 5. Compared with best outcome patients, at baseline those with the worst outcomes were younger ($P = .045$) and had lower NP-NRS ($P = .034$). Minor complications were more common among patients in the best outcome group ($P = .03$). At follow-up the worst outcome group had worse NDI ($P < .001$) and had a modestly but significantly lower CTPA (Table 5).

The best-fit model based on logistic regression analysis included 3 parameters: preoperative NP-NRS ($P = .009$), age ($P = .017$), and posterior-only approach ($P = .038$; Table 6).

Best and Worst Outcomes Based on mJOA

Of the 80 patients in the study cohort, 64 (80.0%) had a baseline and follow-up mJOA documented. For these 64 patients, the mean mJOA was 13.6 (SD = 2.6) and the mean follow-up mJOA was 14.3 (SD = 2.7). Based on the mJOA, the best outcome group (mJOA score improved by ≥ 2 points) consisted of 16 patients, and the worst outcome group (mJOA worsened by ≥ 1 point) consisted of 18 patients (Figure 3). The best outcome group had a mean baseline mJOA of 12.3 (SD = 2.2) that improved to a mean of 16.4 (SD = 1.7) at




Figure 1. Change in Neck Disability Index (NDI) scores from preoperative baseline to minimum 1-year follow-up for 77 patients surgically treated for adult cervical deformity. Each patient had a preoperative baseline NDI score of at least 20 points. Cutoffs for best and worst outcomes are indicated.

Table 3. Univariate Analysis of Baseline, Operative, and Follow-up Parameters Between Patients With the Best Versus Worst Outcomes Based on NDI Following Surgery for Adult Cervical Deformity.^a

Parameter	Worst (n = 32)	Best (n = 28)	P Value
Baseline			
Mean age, years (SD)	60.7 (11.5)	61.9 (9.2)	.68
M/F	14:18	11:17	.80
Mean BMI (SD)	30.7 (9.4)	28.4 (6.7)	.29
Smoker (%)	3.1	3.8	1.00
Depression/anxiety (%)	21.9	32.1	.40
Mean CCI (SD)	1.1 (1.3)	0.5 (0.8)	.009^b
Diagnosis			.58
Degenerative kyphosis (%)	53.1	46.4	
Iatrogenic kyphosis (%)	21.9	21.4	
Degenerative kyphoscoliosis (%)	6.3	3.6	
Degenerative scoliosis (%)	0.0	7.1	
Traumatic kyphosis (%)	0.0	3.6	
Congenital kyphosis (%)	3.1	0.0	
Other (%)	15.6	17.9	
Previous cervical spine surgery (%)	46.9	34.6	.43
Mean NDI (SD)	48.5 (14.4)	49.7 (14.9)	.77
Mean mJOA score (SD)	13.5 (2.6)	13.5 (2.8)	1.00
Mean NP-NRS score (SD)	6.8 (2.4)	6.8 (2.4)	.95 ^b
Mean C2-C7 lordosis, ° (SD)	3.0 (18.5)	11.2 (20.7)	.13
Mean C2-C7 SVA, mm (SD)	48.3 (26.7)	48.7 (23.3)	.95
Mean C7-S1 SVA, mm (SD)	14.3 (81.8)	15.3 (80.3)	.96
Mean CTPA, ° (SD)	4.7 (2.7)	4.9 (2.8)	.81
Mean PT, ° (SD)	13.9 (11.4)	22.7 (10.0)	.003
Mean PI-LL mismatch, ° (SD)	-4.1 (21.6)	6.0 (15.2)	.012^b
Operative			
Mean estimated blood loss, L (SD)	0.8 (0.7)	0.7 (0.8)	.22 ^b
Mean operative time, hours (SD)	5.7 (2.6)	6.6 (5.2)	.85 ^b
Mean number of fusion levels (SD)	8.2 (3.9)	6.4 (2.8)	.058 ^b
Surgical approach			
Anterior-only (%)	9.4	25.0	.17
Posterior-only (%)	56.3	28.6	.039
Combined (%)	34.4	46.4	.43
High-grade posterior osteotomy (%) ^c	25.0	0.0	.004
Any major complication (%)	59.4	21.4	.004
Any minor complication (%)	21.9	42.9	.10
Any complication (%)	71.9	53.6	.18
Follow-up			
Mean NDI (SD)	50.3 (14.7)	21.4 (14.1)	<.001
Mean mJOA score (SD)	13.6 (3.0)	15.3 (2.4)	.042^b
Mean NP-NRS score (SD)	5.9 (2.5)	2.4 (2.7)	<.001
Mean C2-C7 lordosis, ° (SD)	-7.4 (16.8)	-10.4 (11.9)	.23 ^b
Mean C2-C7 SVA, mm (SD)	44.8 (19.4)	37.8 (15.5)	.20
Mean C7-S1 SVA, mm (SD)	51.3 (69.6)	13.4 (40.8)	.027
Mean CTPA, ° (SD)	4.3 (2.0)	4.3 (1.9)	.99
Mean PT, ° (SD)	14.0 (11.2)	20.0 (12.0)	.081
Mean PI-LL mismatch, ° (SD)	-1.8 (18.6)	2.8 (15.6)	.36
Change from baseline to follow-up			
Mean C2-C7 lordosis, ° (SD)	8.2 (17.0)	19.8 (17.5)	.041
Mean C2-C7 SVA, mm (SD)	1.0 (19.0)	-7.8 (20.0)	.16
Mean C7-S1 SVA, mm (SD)	26.8 (48.4)	8.5 (47.4)	.34
Mean CTPA, ° (SD)	-0.6 (1.7)	0.0 (2.1)	.44
Mean PT, ° (SD)	-0.7 (8.2)	-2.2 (6.2)	.61
Mean PI-LL mismatch, ° (SD)	2.0 (9.0)	-0.8 (8.8)	.41

Abbreviations: SD, standard deviation; BMI, body mass index; CCI, Charlson Comorbidity Index; NDI, Neck Disability Index; mJOA, modified Japanese Orthopaedic Association; NP-NRS, Neck Pain Numeric Rating Scale; SVA, sagittal vertical axis; CTPA, cervical-thoracic pelvic angle; PT, pelvic tilt; PI, pelvic incidence; LL, lumbar lordosis; A/P/C, anterior-only/posterior-only/combined approach.

^aSignificant *p*-values are shown in boldface type.

^bMann-Whitney *U* test.

^cHigh-grade osteotomies included pedicle subtraction osteotomy, vertebral column resection, and opening wedge 3-column osteotomy.

follow-up. In contrast, the worst outcome group had a mean baseline mJOA of 14.1 (SD = 2.2) that worsened to a mean follow-up mJOA of 11.9 (SD = 1.8).

Univariate assessments between the best and worst outcomes groups based on mJOA are summarized in Table 7. Compared with best outcome patients, at baseline those with the worst outcomes had better mJOA ($P = .030$). The occurrence of major complications was more common among the patients with the worst outcomes ($P = .007$). Except for differences in mJOA, at follow-up the best and worst outcome groups did not differ based on radiographic measures or outcomes measures (Table 7).

The best-fit model based on logistic regression analysis only included occurrence of a major complication ($P = .008$; Table 8).

Discussion

The present study provides a prospective assessment of the clinical improvement following surgical treatment for

Table 4. Multivariate Analysis of Factors Distinguishing Between Patients With the Best Versus Worst Outcomes Based on NDI Following Surgery for Adult Cervical Deformity^a.

Parameter	OR	95% CI	P Value
Posterior-only surgical approach	0.028	0.002-0.366	.006
Occurrence of major complication	0.016	0.001-0.232	.002
C7-S1 SVA at last follow-up	0.969 ^b	0.946-0.993	.012

Abbreviations: NDI, Neck Disability Index; OR, odds ratio; CI, confidence interval; SVA, sagittal vertical axis.

^aStepwise binary logistic regression; results of best-fit model presented.

^bOR for C7-S1 SVA is per millimeter.

80 ACSD patients at a minimum 1-year follow-up based on standardized outcomes measures of disability (NDI), pain (NP-NRS), and myelopathy (mJOA). The overall patient cohort demonstrated significant improvement in pain and disability and had a modest improvement in myelopathy scores that bordered on statistical significance. Although the outcome score averages reflect overall improvement from the time of surgery to last follow-up, the amount of change in each measure at the individual patient level varied considerably. Comparing the best and worst outcomes groups demonstrated distinguishing factors that included a mix of patient (age, baseline NP-NRS, Charlson Comorbidity Index [CCI]), surgical (complications and surgical approach), and radiographic (follow-up C7-S1 SVA) parameters. Collectively, these findings demonstrate the broad range in outcomes following surgery for ACSD and suggest that the factors that may account for this variation are complex and multifactorial.

Predictors of the best versus worst outcomes varied based on the 3 different outcome metrics, likely due to differences in what the 3 metrics assess. The NP-NRS is a direct measure of pain, while the NDI assesses how neck pain affects a patient's daily life and activities.^{31,32} The mJOA provides a focused assessment of functional disability related to cervical myelopathy.³³

On univariate analysis, the occurrence of complications was a significant distinguishing factor between patients with the best and worst outcomes for all 3 measures assessed. For 2 of the measures (NDI and mJOA), the occurrence of a major complication was incorporated into the best-fit multivariate model. Although this association has not been previously reported for ACSD, authors have previously assessed the impact of complications on outcome for adult thoracolumbar




Figure 2. Change in Neck Pain Numeric Rating Scale (NP-NRS) scores from preoperative baseline to minimum 1-year follow-up for 74 patients surgically treated for adult cervical deformity. Each patient had a preoperative baseline NP-NRS score of at least 3. Cutoffs for best and worst outcomes are indicated.

Table 5. Univariate Analysis of Baseline, Operative, and Follow-up Parameters Between Patients With the Best Versus Worst Outcomes Based on NP-NRS Following Surgery for Adult Cervical Deformity.^a

Parameter	Worst (n = 18)	Best (n = 26)	P Value
Baseline			
Mean age, years (SD)	60.3 (9.7)	66.2 (8.9)	.045
Male/female	5:13	11:15	.36
Mean BMI (SD)	30.3 (11.6)	28.5 (5.4)	.55
Smoker (%)	5.6	0.0	.42
Depression/anxiety (%)	16.7	38.5	.18
Mean CCI (SD)	0.8 (0.9)	0.5 (0.8)	.27 ^a
Diagnosis			.10
Degenerative kyphosis (%)	44.4	61.5	
Iatrogenic kyphosis (%)	22.2	7.7	
Degenerative kyphoscoliosis (%)	16.7	0.0	
Congenital kyphosis (%)	0.0	3.8	
Other (%)	16.7	26.9	
Previous cervical spine surgery (%)	50.0	32.0	.34
Mean NDI (SD)	53.2 (12.7)	47.2 (19.1)	.22
Mean mJOA score (SD)	12.7 (3.1)	13.9 (2.6)	.22
Mean NP-NRS score (SD)	6.3 (1.8)	7.6 (1.4)	.034^b
Mean C2-C7 lordosis, ° (SD)	5.8 (21.5)	9.9 (15.3)	.49
Mean C2-C7 SVA, mm (SD)	50.9 (25.1)	47.7 (26.7)	.70
Mean C7-S1 SVA, mm (SD)	21.0 (88.3)	18.1 (62.4)	.91
Mean CTPA, ° (SD)	5.2 (2.6)	4.9 (3.1)	.73
Mean PT, ° (SD)	16.8 (12.7)	21.8 (10.2)	.15
Mean PI-LL mismatch, ° (SD)	-2.9 (21.3)	4.8 (14.3)	.12 ^b
Operative			
Mean estimated blood loss, L (SD)	1.0 (0.6)	0.7 (0.9)	.069 ^b
Mean operative time, hours (SD)	5.6 (2.6)	6.1 (3.6)	.90 ^b
Mean number of fusion levels (SD)	7.9 (3.2)	7.0 (3.7)	.18 ^b
Surgical approach			
Anterior-only (%)	0.0	23.1	.067
Posterior-only (%)	66.7	34.6	.065
Combined (%)	33.3	42.3	.75
High-grade posterior osteotomy (%) ^c	33.3	11.5	.13
Any major complication (%)	50.0	34.6	.36
Any minor complication (%)	16.7	50.0	.030
Any complication (%)	66.7	61.5	.76
Follow-up			
Mean NDI (SD)	52.0 (14.7)	26.3 (19.4)	<.001
Mean mJOA score (SD)	13.7 (2.5)	14.6 (3.2)	.35
Mean NP-NRS score (SD)	7.6 (1.7)	1.3 (1.4)	<.001^b
Mean C2-C7 lordosis, ° (SD)	-5.8 (17.1)	-9.6 (16.0)	.51
Mean C2-C7 SVA, mm (SD)	45.3 (18.4)	43.8 (18.9)	.81
Mean C7-S1 SVA, mm (SD)	61.4 (75.8)	36.9 (46.7)	.27
Mean CTPA, ° (SD)	4.3 (2.0)	4.8 (2.1)	.52
Mean PT, ° (SD)	16.7 (11.9)	20.8 (13.1)	.32
Mean PI-LL mismatch, ° (SD)	-1.7 (17.6)	4.5 (16.3)	.28
Change from baseline to follow-up			
Mean C2-C7 lordosis, ° (SD)	10.5 (21.1)	19.2 (18.6)	.23
Mean C2-C7 SVA, mm (SD)	-3.1 (26.4)	2.5 (17.1)	.43 ^b
Mean C7-S1 SVA, mm (SD)	34.3 (44.8)	30.5 (57.8)	.87
Mean CTPA, ° (SD)	-1.4 (2.1)	0.4 (1.7)	.035
Mean PT, ° (SD)	-0.2 (9.5)	-0.2 (6.9)	1.00
Mean PI-LL mismatch, ° (SD)	2.3 (9.6)	4.4 (12.3)	.67

Abbreviations: SD, standard deviation; BMI, body mass index; CCI, Charlson Comorbidity Index; NDI, Neck Disability Index; mJOA, modified Japanese Orthopaedic Association; NP-NRS, Neck Pain Numeric Rating Scale; SVA, sagittal vertical axis; CTPA, cervical-thoracic pelvic angle; PT, pelvic tilt; PI, pelvic incidence; LL, lumbar lordosis; A/P/C, anterior-only/posterior-only/combined approach.

^aSignificant *p*-values are shown in boldface type.

^bMann-Whitney *U* test.

^cHigh-grade osteotomies included pedicle subtraction osteotomy, vertebral column resection, and opening wedge 3-column osteotomy.

deformity. Smith and colleagues performed a risk-benefit assessment of surgery for adult thoracolumbar deformity with stratification based on age.³⁴ They reported that, despite having a 4-fold higher complication rate compared with the young patient group, the elderly had greater magnitudes of improvement for pain and disability, suggesting that globally the impact of complications may not preclude significant clinical improvement following surgery. Bridwell and colleagues also reported outcomes for adult thoracolumbar deformity surgery and noted that there was a trend toward a smaller incremental improvement at 2-year follow-up in those patients who had a major complication.³⁵ With 3- to 5-year follow-up of the same patient population, these authors reported a significant impact of complications on outcome.³⁶ The present study, in combination with the findings of Bridwell and colleagues, suggests that complications do have the potential to significantly impact patient outcome, especially when comparing patients at the extremes (best versus worst outcomes). It must be recognized that, despite the best of care, there are inherent risks of complications associated with these often complex procedures.

Nevertheless, these findings further emphasize that efforts to reduce complications associated with these procedures are warranted.

Posterior-only surgical approach was found to be significantly more common among patients with the worst outcomes. This association was apparent in the multivariate modeling for 2 of the outcomes measures (NDI and NP-NRS). There are many potential explanations for these associations. High-grade osteotomies are typically performed through a posterior-only approach.¹⁶ These osteotomies are often reserved for the most severe deformities and are among the most aggressive of correction techniques. It is possible that the posterior-only approach may be disproportionately represented among the patients with the worst outcomes due to its favored use in the most severe and extensive of deformities, especially those requiring a 3-column osteotomy. Notably, for the univariate assessment of factors associated with best versus worst outcomes based on NDI, use of a high-grade osteotomy was significantly associated with the worst outcomes. The substantially greater muscle disruption associated with a posterior approach, compared with an anterior approach, may also partially account for the posterior approach being associated with poorer outcomes based on neck pain and disability. In addition, the posterior approaches used for some patients for deformity correction were extensive, with some necessitating extension of instrumentation and arthrodesis into the lower thoracic or lumbar spine, which could also affect clinical outcomes. These findings certainly do not suggest that the posterior-only approach is always associated with poor outcomes or that it should be abandoned. Selection of surgical approach is often driven by deformity type, history of previous procedures, and the goals of correction. For treatment of many cervical deformities, an anterior-only approach may not be feasible.

Table 6. Multivariate Analysis of Factors Distinguishing Between Patients With the Best Versus Worst Outcomes Based on NP-NRS Following Surgery for Adult Cervical Deformity^a.

Parameter	OR	95% CI	P Value
Preoperative NP-NRS	2.508 ^b	1.254-5.014	.009
Patient age (years)	1.114 ^c	1.019-1.217	.017
Posterior-only surgical approach	0.166	0.030-0.903	.038

Abbreviations: NP-NRS, Neck Pain Numeric Rating Scale; OR, odds ratio; CI, confidence interval.

^aStepwise binary logistic regression; results of best-fit model presented.

^bOR for NP-NRS is per point.

^cOR for patient age is per year.




Figure 3. Change in modified Japanese Orthopaedic Association (mJOA) scores from preoperative baseline to minimum 1-year follow-up for 64 patients surgically treated for adult cervical deformity. Cutoffs for best and worst outcomes are indicated.

Table 7. Univariate Analysis of Baseline, Operative, and Follow-up Parameters Between Patients With the Best Versus Worst Outcomes Based on mJOA Following Surgery for Adult Cervical Deformity.^a

Parameter	Worst (n = 18)	Best (n = 16)	P Value
Baseline			
Mean age, years (SD)	61.5 (11.5)	58.8 (8.0)	.44
Male/female	8:10	4:12	.30
Mean BMI (SD)	28.0 (7.2)	26.7 (7.9)	.64
Smoker (%)	0.0	6.7	.46
Depression/anxiety (%)	33.3	31.3	1.00
Mean CCI (SD)	0.9 (1.6)	0.6 (0.9)	.80 ^b
Diagnosis			.86
Degenerative kyphosis (%)	55.6	43.8	
Iatrogenic kyphosis (%)	22.2	25.0	
Degenerative kyphoscoliosis (%)	5.6	12.5	
Other (%)	16.7	18.8	
Previous cervical spine surgery (%)	55.6	33.3	.30
Mean NDI (SD)	49.0 (18.0)	51.3 (13.7)	.69
Mean mJOA score (SD)	14.1 (2.2)	12.3 (2.2)	.030
Mean NP-NRS score (SD)	6.3 (3.2)	7.6 (1.9)	.33 ^b
Mean C2-C7 lordosis, ° (SD)	5.2 (26.3)	6.8 (15.8)	.91 ^b
Mean C2-C7 SVA, mm (SD)	50.8 (26.4)	44.9 (21.2)	.49
Mean C7-S1 SVA, mm (SD)	7.5 (75.1)	15.5 (46.4)	.72
Mean CTPA, ° (SD)	5.2 (2.7)	4.6 (2.9)	.59
Mean PT, ° (SD)	19.4 (9.9)	23.0 (14.9)	.42
Mean PI-LL mismatch, ° (SD)	0.96 (14.4)	7.3 (16.9)	.25
Operative			
Mean estimated blood loss, L (SD)	0.9 (0.7)	0.7 (0.9)	.10 ^b
Mean operative time, hours (SD)	5.5 (2.5)	5.3 (3.3)	.57 ^b
Mean number of fusion levels (SD)	7.8 (3.2)	6.8 (3.7)	.25 ^b
Surgical approach			
Anterior-only (%)	5.6	25.0	.16
Posterior-only (%)	50.0	50.0	1.00
Combined (%)	44.4	25.0	.30
High-grade posterior osteotomy (%) ^c	22.2	12.5	.66
Any major complication (%)	66.7	18.8	.007
Any minor complication (%)	16.7	50.0	.066
Any complication (%)	77.8	62.5	.46
Follow-up			
Mean NDI (SD)	40.3 (21.3)	30.0 (16.9)	.13
Mean mJOA score (SD)	11.9 (1.8)	16.4 (1.7)	<.001
Mean NP-NRS score (SD)	4.4 (3.2)	4.1 (3.1)	.77
Mean C2-C7 lordosis, ° (SD)	-14.2 (22.5)	-7.0 (9.5)	.30
Mean C2-C7 SVA, mm (SD)	42.0 (19.5)	41.2 (18.6)	.91
Mean C7-S1 SVA, mm (SD)	43.0 (62.9)	11.5 (47.0)	.14
Mean CTPA, ° (SD)	4.3 (2.1)	4.6 (2.5)	.78
Mean PT, ° (SD)	18.2 (10.3)	20.6 (15.2)	.65
Mean PI-LL mismatch, ° (SD)	2.6 (14.1)	3.4 (18.8)	.90
Change from baseline to follow-up			
Mean C2-C7 lordosis, ° (SD)	19.9 (21.0)	13.1 (12.7)	.32
Mean C2-C7 SVA, mm (SD)	-9.3 (20.8)	-2.49 (11.9)	.31
Mean C7-S1 SVA, mm (SD)	38.9 (47.1)	0.9 (45.1)	.091
Mean CTPA, ° (SD)	-1.2 (1.5)	-0.2 (1.2)	.14
Mean PT, ° (SD)	-0.6 (6.1)	-4.3 (6.6)	.21
Mean PI-LL mismatch, ° (SD)	4.0 (8.5)	-4.7 (11.1)	.072

Abbreviations: SD, standard deviation; BMI, body mass index; CCI, Charlson Comorbidity Index; NDI, Neck Disability Index; mJOA, modified Japanese Orthopaedic Association; NP-NRS, Neck Pain Numeric Rating Scale; SVA, sagittal vertical axis; CTPA, cervical-thoracic pelvic angle; PT, pelvic tilt; PI, pelvic incidence; LL, lumbar lordosis; A/P/C, anterior-only/posterior-only/combined approach.

^aSignificant *p*-values are shown in boldface type.

^bMann-Whitney *U* test.

^cHigh-grade osteotomies included pedicle subtraction osteotomy, vertebral column resection, and opening wedge 3-column osteotomy.

Table 8. Multivariate Analysis of Factors Distinguishing Between Patients With the Best Versus Worst Outcomes Based on mJOA Following Surgery for Adult Cervical Deformity^a.

Parameter	OR	95% CI	P Value
Occurrence of major complication	0.115	0.023-0.567	.008

Abbreviations: mJOA, modified Japanese Orthopaedic Association; OR, odds ratio; CI, confidence interval.

^aStepwise binary logistic regression; results of best-fit model presented.

On both univariate and multivariate analysis, residual global sagittal malalignment (C7-S1 SVA) was significantly associated with the worst outcome group based on the NDI. The negative impact of positive global sagittal malalignment on pain and disability is well recognized for thoracolumbar deformity.^{37,38} Although the C7-S1 SVA did not differ preoperatively between the patients that would ultimately have the best and worst outcomes, it is possible that changes in alignment induced by the cervical deformity correction may have altered the global alignment through changes in compensatory measures.^{39,40}

Patient age was a significant distinguishing factor between the best and worst outcomes based on the NP-NRS univariate and multivariate analyses. This association may seem counterintuitive, since the older patients were favored to have the best outcome. The reason for this association is unclear but may relate to unaccounted for differences in deformity types and differences in baseline pain and disability across age groups.

Smith and colleagues have assessed best versus worst outcomes for adult thoracolumbar deformity surgery based on the Oswestry Disability Index and Scoliosis Research Society 22r (SRS-r). Similar to the present study, they reported that the factors that distinguished the best and worst outcomes were a mix of patient factors (baseline depression, BMI, comorbidities, and disability), radiographic factors (residual global sagittal malalignment as assessed by the C7-S1 SVA), and the occurrence of complications.^{20,21} Although baseline depression and BMI were not distinguishing factors in the present study, it is notable that the remaining factors overlap between the 2 studies and suggest that the factors that affect outcomes of adult thoracolumbar and cervical deformity surgeries may have more similarities than differences.

The primary strength of the present study is the prospective multicenter design with standardized data collection. In addition, the patient cohort is heterogeneous and represents a broad spectrum of cervical deformities treated by multiple surgeons across multiple institutions, which enhances the generalizability of the findings. The primary limitation of the present study relates to the number of patients. Although 80 is a relatively large number of patients for ACSD, the limited numbers of patients, especially when focusing on those at the extreme ends of the outcomes spectrum, does impact the ability to perform granular assessments of factors associated with outcomes. In addition, although multiple statistical comparisons were performed in the present study, we intentionally did not perform a

Bonferroni correction, since we would rather commit a Type I error than miss potentially important associations, especially in the setting of limited numbers of patients. Another limitation of the present study relates to the distribution of the patient population enrolled in the database. Since the mean patient age was 62 years and the youngest patient was 36.5 years, the findings of the present study may be most relevant to the older ACSD patient population. Last, there remains no ACSD-specific outcome measure, which necessitated the use of more generalized outcomes measures that cover neck disability, neck pain, and cervical myelopathy.

Conclusions

Based on a prospectively collected multicenter series of ACSD patients, this study provides assessment of the clinical improvement following surgical treatment for 80 ACSD patients at a minimum 1-year follow-up based on standardized outcomes measures of disability, pain, and myelopathy. Although the overall patient cohort demonstrated significant improvement in outcomes, there was substantial variation at the individual patient level. The factors that distinguished between the best and worst outcomes groups included a mix of patient (age, baseline NP-NRS, CCI), surgical (complications and surgical approach), and radiographic (follow-up C7-S1 SVA) parameters. Collectively, these findings demonstrate the broad range in outcomes following surgery for ACSD and suggest that the factors that may account for this variation are complex and multifactorial. These findings suggest areas that may warrant greater awareness to help optimize patient selection, treatment strategies, and preoperative patient counseling.


Declaration of Conflicting Interests

The author(s) declared no potential conflicts of interest with respect to the research, authorship, and/or publication of this article.

Funding

The author(s) disclosed receipt of the following financial support for the research, authorship, and/or publication of this article: The International Spine Study Group (ISSG) is funded through research grants from DePuy Synthes and individual donations.

ORCID iD

Michael Kelly, MD  <https://orcid.org/0000-0001-6221-7406>

References

1. Bovill EG Jr. Osteotomy of cervical part of the spine in ankylosing spondylitis with severe deformity. *Calif Med.* 1965; 102:142-144.
2. Simmons EH. The surgical correction of flexion deformity of the cervical spine in ankylosing spondylitis. *Clin Orthop Relat Res.* 1972;86:132-143.
3. Simmons EH, Brown ME. Surgery for kyphosis in ankylosing spondylitis. *Can Nurse.* 1972;68:24-29.

4. Urist MR. Osteotomy of the cervical spine; report of a case of ankylosing rheumatoid spondylitis. *J Bone Joint Surg Am.* 1958; 40A:833-843.
5. Ames CP, Blondel B, Scheer JK, et al. Cervical radiographical alignment: comprehensive assessment techniques and potential importance in cervical myelopathy. *Spine (Phila Pa 1976).* 2013;38(22 suppl 1):S149-S160.
6. Caruso L, Barone G, Farneti A, Caraffa A. Pedicle subtraction osteotomy for the treatment of chin-on-chest deformity in a post-radiotherapy dropped head syndrome: a case report and review of literature. *Eur Spine J.* 2014;23(suppl 6):634-643.
7. Etame AB, Than KD, Wang AC, La Marca F, Park P. Surgical management of symptomatic cervical or cervicothoracic kyphosis due to ankylosing spondylitis. *Spine (Phila Pa 1976).* 2008;33: E559-E564.
8. Etame AB, Wang AC, Than KD, La Marca F, Park P. Outcomes after surgery for cervical spine deformity: review of the literature. *Neurosurg Focus.* 2010;28:E14.
9. Grosso MJ, Hwang R, Krishnaney AA, Mroz TE, Benzel EC, Steinmetz MP. Complications and outcomes for surgical approaches to cervical kyphosis. *J Spinal Disord Tech.* 2015;28: E385-E393.
10. Han K, Lu C, Li J, et al. Surgical treatment of cervical kyphosis. *Eur Spine J.* 2011;20:523-536.
11. O'Shaughnessy BA, Liu JC, Hsieh PC, Koski TR, Ganju A, Ondra SL. Surgical treatment of fixed cervical kyphosis with myelopathy. *Spine (Phila Pa 1976).* 2008;33:771-778.
12. Scheer JK, Tang JA, Smith JS, et al. Cervical spine alignment, sagittal deformity, and clinical implications: a review. *J Neurosurg Spine.* 2013;19:141-159.
13. Tang JA, Scheer JK, Smith JS, et al. The impact of standing regional cervical sagittal alignment on outcomes in posterior cervical fusion surgery. *Neurosurgery.* 2012;71:662-669.
14. Smith JS, Shaffrey CI, Bess S, et al. Recent and emerging advances in spinal deformity. *Neurosurgery.* 2017;80(3 suppl): S70-S85.
15. Ames CP, Smith JS, Eastlack R, et al. Reliability assessment of a novel cervical spine deformity classification system. *J Neurosurg Spine.* 2015;23:673-683.
16. Ames CP, Smith JS, Scheer JK, et al. A standardized nomenclature for cervical spine soft-tissue release and osteotomy for deformity correction: clinical article. *J Neurosurg Spine.* 2013;19: 269-278.
17. Smith JS, Klineberg E, Shaffrey CI, et al. Assessment of surgical treatment strategies for moderate to severe cervical spinal deformity reveals marked variation in approaches, osteotomies, and fusion levels. *World Neurosurg.* 2016;91:228-237.
18. Smith JS, Klineberg E, Lafage V, et al. Prospective multicenter assessment of perioperative and minimum 2-year postoperative complication rates associated with adult spinal deformity surgery. *J Neurosurg Spine.* 2016;25:1-14.
19. Smith JS, Ramchandran S, Lafage V, et al. Prospective multicenter assessment of early complication rates associated with adult cervical deformity surgery in 78 patients. *Neurosurgery.* 2016;79:378-388.
20. Smith JS, Shaffrey CI, Glassman SD, et al. Clinical and radiographic parameters that distinguish between the best and worst outcomes of scoliosis surgery for adults. *Eur Spine J.* 2013;22: 402-410.
21. Smith JS, Shaffrey CI, Lafage V, et al. Comparison of best versus worst clinical outcomes for adult spinal deformity surgery: a retrospective review of a prospectively collected, multicenter database with 2-year follow-up. *J Neurosurg Spine.* 2015;23:349-359.
22. Ames CP, Smith JS, Scheer JK, et al. Impact of spinopelvic alignment on decision making in deformity surgery in adults: a review. *J Neurosurg Spine.* 2012;16:547-564.
23. Protopsaltis T, Bronsard N, Soroceanu A, et al. Cervical sagittal deformity develops after PJK in adult thoracolumbar deformity correction: radiographic analysis utilizing a novel global sagittal angular parameter, the CTPA. *Eur Spine J.* 2017;26:1111-1120.
24. Champain S, Benchikh K, Nogier A, Mazel C, Guise JD, Skalli W. Validation of new clinical quantitative analysis software applicable in spine orthopaedic studies. *Eur Spine J.* 2006;15: 982-991.
25. Rillardon L, Levassor N, Guigui P, et al. Validation of a tool to measure pelvic and spinal parameters of sagittal balance [in French]. *Rev Chir Orthop Reparatrice Appar Mot.* 2003;89: 218-227.
26. Fehlings MG, Wilson JR, Kopjar B, et al. Efficacy and safety of surgical decompression in patients with cervical spondylotic myelopathy: results of the AOSpine North America prospective multi-center study. *J Bone Joint Surg Am.* 2013;95:1651-1658.
27. Fairbank JC, Couper J, Davies JB, O'Brien JP. The Oswestry low back pain disability questionnaire. *Physiotherapy.* 1980;66: 271-273.
28. Fairbank JC, Pynsent PB. The Oswestry Disability Index. *Spine (Phila Pa 1976).* 2000;25:2940-2952.
29. Tetreault L, Nouri A, Kopjar B, Cote P, Fehlings MG. The minimum clinically important difference of the modified Japanese Orthopaedic Association Scale in patients with degenerative cervical myelopathy. *Spine (Phila Pa 1976).* 2015;40:1653-1659.
30. Auffinger B, Lam S, Shen J, Roitberg BZ. Measuring surgical outcomes in subaxial degenerative cervical spine disease patients: minimum clinically important difference as a tool for determining meaningful clinical improvement. *Neurosurgery.* 2014;74: 206-213.
31. Carreon LY, Glassman SD, Campbell MJ, Anderson PA. Neck Disability Index, short form-36 physical component summary, and pain scales for neck and arm pain: the minimum clinically important difference and substantial clinical benefit after cervical spine fusion. *Spine J.* 2010;10:469-474.
32. Vernon H, Mior S. The Neck Disability Index: a study of reliability and validity. *J Manipulative Physiol Ther.* 1991;14: 409-415.
33. Benzel EC, Lancon J, Kesterson L, Hadden T. Cervical laminectomy and dentate ligament section for cervical spondylotic myelopathy. *J Spinal Disord.* 1991;4:286-295.
34. Smith JS, Shaffrey CI, Glassman SD, et al. Risk-benefit assessment of surgery for adult scoliosis: an analysis based on patient age. *Spine (Phila Pa 1976).* 2011;36:817-824.

35. Bridwell KH, Glassman S, Horton W, et al. Does treatment (non-operative and operative) improve the two-year quality of life in patients with adult symptomatic lumbar scoliosis: a prospective multicenter evidence-based medicine study. *Spine (Phila Pa 1976)*. 2009;34:2171-2178.
36. Bridwell KH, Baldus C, Berven S, et al. Changes in radiographic and clinical outcomes with primary treatment adult spinal deformity surgeries from two years to three- to five-years follow-up. *Spine (Phila Pa 1976)*. 2010;35:1849-1854.
37. Glassman SD, Berven S, Bridwell K, Horton W, Dimar JR. Correlation of radiographic parameters and clinical symptoms in adult scoliosis. *Spine (Phila Pa 1976)*. 2005;30:682-688.
38. Schwab FJ, Blondel B, Bess S, et al. Radiographical spinopelvic parameters and disability in the setting of adult spinal deformity: a prospective multicenter analysis. *Spine (Phila Pa 1976)*. 2013;38:E803-E812.
39. Klineberg E, Schwab F, Ames C, et al. Acute reciprocal changes distant from the site of spinal osteotomies affect global postoperative alignment. *Adv Orthop*. 2011;2011:415946.
40. Smith JS, Shaffrey CI, Lafage V, et al. Spontaneous improvement of cervical alignment after correction of global sagittal balance following pedicle subtraction osteotomy. *J Neurosurg Spine*. 2012;17:300-307.