

# UCLA

## UCLA Previously Published Works

**Title**

Is the Oral Microbiome Important in HIV-Associated Inflammation?

**Permalink**

<https://escholarship.org/uc/item/5sx3x17s>

**Journal**

mSphere, 5(1)

**ISSN**

1556-6811

**Author**

Fulcher, Jennifer A

**Publication Date**

2020-02-26

**DOI**

10.1128/msphere.00034-20

Peer reviewed



# Is the Oral Microbiome Important in HIV-Associated Inflammation?

Jennifer A. Fulcher<sup>a,b</sup>

<sup>a</sup>Division of Infectious Diseases, Department of Medicine, David Geffen School of Medicine at UCLA, Los Angeles, California, USA

<sup>b</sup>VA Greater Los Angeles Healthcare System, Los Angeles, California, USA

**ABSTRACT** Alterations in the gut microbiome during HIV infection have been implicated in chronic inflammation, but the role of the oral microbiome in this process is less clear. The article by M. K. Annavajhala, S. D. Khan, S. B. Sullivan, J. Shah, et al. (mSphere 5:e00798-19, 2020, <https://doi.org/10.1128/mSphere.00798-19>) investigated the relationship between oral and gut microbiome diversity and immune activation in patients with HIV on antiretroviral therapy. In this study, oral microbiome diversity was inversely associated with inflammatory markers such as soluble CD14 (sCD14), but surprisingly similar associations were not seen with gut microbiome diversity. Oral microbiome diversity was also associated with periodontitis in these patients. This study highlights the importance of continuing multisite examinations in studying the gastrointestinal tract microbiome and also stimulates important directions for future research defining the role of the oral-gut axis in HIV-associated inflammation.

**KEYWORDS** human immunodeficiency virus, inflammation, oral microbiology

The gastrointestinal tract contains trillions of commensal microbes that are integral to immune development and function. The microbial composition, or microbiome, is shaped by the local microenvironment and varies greatly by anatomic site from the oral cavity to the large intestine. Most research has focused on the intestinal microbiome, but other sites such as the oral cavity have increasingly recognized roles in health and disease. We are now learning that the microbiome in the gastrointestinal tract is not as compartmentalized as once thought. Recently, Schmidt et al. showed that there is ongoing direct migration of bacteria from the mouth to the gut even in healthy individuals (1). What does this mean? In most individuals, there is no apparent clinical consequence of oral-to-gut bacterial migration, and this may be a necessary mechanism to shape the gut microbiome. However, some oral bacteria appear to cause gut inflammation (2–4), and multiple clinical studies have associated the presence of “oral” bacteria in the gut with inflammatory diseases (5–7), including HIV (8, 9).

The gut microbiome has been implicated in driving systemic inflammation in chronic HIV-1 infection. The loss of the critical barrier regulating Th17 cells during acute HIV-1 infection increases microbial translocation and inflammation (10–12), and indicators of microbial translocation have been repeatedly associated with systemic inflammation in clinical studies (13–15). Gut bacteria may directly contribute to HIV-related inflammation via translocation (13) or direct interaction with mucosal immune cells (16). How the oral microbiome contributes to this cascade has received relatively little attention.

Annavajhala et al. (17) address this question by comparing oral and gut microbiome diversity among persons living with HIV on suppressive antiretroviral therapy. The authors examined the oral and gut microbiomes in 52 women and men with HIV, including longitudinal sampling in a portion of the cohort. Low CD4 nadir was associated with lower gut bacterial diversity, consistent with other studies (18), but this was

**Citation** Fulcher JA. 2020. Is the oral microbiome important in HIV-associated inflammation? mSphere 5:e00034-20. <https://doi.org/10.1128/mSphere.00034-20>.

**Copyright** © 2020 Fulcher. This is an open-access article distributed under the terms of the [Creative Commons Attribution 4.0 International license](https://creativecommons.org/licenses/by/4.0/).

Address correspondence to [jfulcher@mednet.ucla.edu](mailto:jfulcher@mednet.ucla.edu).

For the article discussed, see <https://doi.org/10.1128/mSphere.00798-19>.

*The views expressed in this article do not necessarily reflect the views of the journal or of ASM.*

**Published** 5 February 2020

not seen in the oral microbiome. When examining systemic markers of inflammation, soluble CD14 (sCD14) and interleukin 6 (IL-6) were inversely associated with saliva bacterial diversity, yet no associations were found with gut bacterial diversity. This is in contrast to many other studies which have linked the gut microbiome to systemic inflammation (13–15), though none of these prior studies simultaneously examined the oral compartment. Whether this suggests a more important role of the oral microbiome in systemic inflammation or is simply a product of differences in study design, populations, and sample size is not known. It is also possible that the oral microbiome plays an important role only in certain stages of disease, and these may not coincide with the gut microbiome.

Though compelling, these findings should be considered in the context of a few important limitations. Most notably, this study is limited to people living with HIV, so it is not known how these findings compare to those without HIV. In other settings, the oral microbiome has been associated with periodontitis and systemic inflammation (19); how might HIV augment this relationship? Another important consideration is the contribution of periodontitis to these findings. In this study, saliva bacterial diversity was also associated with moderate to severe periodontitis, and specific taxa including *Prevotella melaninogenica* and *Rothia mucilaginosa* were associated with periodontitis. Interestingly, these same species were associated with HIV disease in a separate study of HIV-infected and uninfected women (20). Finally, though the study by Annavajhala et al. (17) is among the larger studies published so far investigating the role of the oral microbiome in HIV-associated inflammation, it is still limited in size. This highlights the critical need for larger, longitudinal studies with multiple site sampling for comparative analyses of microbial composition and evolution in the context of HIV and inflammation.

Importantly, the study by Annavajhala et al. (17) is one of the few published studies that has included women. No differences in bacterial diversity were seen by sex at either site in this study, but the small sample size and uneven distribution of men and women may have limited the ability to detect differences. Moving forward, a focused effort on better understanding of the role of sex on both the oral and gut microbiome in HIV will be critical.

This study adds further evidence that the effects of the microbiome on mucosal and systemic immunity are not limited to the intestinal tract. The authors challenge the notion that alterations in the gut microbiome are most influential in HIV-associated inflammation and provide data suggesting that oral microbiome diversity may play an equal or even greater role in systemic inflammation. Given the evidence that the oral and gut microbiomes interact, it is unlikely that either site functions in isolation. Defining the dynamics between different microbiome compartments, and how these populations interact, will better our understanding of how commensal microbiota influence disease. What is the timing of oral microbiome changes related to gut dysbiosis? What characteristics of the oral bacterial species lead to inflammation when migrated to a distal site? It is likely that bacteria evolve and adapt to their specific niche, and when placed in another environment cause disruption of the homeostatic ecosystem which has existed. Understanding what is different about these translocated species may give insight into the immune environment perturbations that promote dysbiosis, and potentially provide therapeutic targets to dampen this process.

## ACKNOWLEDGMENTS

This work received no specific grant funding from any public, commercial, or other agency.

I declare that I have no conflicts of interest.

## REFERENCES

1. Schmidt TS, Hayward MR, Coelho LP, Li SS, Costea PI, Voigt AY, Wirbel J, Maistrenko OM, Alves RJ, Bergsten E, de Beaufort C, Sobhani I, Heintz-Buschart A, Sunagawa S, Zeller G, Wilmes P, Bork P. 2019. Extensive transmission of microbes along the gastrointestinal tract. *Elife* 8:e42693. <https://doi.org/10.7554/eLife.42693>.
2. Atarashi K, Suda W, Luo C, Kawaguchi T, Motoo I, Narushima S, Kiguchi Y, Yasuma K, Watanabe E, Tanoue T, Thaiss CA, Sato M, Toyooka K, Said HS, Yamagami H, Rice SA, Gevers D, Johnson RC, Segre JA, Chen K, Kolls JK, Elinav E, Morita H, Xavier RJ, Hattori M, Honda K. 2017. Ectopic colonization of oral bacteria in the intestine drives TH1 cell induction

- and inflammation. *Science* 358:359–365. <https://doi.org/10.1126/science.aan4526>.
3. Arimatsu K, Yamada H, Miyazawa H, Minagawa T, Nakajima M, Ryder MI, Gotoh K, Motooka D, Nakamura S, Iida T, Yamazaki K. 2014. Oral pathobiont induces systemic inflammation and metabolic changes associated with alteration of gut microbiota. *Sci Rep* 4:4828. <https://doi.org/10.1038/srep04828>.
  4. Kato T, Yamazaki K, Nakajima M, Date Y, Kikuchi J, Hase K, Ohno H, Yamazaki K. 2018. Oral administration of *Porphyromonas gingivalis* alters the gut microbiome and serum metabolome. *mSphere* 3:e00460-18. <https://doi.org/10.1128/mSphere.00460-18>.
  5. Gevers D, Kugathasan S, Denson LA, Vazquez-Baeza Y, Van Treuren W, Ren B, Schwager E, Knights D, Song SJ, Yassour M, Morgan XC, Kostic AD, Luo C, Gonzalez A, McDonald D, Haberman Y, Walters T, Baker S, Rosh J, Stephens M, Heyman M, Markowitz J, Baldassano R, Griffiths A, Sylvester F, Mack D, Kim S, Crandall W, Hyams J, Huttenhower C, Knight R, Xavier RJ. 2014. The treatment-naive microbiome in new-onset Crohn's disease. *Cell Host Microbe* 15:382–392. <https://doi.org/10.1016/j.chom.2014.02.005>.
  6. Zhang X, Zhang D, Jia H, Feng Q, Wang D, Liang D, Wu X, Li J, Tang L, Li Y, Lan Z, Chen B, Li Y, Zhong H, Xie H, Jie Z, Chen W, Tang S, Xu X, Wang X, Cai X, Liu S, Xia Y, Li J, Qiao X, Al-Aama JY, Chen H, Wang L, Wu QJ, Zhang F, Zheng W, Li Y, Zhang M, Luo G, Xue W, Xiao L, Li J, Chen W, Xu X, Yin Y, Yang H, Wang J, Kristiansen K, Liu L, Li T, Huang Q, Li Y, Wang J. 2015. The oral and gut microbiomes are perturbed in rheumatoid arthritis and partly normalized after treatment. *Nat Med* 21:895–905. <https://doi.org/10.1038/nm.3914>.
  7. Qin N, Yang F, Li A, Prifti E, Chen Y, Shao L, Guo J, Le Chatelier E, Yao J, Wu L, Zhou J, Ni S, Liu L, Pons N, Batto JM, Kennedy SP, Leonard P, Yuan C, Ding W, Chen Y, Hu X, Zheng B, Qian G, Xu W, Ehrlich SD, Zheng S, Li L. 2014. Alterations of the human gut microbiome in liver cirrhosis. *Nature* 513:59–64. <https://doi.org/10.1038/nature13568>.
  8. Lozupone CA, Li M, Campbell TB, Flores SC, Linderman D, Gebert MJ, Knight R, Fontenot AP, Palmer BE. 2013. Alterations in the gut microbiota associated with HIV-1 infection. *Cell Host Microbe* 14:329–339. <https://doi.org/10.1016/j.chom.2013.08.006>.
  9. Vujkovic-Cvijin I, Dunham RM, Iwai S, Maher MC, Albright RG, Broadhurst MJ, Hernandez RD, Lederman MM, Huang Y, Somsouk M, Deeks SG, Hunt PW, Lynch SV, McCune JM. 2013. Dysbiosis of the gut microbiota is associated with HIV disease progression and tryptophan catabolism. *Sci Transl Med* 5:193ra91. <https://doi.org/10.1126/scitranslmed.3006438>.
  10. Chege D, Sheth PM, Kain T, Kim CJ, Kovacs C, Loutfy M, Halpenny R, Kandel G, Chun TW, Ostrowski M, Kaul R, Toronto Mucosal Immunology Group. 2011. Sigmoid Th17 populations, the HIV latent reservoir, and microbial translocation in men on long-term antiretroviral therapy. *AIDS* 25:741–749. <https://doi.org/10.1097/QAD.0b013e328344cefb>.
  11. Kim CJ, McKinnon LR, Kovacs C, Kandel G, Huibner S, Chege D, Shahabi K, Benko E, Loutfy M, Ostrowski M, Kaul R. 2013. Mucosal Th17 cell function is altered during HIV infection and is an independent predictor of systemic immune activation. *J Immunol* 191:2164–2173. <https://doi.org/10.4049/jimmunol.1300829>.
  12. Klatt NR, Estes JD, Sun X, Ortiz AM, Barber JS, Harris LD, Cervasi B, Yokomizo LK, Pan L, Vinton CL, Tabb B, Canary LA, Dang Q, Hirsch VM, Alter G, Belkaid Y, Lifson JD, Silvestri G, Milner JD, Paiardini M, Haddad EK, Brenchley JM. 2012. Loss of mucosal CD103+ DCs and IL-17+ and IL-22+ lymphocytes is associated with mucosal damage in SIV infection. *Mucosal Immunol* 5:646–657. <https://doi.org/10.1038/mi.2012.38>.
  13. Brenchley JM, Price DA, Schacker TW, Asher TE, Silvestri G, Rao S, Kazza Z, Bornstein E, Lambotte O, Altmann D, Blazar BR, Rodriguez B, Teixeira-Johnson L, Landay A, Martin JN, Hecht FM, Picker LJ, Lederman MM, Deeks SG, Douek DC. 2006. Microbial translocation is a cause of systemic immune activation in chronic HIV infection. *Nat Med* 12:1365–1371. <https://doi.org/10.1038/nm1511>.
  14. Dillon SM, Lee EJ, Kotter CV, Austin GL, Dong Z, Hecht DK, Gianella S, Siewe B, Smith DM, Landay AL, Robertson CE, Frank DN, Wilson CC. 2014. An altered intestinal mucosal microbiome in HIV-1 infection is associated with mucosal and systemic immune activation and endotoxemia. *Mucosal Immunol* 7:983–994. <https://doi.org/10.1038/mi.2013.116>.
  15. Mutlu EA, Keshavarzian A, Losurdo J, Swanson G, Siewe B, Forsyth C, French A, Demarais P, Sun Y, Koenig L, Cox S, Engen P, Chakradeo P, Abbasi R, Gorenz A, Burns C, Landay A. 2014. A compositional look at the human gastrointestinal microbiome and immune activation parameters in HIV infected subjects. *PLoS Pathog* 10:e1003829. <https://doi.org/10.1371/journal.ppat.1003829>.
  16. Dillon SM, Lee EJ, Kotter CV, Austin GL, Gianella S, Siewe B, Smith DM, Landay AL, McManus MC, Robertson CE, Frank DN, McCarter MD, Wilson CC. 2016. Gut dendritic cell activation links an altered colonic microbiome to mucosal and systemic T-cell activation in untreated HIV-1 infection. *Mucosal Immunol* 9:24–37. <https://doi.org/10.1038/mi.2015.33>.
  17. Annavajhala MK, Khan SD, Sullivan SB, Shah J, Pass L, Kister K, Kunen H, Chiang V, Monnot GC, Ricupero CL, Mazur RA, Gordon P, de Jong A, Wadhwa S, Yin MT, Demmer RT, Uhlemann A-C. 2020. Oral and gut microbial diversity and immune regulation in patients with HIV on antiretroviral therapy. *mSphere* 5:e00798-19. <https://doi.org/10.1128/mSphere.00798-19>.
  18. Li SX, Armstrong A, Neff CP, Shaffer M, Lozupone CA, Palmer BE. 2016. Complexities of gut microbiome dysbiosis in the context of HIV infection and antiretroviral therapy. *Clin Pharmacol Ther* 99:600–611. <https://doi.org/10.1002/cpt.363>.
  19. Yamashita Y, Takeshita T. 2017. The oral microbiome and human health. *J Oral Sci* 59:201–206. <https://doi.org/10.2334/josnurd.16-0856>.
  20. Lewy T, Hong BY, Weiser B, Burger H, Tremain A, Weinstock G, Anastos K, George MD. 2019. Oral microbiome in HIV-infected women: shifts in the abundance of pathogenic and beneficial bacteria are associated with aging, HIV load, CD4 count, and antiretroviral therapy. *AIDS Res Hum Retroviruses* 35:276–286. <https://doi.org/10.1089/AID.2017.0200>.