UC San Diego

UC San Diego Previously Published Works

Title

Documentation of ENDS Use in the Veterans Affairs Electronic Health Record (2008-2014)

Permalink

https://escholarship.org/uc/item/5tz6m9n0

Journal

American Journal of Preventive Medicine, 56(3)

ISSN

0749-3797

Authors

Conway, Mike Mowery, Danielle L South, Brett R et al.

Publication Date

2019-03-01

DOI

10.1016/j.amepre.2018.10.019

Peer reviewed

Documentation of ENDS Use in the Veterans Affairs Electronic Health Record (2008-2014)

Mike Conway, PhD,¹ Danielle L. Mowery, PhD,^{1,2} Brett R. South, PhD,^{1,2} Gregory J. Stoddard, MPH,³ Wendy W. Chapman, PhD,^{1,2} Olga V. Patterson, PhD,^{1,3} Shu-Hong Zhu, PhD⁴

From the ¹Department of Biomedical Informatics, University of Utah, Salt Lake City, Utah; ²Informatics, Decision-Enhancement, and Analytic Sciences Center (IDEAS 2.0), Veterans Affairs Salt Lake City Health Care System, Salt Lake City, Utah; ³Department of Internal Medicine, University of Utah, Salt Lake City, Utah; and ⁴Department of Family Medicine and Public Health, University of California San Diego, San Diego, California

Address correspondence to: Mike Conway, PhD, Department of Biomedical Informatics, 421 Wakara Way, Salt Lake City UT 84108. E-mail: mike.conway@utah.edu.

INTRODUCTION

The Department of Veterans Affairs (VA) health system is the largest health system in the U.S., and the only system with a nationwide presence, with more than 1,700 healthcare facilities serving 8.7 million veterans per year.¹ Smoking rates among U.S. Veterans are substantially higher than in the general population, with 20.1% of Veterans smoking, compared with 15.5% of the general U.S. population.² The VA health system and its associated electronic health record (EHR) provides a unique opportunity in the U.S. context for investigating ENDS documentation practices using clinical data. However, the VA EHR, like other widely used EHR, lacks a structured data field for documenting ENDS use. Typically, ENDS documentation, if it exists at all, is recorded in narrative clinical text, and hence requires the use of text-mining methods to extract relevant data. Apart from two studies focused on ENDS documentation in regional health systems in the U.S., 3,4 little is currently known regarding the frequency with which clinicians document ENDS use.

This letter reports on work utilizing EHR data derived from a cohort of 20,000 documented smokers in the VA system with the goal of investigating the extent to which ENDS documentation frequency has changed between 2008 and 2014.

METHODS

The VA Clinical Corporate Data Warehouse was queried to create a nationally representative, randomly sampled cohort of 20,000 patients (8,806,352 clinical notes in total) from Veterans who were documented smokers consistently throughout the years 2008 to 2014. Patients with more than one smoking status class documented in any year (e.g., former smoker, current smoker) were excluded as the primary goal of this study was to understand ENDS documentation changes over time among current smokers. Patient ages in 2008 ranged from 17 to 102 years (mean: 65), with women constituting 5.2% of the sample. To identify ENDS-relevant clinical notes the authors used a list of 14 high precision (i.e., high positive predictive value) keywords (e.g., e cig, e-cig, ecig; Figure 1 provides a complete list) identified and verified in pilot work⁵ to perform a keyword search within the patient clinical notes. High precision keywords were used due to ambiguity associated with ENDS-related keywords, such as "vaporizer," which in this data set refers to nicotine use as opposed to marijuana or medical nebulizer use, <20% of the time. Year-on-year (2008–2014) change in frequency of ENDS mentions at the patient level were then determined. The percentage of patients with an ENDS mention was used as the outcome variable (n=7, for the 7 years), with year as the predictor variable, and modeled using a generalized linear regression model with log link (log transformation of percentage), family Gaussian, and robust SE to account for time-series autocorrelation.

RESULTS

A statistically significant increase (p<0.001) in the number of patients with at least one documented ENDS mention was observed (0 patients in 2008, 174 patients in 2014, i.e., increasing from 0% to 0.87% of patients). In total, 631 ENDS mentions from 291 unique patients were observed, with the greatest year-on-year increase in patients occurring between 2013 and 2014 (Figure 1). Interestingly, the percentage of patients with an ENDS mention increased exponentially, doubling each year from the previous year (1.95-fold increase per year, 95% CI=1.76, 2.15, p<0.001).

DISCUSSION

Analysis of the VA EHR dataset revealed that — consistent with results reported using EHR data derived from a large regional health system³ — the number of VA patients with ENDS mentions increased in frequency significantly over time, with 0 patients in 2008 and 174 patients in 2014. This result indicates that clinicians are increasingly likely to document their ENDS-related discussions with patients who smoke. However, there remains a striking discrepancy between the frequency of patients with ENDS use documentation in the VA EHR (0.87% of patients in 2014) and the much greater prevalence—estimated to be 11.5% (10.4% in smokers aged 45-64 years, and 8.3% in smokers aged 65 years and older)—of ENDS use among U.S. smokers generally, suggesting that ENDS use is currently massively under-documented by clinicians in the VA system, and that there is an acute

need for the inclusion of an ENDS-related structured data field in the VA EHR.

ACKNOWLEDGMENTS

The views expressed in this article are those of the authors and do not necessarily reflect the position or policy of the Department of Veterans

Affairs or the U.S. government. This study was approved by the University of Utah's IRB (IRB 00088382).

MC conceptualized the study and wrote the first draft. DM conducted the analysis and contributed substantially towards writing the first draft. BS performed the initial data pull and contributed substantially towards writing the first draft. WC and OP provided expertise on working with Veterans Affairs clinical notes and contributed to the first draft. GS provided statistical support. SZ provided expertise in the broad area of ENDS use and its relationship to tobacco use. All authors contributed to the writing and approved the final version of the manuscript.

No financial disclosures were reported by the authors of this paper.

REFERENCES


- U.S. Department of Veterans Affairs. Tobacco use in the VA infographic. <u>www.publichealth.va.gov/smoking/professionals/tobacco-use-infographic.asp</u>. Published 2016. Accessed July 1, 2018.
- CDC. Current cigarette smoking among adults in the United States: fast facts and fact sheet.
 www.cdc.gov/tobacco/data_statistics/fact_sheets/adult_data/cig_smoking /index.htm. Published 2018. Accessed July 1, 2018.
- Young-Wolff KC, Klebaner D, Folck B, et al. Do you vape? Leveraging electronic health records to assess clinician documentation of electronic nicotine delivery system use among adolescents and adults. *Prev Med*. 2017;105:32–36. https://doi.org/10.1016/j.ypmed.2017.08.009.
- Winden TJ, Chen ES, Wang Y, et al. Towards the standardized documentation of e-cigarette use in the electronic health record for population health surveillance and research. AMIA Jt Summits Transl Sci Proc. 2015;2015:199–203.
- 5. Mowery DL, South BR, Patterson OV, Zhu SH, Conway M. Investigating the documentation of electronic cigarette use in the Veteran Affairs electronic health record: a pilot study. In: *Proceedings of the Workshop on Current Trends in Biomedical Natural Language Processing (BioNLP 2017)*. Vancouver; 2017. https://doi.org/10.18653/v1/W17-2335.
- 6. Zhu SH, Zhuang YL, Wong S, Cummins SE, Tedeschi GJ. E-cigarette use

and associated changes in population smoking cessation: evidence from U.S. current population surveys. *BMJ*. 2017;358:j3262.

https://doi.org/10.1136/bmj.j3262.

LIST OF FIGURES

Figure 1. Increase in Frequency of ENDS mentions by year in a cohort of 20,000 VA smokers.



electronic_cig
electrical_cig
liquid_tobacco ecig

electric_tobacco
liquid_nicotine
electric_cig
electronic_tobacco
ecigarette
electronic_nicotine