UC Irvine

UC Irvine Previously Published Works

Title

Using mitochondrial and nuclear markers to evaluate the degree of genetic cohesion among Echinococcus populations.

Permalink

<https://escholarship.org/uc/item/5v36t73b>

Journal

Experimental parasitology, 119(4)

ISSN

0014-4894

Authors

Badaraco, JL
Ayala, FJ
Bart, J-M
et al.

Publication Date

2008-08-01

DOI

10.1016/j.exppara.2008.02.004

Copyright Information

This work is made available under the terms of a Creative Commons Attribution License, available at <https://creativecommons.org/licenses/by/4.0/>

Peer reviewed

Using mitochondrial and nuclear markers to evaluate the degree of genetic cohesion among *Echinococcus* populations

J.L. Badaraco^a, F.J. Ayala^b, J.-M. Bart^c, B. Gottstein^d, K.L. Haag^{a,c,*}

^a Programa de Pós Graduação em Genética e Biologia Molecular, Universidade Federal do Rio Grande do Sul, Porto Alegre, RS, CEP 91501-970, Brazil

^b Department of Ecology and Evolutionary Biology, University of California, Irvine, CA 92697, USA

^c Laboratoire de Parasitologie, Faculté de Sciences Médicales et Pharmaceutiques de Besançon, Université de Franche-Comté, Besançon, France

^d Institute of Parasitology, Faculty of Medicine and Veterinary Medicine, University of Berne, Berne CH-3001, Switzerland

^e Departamento de Genética, Instituto de Biociências, Universidade Federal do Rio Grande do Sul, Porto Alegre, 91501-970 RS, Brazil

Received 23 November 2007; received in revised form 7 February 2008; accepted 12 February 2008

Available online 19 February 2008

Abstract

Based on the distinctiveness of their mitochondrial haplotypes and other biological features, several recent publications have proposed that some *Echinococcus granulosus* strains should be regarded as separate species. However, the genetic cohesion of these species has not been extensively evaluated using nuclear markers. We assess the degree of polymorphism of the partial mitochondrial *cox1* (366 bp), the nuclear *mdh* (214 bp) and *EgAgB4* (281–283 bp) genes of *E. granulosus sensu lato* isolates collected from areas where different strains occur sympatrically. Five distinct mitochondrial haplotypes were determined by direct sequencing (G1, G2, G5, G6 and G7). The *mdh* genotypes were first screened by SSCP: three alleles were identified (Md1–Md3), which were further confirmed by nucleotide sequencing. For *EgAgB4*, which was analysed by direct sequencing the PCR products, two groups of sequences were found: *EgAgB4*-1 and *EgAgB4*-2. No haplotype-specific *mdh* or *EgAgB4* sequences occur. Nevertheless, alleles Md1 and Md2 and type 1 sequences of *EgAgB4* showed a higher frequency within the group of haplotypes G1–G2, while allele Md3 and *EgAgB4*-2 are most frequent in the G5–G7 cluster. By AMOVA it is shown that 79% of the total genetic variability is found among haplotype groups. These findings are compatible with two not mutually exclusive evolutionary hypotheses: (a) that haplotypes share an ancestral polymorphism, or (b) that the reproductive isolation between parasites with distinct haplotypes is not complete, leading to gene introgression. The biological and epidemiologic consequences of our findings are discussed.

© 2008 Elsevier Inc. All rights reserved.

Index Descriptors and Abbreviations: *Echinococcus*; Molecular epidemiology; Introgression; SSCP

1. Introduction

After a recent taxonomic review, the genus *Echinococcus* Rudolphi, 1801 (Cestoda; Taeniidae) now includes eight recognized species: *Echinococcus granulosus*, *Echinococcus equinus*, *Echinococcus canadensis* and *Echinococcus ortleppi*, which cause cystic echinococcosis (referred here as *E.*

granulosus sensu lato, for convenience); *Echinococcus multilocularis*, causing alveolar echinococcosis; the Neotropical *Echinococcus oligarthrus* and *Echinococcus vogeli* causing polycystic echinococcosis; and *Echinococcus shiquicus*, which was recently identified in the Tibet mountains, and seems more closely related to *E. multilocularis*. The parasite has a complex life cycle, requiring a carnivore definitive host and an herbivore intermediate host, between which transmission occurs through predator–prey interactions. During the larval stage *Echinococcus* spp. proliferate asexually and develop numerous protoscolices, each with a potential to mature into an adult worm. Worms are hermaphrodites, and reproduce mainly by selfing in the

* Corresponding author. Address: Departamento de Genética, Universidade Federal do Rio Grande do Sul, Avenue Bento Gonçalves, 9500 Predio 43312, Caixa Postal 15053-Porto Alegre, RS, CEP 91501-970, Brazil. Fax: +55 51 3308 9823.

E-mail address: karen.haag@ufrgs.br (K.L. Haag).

carnivore intestine, but a low rate of outcrossing has been inferred from genetic polymorphisms (Thompson and Lymbery, 1996; Lymbery et al., 1997; Haag et al., 1999). Nevertheless, due to the clonal origin of the adult population derived from a single metacystode, cross-fertilization occurs most frequently between identical genotypes (geitonogamy), which is equivalent to selfing with regard to the population genetic consequences.

A remarkable feature of *E. granulosus sensu lato* is a large intra-specific diversity in mitochondrial haplotypes and some adult morphologic traits. Mitochondrial genetic variants differing in epidemiologically relevant characters used to be informally designated “strains” (Bowles et al., 1992, 1995). Most of the evidence used to evaluate if strain designations correlate with morphology comes from larval and adult rostellar hook morphometry. For example, the number and length of hooks can be used as diagnostic features to distinguish the camel and the sheep strain (Ahmadi, 2004), but not the sheep and the Tasmanian sheep strain (Hobbs et al., 1990). Now the taxonomic status of some of these strains has been debated. The horse strain (haplotype G4) was the first to be proposed as a new species, *E. equinus*, due to its genetic and developmental distinctiveness (Thompson et al., 1995). Later, the cattle strain (haplotype G5) was also indicated as a separate species (Thompson and McManus, 2002): *E. ortleppi*. Finally, based on phylogenetic analyses of 11 complete mitochondrial genome sequences, it was proposed by Nakao et al. (2007) that some major clades within the *Echinococcus* phylogeny should be considered as distinct species. Combining their original results with other published mitochondrial sequences, and scattered observations on morphology, host specificities and biogeography, the authors suggested the following reclassification: the clusters G1–G3 should be considered *E. granulosus sensu stricto*, G4 *E. equinus*, G5 *E. ortleppi* and G6–G10 (a cluster including the camel, pig and cervid strains) *E. canadensis*.

It must be kept in mind, nevertheless, that the G5–G8 strains belong to a single cluster in the inferred tree, and that the genetic distances between mitochondrial genomes of this cluster average about 5%. Furthermore, although similar amounts of mtDNA divergence correlate with species circumscriptions in other animals, cautiousness is required in interpreting mitochondrial barcodes. For example, several African cichlid fish species, which radiated in the last 15–200,000 years, differ by less than 1% in mitochondrial genes (Meyer et al., 1990). Salamanders from the species *Ensatina eschscholtzii*, on the other hand, form a ring of subspecies whose members diverged more than 5 million years ago, with many subspecies having more than 5% mtDNA divergence from each other. Yet, because of population connectivity based on intensive ecological research, this complex is considered one species (Moritz et al., 1992).

Unfortunately, the genetic evidence behind the taxonomic debate is very limited. Population genetic variability has largely been neglected in *Echinococcus*.

Furthermore, the genetic studies have concentrated on the analysis of mitochondrial DNA, a non-recombining, fast evolving and maternally inherited genome in most animals. Little is known about the degree of polymorphism in nuclear loci, which would be valuable to test for reproductive isolation. In the present study, we verify the pattern and the amount of variation in a 214-bp fragment of the gene encoding malate dehydrogenase (*mdh*) and in a 281–283-bp segment of CDS encoding the fourth antigen B subunit (*EgAgB4*). Samples of *E. granulosus sensu lato* isolates were collected from geographic areas where strains occur sympatrically.

2. Materials and methods

2.1. Parasite samples and strain determination

Echinococcus granulosus hydatid cysts were obtained from different host species and four geographic areas where the distribution of strains is known to overlap (Table 1). Strains were determined with partial sequences (366 bp) of the mitochondrial cytochrome oxidase 1 gene (*cox1*) using the procedures described by Bowles et al. (1992). Sequencing was performed by cycle sequencing and migrated in an ABI 3730XL machine (Applied Biosystems).

2.2. Genotyping the *mdh* locus

We amplified a fragment of 214 bp from the gene encoding malate dehydrogenase (*mdh*—GenBank Accession No. L08894), which includes parts of the second and third exons and the complete second intron. The PCR primers used were 5'-CGCTCCTTCCATTTCCGAAAG (forward) and 5'-TTGGTGACAACGGCGTG

Table 1
Number of *E. granulosus* isolates from different strains, hosts and geographic locations included in the sample

Country	Total	Host	Haplotype	<i>n</i>
Argentina	33	Sheep	G1	7
			G1	14
		Human	G2	4
			G6	8
Algeria	28	Sheep	G1	10
		Human	G1	5
		Cattle	G1	7
		Dromedary	G1	3
			G6	3
Southern Brazil	171	Cattle	G1	115
			G5	55
			G7	1
Romania	27	Sheep	G1	6
		Human	G1	1
		Cattle	G1	12
		Pig	G7	8
Total	259			259

AGAC (reverse). The reactions contained approximately 30 ng of genomic DNA, 1 U Taq polymerase (Invitrogen), 1.5 mM MgCl₂, 10 mM dNTPs and 20 pmol of each primer in a total volume of 50 µl. The amplification was carried out using the following cycling conditions: 94 °C for 4 min, 40 cycles of 94 °C for 1 min, 55 °C for 1 min and 72 °C for 1 min and a final extension step of 72 °C for 10 min. The PCR fragments were denatured at 95 °C for 3 min and migrated in 12% polyacrylamide gels to separate the single stranded DNA according to its conformation (Single Strand Conformation Polymorphism—SSCP) using the GenePhor system (GE Healthcare). Gels were run in “buffer A” (pH 9.0—GE Healthcare) at 12 °C and constant voltage of 200 V for 1 h and 30 min. The gels were silver stained using standard protocols. Genotypes were attributed to the distinct SSCP patterns and confirmed by automated nucleotide sequencing. At least six independent PCR reactions corresponding to each SSCP pattern were sequenced.

2.3. *EgAgB4* sequence analyses

We designed primers to specifically amplify the *EgAgB4* gene, based on the GenBank Accession No. sequence DQ152008. The forward primer PB4F_ol1 (5'-GGAT GGAGTATAAGGAGCAG) anneals within the upstream flanking region, while the reverse primer AgBRev2 (5'-GACATATTTCTTCAACACTTCGTGAAC) anneals inside the second exon, generating a fragment of about 350 bp. The PCR reactions were similar as those described for *mdh*, except for the annealing temperatures, which followed a touchdown procedure, starting with 60 °C and decreasing until 50 °C during the 20 initial cycles. The amplicons were purified through columns (GE Healthcare) and sequenced directly.

2.4. Statistic analyses

Sequence trace quality was assessed with the SeqMan tool from the Lasergene software (version 7.0). The amount of sequence polymorphism in the *mdh* locus was estimated by the number of segregating sites (*s*) and nucleotide diversity (π) using the dnaSP software version 4.0 (Rozas et al., 2003). The Hardy–Weinberg genotype proportions and the population genetic structure were analyzed with Arlequin version 3.1 (Excoffier et al., 2005). For the genetic structure analyses, we used a hierarchical approach, first grouping the isolates by their mitochondrial haplotype (strain) and further subdividing the groups into populations according to the geographic location. This approach was used to decompose the genetic variation with AMOVA (Excoffier et al., 1992). Gene flow between strains in each geographic area was verified through the F_{ST} parameter (Reynolds et al., 1983), which can be used to estimate the number of migrants per generation (N_m) between populations (Slatkin and Maddison, 1989).

3. Results

3.1. Strain polymorphism at the *mdh* locus

Five distinct haplotypes are present in our sample, as determined by the *cox1* mitochondrial marker (G1, G2, G5, G6 and G7; Table 1). The sequence regions with traces of high quality never showed double peaks for *cox1*.

For *mdh*, SSCP discriminated three alleles segregating within our samples (Md1, Md2 and Md3; Fig. 1), and their distinctiveness was confirmed by nucleotide sequencing (GenBank Accession Nos. EF640368, EF640371 and EF640370, respectively). Alleles Md1 and Md2 differ by 3 nucleotide substitutions, while Md3 differs from Md1 and Md2 by 10 and 8 nucleotide substitutions, respectively. The *mdh* genotype and allele frequencies are displayed in Table 2. There are no exclusive alleles in the strains, but Md3 occurs at a higher frequency (at least 0.88) in the cattle, camel and pig strains (haplotypes G5, G6 and G7). Alleles Md1 and Md2, on the other hand, are present at low frequencies in these strains, but at a higher frequency in the sheep and Tasmanian sheep strains (G1 and G2). A single parasite of G1 in Brazil is heterozygous for Md2 and Md3, and two were homozygous for Md3.

Large heterozygote deficiencies were found for the G1 populations from Argentina ($p < 0.01$), Algeria ($p < 0.05$) and Brazil ($p < 0.01$), and for G5 in Brazil ($p < 0.01$). Ten, out of 214 nucleotide sites at the *mdh* locus are

Fig. 1. SSCP gel showing the patterns of the *mdh* genotypes identified in this study: (1) Md1/Md1; (2) Md1/Md2; (3) Md2/Md2; (4) Md2/Md3; (5) Md3/Md3. The bands corresponding to the renatured double-stranded amplicons, including the 214 bp insert plus both primers, are indicated by ds. It is verified that the heterozygous genotypes form heteroduplexes after renaturation.

Table 2
Genotype numbers and allele frequencies for the *mdh* locus

Country	Haplotype	Genotype (%)					Total	Allele frequency		
		Md1/Md1	Md1/Md2	Md2/Md2	Md2/Md3	Md3/Md3		Md1	Md2	Md3
Argentina	G1	1 (4.76)	2 (9.52)	17 (80.95)	—	1 (4.76)	21	0.1	0.86	0.04
	G2	—	1 (25.00)	3 (75.00)	—	—	4	0.12	0.88	—
	G6	—	1 (12.5)	—	—	7 (87.5)	8	0.06	0.06	0.88
	Total	1 (3.03)	4 (12.12)	20 (60.61)	—	8 (24.24)	33	0.09	0.67	0.24
Algeria	G1	5 (20.00)	9 (36.00)	10 (40.00)	—	1 (4.00)	25	0.38	0.58	0.04
	G6	—	—	—	—	3 (100.00)	3	—	—	1.00
	Total	5 (17.86)	9 (32.14)	10 (35.71)	—	4 (14.29)	28	0.52	0.34	0.14
Southern Brazil	G1	25 (21.74)	46 (40.00)	41 (35.65)	1 (0.87)	2 (1.74)	115	0.42	0.56	0.02
	G5	—	3 (5.50)	—	—	52 (94.50)	55	0.03	0.03	0.94
	G7	—	—	—	—	1 (100.00)	1	—	—	1.00
	Total	25 (14.62)	49 (28.65)	41 (23.98)	1 (0.58)	54 (31.58)	171	0.29	0.39	0.32
Romania	G1	2 (10.53)	6 (31.58)	11 (57.89)	—	—	19	0.26	0.74	—
	G7	—	—	—	1 (12.5)	7 (87.5)	8	—	0.06	0.94
	Total	2 (7.41)	6 (22.22)	11 (40.74)	1 (3.7)	7 (25.93)	27	0.19	0.53	0.28

polymorphic in the surveyed populations; overall, the nucleotide diversity within each strain is below 1% (Table 1, Supplementary data). Of the 10 polymorphic sites, 3 are located inside the second exon, 3 in the intron and 4 in the third exon (data not shown).

When the genetic diversity is decomposed in three hierarchical levels using AMOVA, 79.07% of the variation occurs among strains, 1.1% among populations from the same strain and 19.9% is present within populations. The pairwise F_{ST} estimates between populations of the G1–G2 cluster ranged from 0.79 to 0.90 ($p < 0.01$) when compared to the G5–G7 cluster (Table 3). Not significant and low F_{ST} values occur within each cluster, except for the G1 populations of Southern Brazil $\times$ Argentina ($F_{ST} = 0.13$) and Southern Brazil $\times$ Algeria ($F_{ST} = 0.03$), which showed significant genetic differentiation ($p < 0.01$ and 0.05, respectively). Conversely, the estimates of gene

flow are higher for populations of the same strain and for strains of the same cluster, but very low between strains of distinct clusters. Sympatric populations of different clusters seem to show a small amount of gene flow, ranging between 0.08 and 0.13 migrants per generation, as inferred from our results (Table 3).

3.2. Sequence variability for *EgAgB4*

Due to technical difficulties, only a fraction of our sample could be analyzed both for *mdh* and *EgAgB4* sequence diversity. Overall, 176 isolates were typed both for *mdh* and *EgAgB4* (Table 4). Furthermore, due to the high degree of nucleotide sequence variation in the *EgAgB4* upstream flanking region, our analyses focus only on the polymorphisms found within a subset of 281–283 bp after the start codon (see Appendix A, Supplementary data). The

Table 3
Pairwise F_{ST} (below the diagonal) and the estimated number of migrants per generation between populations (N_m)

			Argentina			Algeria		Brazil		Romania	
			G1	G2	G6	G1	G6	G1	G5	G1	G2
Argentina	G1	21	—	∞	0.13	4.32	0.08	3.30	0.07	15.32	0.09
	G2	4	−0.06	—	0.12	8.10	0.02	5.04	0.06	∞	0.06
	G6	8	0.8**	0.80**	—	0.18	∞	0.16	285.70	0.11	∞
Algeria	G1	25	0.10**	0.06	0.73*	—	0.13	∞	0.12	23.34	0.14
	G6	3	0.87**	0.95*	−0.03	0.79**	—	0.09	∞	0.06	∞
Southern Brazil	G1	115	0.13**	0.09	0.76*	−0.01	0.84**	—	0.11	13.46	0.13
	G5	55	0.87**	0.9**	0.00	0.81**	−0.06	0.82**	—	0.07	∞
Romania	G1	19	0.03	−0.02	0.81*	0.02	0.89**	0.03*	0.88**	—	0.08
	G7	8	0.85**	0.89**	−0.04	0.78**	−0.08	0.79**	−0.03	0.86**	—

The N_m value for sympatric strains is displayed in bold letters.

* $p < 0.05$.

** $p < 0.01$.

Table 4
Genotype numbers for the *mdh* locus and *EgAgB4* combined

Country	Haplotype	Genotype	<i>n</i>
Argentina	G1	Md1/Md1; EgAgB4-1	2
		Md1/Md2; EgAgB4-1	2
		Md2/Md2; EgAgB4-1	8
	G2	Md1/Md2; EgAgB4-1	1
		Md2/Md2; EgAgB4-1	2
	G6	Md1/Md2; EgAgB4-1*	1
	Total		16
Algeria	G1	Md1/Md1; EgAgB4-1	5
		Md1/Md2; EgAgB4-1	9
		Md2/Md2; EgAgB4-1	9
		Md3/Md3; EgAgB4-1*	1
	G6	Md3/Md3; EgAgB4-1*	1
		Md3/Md3; EgAgB4-2	1
	Total		26
Southern Brazil	G1	Md1/Md1; EgAgB4-1	19
		Md1/Md2; EgAgB4-1	35
		Md2/Md2; EgAgB4-1	29
		Md2/Md3; EgAgB4-1*	1
		Md3/Md3; EgAgB4-2*	1
	G5	Md1/Md2; EgAgB4-1*	1
		Md3/Md3; EgAgB4-1*	6
		Md3/Md3; EgAgB4-2	26
	G7	Md3/Md3; EgAgB4-2	1
	Total		119
Romania	G1	Md1/Md1; EgAgB4-1	2
		Md1/Md2; EgAgB4-1	3
		Md2/Md2; EgAgB4-1	9
	G7	Md3/Md3; EgAgB4-1*	1
	Total		15
Total			176

* Genotypes suggesting introgression.

sequences are separated in two highly divergent groups ($s = 35$, leading to a divergency of 12%), which we call *EgAgB4-1* (GenBank Accession No. DQ152008) and *EgAgB4-2* (GenBank Accession No. AY569358). Since the *E. granulosus* isolates from both strain clusters always show either a type 1 or a type 2 *EgAgB4* sequence pattern, they are separated accordingly (Appendix A, Supplementary data). However, several segregating sites are found for each group ($s = 3$ within *EgAgB4-1*; $s = 6$ within *EgAgB4-2*, data not shown), including sites with double peaks within regions of high quality traces, which we interpret as polymorphisms. Since we cannot assure that they belong to alleles from the same locus, we ignore these additional polymorphisms at the present stage of our analyses. The majority of isolates from the G1–G2 cluster harbor *EgAgB4-1* sequences (137 out of 138 isolates, or 99%), while 28 out of 38 isolates (74%) belonging to the G5–G7 cluster contain the *EgAgB4-2* pattern. However, as for *mdh*, several recombinant forms are found for this genetic marker as well (Table 4).

4. Discussion

We analyzed the genetic structure of *E. granulosus sensu lato* in four geographic areas where distinct strains occur in sympatry, and therefore could share the same definitive host, using one mitochondrial (*cox1*) and two nuclear markers (*mdh* and *EgAgB4*). The *cox1* sequences were used to identify strains, while the *mdh* genotypes allowed an estimate of gene flow among populations. We already have indications that the *EgAgB* genes are highly redundant in the *Echinococcus* genome (Haag et al., 2006), thus the polymorphism within *EgAgB4-1* and *EgAgB4-2* was ignored, and the marker was not included in the population structure analyses. However, these two groups of *EgAgB4* sequences are highly divergent, and the totality of the 176 isolates analysed by the three genetic markers show either a type 1 or a type 2 pattern. In Table 3, all genotypes showing a recombinant genotype taking into account the three markers (mitochondrial $\times$ nuclear or nuclear $\times$ nuclear markers) are indicated by an asterisk, and correspond to 7.4% ($n = 13$) of our sample.

It has been suggested already that the heterozygote deficiency in *E. granulosus* populations is due to the fact that worms usually develop in the dog intestine in an aggregate manner, increasing the chance of crossing between clones derived from a single hydatid cyst, which is equivalent to selfing (Lymbery et al., 1989). In the present study, *mdh* heterozygote deficiencies were found in populations with statistically representative samples ($n \geq 20$). This differs from our results in a previous study (Haag et al., 1999), which showed an excess of heterozygotes in the European population and equilibrium frequencies for Southern Brazil and Australia, for two polymorphic loci. Since the samples were considerably smaller and collected in different time periods, it is possible that this difference was due to sampling bias. Nevertheless, the previous study showed strong linkage disequilibrium for all populations, in agreement with the conclusion that selfing (or geitonogamy) is the prevalent mode of reproduction in *E. granulosus* populations.

The immediate consequence of such a reproductive mode is the genetic structuring of the population in separate groups, corresponding to the alternative homozygote genotypes and linkage groups (Hartl, 2000). Moreover, if inbreeding is associated with natural selection, such as the adaptation to a new host species, genetic changes in the population can occur rapidly, because the adaptive alleles will remain “arrested” in their linkage groups, increasing the differentiation among sub-populations. Rapid evolutionary changes are facilitated by the asexual amplification of parasite genotypes during the larval stage. This phenomenon was envisioned by Smyth and Smyth (1964), who proposed that “mutations” amplified during protoscolex formation could lead to the formation of new strains. Overall, it means that animal domestication by primitive human communities could have accelerated the *Echinococcus* evolutionary process, generating parasites adapted to their local livestock (Rausch, 1967).

With the beginning of a worldwide animal trade, some *E. granulosus* variants (strains) came in contact. The consequences of this contact, which represents the current situation in several regions such as Argentina (Kamenetzky et al., 2002), are largely unpredictable. In the present study, we show that *mdh* alleles are shared among some *E. granulosus* strains. It indicates that the history described by the mitochondrial genes is only partial, and it illustrates how misleading it can be to rely only on a single kind of marker for population genetic studies (Anderson, 2001). Two hypotheses could explain our findings: (a) that strains share an ancestral polymorphism, and (b) that the reproductive isolation between strains is not complete, leading to gene introgression. Although the two hypotheses are not mutually exclusive, we believe that some gene flow is occurring among strains. For this reason, we are now evaluating how the genetic diversity is distributed among strains in other nuclear loci (including those analysed in our previous study, Haag et al., 1999) and other stages of the parasite life cycle.

If a small amount of gene flow is occurring among strains, the population dynamics of *E. granulosus* in some regions can become very complex, with important epidemiologic consequences. Although the number of migrants per generation estimated with the *mdh* polymorphism is small, it suggests that adaptive alleles can spread from one strain to another, and be amplified in the population as envisioned by Smyth and Smyth (1964). For example, it is thought that the camel strain is less infective to humans (Lymbery and Thompson, 1988), but in Argentina it was found in 4 out of 9 human patients (Rosenzvit et al., 1999), and in 3 out of 33 human patients in Iran (Harandi et al., 2002). McManus and Thompson (2003) speculated that this could have been the result of a mutation in the G6 haplotype, but it could also be explained by introgression of unknown “human infectivity” genes from another strain.

Another implication of our findings concerns the *Echinococcus* taxonomic debate. Evolutionary theory predicts that the speciation process is accomplished if the differentiated populations, or the putative species, maintain their genetic identity when in geographic contact, due to reproductive isolation mechanisms (Mayr, 1942). Accordingly, it was proposed that a genetic yardstick should be applied to define *Echinococcus* species, based on the reciprocal monophyly and the genetic cohesion of the divergent clades on a phylogeny (Lymbery, 1992). The *mdh* locus does not differentiate between the cattle, camel and pig strain populations (haplotypes G5–G7), since their F_{ST} values are not statistically significant. However, our previous study (Haag et al., 1999) showed that in 3 out of 5 nuclear loci the pig and cattle strains segregate distinct alleles. For this reason, we suggest that an analysis of larger number of nuclear loci should be taken into account for evaluating the degree of their genetic cohesion, and, consequently, identifying good *Echinococcus* species.

Acknowledgments

This work was supported by CNPq (Grants 471214/2004-3 and 450386/2007-4), and was part of the M.Sc. dissertation of Jeferson Badaraco obtained with a CNPq fellowship. We are indebted to one anonymous referee for his contribution.

Appendix A. Supplementary data

Supplementary data associated with this article can be found, in the online version, at doi:10.1016/j.exppara.2008.02.004.

References

- Ahmadi, N.A., 2004. Using morphometry of the larval rostellar hooks to distinguish Iranian strains of *Echinococcus granulosus*. *Annals of Tropical Medicine and Parasitology* 98, 211–220.
- Anderson, T.J., 2001. The dangers of using single locus markers in parasite epidemiology: *Ascaris* as a case study. *Trends in Parasitology* 17, 183–188.
- Bowles, J., Blair, D., McManus, D.P., 1992. Genetic variants within the genus *Echinococcus* identified by mitochondrial DNA sequencing. *Molecular and Biochemical Parasitology* 54, 165–173.
- Bowles, J., Blair, D., McManus, D.P., 1995. A molecular phylogeny of the genus *Echinococcus*. *Parasitology* 110, 317–328.
- Excoffier, L., Smouse, P.E., Quattro, J.M., 1992. Analysis of molecular variance inferred from metric distances among DNA haplotypes: application to human mitochondrial DNA restriction data. *Genetics* 131, 479–491.
- Excoffier, L., Laval, G., Schneider, S., 2005. Arlequin ver. 3.0: an integrated software package for population genetics data analysis. *Evolutionary Bioinformatics Online* 1, 47–50.
- Haag, K.L., Araujo, A.M., Gottstein, B., Siles-Lucas, M., Thompson, R.C., Zaha, A., 1999. Breeding systems in *Echinococcus granulosus* (Cestoda: Taeniidae): selfing or outcrossing? *Parasitology* 118, 63–71.
- Haag, K.L., Gottstein, B., Muller, N., Schnorr, A., Ayala, F.J., 2006. Redundancy and recombination in the *Echinococcus* AgB multigene family: is there any similarity with protozoan contingency genes? *Parasitology* 133, 411–419.
- Harandi, M.F., Hobbs, R.P., Adams, P.J., Mobedi, I., Morgan-Ryan, U.M., Thompson, R.C., 2002. Molecular and morphological characterization of *Echinococcus granulosus* of human and animal origin in Iran. *Parasitology* 125, 953–962.
- Hartl, D.L., 2000. *A Primer of Population Genetics*. Sinauer, Sunderland.
- Hobbs, R.P., Lymbery, A.J., Thompson, R.C., 1990. Rostellar hook morphology of *Echinococcus granulosus* (Batsch, 1786) from natural and experimental Australian hosts, and its implications for strain recognition. *Parasitology* 101, 273–281.
- Kamenetzky, L., Gutierrez, A.M., Canova, S.G., Haag, K.L., Guarnera, E.A., Parra, A., Garcia, G.E., Rosenzvit, M.C., 2002. Several strains of *Echinococcus granulosus* infect livestock and humans in Argentina. *Infection Genetics and Evolution* 2, 129–136.
- Lymbery, A.J., 1992. Interbreeding, monophyly and the genetic yardstick: species concepts in parasites. *Parasitology Today* 8, 208–211.
- Lymbery, A.J., Constantine, C.C., Thompson, R.C.A., 1997. Self-fertilization without genomic or population structuring in a parasitic tapeworm. *Evolution* 51, 289–294.
- Lymbery, A.J., Hobbs, R.P., Thompson, R.C., 1989. The dispersion of *Echinococcus granulosus* in the intestine of dogs. *The Journal of Parasitology* 75, 562–570.
- Lymbery, A.J., Thompson, R.C., 1988. Electrophoretic analysis of genetic variation in *Echinococcus granulosus* from domestic hosts in Australia. *International Journal of Parasitology* 18, 803–811.

- Mayr, E., 1942. *Systematics and the Origin of Species*. Columbia University Press, New York.
- Meyer, A., Kocher, T.D., Basasibwaki, P., Wilson, A.C., 1990. Monophyletic origin of Lake Victoria cichlid fishes suggested by mitochondrial DNA sequences. *Nature* 347, 550–553.
- McManus, D.P., Thompson, R.C., 2003. Molecular epidemiology of cystic echinococcosis. *Parasitology* 127 (Suppl.), S37–S51.
- Moritz, C.C., Schneider, C.J., Wake, D.B., 1992. Evolutionary relationships within the *Ensatina eschscholtzii* complex confirm the ring species interpretation. *Systematic Biology* 41, 273–291.
- Nakao, M., McManus, D.P., Schantz, P.M., Craig, P.S., Ito, A., 2007. A molecular phylogeny of the genus *Echinococcus* inferred from complete mitochondrial genomes. *Parasitology* 134, 713–722.
- Rausch, R.L., 1967. On the ecology and distribution of *Echinococcus* spp. (Cestoda: Taeniidae), and characteristics of their development in the intermediate host. *Annales de Parasitologie Humaine et Comparée* 42, 19–63.
- Reynolds, J., Weir, B.S., Cockerham, C.C., 1983. Estimation for the coancestry coefficient: basis for a short-term genetic distance. *Genetics* 105, 767–779.
- Rosenzvit, M.C., Zhang, L.H., Kamenetzky, L., Canova, S.G., Guarnera, E.A., McManus, D.P., 1999. Genetic variation and epidemiology of *Echinococcus granulosus* in Argentina. *Parasitology* 118, 523–530.
- Rozas, J., Sanchez-Del Barrio, J.C., Messeguer, X., Rozas, R., 2003. DnaSP: DNA polymorphism analyses by the coalescent and other methods. *Bioinformatics* 19, 2496–2497.
- Slatkin, M., Maddison, W.P., 1989. A cladistic measure of gene flow inferred from the phylogenies of alleles. *Genetics* 7, 603–613.
- Smyth, J.D., Smyth, M.M., 1964. Natural and experimental hosts of *Echinococcus granulosus* and *E. multilocularis*, with comments on the genetics of speciation in the genus *Echinococcus*. *Parasitology* 54, 493–514.
- Thompson, R.C., Lymbery, A.J., 1996. Genetic variability in parasites and host-parasite interactions. *Parasitology* 112 (Suppl.), S7–S22.
- Thompson, R.C.A., Lymbery, A.J., Constantine, C.C., 1995. Variation in *Echinococcus*: towards a taxonomic revision of the genus. *Advances in Parasitology* 35, 146–176.
- Thompson, R.C., McManus, D.P., 2002. Towards a taxonomic revision of the genus *Echinococcus*. *Trends in Parasitology* 18, 452–457.