UC Irvine

UC Irvine Previously Published Works

Title

Formation and destruction of the heavy-electron ground state in U compounds with AuBe₅ crystal structure.

Permalink

<https://escholarship.org/uc/item/5zm0798s>

Journal

Physical review. B, Condensed matter, 35(3)

ISSN

0163-1829

Authors

Ott, HR
Rudigier, H
Felder, E
et al.

Publication Date

1987

DOI

10.1103/physrevb.35.1452

Copyright Information

This work is made available under the terms of a Creative Commons Attribution License, available at <https://creativecommons.org/licenses/by/4.0/>

Peer reviewed

Formation and destruction of the heavy-electron ground state in U compounds with AuBe₅ crystal structure

H. R. Ott, H. Rudigier, and E. Felder

Laboratorium für Festkörperphysik, Eidgenössische Technische Hochschule Zürich-Hönggerberg, 8093 Zürich, Switzerland

Z. Fisk and J. D. Thompson

Materials Science and Technology Division, Los Alamos National Laboratory, Los Alamos, New Mexico 87545

(Received 6 June 1986; revised manuscript received 12 November 1986)

The strong influence of changes of the chemical composition on the formation of a heavy-electron ground state is demonstrated in uranium intermetallic compounds crystallizing in the cubic AuBe₅ structure. The most spectacular effect is the large enhancement of the low-temperature electronic specific heat in UPt₅ when one Pt atom per formula unit is replaced by Au. UAuPt₄ may be a heavy-electron superconductor.

The formation of a heavy-electron ground state in uranium intermetallic compounds is, so far, observed exclusively in materials where the nearest distance d_{U-U} between uranium atoms in the crystal lattice exceeds 4 Å. The most prominent examples are UBe₁₃ (Ref. 1) and UPt₃,² both heavy-electron superconductors, and U₂Zn₁₇ (Ref. 3) and UCd₁₁,⁴ which order magnetically out of a heavy-electron state. Among these examples the largest electronic contributions to the molar low-temperature specific heat c_p are observed in UBe₁₃ and UCd₁₁, for which d_{U-U} exceeds 5 and 6 Å, respectively. More recently, the formation of a heavy-electron state, indicated by a c_p/T ratio which increases with decreasing temperature, in an already magnetically ordered material was observed in UCu₅,⁵ another uranium compound where d_{U-U} is close to 5 Å. While these observations suggest that a large separation between 5f-electron-carrying atoms is favorable for the formation of a heavy-electron state it seems *a priori* questionable that this condition alone is sufficient, because, as we shall see below, other compounds with U-U separations close to or more than 5 Å do not show heavy-electron characteristics at low temperatures. In a discussion of the low-temperature behavior of uranium compounds with the Cu₃Au crystal structure, Koelling, Dunlap, and Crabtree⁶ have already noted that d_{U-U} alone is not a reliable parameter for predictions of the magnetic properties and that hybridization effects play a crucial role.

Clear indications that even more subtle details of the electronic structure are essential were obtained from investigations of the effect of impurities on the low-temperature properties of UBe₁₃ and UPt₃. Small amounts of impurity atoms replacing U in UBe₁₃ lead to sizeable changes of the c_p/T ratio of UBe₁₃ at very low temperatures⁷ and similar effects were observed by introducing either Pd on the Pt sites⁸ or Th on the U sites⁹ of UPt₃.

Further evidence was obtained from drastic reductions of the low-temperature specific heat of CeAl₃,¹⁰ UBe₁₃,¹¹ and UPt₃ (Ref. 12) under rather moderate external pres-

ures and from the formation and destruction of a heavy-electron state in the U(Sn_{1-x}In_x)₃ series upon varying x .¹³ As we shall see below, small amounts of impurities can also influence the above-mentioned formation of the state with enhanced specific heat in magnetically ordered UCu₅.

In this work we will give distinct examples of how changes of chemical compositions in isostructural compounds may influence their low-temperature properties, with particular emphasis on the formation or destruction of a heavy-electron ground state. For this purpose we choose uranium compounds crystallizing in the cubic AuBe₅ structure¹⁴ shown in Fig. 1. Within the Be sublattice we recognize two inequivalent sites forming large and small tetrahedra as emphasized in Fig. 1. Its obvious simplicity makes this structure attractive for the investigations we have in mind; in addition, in these compounds the above-mentioned requirement of a large separation of the

FIG. 1. The AuBe₅ crystal structure. The large open circles denote the Au sites; the large dotted circles and the small open circles denote the Be sites. The lines are drawn to visualize the individual positions.

uranium atoms is fulfilled.

Both UNi_5 and UCu_5 crystallize in the structure shown in Fig. 1. In the original work of van Daal, Buschow, van Aken, and van Maaren¹⁵ on $\text{UNi}_{5-x}\text{Cu}_x$, it was pointed out that replacing Ni by Cu leads to dramatic changes in the low-temperature behavior of these substances. While UNi_5 has an essentially temperature-independent magnetic susceptibility below room temperature and stays paramagnetic, $\chi(T)$ of UCu_5 increases substantially with decreasing temperature and antiferromagnetic order is observed below 15 K. Moreover, the c_p/T ratio at helium temperatures is about one order of magnitude smaller in UNi_5 than in UCu_5 . For UNi_5 and UCu_5 the nearest U-U distances are 4.796 and 4.973 Å, respectively, and it might be argued that the difference in this separation alone is responsible for the observed different behavior. In our first example we demonstrate that this is only partially true, particularly with respect to the behavior below 4 K, disregarding, however, in this discussion the recently discovered phase transition around 1 K in UCu_5 .

In Fig. 2 we show the temperature dependence of the low-temperature specific heat of UCu_5 and $\text{UCu}_{4.95}\text{Ni}_{0.05}$ in the form of c_p/T vs T^2 plots. It is evident that 1% Ni impurities on the Cu sites do not greatly affect the overall specific heat, but they are sufficient to suppress the c_p enhancement below 4 K. We recall that this enhancement, however, persists when 20% of the Cu sites are occupied by Ag atoms.⁵ Therefore, it seems difficult to ascribe the change of the c_p/T ratio displayed in Fig. 2 to a simple size effect in the sense mentioned above. It is rather the electronic part of the above-mentioned background specific heat which changes upon larger variations of $d_{\text{U-U}}$, as was shown in Ref. 15, and which is most likely due to band-structure effects. Our measurements of the electrical resistivity ρ in this temperature range reveal that $\rho(T)$ is also quite different in UCu_5 and $\text{UCu}_{4.95}\text{Ni}_{0.05}$. At 1.5 K, ρ of the Ni-doped material is at least a factor of 2 larger than that of pure UCu_5 . In contrast, $\rho(T)$ of UAgCu_4 is almost identical to that of UCu_5 between 1.5 and 10 K.⁵

FIG. 2. c_p/T vs T^2 for UCu_5 and $\text{UCu}_{4.95}\text{Ni}_{0.05}$ between 1.5 and 10 K.

In another example we show that changing the chemical composition of AuBe_5 -type uranium compound may lead to drastic changes in the opposite direction, i.e., to the formation of a heavy-electron state, rather than its destruction as demonstrated above. For this purpose we consider UPt_5 and UAuPt_4 . The physical properties of UPt_5 have been investigated before. $\chi(T)$ below 300 K and the room-temperature valence-band photoemission spectrum were reported by Schneider and Laubschat,¹⁶ $c_p(T)$ was measured by Frings¹⁷ between 1.5 and 15 K, and $\rho(T)$ data between 1.5 and 300 K were published by de Visser, Franse, and Menovsky.¹⁸ Although in UPt_5 the U-U separation is 5.25 Å and hence favorable for the formation of a heavy-electron state, the properties mentioned above indicate a much weaker trend to it than is observed in UPt_3 , where the nearest U-U distance is approximately 1 Å less. It is mainly the temperature dependences of c_p and ρ below 6 K which suggest the presence of spin fluctuations in UPt_5 . As Frings points out, a $T^3 \ln T$ term is needed to properly describe $c_p(T)$ below 6 K, and de Visser and co-workers note a T^2 dependence of ρ below 5 K.

Some of the properties we just described change considerably when one Pt atom per formula unit is replaced by Au. Below, we demonstrate this by reporting results of measurements of $\rho(T)$ and $\chi(T)$ of UAuPt_4 between 1.5 and 300 K and of $c_p(T)$ between 0.15 and 15 K, and by displaying them in comparison with available data on UPt_5 .

In Fig. 3 we show $\chi(T)$ for UAuPt_4 and UPt_5 below room temperature, where the latter data are those of Ref. 16. It may be seen that this change in chemical composition enhances the susceptibility by at least a factor of 2 at all temperatures. Above 60 K, $\chi(T)$ of UAuPt_4 may be described by a Curie-Weiss-type straight line corresponding to an effective moment $\mu_{\text{eff}} = 3.21 \mu_B/(\text{U ion})$ and a paramagnetic Curie temperature Θ_p of -135 K. However, this simple interpretation is questionable because, in spite of the large negative value of Θ_p , UAuPt_4 does not order magnetically down to 0.15 K. Quite obvious is also the change in electrical resistivity as is shown in Fig. 4, where

FIG. 3. Inverse magnetic susceptibility $\chi^{-1}(T)$ of UPt_5 and UAuPt_4 between 1.5 and 220 K. The data for UPt_5 were taken from Ref. 14.

FIG. 4. Temperature dependence of the electrical resistivity $\rho(T)$ of UPt_5 and UAuPt_4 between 1.2 and 300 K.

we present $\rho(T)$ between 1.5 and 300 K for both UAuPt_4 and UPt_5 . Both sets of data were obtained in this work and our results for UPt_5 are somewhat at variance with those reported by de Visser *et al.*,¹⁸ the main difference being the sizeably lower values of ρ obtained in our measurements. $\rho(T)$ of UAuPt_4 is clearly enhanced with respect to that of UPt_5 and the enhancement increases with decreasing temperature. The temperature dependence of ρ of UAuPt_4 approaches that of the heavy-electron uranium compounds UCd_{11} (Ref. 4) and U_2Zn_{17} .³

The most dramatic change is observed in the temperature dependence of the low-temperature specific heat. This may be seen from the c_p/T vs T^2 plots for both UAuPt_4 and UPt_5 shown in Fig. 5. The data for UPt_5 are those of Frings.¹⁷ The slight enhancement of c_p of UAuPt_4 with respect to UPt_5 observed at 15 K, increases steadily with decreasing temperature and below 5 K; c_p/T of UAuPt_4 shows the characteristic upturn of heavy-electron materials. From our measurements down to 0.15 K we extrapolate a c_p/T ratio at $T=0$ K of 725 mJ/mole K², indicating that the zero-temperature electronic-specific-heat parameter γ is eight times larger in UAuPt_4 than in UPt_5 . Below 1 K, the specific heat of UAuPt_4 can be described very accurately by considering c_p due to electrons, the lattice, and a spin-fluctuation-type term of the form $T^3 \ln T$. The resulting fit, however, clearly deviates from the experimental data above 2 K and therefore we have not much faith in that sort of interpretation, especially also because the resulting fit parameter of the spin-fluctuation temperature $T_{\text{SF}}=9.5$ K is equal to T_{SF} in UPt_5 .¹⁷ The separation of the U atoms $d_{\text{U-U}}$ in UPt_5 and UAuPt_4 is 5.25 and 5.285 Å, respectively. Although the change of about 0.7% is quite remarkable we tend to ascribe the large enhancement of c_p/T below 5 K again to a many-body effect and not to a size-induced band-structure effect which may, however, cause the specific-heat increase in UAuPt_4 above 7 K.

From the temperature dependence of ρ we suspect, without definite proof yet, that UAuPt_4 will ideally form an ordered compound with all the Au atoms occupying those Be sites of the AuBe_5 structure that form large tetrahedra (see Fig. 1). Remaining deviations from the

FIG. 5. c_p/T vs T^2 for UPt_5 and UAuPt_4 below 15 K. The data for UPt_5 (above 1.5 K) are those of Ref. 15. The data for UAuPt_4 extend down to 0.15 K.

ideal distribution of the atoms on different lattice sites are most likely the cause for the absence of any phase transition in our samples. This statement is supported by the fact that all known heavy-electron uranium compounds undergo a phase transition at low temperatures unless it is suppressed by impurities or other imperfections of the crystal lattice. In other words, we may also state that we doubt that UAuPt_4 is a uranium analogue to CeAl_3 or CeCu_6 , two heavy-electron Ce compounds that show no phase transition down to 20 mK. It is thus of interest to speculate what kind of phase transition is the most likely to occur in perfect UAuPt_4 . In a previous comparison of low-temperature properties of heavy-electron U compounds¹⁹ it became apparent that the superconductors UBe_{13} and UPt_3 have distinctly larger electronic specific heats per unit volume γ_v than all other compounds which order antiferromagnetically. For UAuPt_4 we calculate a γ_v value of 11.5 mJ/cm³ K², very close to those obtained for UPt_3 and UBe_{13} , respectively. Hence, we conjecture that UAuPt_4 may be a heavy-electron superconductor once it can be prepared in perfect enough form. This conjecture is also supported by the fact that the renormalized ratio of the values of the low-temperature magnetic susceptibility and specific heat $(\pi^2 k_B^2 / 3 \mu_B^2) (\chi / \gamma)$ for UAuPt_4 is 1.43, close to the values found for the three known heavy-electron superconductors. This value is also near a phenomenological boundary separating magnets and superconductors, which was recently proposed by DeLong.²⁰

Our last point is of general and fundamental interest. It should be possible to perform band-structure calculations for both UPt_5 and UAuPt_4 .²¹ This would then allow one to test the significance of such calculations in conjunction with heavy-electron behavior because the large difference in the low-temperature specific heat of these two compounds should reflect itself in the calculated energy dependence of the electronic density of states, if band-structure effects were indeed responsible for the obviously different behavior.

Financial support from the Schweizerische Nationalfonds zur Förderung der Wissenschaftlichen Forschung is gratefully acknowledged. Work at Los Alamos was done under the auspices of the U.S. Department of Energy.

-
- ¹H. R. Ott, H. Rudigier, Z. Fisk, and J. L. Smith, *Phys. Rev. Lett.* **50**, 1595 (1983).
 - ²G. R. Stewart, Z. Fisk, J. O. Willis, and J. L. Smith, *Phys. Rev. Lett.* **52**, 679 (1984).
 - ³H. R. Ott, H. Rudigier, P. Delsing, and Z. Fisk, *Phys. Rev. Lett.* **52**, 1551 (1984).
 - ⁴Z. Fisk, G. R. Stewart, J. O. Willis, H. R. Ott, H. Rudigier, and F. Hulliger, *Phys. Rev. B* **30**, 6360 (1984).
 - ⁵H. R. Ott, H. Rudigier, E. Felder, Z. Fisk, and B. Batlogg, *Phys. Rev. Lett.* **55**, 1595 (1985).
 - ⁶D. D. Koelling, B. D. Dunlap, and G. W. Crabtree, *Phys. Rev. B* **31**, 4966 (1985).
 - ⁷H. R. Ott, H. Rudigier, E. Felder, Z. Fisk, and J. L. Smith, *Phys. Rev. B* **33**, 126 (1986).
 - ⁸A. de Visser, J. C. P. Klaasse, M. van Sprang, J. J. M. Franse, and A. Menovsky, *J. Magn. Magn. Mater.* **54–57**, 375 (1986).
 - ⁹A. P. Ramirez, B. Batlogg, A. S. Cooper, and E. Bucher, *Phys. Rev. Lett.* **57**, 1072 (1986).
 - ¹⁰G. E. Brodale, R. A. Fisher, N. E. Phillips, and J. Flouquet, *Phys. Rev. Lett.* **56**, 390 (1986).
 - ¹¹N. E. Phillips, R. A. Fisher, J. Flouquet, A. L. Giorgi, J. A. Olsen, and G. R. Stewart, *J. Magn. Magn. Mater.* (to be published).
 - ¹²G. E. Brodale, R. A. Fisher, N. E. Phillips, G. R. Stewart, and A. L. Giorgi, *Phys. Rev. Lett.* **57**, 234 (1986).
 - ¹³C. L. Lin, L. W. Zhou, J. E. Crow, R. P. Guertin, and G. R. Stewart, *J. Magn. Magn. Mater.* **54–57**, 391 (1986).
 - ¹⁴L. Misch, *Metallwirtsch.* **14**, 897 (1935).
 - ¹⁵H. J. van Daal, K. H. J. Buschow, P. B. van Aken, and M. H. van Maaren, *Phys. Rev. Lett.* **34**, 1457 (1975).
 - ¹⁶W. D. Schneider and C. Laubschat, *Phys. Rev. B* **23**, 997 (1981).
 - ¹⁷P. H. Frings, thesis, University of Amsterdam, 1984.
 - ¹⁸A. de Visser, J. J. M. Franse, and A. Menovsky, *J. Magn. Magn. Mater.* **43**, 43 (1984).
 - ¹⁹Z. Fisk, J. L. Smith, H. R. Ott, and B. Batlogg, *J. Magn. Magn. Mater.* **52**, 79 (1985).
 - ²⁰L. E. DeLong, *Phys. Rev. B* **33**, 3556 (1986).
 - ²¹R. C. Albers and M. Boring (private communication).