

UC San Diego

UC San Diego Previously Published Works

Title

Functional and clinical studies reveal pathophysiological complexity of CLCN4-related neurodevelopmental condition.

Permalink

<https://escholarship.org/uc/item/6039092w>

Journal

Molecular Psychiatry, 28(2)

Authors

Palmer, Elizabeth

Pusch, Michael

Picollo, Alessandra

et al.

Publication Date

2023-02-01

DOI

10.1038/s41380-022-01852-9

Peer reviewed

ARTICLE OPEN



Functional and clinical studies reveal pathophysiological complexity of *CLCN4*-related neurodevelopmental condition

Elizabeth E. Palmer^{1,2,82}, Michael Pusch^{3,82}, Alessandra Picollo³, Caitlin Forwood¹, Matthew H. Nguyen^{2,4}, Vanessa Suckow⁵, Jessica Gibbons⁵, Alva Hoff^{3,6}, Lisa Sigfrid^{3,6}, Andre Megarbane^{7,8}, Mathilde Nizon^{9,10}, Benjamin Cogné^{9,10}, Claire Beneteau⁹, Fowzan S. Alkuraya¹¹, Aziza Chedrawi¹², Mais O. Hashem¹¹, Hannah Stamberger^{13,14}, Sarah Weckhuysen^{13,14,15}, Arnaud Vanlander¹⁶, Bertien Ceulemans¹⁷, Sulekha Rajagopalan⁴, Kenneth Nunn¹⁸, Stéphanie Arpin¹⁹, Martine Raynaud¹⁹, Constance S. Motter²⁰, Catherine Ward-Melver²⁰, Katrien Janssens²¹, Marije Meuwissen²¹, Diane Beysen²², Nicola Dikow²³, Mona Grimmel²⁴, Tobias B. Haack²⁴, Emma Clement²⁵, Amy McTague^{26,27}, David Hunt²⁸, Sharron Townshend²⁹, Michelle Ward²⁹, Linda J. Richards^{30,31}, Cas Simons^{32,33}, Gregory Costain³⁴, Lucie Dupuis³⁴, Roberto Mendoza-Londono³⁴, Tracy Dudding-Byth^{35,36}, Jackie Boyle³⁵, Carol Saunders^{37,38}, Emily Fleming³⁹, Salima El Chehadeh^{40,41}, Marie-Aude Spitz⁴², Amelie Piton⁴³, Bénédicte Gerard⁴³, Marie-Thérèse Abi Warde^{42,44}, Gillian Rea⁴⁵, Caoimhe McKenna⁴⁵, Sofia Douzgou^{46,47}, Siddharth Banka^{47,48}, Cigdem Akman⁴⁹, Jennifer M. Bain⁴⁹, Tristan T. Sands⁴⁹, Golder N. Wilson⁵⁰, Erin J. Silvertooth⁵¹, Lauren Miller⁵², Damien Lederer⁵³, Rani Sachdev^{1,2}, Rebecca Macintosh^{1,2}, Olivier Monestier⁵³, Deniz Karadurmus⁵³, Felicity Collins⁵⁴, Melissa Carter⁵⁵, Luis Rohena^{56,57}, Marjolein H. Willemsen⁵⁸, Charlotte W. Ockeloen⁵⁸, Rolph Pfundt⁵⁸, Sanne D. Kroft⁵⁹, Michael Field³⁵, Francisco E. R. Laranjeira⁶⁰, Ana M. Fortuna⁶¹, Ana R. Soares⁶¹, Vincent Michaud^{62,63}, Sophie Naudion⁶², Sailaja Golla⁶⁴, David D. Weaver⁶⁵, Lynne M. Bird⁶⁶, Jennifer Friedman⁶⁶, Virginia Clowes^{67,68}, Shelagh Joss⁶⁹, Laura Pölsler⁷⁰, Philippe M. Campeau⁷¹, Maria Blazo⁷², Emilia K. Bijlsma⁷³, Jill A. Rosenfeld^{74,75}, Christian Beetz⁷⁶, Zöe Powis⁷⁷, Kirsty McWalter⁷⁸, Tracy Brandt⁷⁸, Erin Torti⁷⁸, Mikaël Mathot⁷⁹, Shekeeb S. Mohammad^{18,80}, Ruth Armstrong⁸¹ and Vera M. Kalscheuer⁵

© The Author(s) 2022

Missense and truncating variants in the X-chromosome-linked *CLCN4* gene, resulting in reduced or complete loss-of-function (LOF) of the encoded chloride/proton exchanger CIC-4, were recently demonstrated to cause a neurocognitive phenotype in both males and females. Through international clinical matchmaking and interrogation of public variant databases we assembled a database of 90 rare *CLCN4* missense variants in 90 families: 41 unique and 18 recurrent variants in 49 families. For 43 families, including 22 males and 33 females, we collated detailed clinical and segregation data. To confirm causality of variants and to obtain insight into disease mechanisms, we investigated the effect on electrophysiological properties of 59 of the variants in *Xenopus* oocytes using extended voltage and pH ranges. Detailed analyses revealed new pathophysiological mechanisms: 25% (15/59) of variants demonstrated LOF, characterized by a “shift” of the voltage-dependent activation to more positive voltages, and nine variants resulted in a toxic gain-of-function, associated with a disrupted gate allowing inward transport at negative voltages. Functional results were not always in line with *in silico* pathogenicity scores, highlighting the complexity of pathogenicity assessment for accurate genetic counselling. The complex neurocognitive and psychiatric manifestations of this condition, and hitherto unrecognized impacts on growth, gastrointestinal function, and motor control are discussed. Including published cases, we summarize features in 122 individuals from 67 families with *CLCN4*-related neurodevelopmental condition and suggest future research directions with the aim of improving the integrated care for individuals with this diagnosis.

Molecular Psychiatry (2023) 28:668–697; <https://doi.org/10.1038/s41380-022-01852-9>

INTRODUCTION

CLCN4 encodes the intracellularly located chloride/proton ion-exchanger CIC-4, and is located on the human X chromosome at Xp22.2. Rare inherited or de novo missense and truncating variants are identified in a growing number of males and females with a range of neurodevelopmental and psychiatric complications. However, the establishment of the pathogenicity of previously unreported rare missense variants remains challenging. As of 22nd May 2022, from the 153 missense *CLCN4* variants listed

in the publicly available database ClinVar, 73% (111) were classified to be of uncertain significance. Without clear establishment of pathogenicity, families remain on a diagnostic odyssey, cannot make fully informed reproductive choices, or benefit from advances in condition-specific management guidelines or targeted therapies.

The first *CLCN4* variant was reported in an infant male with developmental and epileptic encephalopathy and suggested *CLCN4* as a novel candidate disease gene [1]. Three years later,

A full list of author affiliations appears at the end of the paper.

Received: 5 April 2022 Revised: 10 October 2022 Accepted: 21 October 2022

Published online: 16 November 2022

as part of an X chromosomal exome sequencing study, our group demonstrated that truncating and missense variants were associated with a neurocognitive phenotype in males in five unrelated families [2]. Two families had linkage intervals including Xp22: A two generation French family with five affected males with severe to profound intellectual disability (ID) and variable behavioral difficulties was reported by Raynaud et al., in 1996 [3] and a Belgian family with five males spanning two generations with ID, challenging behaviors and autistic features described by Claes et al. [4]. Heterozygous females in those families were neurotypical or had a mild neurocognitive/psychiatric phenotype. Therefore, a phenotypic entity of X-linked recessive ID (Raynaud-Claes Syndrome) was proposed (MIM *300114).

Subsequently, we reported 10 additional families consisting of 29 hemizygous males and 23 heterozygous females [5]. We clarified that all males had a core phenotype of mild to severe ID, with considerable intrafamilial heterogeneity. For the first time, we reported the phenotype in females with *de novo* variants, which overlapped in severity with that of males. Other common clinical features included epilepsy, subtle white matter changes on neuroimaging, autism spectrum disorder, challenging behaviors, and mental health complications including bipolar disorder, depression, and anxiety. More recently, an additional six males with *CLCN4*-related neurodevelopmental condition were reported confirming the core feature of ID and common comorbidities of epilepsy and challenging behaviors [2, 6, 7]. Xu et al., reported on a female with ID, autistic features and brain abnormalities, with a maternally inherited *CLCN4* missense variant where the mother had mild ID [8]. We recently summarized the published genotypic and phenotypic spectrum [9], noting that, to date, all *CLCN4* variants studied in the *Xenopus* expression system demonstrated partial or complete loss-of-function (LOF) [1, 10, 11].

CIC-4 is one of the nine members of the CLC gene family encoding anion-transporting membrane proteins [12]. CLC proteins are divided into two groups: four members (CIC-1, CIC-2, CIC-Ka, and CIC-Kb) are Cl^- channels localized in the plasma membrane, while the remaining CLCs (CIC-3 to -7) are secondary active Cl^-/H^+ antiporters physiologically localized in intracellular endo-/lysosomal membranes; the latter are also called vesicular CLCs (vCLCs). Among the vCLCs, CIC-3 to -5 are highly homologous and are localized to endosomes, while the more distantly related CIC-6 and CIC-7 are localized to late endosomes and lysosomes, respectively [12]. The vesicular Cl^-/H^+ antiport activity is important for ionic homeostasis of endo-/lysosomes by assisting in vesicular acidification and increasing luminal Cl^- concentration. The function of CIC-4 critically depends on the highly related CIC-3 transporter, with which it forms heterodimers [13, 14]. While most CLCs are physiologically homodimeric, CIC-4 appears to preferentially associate with CIC-3, whereas CIC-4 homodimers are biochemically relatively unstable [13, 14].

CIC-4, and other members of this protein family, CIC-3, CIC-6, CIC-7, and Ostm1, an obligatory subunit of CIC-7, are implicated in neurological disorders [2, 5, 12, 15, 16]. This could be postulated to be related to the postmitotic nature of neurons and their heavy reliance on vesicular trafficking. For example, mice lacking late endosomal CIC-6 transporters show signs of lipofuscin accumulation [17], and lacking lysosomal CIC-7 exhibit a severe lysosomal storage phenotype, respectively [18]. Recently a recurrent gain-of-function (GOF) variant reported in *CLCN6* caused the severe neurodegenerative disease CONRIBA (Neurodegeneration, childhood-onset, hypotonia, respiratory insufficiency and brain imaging abnormalities CONRIBA; MIM 619173) [15] while a variant found in a patient with clinical features of late-onset neuronal ceroid lipofuscinosis [17] was found to have greatly reduced functional activity [19]. LOF of CIC-3 in mice leads to neurodegeneration [20] and both GOF and LOF *CLCN3* variants in humans cause severe global developmental delay [16]. Conversely, knock-out mouse models of CIC-4 have no overt phenotype [21], implying a

complex causative mechanism that requires further exploration to understand the pathophysiological basis of *CLCN4*-related neurodevelopmental condition.

Understanding the pathogenicity of missense variation in *CLCN4* both clinically and functionally is therefore the next step [6]. We firstly undertook a collaborative study aiming to further characterize the genotypic and phenotypic spectrum of *CLCN4*-related neurodevelopmental condition in both males and females. Secondly, we studied the functional impact of novel and previously reported missense variants in heterologously expressing *Xenopus* oocytes by employing electrophysiological measurements using extended voltage-protocols.

SUBJECTS AND METHODS

Subjects

We collected de-identified detailed clinical data on 55 individuals from 43 previously unreported families with (presumed) *CLCN4*-related neurodevelopmental condition, including individuals from three families where the proband had a blended clinical phenotype with a second genetic diagnosis. Data were obtained through an international collaborative process wherein clinicians and diagnostic laboratories with variants identified in *CLCN4* contacted our team, and we also contacted the laboratory or clinician who had deposited variants in *CLCN4* in the public databases DECIPHER, ClinVar, and LOVD [22–24]. In each participating center, written informed consent was obtained from the individual's legal guardians before genetic testing as approved by relevant local ethical committees. Clinical information was obtained by review of medical records and examination of affected individuals. Written informed consent for the publication of clinical data and photographs was also obtained from the participants' legal guardians.

Expression construct

The human CIC-4 cDNA was cloned in the pTLN expression vector [25], in which the disease-associated variants were introduced using standard restriction-free mutagenesis. All constructs were verified by Sanger sequencing.

Expression in oocytes

RNA was transcribed using the SP6 mMessageMachine kit (ThermoFisher, Milan, Italy) after linearization with *Mlu*I. *Xenopus laevis* oocytes were injected with ~6 ng of RNA and incubated at 18 °C for 2–5 days prior to measurements as described previously [26].

Two electrode voltage clamp recordings

Recording pipettes were filled with 3 M KCl (resistance about 0.6 MΩ) and currents were recorded using a TEC03 two electrode voltage clamp amplifier (npi electronics, Tamm, Germany). Ground electrodes were connected to the bath via agar bridges. The standard extracellular solution contained 100 mM NaCl, 5 mM MgSO_4 , 10 mM HEPES (pH 7.3). For solutions at pH 6.3 and 5.3, HEPES buffer was replaced by MES (2-(*n*-morpholino) ethanesulfonic acid) buffer. pH was adjusted with NaOH. Currents were acquired using the custom GePulse acquisition program and an itc-16 interface (Instrutech, Colorado, USA), filtering at 5 kHz and sampling at 50 kHz. Two types of stimulation protocols were applied from a holding potential of –30 mV. The first consisted of 10 ms pulses to voltages ranging from +160 to –120 mV (in 20 mV steps) without leak-subtraction. The second protocol consisted of steps ranging from +170 to –10 mV (in 10 mV steps), applying linear leak and capacity subtraction using a 'P/4' leak subtraction protocol from the holding potential –30 mV. For this procedure 4 pulses of ¼ of the regular amplitude were applied towards negative voltages, their response was averaged, adequately scaled, and subtracted. This procedure approximately eliminates linear capacitive currents and 'leak', assuming that CIC-4 is inactive at negative voltages.

Data analysis

To evaluate the relative expression levels of mutant compared to wild-type (WT) CIC-4, currents were measured for >=6 oocytes for each batch of injection of each construct, and the average current-voltage relationship was obtained using the P/4 subtracted protocol. Average currents

from $>=6$ non-injected oocytes from the same batch were subtracted. For the average IV curves, currents were normalized to the current measured for WT from the same batch at 170 mV, and data from at least four injections for each construct were averaged. For the average ratios of mutant versus WT currents at a given voltage, data, currents were normalized to the respective current measured for WT. This procedure highlights possible alterations of the voltage-dependence. A voltage-independent reduction (or increase) in current size would result in a voltage-independent ratio. We interpret alterations of the voltage-dependent rectification as a change of a gating process that depends on both subunits. Such a gating process is clearly present in CIC-6 and CIC-7 transporters [19, 27] and similarly most likely underlies the extreme rectification of CIC-3, CIC-4, and CIC-5 [28]. In agreement with this hypothesis, practically all variants found here that lead to an apparent shift of the voltage-dependence to more positive voltages are located close to the dimer interface.

For data analysis of currents measured at various external pH values, the following leak-subtraction was performed. For each oocyte, currents measured at pH 7.3 were fitted in the range $-120 \text{ mV} \leq V \leq 0 \text{ mV}$ with a straight line. The line was extrapolated to all voltages and subtracted from the current-voltage relationships (IVs) measured in the various conditions, and normalized to the current at pH 7.3, 160 mV. This is because for WT CIC-4 and for most variants, at pH 7.3, currents recorded at voltages $V \leq 0 \text{ mV}$ are very small and indistinguishable from currents in un-injected oocytes and represent a mixture of leak and endogenous currents. Similar to the “voltage-shifted” variants, we interpret the emergence of inward currents at acidic as a partial disruption of the gating process that in WT keeps the transporter inactive at negative voltages, similar to what described for *CLCN3* variants [16]. Error bars in all figures represent SEM. Statistical significance was assessed by Student’s unpaired two-tailed *t*-test. Variance is similar between all groups because the same batches of oocytes were utilized for WT and variant measurements.

RESULTS

Detailed clinical data were analyzed on 55 previously unreported individuals, 22 hemizygous males and 33 female heterozygotes, from 43 previously unreported families, as well as updated clinical information on one previously reported female who was now recognized to have a recurrent variant [5]. The 44 families were divided into five groups (A-E). This includes families with missense variants, who were divided into groups A-D based on the functional results obtained in the *Xenopus laevis* oocyte model for the *CLCN4* missense variants as described below, as well as three additional patients with novel truncating variants (Group E). Demographic details of these affected individuals, the *CLCN4* variants, their frequency in the gnomAD database, in silico pathogenicity predictors, and results of in vitro functional studies in *Xenopus* oocytes are presented in Table 1. Table 1 also includes the details of variants from the public database ClinVar which we investigated with in vitro functional studies in *Xenopus* oocytes but where we were unable to obtain consent to publish clinical data, as well as variants from ClinVar (as of 25th May 2022) and publications which were recurrent with our investigated variants. New ClinVar accession numbers were obtained for any variant with functional data not already listed in ClinVar and added to Table 1. Figure 1 shows the pedigrees of the unreported families with a novel inherited *CLCN4* variant previously unpublished in the peer-reviewed medical literature. Figure 2 shows clinical photographs and MRI brain images. Figure 3A is a schematic drawing of the *CLCN4* gene and CIC-4 protein with all variants of clearly affected males and females with clinical information available (this study and published). More details on the clinical presentation of these previously unreported families are detailed in Supplementary Table 1 and the case reports, and Supplementary Fig. 3 of individuals with blended phenotypes.

To test for possible impact of variants on the electrophysiological properties of the CIC-4 Cl^-/H^+ antiporter, missense changes as present in the affected individuals were introduced in CIC-4 expression constructs and studied by the 2-electrode voltage-clamp recording method in *Xenopus* oocytes. Example recordings

are shown in Fig. 4A and all results in Table 1 and Supplementary Fig. 1. WT CIC-4 shows typical outwardly rectifying currents as described [29, 30]. An overview of all variants investigated functionally in this study are shown in a topology model and in a three-dimensional model in Fig. 3B, C.

Group A consists of 21 previously unreported families with detailed clinical data (see Supplementary Table 1) whose functional studies are part of this study and demonstrated a LOF, reduced function or shift of voltage dependence, as detailed below. These families included 13 with a male proband and eight with a female proband. In Table 1, Group A also includes seven families that we previously reported on with loss or reduced function [5]. It also includes data from 23 families whose variant was recurrent with one in our cohort, for whom only limited data was available in public databases (ClinVar, DECIPHER, LOVD) as of the 25th May 2022, or from publications from other groups [7, 8, 11, 31]. We included these families as further evidence of the pathogenicity of these recurrent variants.

Some variants, for example p.(Val92Met), showed current levels that were barely above those seen in un-injected oocytes (Fig. 4A–C). A similar near complete loss or reduced function was observed for other variants e.g., p.(Lys62Arg), p.(Ser278Arg), p.(Gly342Glu), and p.(Gly484Arg) (for full list see Table 1, Supplementary Fig. 1). As the variant p.Gly731Val affected the last amino acid of exon 12, we analyzed if this variant impacted splicing (Supplementary Fig. 2), but this could not be demonstrated. Little mechanistic insight can be obtained from these LOF variants as we did not analyze for example if protein stability was affected.

Other variants, for example p.(Asn309Ser), showed a reduced expression level, but no sign of altered voltage-dependence (Fig. 4A–C). The lack of altered voltage-dependence is highlighted in Fig. 4C, which shows that the ratio of currents mediated by variant p.(Asn309Ser) and currents of WT CIC-4 has a practically voltage-independent value of ~ 0.25 . A similar, partially reduced function was observed for variants p.(Ile374Thr) and p.(Gln489Lys) (Table 1, Supplementary Fig. 1).

In contrast to the voltage-independent reduction seen in the variants described above, several other variants, including p.(Leu276Phe), showed a “right-shifted” voltage dependence. This is difficult to appreciate by just comparing the raw current traces (Fig. 4A) or the average current-voltage relationship (Fig. 4B) but is clear in Fig. 4C. For p.(Leu276Phe), the ratio of currents compared to WT is small for $V \sim 20 \text{ mV}$ but progressively enlarged at more positive voltages. The reduction of currents at “physiological” voltages is overcome by sufficiently large positive voltages. This essential LOF phenotype likely reflects an effect on the gating process of CIC-4, as detailed in Subjects and methods. Similar LOF by apparently right-shifted gating was observed to various degrees for variants p.(Gly78Ser), p.(Val212Gly), p.(Gly269Asp), p.(Ile272Val), p.(Val275Leu), p.(Val275Met), p.(Phe319Ser), and p.(Arg718Trp) (Table 1, Supplementary Fig. 1).

Group B. This group includes nine missense *CLCN4* variants from 17 independent families: 14 were previously unreported, one, a female with the *de novo* variant p.(Ala555Val), was previously reported by our group [5], and four families with the same variants or amino acid mutated, that were included in public databases but for whom we could not obtain detailed clinical data. These variants were grouped, as they showed compelling clinical evidence for pathogenicity (rarity, *de novo* status, matching clinical phenotype, and recurrence across unrelated families), without gross effect at the regular recording conditions at pH 7.3. However, p.(Ile549Asn) and some other variants exhibited a characteristic alteration (Fig. 4C): the ratio of currents compared to WT became progressively larger towards more negative voltages. This behavior is reminiscent of a GOF effect described for *CLCN3* variants [16]. Indeed, closer inspection of this and other variants revealed a dramatic GOF that is apparent particularly at acidic

Table 1. Summary of rare *CLCN4* variants reported in this study and, if recurrent, in previous literature or public databases.

	GROUP A: Missense variants LOF					
Families	A1	A2	A3	A4	A5	A6
Genomic position and variant, (GRCh38), NC_000023.10	X:10187555A>G	X:10187602 G>A	X:10194940 G>A	X:10206410 C>T	X:10206437 T>G	X:10206479 C>T
Exon number	4	4	5	7	7	7
c.DNA change, NM_001830.4(CLCN4):	c.185A>G	c.232G>A	c.274G>A	c.608C>T	c.635T>G	c.677C>T
Protein change, NP_001821	p.(Lys62Arg)	p.(Gly78Ser)	p.(Val92Met)	p.(Thr203Ile)	p.(Val212Gly)	p.(Pro226Leu)
Protein domain* for missense variants	N-term, intracellular	Helix B, transmembrane	Helix B, transmembrane	Helix E, intramembrane	Helix E, intramembrane	Helix F, intramembrane
Source :This study = families previously unreported in the medical literature with detailed clinical data; ClinVAR; LOVD and/ or DECIPHER	<i>This study; ClinVAR SCV002525715</i>	<i>Hu et al., 2016; Palmer et al., 2018; ClinVar SCV000297912.2</i>	<i>This study; ClinVar SCV000920556.1</i>	<i>This study; ClinVAR SCV002525716</i>	<i>Hu et al., 2016; Palmer et al., 2018; ClinVar SCV000245780.1</i>	<i>This study; ClinVAR SCV002525717</i>
Gender of proband. Others with variant in family.	1 affected male, mother unaffected	3 affected males	1 affected female proband, 1 affected male (father)	1 affected male, mother unaffected	2 affected males (3 other males in family with ID not tested).	1 affected male
Inheritance	Maternally inherited	Maternally inherited	Paternally inherited	Maternally inherited	Maternally inherited	<i>De novo</i>
Recurrent in unrelated families	No	No	No	No	No	No
Assessment of pathogenicity according to ACMG criteria on ClinVAR, or as assessed by authors using Varsome	VOUS	P	VOUS	LP	P	VOUS
SIFT (dbNSFP version 4.2); converted rankscore	Tolerated (0.48)	Deleterious (0)	Deleterious (0)	Deleterious (0)	Deleterious (0)	Deleterious (0)
PolyPhen	Benign (0.005)	Possibly damaging (0.817)	Benign (0.03)	Probably damaging (0.992)	Probably damaging (0.925)	Probably damaging (0.997)
CADD	21.2	25.4	21.8	25.1	27.2	25.1
REVEL	0.466	0.937	0.608	0.969	0.944	0.967
SpliceAI	≤ 0.2	≤ 0.2	≤ 0.2	≤ 0.2	≤ 0.2	≤ 0.2
Frequency in heterozygotes (gnomAD)	0	0	0	0	0	0
Frequency in hemizygotes (gnomAD)	0	0	0	0	0	0
Functional impact in <i>Xenopus</i> oocyte model	LOF	LOF by shift of voltage dependence	Almost complete LOF	Almost complete LOF	LOF by shift of voltage dependence	LOF
Severe functional impact in <i>Xenopus</i> oocyte model	No	Yes	No	No	Yes	No
Blended phenotype?	No	No	No	No	No	No
Genetic test	Trio exome sequencing	X-chromosome exome	Singleton exome sequencing	Trio exome sequencing	Targeted X-exome sequencing	Trio exome sequencing

Families	A7	A8	A9	A10	A11	A12
Genomic position and variant, (GRCh38), NC_000023.10	X:10206739 G>A	X:10206739 G>A	X:10206747 A>G	X:10206756 G>C	X:10206756 G>A	X:10206756 G>A
Exon number	8	8	8	8	8	8
c.DNA change, NM_001830.4(CLCN4):	c.806G>A	c.806G>A	c.814A>G	c.823G>C	c.823G>A	c.823G>A
Protein change, NP_001821	p.(Gly269Asp)	p.(Gly269Asp)	p.(Ile272Val)	p.(Val275Leu)	p.(Val275Met)	p.(Val275Met)
Protein domain* for missense variants	Helix G, intramembrane	Helix G, intramembrane	Helix H, intramembrane	Helix H, intramembrane	Helix H, intramembrane	Helix H, intramembrane
Source :This study = families previously unreported in the medical literature with detailed clinical data; ClinVAR; LOVD and/ or DECIPHER	<i>This study; ClinVar SCV000607256.1</i>	<i>ClinVar SCV000582636.4</i>	<i>ClinVar SCV000742044.2</i>	<i>This study; ClinVar SCV0002525718</i>	<i>Palmer et al., 2018; ClinVar SCV000245786.1</i>	<i>ClinVar SCV000577686.4</i>
Gender of proband. Others with variant in family.	1 affected female	NR	NR	1 affected male, mother unaffected	1 affected female	NR
Inheritance	<i>De novo</i>	NR	NR	Maternally inherited. <i>De novo</i> in mother	<i>De novo</i>	NR
Recurrent in unrelated families	Yes	Yes	No	No	Yes	Yes
Assessment of pathogenicity according to ACMG criteria on ClinVAR, or as assessed by authors using Varsome	LP	LP	VOUS	VOUS	P	P
SIFT (dbNSFP version 4.2); converted rankscore	Deleterious (0)	Deleterious (0)	Tolerated (0.16)	Deleterious (0)	Deleterious (0)	Deleterious (0)
PolyPhen	Probably damaging (1)	Probably damaging (1)	Benign (0.173)	Probably damaging (0.919)	Probably damaging (0.971)	Probably damaging (0.971)
CADD	26.2	26.2	18.6	25.7	26.7	26.7
REVEL	0.946	0.946	0.564	0.92	0.925	0.925
SpliceAI	≤ 0.2	≤ 0.2	≤ 0.2	≤ 0.2	≤ 0.2	≤ 0.2
Frequency in heterozygotes (gnomAD)	0	0	0	0	0	0
Frequency in hemizygotes (gnomAD)	0	0	0	0	0	0
Functional impact in <i>Xenopus</i> oocyte model	LOF by shift of voltage-dependence	LOF by shift of voltage-dependence	LOF by shift of voltage-dependence	LOF by shift of voltage-dependence	LOF by shift of voltage-dependence	LOF by shift of voltage-dependence
Severe functional impact in <i>Xenopus</i> oocyte model	Yes	Yes	No	No	No	No
Blended phenotype?	No	No	NR	No	No	NR
Genetic test	Trio exome sequencing	NR	NR	Singleton exome sequencing	Trio exome	Exome

Families	A13	A14	A15	A16	A17	A18
Genomic position and variant, (GRCh38), NC_000023.10	X:10206759 C>T	X:10206759 C>T	X:10206759 C>T	X:10206768 C>G	X:10206773 A>T	X:10208049 G>A
Exon number	8	8	8	8	8	9
c.DNA change, NM_001830.4(CLCN4):	c.826C>T	c.832A>C	c.832A>C	c.835C>G	c.840A>T	c.848G>A
Protein change, NP_001821	p.(Leu276Phe)	p.(Ser278Arg)	p.(Ser278Arg)	p.(Leu279Val)	p.(Glu280Asp)	p.(Ser283Asn)
Protein domain* for missense variants	Helix H, intramembrane	Helix H, intramembrane	Helix H, intramembrane	Helix H, intramembrane	Helix H, intramembrane	Loop H-I, intracellular
Source :This study = families previously unreported in the medical literature with detailed clinical data; ClinVAR; LOVD and/ or DECIPHER	<i>This study; ClinVAR SCV002525719</i>	<i>ClinVar SCV000549940.2</i>	<i>ClinVar SCV001542314.1</i>	<i>This study; ClinVAR SCV002525720</i>	<i>This study; ClinVAR SCV002525721</i>	<i>This study; ClinVAR SCV002525722</i>
Gender of proband. Others with variant in family.	1 affected male, mother unaffected	NR	NR	1 affected female	1 affected male	1 affected female
Inheritance	Maternally inherited	NR	NR	<i>De novo</i>	<i>De novo</i>	<i>De novo</i>
Recurrent in unrelated families	No	Yes	Yes	No	No	No
Assessment of pathogenicity according to ACMG criteria on ClinVAR, or as assessed by authors using Varsome	VOUS	VOUS	VOUS	LP	LP	VOUS
SIFT (dbNSFP version 4.2); converted rankscore	Deleterious (0)	Deleterious (0)	Deleterious (0)	Deleterious (0)	Deleterious (0)	Deleterious (0)
PolyPhen	Probably damaging (0.94)	Probably damaging (0.99)	Probably damaging (0.99)	Probably damaging (0.964)	Probably damaging (0.998)	Probably damaging (0.964)
CADD	26.2	27.1	27.1	23.8	23.5	25.4
REVEL	0.945	0.985	0.985	0.801	0.888	0.915
SpliceAI	≤ 0.2	≤ 0.2	≤ 0.2	≤ 0.2	≤ 0.2	≤ 0.2
Frequency in heterozygotes (gnomAD)	0	0	0	0	0	0
Frequency in hemizygotes (gnomAD)	0	0	0	0	0	0
Functional impact in Xenopus oocyte model	LOF by shift of voltage-dependence	Almost complete LOF	Almost complete LOF	LOF	LOF	LOF
Severe functional impact in <i>Xenopus</i> oocyte model	Yes	No	No	Yes	No	No
Blended phenotype?	No	NR	NR	No	No	No
Genetic test	Trio exome sequencing	NR	NR	Trio exome sequencing	Trio exome sequencing	Trio whole genome sequencing

Families	A19	A20	A21	A22	A23	A24
Genomic position and variant, (GRCh38), NC_000023.10	X:10208127 A>G	X:10208127 A>G	X:10208157 T>C	X:10208226 G>A	X:10208226 G>A	X:10208279 C>A
Exon number	9	9	9	9	9	9
c.DNA change, NM_001830.4(CLCN4):	c.926A>G	c.926A>G	c.956T>C	c.1025G>A	c.1025G>A	c.1078C>A
Protein change, NP_001821	p.(Asn309Ser)	p.(Asn309Ser)	p.(Phe319Ser)	p.(Gly342Glu)	p.(Gly342Glu)	p.(Arg360Ser)
Protein domain* for missense variants	Loop I-J, extracellular	Loop I-J, extracellular	Loop I-J, extracellular	Helix J, transmembrane	Helix J, transmembrane	Helix J, transmembrane
Source :This study = families previously unreported in the medical literature with detailed clinical data; ClinVAR; LOVD and/ or DECIPHER	<i>This study; ClinVAR SCV002525723</i>	<i>ClinVAR SCV002032467.1</i>	<i>This study; ClinVAR SCV002525724</i>	<i>This study; ClinVAR SCV002525725</i>	<i>ClinVAR SCV002163577.1</i>	<i>This study; ClinVAR SCV002525726</i>
Gender of proband. Others with variant in family.	2 affected brothers, mother mildly affected	NR	1 affected male, mother unaffected	1 affected male, mother unaffected	NR	1 affected male, mother unaffected
Inheritance	Maternally inherited	NR	Maternally inherited	Maternally inherited. Mosaic in mother	NR	Maternally inherited. <i>De novo</i> in mother
Recurrent in unrelated families	Yes	Yes	No	Yes	Yes	No
Assessment of pathogenicity according to ACMG criteria on ClinVAR, or as assessed by authors using Varsome	VOUS	VOUS	VOUS	VOUS	VOUS	VOUS
SIFT (dbNSFP version 4.2); converted rankscore	Deleterious (0)	Deleterious (0)	Deleterious (0)	Deleterious (0)	Deleterious (0)	Deleterious (0)
PolyPhen	Probably damaging (0.943)	Probably damaging (0.943)	Probably damaging (0.996)	Probably damaging (0.998)	Probably damaging (0.998)	Probably damaging (0.997)
CADD	24.6	24.6	25.7	26.6	26.6	25.3
REVEL	0.821	0.821	0.971	0.966	0.966	0.905
SpliceAI	≤ 0.2	≤ 0.2	≤ 0.2	≤ 0.2	≤ 0.2	≤ 0.2
Frequency in heterozygotes (gnomAD)	0	0	0	0	0	0
Frequency in hemizygotes (gnomAD)	0	0	0	0	0	0
Functional impact in <i>Xenopus</i> oocyte model	Reduced function	Reduced function	LOF by shift of voltage-dependence	LOF	LOF	LOF
Severe functional impact in <i>Xenopus</i> oocyte model	No	No	Yes	Yes	Yes	No
Blended phenotype?	No	No	No	No	No	No
Genetic test	Quad exome sequencing (research)	NR	Singleton exome sequencing (research)	Targeted MPS gene panel (epilepsy)	NR	Trio exome sequencing

Families	A25	A26	A27	A28	A29	A30
Genomic position and variant, (GRCh38), NC_000023.10	X:10208322 T>C	X:10208322 T>C	X:10212527 G>A	X:10212542 C>A	X:10212653 G>A	X:10212653 G>A
Exon number	9	9	10	10	10	10
c.DNA change, NM_001830.4(CLCN4):	c.1121T>C	c.1121T>C	c.1450G>A	c.1465C>A	c.1576G>A	c.1576G>A
Protein change, NP_001821	p.(Ile374Thr)	p.(Ile374Thr)	p.(Gly484Arg)	p.(Gln489Lys)	p.(Gly526Ser)	p.(Gly526Ser)
Protein domain* for missense variants	Helix K, intramembrane	Helix K, intramembrane	Helix N, intramembrane	Helix N, intramembrane	Helix O, intramembrane	Helix O, intramembrane
Source :This study = families previously unreported in the medical literature with detailed clinical data; ClinVAR; LOVD and/ or DECIPHER	<i>ClinVar</i> SCV000577573.3	<i>ClinVar</i> SCV002200551.1	LOVD#0000346105	<i>This study;</i> <i>ClinVar</i> SCV000589760.3	<i>This study;</i> <i>ClinVar</i> SCV000693819.1	<i>ClinVar</i> SCV000942548.4
Gender of proband. Others with variant in family.	NR	NR	NR	1 affected female	1 affected male, 1 affected brother and maternal uncle (not tested)	NR
Inheritance	NR	NR	NR	<i>De novo</i>	Maternally inherited	NR
Recurrent in unrelated families	Yes - but also present in gnomAD	Yes - but also present in gnomAD	No	No	Yes	Yes
Assessment of pathogenicity according to ACMG criteria on ClinVAR, or as assessed by authors using Varsome	VOUS	VOUS	LP	LP	LP	VOUS
SIFT (dbNSFP version 4.2); converted rankscore	Tolerated	Tolerated	Deleterious (0)	Tolerated (0.06)	Deleterious (0.03)	Deleterious (0.03)
PolyPhen	Benign (0.04)	Benign (0.04)	Probably damaging (0.999)	Benign (0.221)	Possibly damaging (0.73)	Possibly damaging (0.73)
CADD	21.2	21.2	28.1	22.3	33	33
REVEL	0.733	0.733	0.975	0.806	0.913	0.913
SpliceAI	≤ 0.2	≤ 0.2	≤ 0.2	≤ 0.2	ΔS donor gain 0.54	ΔS donor gain 0.54
Frequency in heterozygotes (gnomAD)	1.02 × 10 ⁻⁵	1.02 × 10 ⁻⁵	0	0	0	0
Frequency in hemizygotes (gnomAD)	0	0	0	0	0	0
Functional impact in <i>Xenopus</i> oocyte model	Reduced function	Reduced function	LOF	Reduced function	LOF by shift of voltage-dependence	LOF by shift of voltage-dependence
Severe functional impact in <i>Xenopus</i> oocyte model	No	No	No	No	No	No
Blended phenotype?	NR	NR	NR	No	No	NR
Genetic test	NR	NR	NR	Trio exome sequencing	Exome	NR

Families	A31	A32	A33	A34	A35	A36
Genomic position and variant, (GRCh38), NC_000023.10	X:10213701 G>A	X:10213710 G>A	X:10213710 G>A	X:10213734 G>C	X:10213734 G>C	X:10213738 G>A
Exon number	11	11	11	11	11	11
c.DNA change, NM_001830.4(CLCN4):	c.1597G>A	c.1606G>A	c.1606G>A	c.1630G>A	c.1630G>A	c.1633G>A
Protein change, NP_001821	p.(Val533Met)	p.(Val536Met)	p.(Val536Met)	p.(Gly544Arg)	p.(Gly544Arg)	p.(Gly545Ser)
Protein domain* for missense variants	Helix P, intramembrane	Helix P, intramembrane	Helix P, intramembrane	Loop P-Q, intramembrane	Loop P-Q, intramembrane	Loop P-Q, intramembrane
Source :This study = families previously unreported in the medical literature with detailed clinical data; ClinVAR; LOVD and/ or DECIPHER	<i>This study; ClinVar SCV002525727</i>	<i>Hu et al., 2016; Palmer et al., 2018; ClinVar SCV000297914.2</i>	<i>ClinVar SCV001847703.1</i>	<i>Veeramah et al., 2013; ClinVar SCV000120005.3</i>	<i>Palmer et al., 2018; ClinVar SCV000245787.1</i>	<i>ClinVar SCV000570417.4</i>
Gender of proband. Others with variant in family.	1 affected male	7 affected males, two affected females (one severely affected)	NR	1 affected male	1 affected male	NR
Inheritance	Maternally inherited	Maternally inherited	Maternally inherited	<i>De novo</i>	<i>Mosaic de novo</i>	NR
Recurrent in unrelated families	No	Yes	Yes	Yes	Yes	No
Assessment of pathogenicity according to ACMG criteria on ClinVAR, or as assessed by authors using Varsome	VOUS	P	P	P	P	VOUS
SIFT (dbNSFP version 4.2); converted rankscore	Deleterious (0.02)	Deleterious (0)	Deleterious (0)	Deleterious (0)	Deleterious (0)	Tolerated (0.17)
PolyPhen	Probably damaging (0.924)	Probably damaging (0.997)	Probably damaging (0.997)	Probably damaging (0.999)	Probably damaging (0.999)	Benign (0.402)
CADD	26.3	26.8	26.8	27.1	27.1	22.8
REVEL	0.871	0.907	0.907	0.884	0.884	0.648
SpliceAI	≤ 0.2	≤ 0.2	≤ 0.2	≤ 0.2	≤ 0.2	≤ 0.2
Frequency in heterozygotes (gnomAD)	0	0	0	0	0	0
Frequency in hemizygotes (gnomAD)	0	0	0	0	0	0
Functional impact in <i>Xenopus</i> oocyte model	LOF by shift of voltage-dependence	LOF by shift of voltage-dependence	LOF by shift of voltage-dependence	LOF by shift of voltage-dependence	LOF by shift of voltage-dependence	Almost complete LOF
Severe functional impact in <i>Xenopus</i> oocyte model	No	No	No	Yes	Yes	No
Blended phenotype?	No	No	No	No	NR	NR
Genetic test	Exome	X-chromosome exome	NR	X-chromosome exome	NR	NR

Families	A37	A38	A39	A40	A41	A42
Genomic position and variant, (GRCh38), NC_000023.10	X:10181778 G>A	X:10213749A>C	X:10213782A>G	X:10214008 C>G	X:10214010 G>A	X:10220837 C>T
Exon number	11	11	11	11	11	12
c.DNA change, NM_001830.4(CLCN4):	c.1634G>A	c.1645A>C	c.1678A>G	c.1904C>G	c.1906G>A	c.2152C>T
Protein change, NP_001821	p.(Gly545Asp)	p.(Ile549Leu)	p.(Lys560Glu)	p.(Pro635Arg)	p.(Val636Met)	p.(Arg718Trp)
Protein domain* for missense variants	Loop P-Q, intramembrane	Helix Q, transmembrane	Helix Q, transmembrane	CBS1, intracellular	CBS1, intracellular	CBS2, intracellular
Source :This study = families previously unreported in the medical literature with detailed clinical data; ClinVAR; LOVD and/ or DECIPHER	ClinVar SCV000741711.2	ClinVar SCV001986448.1	ClinVar SCV000780957.12	This study; ClinVAR SCV002525728	This study; ClinVar SCV001572293.1	This study; ClinVAR SCV002525729
Gender of proband. Others with variant in family.	NR	NR	NR	1 affected female, mother mildly affected	1 affected female, mother unaffected	1 affected female
Inheritance	NR	NR	NR	Maternally inherited. <i>De novo</i> in mother	Maternally inherited	<i>De novo</i>
Recurrent in unrelated families	No	No	No	No	No	Yes
Assessment of pathogenicity according to ACMG criteria on ClinVAR, or as assessed by authors using Varsome	LP	VOUS	VOUS	VOUS	VOUS	P
SIFT (dbNSFP version 4.2); converted rankscore	Deleterious (0)	Deleterious (0)	Deleterious (0)	Deleterious (0.02)	Deleterious (0.01)	Deleterious (0)
PolyPhen	Possibly damaging (0.877)	Benign (0.17)	Probably damaging (0.993)	Probably damaging (0.998)	Probably damaging (0.914)	Probably damaging (0.999)
CADD	25.7	23.4	26.8	24.9	26.2	26
REVEL	0.828	0.675	0.962	0.954	0.9	0.874
SpliceAI	≤ 0.2	≤ 0.2	≤ 0.2	≤ 0.2	≤ 0.2	≤ 0.2
Frequency in heterozygotes (gnomAD)	0	0	0	0	0	0
Frequency in hemizygotes (gnomAD)	0	0	0	0	0	0
Functional impact in <i>Xenopus</i> oocyte model	LOF by shift of voltage-dependence	LOF by shift of voltage-dependence	LOF	LOF	Reduced function	LOF by shift of voltage-dependence
Severe functional impact in <i>Xenopus</i> oocyte model	Yes	Yes	No	No	No	No
Blended phenotype?	NR	NR	No	No	No	No
Genetic test	NR	NR	NR	NR	Exome sequencing	Targeted MPS gene panel

Families	A43	A44	A45	A46	A47	A48
Genomic position and variant, (GRCh38), NC_000023.10	X:10220837 C>T	X:10220837 C>T	X:10220837 C>T	X:10220837 C>T	X:10220837 C>T	X:10220837 C>T
Exon number	12	12	12	12	12	12
c.DNA change, NM_001830.4(CLCN4):	c.2152C>T	c.2152C>T	c.2152C>T	c.2152C>T	c.2152C>T	c.2152C>T
Protein change, NP_001821	p.(Arg718Trp)	p.(Arg718Trp)	p.(Arg718Trp)	p.(Arg718Trp)	p.(Arg718Trp)	p.(Arg718Trp)
Protein domain* for missense variants	CBS2, intracellular	CBS2, intracellular	CBS2, intracellular	CBS2, intracellular	CBS2, intracellular	CBS2, intracellular
Source :This study = families previously unreported in the medical literature with detailed clinical data; ClinVAR; LOVD and/ or DECIPHER	Palmer et al., 2018; ClinVar SCV000245785.1	He et al., 2021	Zhou et al., 2018	ClinVar SCV002069088.1	ClinVar SCV002058687.1	ClinVar SCV001976771.1
Gender of proband. Others with variant in family.	1 affected female	1 affected male, unaffected mother	1 affected male	NR	1 affected female	NR
Inheritance	<i>De novo</i>	Maternal (n.b. mother has a karyotype 47,XXX/46,XX)	<i>De novo</i>	NR	NR	NR
Recurrent in unrelated families	Yes	Yes	Yes	Yes	Yes	Yes
Assessment of pathogenicity according to ACMG criteria on ClinVAR, or as assessed by authors using Varsome	P	P	P	P	P	P
SIFT (dbNSFP version 4.2); converted rankscore	Deleterious (0)	Deleterious (0)	Deleterious (0)	Deleterious (0)	Deleterious (0)	Deleterious (0)
PolyPhen	Probably damaging (0.999)	Probably damaging (0.999)	Probably damaging (0.999)	Probably damaging (0.999)	Probably damaging (0.999)	Probably damaging (0.999)
CADD	26	26	26	26	26	26
REVEL	0.874	0.874	0.874	0.874	0.874	0.874
SpliceAI	≤ 0.2	≤ 0.2	≤ 0.2	≤ 0.2	≤ 0.2	≤ 0.2
Frequency in heterozygotes (gnomAD)	0	0	0	0	0	0
Frequency in hemizygotes (gnomAD)	0	0	0	0	0	0
Functional impact in <i>Xenopus</i> oocyte model	LOF by shift of voltage-dependence	LOF by shift of voltage-dependence	LOF by shift of voltage-dependence	LOF by shift of voltage-dependence	LOF by shift of voltage-dependence	LOF by shift of voltage-dependence
Severe functional impact in <i>Xenopus</i> oocyte model	No	No	No	No	No	No
Blended phenotype?	No	No	No	No	No	No
Genetic test	WGS	Trio exome	Targeted MPS gene panel	NR	NR	NR

	GROUP B: Missense variants GOF					
Families	A49	A50	A51	B1	B2	B3
Genomic position and variant, (GRCh38), NC_000023.10	X:10220837 C>T	X:10220876 G>A	X:10220877 G>T	X:10194931 G>A	X:10194931 G>A	X:10206737 T>G
Exon number	12	12	12	5	5	8
c.DNA change, NM_001830.4(CLCN4):	c.2152C>T	c.2191G>A	c.2192G>T	c.265G>A	c.265G>A	c.804T>G
Protein change, NP_001821	p.(Arg718Trp)	p.(Gly731Arg)	p.Gly731Val	p.(Asp89Asn)	p.(Asp89Asn)	p.(Phe268Leu)
Protein domain* for missense variants	CBS2, intracellular	CBS2, intracellular	CBS2, intracellular	Helix B, transmembrane	Helix B, transmembrane	Helix G, intramembrane
Source :This study = families previously unreported in the medical literature with detailed clinical data; ClinVAR; LOVD and/ or DECIPHER	ClinVar SCV000957439.2	Palmer et al., 2018; ClinVar SCV001582304.2	This study; ClinVar SCV002525730	This study; ClinVar SCV000569027.4	ClinVar SCV001468990.1	This study; ClinVar SCV000589740.2
Gender of proband. Others with variant in family.	NR	3 affected males	1 affected male	1 affected female	NR	1 affected female
Inheritance	<i>De novo</i>	Maternally inherited	Maternally inherited	<i>De novo</i>	NR	<i>De novo</i>
Recurrent in unrelated families	Yes	No	No	Yes	Yes	No
Assessment of pathogenicity according to ACMG criteria on ClinVAR, or as assessed by authors using Varsome	P	P	VOUS	VOUS	LP	LP
SIFT (dbNSFP version 4.2); converted rankscore	Deleterious (0)	Deleterious (0.03)	Deleterious (0)	Deleterious (0)	Deleterious (0)	Deleterious (0)
PolyPhen	Probably damaging (0.999)	Possibly damaging (0.798)	Probably damaging (0.993)	Probably damaging (0.918)	Probably damaging (0.918)	Probably damaging (0.996)
CADD	26	32	33	24.7	24.7	23.2
REVEL	0.874	0.878	0.926	0.685	0.685	0.934
SpliceAI	≤ 0.2	≤ 0.2	≤ 0.2	≤ 0.2	≤ 0.2	≤ 0.2
Frequency in heterozygotes (gnomAD)	0	0	0	0	0	0
Frequency in hemizygotes (gnomAD)	0	0	0	0	0	0
Functional impact in <i>Xenopus</i> oocyte model	LOF by shift of voltage-dependence	LOF	LOF	GOF	GOF	GOF
Severe functional impact in <i>Xenopus</i> oocyte model	No	No	No	No	No	No
Blended phenotype?	No	No	No	NR	No	No
Genetic test	NR	X-chromosome exome sequencing	Exome sequencing	Exome sequencing	NR	Exome sequencing

Families	B4	B5	B6	B7	B8	B9
Genomic position and variant, (GRCh38), NC_000023.10	X:10208129 C>T	X:10208150 G>A	X:10208150 G>A	X:10208150 G>A	X:10208150 G>T	X:10208386 C>G
Exon number	9	9	9	9	9	9
c.DNA change, NM_001830.4(CLCN4):	c.928C>T	c.949G>A	c.949G>A	c.949G>A	c.949G>T	c.1185C>G
Protein change, NP_001821	p.(Pro310Ser)	p.(Val317Ile)	p.(Val317Ile)	p.(Val317Ile)	p.(Val317Phe)	p.(Ser395Arg)
Protein domain* for missense variants	Loop I-J, extracellular	Loop I-J, extracellular	Loop I-J, extracellular	Loop I-J, extracellular	Loop I-J, extracellular	Loop K-L, intramembrane
Source :This study = families previously unreported in the medical literature with detailed clinical data; ClinVAR; LOVD and/ or DECIPHER	<i>This study; ClinVar SCV002525731</i>	<i>This study; ClinVar SCV001437777.1</i>	<i>This study; ClinVar SCV000572387.4</i>	<i>This study, DECIPHER Patient 279296</i>	<i>ClinVar SCV000621815.2</i>	<i>This study; ClinVar SCV002525733</i>
Gender of proband. Others with variant in family.	1 affected female	1 affected male	1 affected male, mother mildly affected	1 affected male	NR	1 affected female
Inheritance	<i>De novo</i>	<i>De novo</i>	Maternally inherited. Mosaic in mother	<i>De novo</i>	NR	<i>De novo</i>
Recurrent in unrelated families	No	Yes	Yes	Yes	No	No
Assessment of pathogenicity according to ACMG criteria on ClinVAR, or as assessed by authors using Varsome	VOUS	VOUS	VOUS	LP	VOUS	VOUS
SIFT (dbNSFP version 4.2); converted rankscore	Deleterious (0)	Tolerated (0.15)	Tolerated (0.15)	Tolerated (0.15)	Deleterious (0)	Deleterious (0)
PolyPhen	Possibly damaging (0.883)	Possibly damaging (0.612)	Possibly damaging (0.612)	Possibly damaging (0.612)	Probably damaging (0.993)	Probably damaging (0.933)
CADD	24.5	22.8	22.8	22.8	25	14.46
REVEL	0.931	0.525	0.525	0.525	0.924	0.732
SpliceAI	≤ 0.2	≤ 0.2	≤ 0.2	≤ 0.2	≤ 0.2	≤ 0.2
Frequency in heterozygotes (gnomAD)	0	0	0	0	0	0
Frequency in hemizygotes (gnomAD)	0	0	0	0	0	0
Functional impact in Xenopus oocyte model	GOF	GOF	GOF	GOF	GOF	Reduced outward currents, and slight GOF
Severe functional impact in Xenopus oocyte model	No	No	No	No	No	No
Blended phenotype?	No	No	No	No	NR	No
Genetic test	Exome sequencing	Singleton exome sequencing	Duo exome sequencing (proband and mother)	DDD project (whole genome sequencing)	NR	Targeted MPS gene panel

Families	B10	B11	B12	B13	B14	B15
Genomic position and variant, (GRCh38), NC_000023.10	X:10213750 T>A	X:10213752 G>C	X:10213768 C>T	X:10213768 C>T	X:10213768 C>T	X:10213768 C>T
Exon number	11	11	11	11	11	11
c.DNA change, NM_001830.4(CLCN4):	c.1646T>A	c.1648G>C	c.1664C>T	c.1664C>T	c.1664C>T	c.1664C>T
Protein change, NP_001821	p.(Ile549Asn)	p.(Val550Leu)	p.(Ala555Val)	p.(Ala555Val)	p.(Ala555Val)	p.(Ala555Val)
Protein domain* for missense variants	Helix Q, transmembrane	Helix Q, transmembrane	Helix Q, transmembrane	Helix Q, transmembrane	Helix Q, transmembrane	Helix Q, transmembrane
Source :This study = families previously unreported in the medical literature with detailed clinical data; ClinVAR; LOVD and/ or DECIPHER	<i>This study; ClinVar SCV002525734</i>	<i>This study; ClinVar SCV002525735</i>	<i>Palmer et al., 2018; ClinVar SCV000245784.1</i>	<i>This study; ClinVar SCV002525736</i>	<i>This study; ClinVar SCV000511380.1</i>	<i>This study; ClinVar SCV000490472.1</i>
Gender of proband. Others with variant in family.	1 affected female	1 affected female	1 affected female	1 affected female	1 affected female	1 affected female
Inheritance	<i>De novo</i>	<i>De novo</i>	<i>De novo</i>	<i>De novo</i>	<i>De novo</i>	<i>De novo</i>
Recurrent in unrelated families	No	No	Yes	Yes	Yes	Yes
Assessment of pathogenicity according to ACMG criteria on ClinVAR, or as assessed by authors using Varsome	VOUS	VOUS	P	P	P	P
SIFT (dbNSFP version 4.2); converted rankscore	Deleterious (0)	Tolerated (0.29)	Deleterious (0)	Deleterious (0)	Deleterious (0)	Deleterious (0)
PolyPhen	Probably damaging (0.964)	Benign (0.279)	Possibly damaging (0.627)	Possibly damaging (0.627)	Possibly damaging (0.627)	Possibly damaging (0.627)
CADD	26.3	21.7	23.1	23.1	23.1	23.1
REVEL	0.926	0.694	0.734	0.734	0.734	0.734
SpliceAI	≤ 0.2	≤ 0.2	≤ 0.2	≤ 0.2	≤ 0.2	≤ 0.2
Frequency in heterozygotes (gnomAD)	0	0	0	0	0	0
Frequency in hemizygotes (gnomAD)	0	0	0	0	0	0
Functional impact in Xenopus oocyte model	GOF	GOF	GOF	GOF	GOF	GOF
Severe functional impact in Xenopus oocyte model	No	No	No	No	No	No
Blended phenotype?	No	No	No	No	No	No
Genetic test	Trio exome sequencing	Targeted MPS gene panel (intellectual disability)	Trio exome sequencing	Targeted MPS gene panel (intellectual disability)	Targeted MPS gene panel (epilepsy)	Trio exome sequencing

	GROUP C: Missense variants LOF: blended phenotype					
Families	<i>B16</i>	<i>B17</i>	<i>C1</i>	<i>C2</i>	<i>C3</i>	<i>C4</i>
Genomic position and variant, (GRCh38), NC_000023.10	X:10213768 C>T	X:10213768 C>T	X:10185132 G>A	X:10187576 C>T	X:10213768 C>T	X:10213768 C>T
Exon number	11	11	3	4	9	9
c.DNA change, NM_001830.4(CLCN4):	c.1664C>T	c.1664C>T	c.100G>A	c.206C>T	c.1106C>T	c.1106C>T
Protein change, NP_001821	p.(Ala555Val)	p.(Ala555Val)	p.(Asp34Asn)	p.(Ser69Leu)	p.(Pro369Leu)	p.(Pro369Leu)
Protein domain* for missense variants	Helix Q, transmembrane	Helix Q, transmembrane	N term, intracellular	N term, intracellular	Helix K, intramembrane	Helix K, intramembrane
Source :This study = families previously unreported in the medical literature with detailed clinical data; ClinVAR; LOVD and/ or DECIPHER	<i>ClinVar</i> SCV000740693.2	<i>ClinVar</i> SCV002242006.1	<i>This Study; DECIPHER</i> <i>Patient 277726; ClinVar</i> SCV002525737	<i>This study; ClinVar</i> SCV002525738	<i>This study; ClinVar</i> SCV002525739	<i>ClinVar</i> SCV001503010.2
Gender of proband. Others with variant in family.	NR	NR	1 affected male, mother and sister unaffected neurodevelopmentally	1 affected male, mother unaffected	1 affected male	NR
Inheritance	NR	NR	Maternally inherited.	Maternally inherited	<i>De novo</i>	NR
Recurrent in unrelated families	Yes	Yes	No	No	Yes	Yes
Assessment of pathogenicity according to ACMG criteria on ClinVAR, or as assessed by authors using Varsome	VOUS	P	VOUS	VOUS	VOUS	VOUS
SIFT (dbNSFP version 4.2); converted rankscore	Deleterious (0)	Deleterious (0)	Deleterious (0)	Deleterious (0)	Deleterious (0)	Deleterious (0)
PolyPhen	Possibly damaging (0.627)	Possibly damaging (0.627)	Probably damaging (1)	Benign (0.332)	Probably damaging (0.925)	Probably damaging (0.925)
CADD	23.1	23.1	27.7	24.2	25.2	25.2
REVEL	0.734	0.734	0.778	0.869	0.896	0.896
SpliceAI	≤ 0.2	≤ 0.2	≤ 0.2	≤ 0.2	≤ 0.2	≤ 0.2
Frequency in heterozygotes (gnomAD)	0	0	1	0	0	0
Frequency in hemizygotes (gnomAD)	0	0	0	0	0	0
Functional impact in <i>Xenopus</i> oocyte model	GOF	GOF	LOF	Almost complete LOF	LOF	LOF
Severe functional impact in <i>Xenopus</i> oocyte model	No	No	No	No	No	No
Blended phenotype?	No	No	Yes with Desbuquois syndrome	Yes with SOX11-related condition	Yes a clinical diagnosis of Donnai-Barrow syndrome	NR
Genetic test	NR	NR	DDD project (WGS)	Trio exome sequencing	Trio whole genome sequencing	NR

	GROUP D: Functional studies like wild type					
Families	D1	D2	D3	D4	D5	D6
Genomic position and variant, (GRCh38), NC_000023.10	X:10185119 C>G	X:10194980 C>G	X:10194980 C>G	X:10206514 T>C	X:10208145 G>A	X:10208145 G>A
Exon number	3	5	5	7	9	9
c.DNA change, NM_001830.4(CLCN4);	c.87C>G	c.314C>G	c.314C>G	c.712T>C	c.944G>A	c.944G>A
Protein change, NP_001821	p.(Asp29Glu)	p.(Ser105Cys)	p.(Ser105Cys)	p.(Phe238Leu)	p.(Arg315His)	p.(Arg315His)
Protein domain* for missense variants	N-term	Loop B-C, extracellular	Loop B-C, extracellular	Helix F, intramembrane	Loop I-J, extracellular	Loop I-J, extracellular
Source :This study = families previously unreported in the medical literature with detailed clinical data; ClinVAR; LOVD and/ or DECIPHER	<i>This study; ClinVar SCV002525740</i>	<i>ClinVar SCV002003533.1</i>	<i>ClinVar SCV000549937.4</i>	<i>ClinVar SCV000570777.4</i>	<i>This study; ClinVar SCV002525741</i>	<i>ClinVar SCV001480412.1</i>
Gender of proband. Others with variant in family.	2 affected males, 1 mildly affected female	NR	NR	NR	1 affected female	NR
Inheritance	Maternally inherited	NR	NR	NR	<i>De novo</i>	Maternally inherited
Recurrent in unrelated families	No	Yes - but also present in gnomAD	Yes - but also present in gnomAD	No	Yes - but also present in gnomAD	Yes - but also present in gnomAD
Assessment of pathogenicity according to ACMG criteria on ClinVAR, or as assessed by authors using Varsome	VOUS	VOUS	VOUS	VOUS	LP	VOUS
SIFT (dbNSFP version 4.2); converted rankscore	Deleterious (0.04)	Deleterious (0.03)	Deleterious (0.03)	Tolerated (1)	Tolerated (1)	Tolerated (1)
PolyPhen	Probably damaging (0.983)	Benign (0.246)	Benign (0.246)	Benign (0.013)	Benign (0.01)	Benign (0.01)
CADD	23.9	22.6	22.6	20.9	20.5	20.5
REVEL	0.852	0.494	0.494	0.583	0.575	0.575
SpliceAI	≤ 0.2	≤ 0.2	≤ 0.2	≤ 0.2	≤ 0.2	≤ 0.2
Frequency in heterozygotes (gnomAD)	0	1	1	0	0	0
Frequency in hemizygotes (gnomAD)	0	0	0	0	1	1
Functional impact in <i>Xenopus</i> oocyte model	WT	WT	WT	WT	WT	WT
Severe functional impact in <i>Xenopus</i> oocyte model	No	No	No	No	No	No
Blended phenotype?	No	NR	NR	NR	No	No
Genetic test	Trio whole genome sequencing	NR	NR	NR	Trio exome sequencing	NR

Families	D7	D8	D9	D10	D11	D12
Genomic position and variant, (GRCh38), NC_000023.10	X:10208145 G>A	X:10208291 A>G	X:10208496 G>A	X:10213966 A>G	X:10213990 C>T	X:10214041 T>C
Exon number	9	9	9	11	11	11
c.DNA change, NM_001830.4(CLCN4):	c.944G>A	c.1090A>G	c.1295G>A	c.1862A>G	c.1886C>T	c.1937T>C
Protein change, NP_001821	p.(Arg315His)	p.(Arg364Gly)	p.(Arg432Gln)	p.(Asp621Gly)	p.(Thr629Ile)	p.(Ile646Thr)
Protein domain* for missense variants	Loop I-J, extracellular	Loop J-K, intracellular	Loop L-M, extracellular	CBS1, intracellular	CBS1, intracellular	CBS1, intracellular
Source :This study = families previously unreported in the medical literature with detailed clinical data; ClinVAR; LOVD and/ or DECIPHER	<i>ClinVar</i> SCV002250535.1	<i>This study;</i> <i>ClinVar</i> SCV002525742	<i>ClinVar</i> SCV000594131.1	<i>ClinVar</i> SCV000620779.1	<i>This study;</i> <i>ClinVar</i> SCV002525743	<i>ClinVar</i> SCV000741977.2
Gender of proband. Others with variant in family.	NR	1 affected male	NR	NR	1 affected female	NR
Inheritance	NR	NR	NR	NR	Maternally inherited	NR
Recurrent in unrelated families	Yes - but also present in gnomAD	No	No	No	No	Yes - but also present in gnomAD
Assessment of pathogenicity according to ACMG criteria on ClinVAR, or as assessed by authors using Varsome	VOUS	VOUS	VOUS	VOUS	VOUS	VOUS
SIFT (dbNSFP version 4.2); converted rankscore	Tolerated (1)	Tolerated (0.18)	Tolerated (0.23)	Deleterious (0.05)	Deleterious (0.01)	Deleterious (0.03)
PolyPhen	Benign (0.01)	Benign (0.062)	Benign (0.037)	Benign (0.044)	Possibly damaging (0.851)	Benign (0.158)
CADD	20.5	18.99	22.7	23.1	24.4	23.8
REVEL	0.575	0.366	0.576	0.72	0.888	0.868
SpliceAI	≤ 0.2	≤ 0.2	≤ 0.2	≤ 0.2	≤ 0.2	≤ 0.2
Frequency in heterozygotes (gnomAD)	0	0	2	0	0	11
Frequency in hemizygotes (gnomAD)	1	0	0	0	0	4
Functional impact in <i>Xenopus</i> oocyte model	WT	WT	WT	WT	WT	WT
Severe functional impact in <i>Xenopus</i> oocyte model	No	No	No	No	No	No
Blended phenotype?	No	No	NR	NR	NR	NR
Genetic test	NR	Intellectual disability gene panel off exome backbone	NR	NR	Singleton exome sequencing	NR

Families	D13	D14	D15	D16	D17	D18
Genomic position and variant, (GRCh38), NC_000023.10	X:10214041 T>C	X:10214059 G>C	X:10214067 A>G	X:10220838 G>A	X:10220838 G>A	X:10220838 G>A
Exon number	11	11	11	12	12	12
c.DNA change, NM_001830.4(CLCN4):	c.1937T>C	c.1955G>C	c.1963A>G	c.2153G>A	c.2153G>A	c.2153G>A
Protein change, NP_001821	p.(Ile646Thr)	p.(Arg652Thr)	p.(Ile655Val)	p.(Arg718Gln)	p.(Arg718Gln)	p.(Arg718Gln)
Protein domain* for missense variants	CBS1, intracellular	CBS1, intracellular	CBS1, intracellular	CBS2, intracellular	CBS2, intracellular	CBS2, intracellular
Source :This study = families previously unreported in the medical literature with detailed clinical data; ClinVAR; LOVD and/ or DECIPHER	<i>ClinVar</i> SCV000549938.4	<i>ClinVar</i> SCV000569420.4	<i>ClinVar</i> SCV000549939.4	<i>ClinVar</i> SCV000742750.2	<i>ClinVar</i> SCV001622680.1	<i>ClinVar</i> SCV002139072.1
Gender of proband. Others with variant in family.	NR	NR	NR	NR	1 affected male	NR
Inheritance	NR	NR	NR	NR	Maternally inherited	NR
Recurrent in unrelated families	Yes - but also present in gnomAD	No	No	Yes - but also present in gnomAD	Yes - but also present in gnomAD	Yes - but also present in gnomAD
Assessment of pathogenicity according to ACMG criteria on ClinVAR, or as assessed by authors using Varsome	Benign	VOUS	VOUS	VOUS	LP	VOUS
SIFT (dbNSFP version 4.2); converted rankscore	Deleterious (0.03)	Deleterious (0.04)	Tolerated (0.37)	Tolerated (0.09)	Tolerated (0.09)	Tolerated (0.09)
PolyPhen	Benign (0.158)	Benign (0.224)	Benign (0)	Possibly damaging (0.658)	Possibly damaging (0.658)	Possibly damaging (0.658)
CADD	23.8	23.7	17.72	25.2	25.2	25.2
REVEL	0.868	0.857	0.271	0.779	0.779	0.779
SpliceAI	≤ 0.2	≤ 0.2	≤ 0.2	≤ 0.2	≤ 0.2	≤ 0.2
Frequency in heterozygotes (gnomAD)	11	0	1	3	3	3
Frequency in hemizygotes (gnomAD)	4	0	0	2	2	2
Functional impact in <i>Xenopus</i> oocyte model	WT	WT	WT	WT	WT	WT
Severe functional impact in <i>Xenopus</i> oocyte model	No	No	No	No	No	No
Blended phenotype?	NR	NR	NR	NR	NR	NR
Genetic test	NR	NR	NR	NR	NR	NR

	GROUP E: Truncating variants		
Families	E1	E2	E3
Genomic position and variant, (GRCh38), NC_000023.10	X:10208122 CATCA>C	X:10220667 CCAGA>C	X:10220710 C>G
Exon number	9	9	9
c.DNA change, NM_001830.4(CLCN4):	c.925_928del	c.1987_1990del	c.2025C>G
Protein change, NP_001821	p.(Asn309Profs)	p.(Gln663Glyfs)	p.(Tyr675Ter)
Protein domain* for missense variants	FRAMESHIFT	FRAMESHIFT	NONSENSE
Source :This study = families previously unreported in the medical literature with detailed clinical data; ClinVAR; LOVD and/ or DECIPHER	<i>This study; ClinVar SCV002525743</i>	<i>This study; ClinVar SCV002525744</i>	<i>This study; ClinVar SCV002525745</i>
Gender of proband. Others with variant in family.	1 affected male, mother unaffected	1 affected male, 1 affected maternal uncle, mother unaffected	1 affected male
Inheritance	Maternally inherited	Maternally inherited.	<i>De novo</i>
Recurrent in unrelated families	No	No	No
Assessment of pathogenicity according to ACMG criteria on ClinVAR, or as assessed by authors using Varsome	P	LP	P
SIFT (dbNSFP version 4.2); converted rankscore	NA-frameshift	NA-frameshift	NA-nonsense
PolyPhen	NA-frameshift	NA-frameshift	NA-nonsense
CADD	NA-frameshift	NA-frameshift	36
REVEL	NA-frameshift	NA-frameshift	NA-nonsense
SpliceAI	NA-frameshift	NA-frameshift	≤ 0.2
Frequency in heterozygotes (gnomAD)	0	0	0
Frequency in hemizygotes (gnomAD)	0	0	0
Functional impact in <i>Xenopus</i> oocyte model	Frameshift	Frameshift	Nonsense
Severe functional impact in <i>Xenopus</i> oocyte model	Not tested	Not tested	Not tested
Blended phenotype?	No	No	No
Genetic test	Intellectual disability gene panel of exome backbone	Singleton whole genome sequencing	Trio exome sequencing

Data presented include genomic coordinates, reporting laboratory assessment of pathogenicity using ACMG criteria as reported in ClinVar or determined by the authors using VARSOME prior to functional studies were conducted, demographic details, inheritance, recurrence within families, recurrence across families, including public databases as of 25th May 2022, selected in silico pathogenicity scores, frequency in gnomAD database, functional impact in *Xenopus* oocyte model, and if the individual has more than one genetic diagnosis (blended phenotype). Data which are supportive of pathogenicity is color-coded orange (with darker orange for most supportive data), data which are not supportive of pathogenicity are coded green. ACMG American College of Medical Genetics and Genomics, CBS cystathionine β-synthase, NA not applicable, NR not reported; N-term N terminus, GOF gain of function, LOF loss-of-function, LOVD Leiden Open Variation Database, MPS massively parallel sequencing, WT wild type. In silico scores include PolyPhen, CADD, REVEL and SpliceAI.





Fig. 1 Pedigrees of all previously unreported families with inherited *CLCN4* variants. Filled square/circle = affected individual, lightly shaded circle/square = mildly affected individual, *familial **CLCN4* variant present in affected males, – familial *CLCN4* variant absent in male, */– familial *CLCN4* variant present in female, –/– familial *CLCN4* variant absent in female. Pedigrees of families with a *de novo* variant are not shown.

extracellular pH (Fig. 4E–G). While outward currents of WT CIC-4 were slightly inhibited at acidic pH and inward currents remained undetectable, comparably large inward currents became visible at pH 6.3 and 5.3 for variant p.(Ile549Asn) (Fig. 4D, E). A quantitative analysis revealed that similarly large inward currents were seen for variants p.(Phe268Leu) and p.(Ala555Val) (Fig. 4B, C). Smaller, but highly significant inward currents were also detected for variants p.(Asp89Asn), p.(Pro310Ser), p.(Val317Ile), p.(Val317Phe), p.(Ser395Arg), and p.(Ala550Leu) (Fig. 4F, G and Supplementary Fig. 1). Evidently, these variants partially disrupted the gating process of CIC-4 that normally prevents inward currents even at very acidic pH. For variants p.(Phe268Leu) and p.(Ile549Asn) inward currents were large enough to estimate reversal potentials at pH 6.3 and pH 5.3. The fact that the reversal potential in these conditions differed by about 12.5 mV for both variants (Fig. 4H)

demonstrates that the inward currents carried by these variants are at least partially mediated by H^+ transport. However, the difference falls short of the expected value of ~20 mV for a coupled $2Cl^-/1H^+$ antiporter [32], suggesting that currents mediated by the variants are at least partially uncoupled. More detailed studies will however be needed to determine precise transport stoichiometry for these variants as well as for WT CIC-4.

Group C consisted of families with variants p.(Asp34Asn), p.(Ser69Leu) and p.(Pro369Leu). Although these variants all showed a functional LOF similar to those in Group A (Supplementary Fig. 1), the affected individuals had more complex clinical presentations (Supplementary Fig. 3) and an additional genetic condition was proven or strongly suspected, consistent with a blended phenotype. Consequently, they were separated from the other groups, and not included in the clinical summary in Table 2.

A



B




Fig. 2 Clinical photographs of individuals with previously unreported variants in *CLCN4*, and representative neuroimaging. A Clinical photographs demonstrate that some males and females have progressive lengthening of their face and 'squaring' of the jaw with age. LOF loss-of-function, GOF gain-of-function, ROF reduction of function, m months, y years. **B** Neuroimaging (T1 mid-sagittal view) from affected probands. In all individuals there are abnormalities of the corpus callosum. The proband of Family A10 has a dysplastic corpus callosum: it is of normal length but globally hypoplastic. Family A19: two affected brothers both display complete agenesis of the corpus callosum with colpocephaly. Family B5: the proband has partial agenesis of the corpus callosum (affecting the posterior part and splenium), colpocephaly and mild dilatation of the 3rd ventricle. Family B6: the proband has a dysplastic corpus callosum, and mildly small optic chiasm and optic nerves bilaterally.





Fig. 3 Mapping of all *CLCN4* variants functionally investigated in this study. A Schematic of the *CLCN4* gene and CIC4-protein with position of variants from newly identified families with clearly affected males and females depicted above the schematic, and position of variants published to date shown below the schematic. **B** Position of the investigated missense variants in a CLC topology model. Altered residues are shown as circles and functional effects are color-coded as indicated in the figure. **C** Three-dimensional homology model of the human CIC-4 protein based on the structure of the CmCIC homodimer (Protein Data Bank: 3ORG). The view from within the membrane delimited by dashed lines. The two subunits forming the homodimer are shown in dark and light grey. Mutated residues are shown as spheres colored as in B. Right 3D model viewed from the extracellular site.



Group D consisted of 18 families with rare missense variants with supportive *in silico* pathogenicity scores and/or clinical features suggestive of *CLCN4*-related condition, but for which no functional impact in the *Xenopus* expression system could be demonstrated. This group included variants p.(Asp29Glu),

p.(Ser105Cys), p.(Phe238Leu), p.(Arg315His), p.(Arg364Gly), p.(Arg432Gln), as well as variants located in the intracellular CBS1 and CBS2 domains: the variants p.(Asp621Gly), p.(Thr629Ile), p.(Ile646Thr), p.(Arg652Thr), p.(Ile655Val), and p.(Arg718Gln) (Table 1, Supplementary Fig. 1). It is plausible that these variants

Fig. 4 Expression of *CLCN4* variants in *Xenopus* oocytes. Panel **A** shows example recordings of the indicated constructs evoked by the voltage-clamp protocol indicated in the inset and using a “P/4” leak subtraction protocol (see Methods). Scale bars apply to all constructs. **B** shows average normalized IV relationships of the same variants. Currents are normalized to that of WT at 170 mV as described in Methods. In **C**, currents are normalized to the current of WT at the same voltage (see Methods). Data points are significantly different at practically all voltages from the value of 1 (i.e., WT) for all indicated variants ($p < 0.05$). Panel **D** shows typical current traces recorded without leak subtraction of WT *CIC-4* in the presence of neutral and acidic extracellular pH, with outward currents being inhibited and inward currents remaining at a negligible level [29]. **E** illustrates the pH response of variant p.(Ile549Asn), which shows the activation of relatively large inward currents at acidic pH. **F, G** quantitative analysis of pH dependence of indicated GOF variants. Currents recorded at pH 5.3 were normalized to values measured at pH 7.3 as described in methods. The GOF effect of variants p.(Val317Ile) and p.(Ser395Arg) becomes apparent in panel **G** that shows the same data as panel F at a magnified scale. Data points are significantly different at voltages ≤ -40 mV from WT for all indicated variants ($p < 0.05$). Panel **H** shows differences in reversal potential measured at pH 6.3 and pH 5.3 for variants p.(Phe268Leu) and p.(Ile549Asn). The red line indicates the value expected for a stoichiometrically coupled 2 $\text{Cl}^-/1 \text{H}^+$ antiporter.

are pathogenic by a mechanism not modelled in our cellular system, but given the lack of evidence on their pathogenicity, the families with Group D variants were not included in the summary Table 2.

Group E consisted of three individuals with a frameshift or nonsense variant in *CLCN4* for whom detailed clinical data were available.

This study thus brings the total number of individuals with (likely) pathogenic variants in *CLCN4* to a total of 122: 58 males and 64 females. For 20 of the females, parental studies demonstrated the variant to be *de novo*, while the other 44 females were identified as being heterozygous for a *CLCN4* variant only after a relative (usually a son, but on two occasions a daughter) was identified in their family to have *CLCN4*-related condition [1, 2, 5, 7, 11, 31]. The clinical features of this expanded cohort are summarized in Table 2.

DISCUSSION

Our study addresses the interpretation of novel missense variants, a common clinical conundrum across clinical genetic practice [33]. We robustly demonstrate a much wider range of functional impacts of *CLCN4* variants in the *Xenopus* oocyte model than had been previously demonstrated. In addition, we provide new insights into the common clinical features of *CLCN4*-related neurodevelopmental condition, which have enabled us to provide updated clinical management advice to clinicians [9] and improved patient and family education via the patient advocacy group CureCLCN4.

We confirm that cognitive disability is the most common clinical feature in males, most commonly in the moderate or severe/profound range (Table 2 and Supplementary Table 1). For the first time, however, we report a male with a verbal IQ in the normal range. This 12-year-old male (Family A21; p.(Phe319Ser)mat; functional studies: LOF by shifted voltage dependence) had a verbal IQ of 90 on formal psychometric testing (WISC-II-NL) but did have a lower performance IQ (61: within the mild ID range) and significant comorbidities with delayed language acquisition, articulation difficulties, severe treatment-resistant epilepsy, autistic features, and hyperactivity. We also report the first observation of a male with *CLCN4*-related condition and mild ID who has had a family (Family A3; p.(Val92Met); functional studies: almost complete LOF). He has two daughters who are obligate heterozygotes, one with mild ID and the other with specific learning disabilities.

Phenotypic prediction of cognitive function in females with *CLCN4*-related condition is very difficult with a wide spectrum of severity of neurodevelopmental and medical issues, including about half of heterozygous female carriers being apparently completely unaffected (Table 2). In general, females with a *de novo* variant had a more severe phenotype than those with an inherited variant. However, this observation is far from absolute, as evidenced by several female individuals in the cohort. For example, in families A10 and A24, the mother of a severely

affected male had a *de novo* variant and yet was completely unaffected. On the other hand, we report females with inherited variants who have severe phenotypes: for example, the proband in Family A40 had moderate ID and a missense variant (p.(Pro635Arg); functional studies: LOF), which she inherited from a mother with mild ID. The proband had no additional genetic condition identified by WGS, and there was no evidence of mosaicism in the unaffected mother. We have previously reported that X-inactivation status does not correspond to clinical severity [5], and, as demonstrated across this and previous studies [8, 31], female-to-female inheritance from a very mildly, or even apparently non-affected mother does not ensure a mild phenotype in the daughter. Clinically, were a *de novo* missense *CLCN4* variant to be detected on a prenatal exome in a female embryo, there would remain a degree of uncertainty whether there would be a neurodevelopmental phenotype postnatally. From our previous study, it was apparent that females with a *CLCN4* frameshift or nonsense variant or a small intragenic chromosomal deletion of *CLCN4* are typically unaffected [5]. This was confirmed in the current study: both female carriers in the two families with inherited truncating variants in Group E were unaffected (Family E1 and E2). This observation may signify that the impact of missense variants in females could lead to a toxic gain-of function or a LOF that could be at least partially imparted also on *CIC-3/CIC-4* heterodimers.

Behavioral and mental health disorders are the next most common clinical features. The four most common conditions were attention deficit hyperactivity disorder (ADHD) or significant hyperactivity, impulsiveness, or restlessness affecting 59% of all males and 46.7% of females with *de novo* variants; autism spectrum disorder (or autistic behavior) affecting 54.5% of all males and 40% of females with *de novo* variants; angry outbursts or challenging behaviors, affecting 36.4% of males and 26.7% of females, and lastly anxiety, affecting 27.2% of all males, 53% of females with *de novo* variants and 10.5% of females with inherited variants or variants with unknown inheritance. The mental health conditions were reported to significantly impact the affected individual's ability to learn and their quality of life. Less frequent mental health disorders included obsessive compulsive disorder and depression/ bipolar disorder, which commonly had onset in late teenage years or early adulthood and caused a significant deterioration in quality of life. This highlights the need for close monitoring of all individuals for psychiatric complications, with appropriate referral to a psychiatrist skilled in the management of individuals with neurodevelopmental conditions.

Epilepsy is also confirmed as a significant feature of *CLCN4*-related neurodevelopmental condition, affecting 59% of all males and 20% of females. Most individuals with epilepsy had seizure onset within the first three years of life, although two were diagnosed at age 13, highlighting the need for ongoing seizure surveillance beyond childhood. Seizure semiologies were broad, including generalized absence and tonic-clonic seizures and focal onset seizures, as evidenced by EEG showing focal onset in some and generalized onset in others. Epilepsy can be severe, consistent

Table 2. Summarized clinical features, presented with HPO (Human Phenotype Ontology) nomenclature, of all individuals with a *CLCN4*-related neurodevelopmental condition from this study and from previous reports (in the case that detailed clinical data were available).

Feature	HPO term	Males				Females				Males and females total							
		All variants				De novo variants				Inherited or inheritance unknown variants				All variants			
		Previously reported positive informative/total	This study positive informative/total	Total positive/total	Previously reported positive informative/total	This study positive informative/total	Total positive/total	Previously reported positive informative/total	This study positive informative/total	Total positive/total	Previously reported positive informative/total	This study positive informative/total	Total positive/total	Previously reported positive informative/total	This study positive informative/total	Total positive/total	
Neurodevelopment	Unaffected	0/36 (0%)	0/22 (0%)	0/58 (0%)	2/15 (13.3%)	2/20 (10%)	2/25 (8%)	13/19 (68.4%)	35/44 (79.5%)	37/64 (57.8%)	37/122 (30.3%)						
	Borderline intellectual disability	1/36 (2.8%)	0/22 (0%)	1/58 (1.7%)	0/15 (0%)	1/20 (5%)	0/25 (0%)	0/19 (0%)	0/44 (0%)	1/64 (1.6%)	2/122 (1.6%)						
	Mild intellectual disability	9/36 (25%)	4/22 (18.2%)	13/58 (22.4%)	6/15 (40%)	6/20 (30%)	1/25 (4%)	5/19 (26.3%)	6/44 (13.6%)	12/64 (18.8%)	25/122 (20.5%)						
	Moderate intellectual disability	9/36 (25%)	9/22 (41%)	18/58 (31%)	5/15 (33.3%)	7/20 (35%)	0/25 (0%)	1/19* (5.3%)	1/44 (2.3%)	8/64 (12.5%)	26/122 (21.3%)						
	Severe/profound intellectual disability	17/36 (47.2%)	7/22 (31.8%)	24/58 (41.4%)	2/15 (13.3%)	4/20 (20%)	2/25 (8%)	0/19 (0%)	2/44 (4.5%)	6/64 (9.4%)	30/122 (24.6%)						
	Specific learning disability	0/36 (0%)	2/22 (9.1%)	2/58 (3.4%)	1/15 (6.7%)	1/20 (5%)	0/25 (0%)	0/19 (0%)	0/44 (0%)	1/64 (1.6%)	3/122 (2.5%)						
	Delayed speech and language	36/36 (100%)	22/22 (100%)	58/58 (100%)	14/15 (93.3%)	19/20 (95%)	NA	6/19 (31.6%)	6/44 (13.6%)	25/64 (39%)	83/122 (68%)						
	Epilepsy	22/36 (61.1%)	13/22 (59.1%)*	35/58 (60.3%)	3/15 (20%)	5/20 (25%)	1/25 (4%)	1/19 (5.3%)	2/44 (4.5%)	7/64 (10.9%)	42/122 (34.4%)						
	Well-controlled epilepsy	8/22 (36.4%)	10/13 (77%)	18/35 (51.4%)	3/3 (100%)	4/5 (80%)	0/1 (0%)	1/1 (100%)	1/2 (50%)	5/7 (71.4%)	23/42 (54.8%)						
	Treatment-resistant epilepsy	12/22 (54.5%)	3/13 (23.1%)	15/35 (42.9%)	0/3 (0%)	1/5 (20%)	1/1 (100%)	0/1 (0%)	1/2 (50%)	2/7 (28.6%)	17/42 (40.5%)						
Neurology	Information about seizure control not available	2/22 (9.1%)	0/13 (0%)	2/35 (5.7%)	0/3 (0%)	0/5 (0%)	0/1 (0%)	0/1 (0%)	0/2 (0%)	0/7 (0%)	2/42 (4.8%)						
	Infantile hypotonia/ neonatal hypotonia	11/36 (31%)	12/22 (54.5%)	23/58 (39.6%)	8/15 (53.3%)	11/20 (55%)	1/18 (5.6%)	2/19 (10.5%)	3/37 (8.1%)	14/57 (24.6%)	37/115 (32.2%)						
	Progressive neurological manifestations	8/36 (22.2%)	7/22 (31.8%)	15/58 (25.9%)	6/15 (40%)	7/20 (35%)	2/18 (11.1%)	1/19 (5.3%)	3/37 (8.1%)	10/57 (17.5%)	25/115 (21.7%)						
	Abnormality of the brain	14/18 (77.8%)	10/17 (58.8%)	24/35 (68.6%)	8/13 (61.5%)	10/17 (58.8%)	1/1 (100%)	2/2 (100%)	3/3 (100%)	13/20 (65%)	37/55 (67.3%)						
	Abnormality of white matter (e.g. white matter hyperintensities/ periventricular leukomalacia/ delayed or abnormal myelination)	11/18 (61.1%)	9/17 (52.9%)	20/35 (57.1%)	3/13 (23.1%)	5/17 (29.4%)	1/1 (100%)	0/2 (0%)	1/3 (33.3%)	6/20 (30%)	26/55 (47.3%)						
	Abnormality of the corpus callosum	6/18 (33%)	8/17 (47%)	14/35 (40%)	3/13 (23.1%)	4/17 (23.5%)	1/1 (100%)	0/2 (0%)	1/3 (33.3%)	5/20 (25%)	19/55 (34.5%)						
	Cerebral and/ or cerebellar atrophy	6/18 (33%)	2/17 (11.8%)	8/35 (22.8%)	5/13 (38.5%)	7/17 (41.2%)	1/1 (100%)	0/2 (0%)	1/3 (33.3%)	8/20 (40%)	16/55 (29.1%)						
	Other abnormality of the brain, e.g. Cortical dysplasia/ sclerosis/ cortical hyperintensities	1/18 (5.5%)	1/17 (5.9%)	2/35 (5.7%)	3/13 (23.1%)	4/17 (23.5%)	0/1 (0%)	2/2 (100%)	2/3 (66.7%)	6/20 (30%)	8/55 (14.5%)						
	Autism spectrum disorder or autistic behavior	2/36 (5.5%)	12/22 (54.5%)	14/58 (24.1%)	6/15 (40%)	6/20 (30%)	1/18 (5.5%)	4/19 (21%)	5/37 (13.5%)	11/57 (19.3%)	25/115 (21.7%)						
	Depression/ bipolar disorder	4/36 (11.1%)	1/22 (4.5%)	5/58 (8.6%)	1/15 (6.7%)	1/20 (5%)	1/18 (5.5%)	2/19 (10.5%)	3/37 (8.1%)	4/57 (7%)	9/115 (7.8%)						
Psychiatry	Anxiety	4/36 (11.1%)	6/22 (27.3%)	10/58 (17.2%)	8/15 (53.3%)	9/20 (45%)	1/18 (5.5%)	2/19 (10.5%)	3/37 (8.1%)	12/57 (21.1%)	22/115 (19.1%)						
	Obsessive and or compulsive behaviors	2/36 (5.6%)	4/22 (18.2%)	6/58 (10.3%)	2/15 (13.3%)	2/20 (10%)	0/18 (0%)	1/19 (5.3%)	1/37 (2.7%)	3/57 (5.3%)	9/115 (7.8%)						
	Attention Deficit Hyperactivity Disorder/ or significant hyperactivity/ restlessness/ impulsivity	4/36 (11.1%)	13/22 (59.1%)	17/58 (29.3%)	7/15 (46.7%)	7/20 (35%)	0/18 (0%)	3/19 (15.8%)	3/37 (8.1%)	10/57 (17.5%)	27/115 (23.5%)						

Table 2. continued

Feature	HPO term	Males		Females				Males and females total	
		All variants		De novo variants		Inherited or inheritance unknown variants		All variants	
		Previously reported positive informative/total	This study positive informative/total	Previously reported positive informative/total	This study positive informative/total	Previously reported positive informative/total	This study positive informative/total	Total positive/total informative	Total positive/total informative
Psychotic disorder	HP:0000709	0/36 (0%)	1/22 (4.5%)	0/5 (0%)	0/15 (0%)	0/18 (0%)	0/19 (0%)	0/37 (0%)	2/115 (1.7%)
	HP:0000718	8/36 (22.2%)	8/22 (36.4%)	2/5 (40%)	4/15 (26.7%)	0/18 (0%)	1/19 (5.3%)	1/37 (2.7%)	23/115 (20%)
	HP:0002020	1/36* (2.8%)	8/22 (36.4%)	0/5 (0%)	5/15 (33.3%)	0/18 (0%)	1/19 (5.3%)	1/37 (2.7%)	15/115 (13%)
Gastrointestinal and growth	HP:0002019	0/36 (0%)	8/22 (36.4%)	1/5 (20%)	6/15 (40%)	0/18 (0%)	1/19 (5.3%)	1/37 (2.7%)	16/115 (13.9%)
	HP:0011968	2/36 (5.6%)	8/22 (36.4%)	3/5 (60%)	8/15 (53.3%)	0/18 (0%)	3/19 (15.8%)	3/37 (8.1%)	24/115 (20.9%)
	HP:0005484	5/28 (17.8%)	4/20 (20%)	2/5 (40%)	9/11 (81.8%)	0/18 (0%)	1/19 (5.3%)	1/37 (2.7%)	21/115 (18.3%)
Failure to thrive	HP:0001508	2/27 (7.4%)	4/18 (22.2%)	0/5 (0%)	5/13 (38.5%)	0/18 (0%)	1/19 (5.3%)	1/37 (2.7%)	12/100 (12%)
	HP:0004322	1/27 (3.7%)	6/22 (27.2%)	1/4 (20%)	4/13 (30.7%)	0/18 (0%)	1/19 (5.3%)	1/37 (2.7%)	13/103 (12.6%)
	HP:0002360	0/36 (0%)	2/22 (9.1%)	1/5 (20%)	3/13 (23.1%)	0/18 (0%)	1/19 (5.3%)	1/37 (2.7%)	7/113 (6.2%)
Scoliosis/ kyphosis	HP: 0010674	3/36 (8.3%)	1/22 (4.5%)	1/5 (20%)	0/13 (0%)	0/18(0%)	0/19 (0%)	0/37 (0%)	5/113 (4.4%)
	HP:0001763;	1/36 (2.8%)	7/22 (31.8%)	2/5 (40%)	1/13 (7.7%)	0/18 (0%)	0/19 (0%)	0/37 (0%)	11/113 (9.7%)
	HP:0030984;								
Hearing impairment	HP:0000403;	0/36 (0%)	6/22 (27.3%)	0/5 (0%)	2/15 (13.3%)	0/18 (0%)	1/19 (5.3%)	0/37 (0%)	8/113 (7.1%)
	HP:0000407								
	HP:0000486;	0/36 (0%)	6/22 (27.3%)	0/5 (0%)	3/18 (16.7%)	0/18 (0%)	2/19 (10.5%)	0/37 (0%)	9/113 (8%)
Vision impairment	HP:0000558;								
	HP:0000545;								
	HP:0000505								

This table excludes patients with rare missense variants from Group D, for whom the functional studies were similar to wild type, and patients from Group C that had a more severe phenotype due to an additional monogenic condition.

with a developmental and epileptic encephalopathy as highlighted in recent reports [1, 11]. The severity of epilepsy, however, does not necessarily correlate with the severity of cognitive impairment. Due to *CLCN4* being an antiporter of protons and chloride, which may be important in acid-base balance, acetazolamide has been trialed without any clear evidence of improvement in seizure control. Indeed, no specific anti-seizure medications have been demonstrated to best correlate with epilepsy control.

Neuroimaging showed abnormalities in 58.8% of males and 61.5% of females, most commonly of the white matter. Two brothers and one female had complete agenesis of the corpus callosum. This suggests that *CLCN4* should be added to panels of genes interrogated in individuals with corpus callosum abnormalities [34, 35].

Infantile hypotonia was reported in about half of all males and of females with *de novo* variants in this cohort. Progressive microcephaly was more common in females with *de novo* variants (80.8%) compared to males (20%). 31.8% of males and 40% of females with *de novo* variants had later onset neurological symptoms including tremor, ataxia, hyperkinesia or stereotypical movements, changes in gait such as walking with a stooped posture, or progressive spasticity.

Functional gastrointestinal symptoms, such as gastroesophageal reflux and constipation, were common in females with *de novo* variants (33.3% had gastroesophageal reflux and 40% had constipation) and impacted also a significant proportion of males (36.4% had gastroesophageal reflux and 36.4% had constipation). A small proportion of individuals, particularly those with GOF variants, have a striking growth phenotype. All four females with the recurrent *de novo* p.(Ala555Val) variant, for whom clinical data were available (Families B12-B15) had severe symmetrical growth restriction and feeding difficulties, two requiring gastrostomy feeds. The female proband from Family B13 was investigated by a pediatric endocrinologist, without evidence of growth hormone deficiency. The cause of this growth restriction requires further study but may reflect roles of the CIC-4 protein in fundamental growth processes or impact on enteric neurological function. Our findings underscore the importance of involving neurogastroenterology specialists in the comprehensive management of children with neurodevelopmental conditions, due to the significant impact on quality of life of underrecognized and untreated functional gastrointestinal comorbidities [36].

Other, less commonly noted clinical features include scoliosis, pes planus and/or lax joints, sleep disorders, otitis media with effusions, and strabismus. However, to date, our and other studies suggest that non-neurological congenital anomalies outside of the neurological system are not core features of *CLCN4*-related condition. With age, as previously described, there is a progressive lengthening of the face in males and females, with some males having a relatively 'square' jaw [5] (Fig. 2). Facial features in infancy and childhood are variable, without a recognizable 'gestalt'.

With a larger cohort now functionally characterized, we examined whether distinct functional impacts of the CIC-4 variants correlated with phenotypic features. Some early observations could be made. Firstly, the GOF variants (Group B) were commonly associated with a severe growth, feeding and/or functional gastrointestinal component. Secondly, they had a higher female:male ratio; 73% of the affected individuals in Group A (LOF) were male, compared to only 41% in Group B (GOF). Thirdly, all three males with GOF variants had the same variant (p.(Val317Ile)): in two of these families the variant was *de novo*, in one maternally inherited. These males had similar clinical phenotypes including moderate to severe global developmental delay or ID, visual impairment (two were proven to have optic atrophy) and abnormalities of the corpus callosum. The functional impact of this variant was milder than that of the other GOF variants present

in females. A possibility is that a severe GOF variant may not be compatible with life in a hemizygous male.

We cannot yet discount the pathogenicity of variants which performed like WT in our cellular model, as this is far from a complete model of the complexity of CIC-4 in animals in vivo, and, more specifically in the developing human. For example, variants that behaved like WT included the rare p.(Arg315His) *de novo* variant in a female (Family D5), who had clinical features entirely consistent with the spectrum seen in *CLCN4*-associated neurodevelopmental condition: however, this variant has also been reported in two other unrelated families in gnomAD. We also could not demonstrate a functional impact for several variants in the distal CBS domain, although it is possible that these variants may impact protein sorting or other mechanisms unable to be evaluated with the current *Xenopus* oocyte model.

In a structural model of a homodimeric CIC-4 protein, most variants characterized by a LOF with "rightward shifted voltage dependence" are localized at or near the dimer interface. This observation agrees with the hypothesis that voltage-dependent gating of CIC-4 is associated with a rearrangement of the dimer interface, as has been proposed for gating of the lysosomal CIC-7 [37]. Similarly, most GOF variants cluster at the dimer interface, mostly close to the luminal side. These mutants appear to partially destabilize the gate of the transporter that evidently must be tightly closed at negative voltages for proper function in endosomes. Interestingly, the isoleucine mutated in variant p.(Ile549Asn) (Family B10, severely affected female) that shows a particularly large GOF corresponds with Ile607 in the highly homologous CIC-3 protein; a variant at this position in *CLCN3* (p.(Ile607Thr)) similarly caused a dramatic GOF and the affected individual died within the first month of life. It is important to note that CIC-4 most likely forms heterodimeric complexes with CIC-3 [13]. Overall, the disease phenotypes caused by *CLCN3* and *CLCN4* variants are quite different, demonstrating that the two genes have overlapping but not identical functions. Our previous investigations on *CLCN4* missense variants which were found in heterozygous females did not support a potential dominant negative effect when equal amounts of WT and mutant CIC-4 were co-expressed in *Xenopus* oocytes [5]. However, the effect of voltage-gated shifted variants as well as GOF variants in heterodimeric CIC-3/CIC-4 complexes remains to be investigated [14]. Interestingly, the recurrent GOF variant p.(Tyr553Cys) in the late-endosomal CIC-6 causes a marked leftward-shift of the gating process [19, 38]. The corresponding tyrosine residue in CIC-4 is located just one residue away from Ile549. Both residues are in the linker connecting helices P and Q. The dramatic functional alterations of these variants provide additional evidence for a critical role of the linker P-Q in CLC transporter gating and corroborate the hypothesis that the GOF variants of vesicular CLCs are associated with a disrupted gating process.

We attempted to look at the possible impact of mosaicism on the phenotypic severity of *CLCN4* variants, but data are too scarce to robustly conclude that mosaicism for a *CLCN4* variant is predictive of phenotypic expression in females or males. This may be due to the lack of knowledge between the level of mosaicism in blood to that in the brain. For example, the variant p.(Arg718Trp), in the CBS2 domain, has now been reported *de novo* in both males and females with a severe phenotype (Table 1, Supplementary information), as well as in one unaffected mother, reported by He et al. [11]. However, we do note that this unaffected mother had a mosaic karyotype (47,XXX/46,XX) and it is possible that the 'extra' X chromosome may have somewhat moderated her phenotypic expression, as we considered for the unaffected male with Klinefelter syndrome, with an inherited *CLCN4* variant which resulted in a severe phenotype in his male relatives [5].

We report on four individuals (C1-C4) with a *de novo* or inherited missense *CLCN4* variant and supportive functional

studies, but a more complex clinical phenotype, which we could attribute to a likely, or confirmed, blended genotype due to two monogenic conditions. For example, the male proband in Family C1 (p.(Asp34Asn)); whose functional studies were consistent with a LOF of *CIC-4*, has short stature and distinctive skeletal and facial features consistent with a diagnosis of Desbuquois dysplasia (*XYLT1*-related) that he shares with his sister. However, he has significant ID, epilepsy, and autism spectrum disorder, which are atypical for Desbuquois syndrome, and thus most likely has a blended phenotype of Desbuquois dysplasia and *CLCN4*-related neurodevelopmental condition. The finding of four patients with a blended phenotype due to suspected or proven multi-locus pathogenic variation in a total cohort of 122 individuals with *CLCN4* variants (4/122: 3.3%) is consistent with other studies estimating this phenomenon occurs in about 5% of individuals with an identified diagnosis after unbiased sequencing [39]. It emphasizes that for affected individuals, whose clinical features are not entirely consistent with their diagnosed monogenic condition, broadening the scope of genomic sequencing to an unbiased exome or whole genome sequencing approach may be appropriate to look for additional pathogenic findings

In summary, our study considerably expands our knowledge of the range of phenotypic and genotypic variation in *CLCN4*-related condition and for the first time robustly demonstrates a range of functional impacts, including gain of function. Variant classification still remains a nuanced art, rather than a precise science [40]. Fully informed genetic counselling is required to guide families through the diagnostic limitations and uncertainties inherent in genetic testing for neurodevelopmental conditions [41]. Several research priorities remain. We need to better ascertain the causality of all rare missense variants to elucidate targeted treatments. Establishment of a robust animal model is an urgent priority. This could potentially be a rat model, given that *Cln4* is on the X chromosome in the rat, as opposed to in the mouse where it is autosomal [42]. A high throughput functional and therapeutic assay system, such as neuronal micro-electrode assays, which have been successfully applied in other neurodevelopmental conditions [43], would also be very helpful. With recent inclusion of *CLCN4* in the SFARI gene project [44] scientists and clinicians are working together to better understand and manage this condition.

REFERENCES

- Veeramah KR, Johnstone L, Karafet TM, Wolf D, Sprissler R, Salogiannis J, et al. Exome sequencing reveals new causal mutations in children with epileptic encephalopathies. *Epilepsia* 2013;54:1270–81.
- Hu H, Haas SA, Chelly J, Van Esch H, Raynaud M, de Brouwer AP, et al. X-exome sequencing of 405 unresolved families identifies seven novel intellectual disability genes. *Mol Psychiatry*. 2016;21:133–48.
- Raynaud M, Gendrot C, Dessay B, Moncla A, Ayrault AD, Moizard MP, et al. X-linked mental retardation with neonatal hypotonia in a French family (MRX15): gene assignment to Xp11.22-Xp21.1. *Am J Med Genet*. 1996;64:97–106.
- Claes S, Vogels A, Holvoet M, Devriendt K, Raeymaekers P, Cassiman JJ, et al. Regional localization of two genes for nonspecific X-linked mental retardation to Xp22.3-p22.2 (MRX49) and Xp11.3-p11.21 (MRX50). *Am J Med Genet*. 1997;73:474–9.
- Palmer EE, Stuhlmann T, Weinert S, Haan E, Van Esch H, Holvoet M, et al. De novo and inherited mutations in the X-linked gene *CLCN4* are associated with syndromic intellectual disability and behavior and seizure disorders in males and females. *Mol Psychiatry*. 2018;23:222–30.
- Dingemans AJM, Stremmelaar DE, Vissers L, Jansen S, Nabais Sa MJ, van Remortele A, et al. Human disease genes website series: An international, open and dynamic library for up-to-date clinical information. *Am J Med Genet A* 2021;185:1039–46.
- Guo YX, Ma HX, Zhang YX, Chen ZH, Zhai QX. Whole-exome sequencing for identifying genetic causes of intellectual developmental disorders. *Int J Gen Med*. 2021;14:1275–82.
- Xu X, Lu F, Zhang L, Li H, Du S, Tang J. Novel *CLCN4* variant associated with syndromic X-linked intellectual disability in a Chinese girl: a case report. *BMC Pediatr*. 2021;21:384.
- Adam MP, Ardinger HH, Pagon RA, Wallace SE, Bean LJM, Gripp KW, et al. *GeneReviews*. 1993.
- Palmer EE, Sachdev R, Macintosh R, Melo US, Mundlos S, Righetti S, et al. Diagnostic yield of whole genome sequencing after nondiagnostic exome sequencing or gene panel in developmental and epileptic encephalopathies. *Neurology* 2021;96:e1770–e82.
- He H, Guzman RE, Cao D, Sierra-Marquez J, Yin F, Fahlke C, et al. The molecular and phenotypic spectrum of *CLCN4*-related epilepsy. *Epilepsia* 2021;62:1401–15.
- Jentsch TJ, Pusch M. CLC chloride channels and transporters: structure, function, physiology, and disease. *Physiol Rev*. 2018;98:1493–590.
- Guzman RE, Bungert-Plumke S, Franzen A, Fahlke C. Preferential association with *CIC-3* permits sorting of *CIC-4* into endosomal compartments. *J Biol Chem*. 2017;292:19055–65.
- Weinert S, Gimber N, Deuschel D, Stuhlmann T, Puchkov D, Farsi Z, et al. Uncoupling endosomal CLC chloride/proton exchange causes severe neurodegeneration. *EMBO J*. 2020;39:e103358.
- Polovitskaya MM, Barbini C, Martinelli D, Harms FL, Cole FS, Calligari P, et al. A Recurrent Gain-of-Function mutation in *CLCN6*, encoding the *CIC-6* Cl(-)/H(+)-Exchanger, causes early-onset neurodegeneration. *Am J Hum Genet*. 2020;107:1062–77.
- Duncan AR, Polovitskaya MM, Gaitan-Penas H, Bertelli S, VanNoy GE, Grant PE, et al. Unique variants in *CLCN3*, encoding an endosomal anion/proton exchanger, underlie a spectrum of neurodevelopmental disorders. *Am J Hum Genet*. 2021;108:1450–65.
- Poet M, Kornak U, Schweizer M, Zdebek AA, Scheel O, Hoelzer S, et al. Lysosomal storage disease upon disruption of the neuronal chloride transport protein *CIC-6*. *Proc Natl Acad Sci USA*. 2006;103:13854–9.
- Kasper D, Planells-Cases R, Fuhrmann JC, Scheel O, Zeitz O, Ruether K, et al. Loss of the chloride channel *CIC-7* leads to lysosomal storage disease and neurodegeneration. *EMBO J*. 2005;24:1079–91.
- Zifarelli G, Pusch M, Fong P. Altered voltage-dependence of slowly activating chloride-proton antiporter by late endosomal *CIC-6* explains distinct neurological disorders. *J Physiol*. 2022;600:2147–64.
- Stobrawa SM, Breiderhoff T, Takamori S, Engel D, Schweizer M, Zdebek AA, et al. Disruption of *CIC-3*, a chloride channel expressed on synaptic vesicles, leads to a loss of the hippocampus. *Neuron* 2001;29:185–96.
- Rickheit G, Wartosch L, Schaffer S, Stobrawa SM, Novarino G, Weinert S, et al. Role of *CIC-5* in renal endocytosis is unique among *CIC* exchangers and does not require PY-motif-dependent ubiquitylation. *J Biol Chem*. 2010;285:17595–603.
- Firth HV, Richards SM, Bevan AP, Clayton S, Corpas M, Rajan D, et al. DECIPHER: Database of chromosomal imbalance and phenotype in humans using ensembl resources. *Am J Hum Genet*. 2009;84:524–33.
- Landrum MJ, Kattman BL. *ClinVar* at five years: Delivering on the promise. *Hum Mutat*. 2018;39:1623–30.
- Fokkema IF, Taschner PE, Schaafsma GC, Celli J, Laros JF, den Dunnen JT. *LOVD v.2.0*: the next generation in gene variant databases. *Hum Mutat*. 2011;32:557–63.
- Lorenz C, Pusch M, Jentsch TJ. Heteromultimeric CLC chloride channels with novel properties. *Proc Natl Acad Sci USA*. 1996;93:13362–6.
- Traverso S, Zifarelli G, Aiello R, Pusch M. Proton sensing of CLC-0 mutant E166D. *J Gen Physiol*. 2006;127:51–65.
- Leisle L, Ludwig CF, Wagner FA, Jentsch TJ, Stauber T. *CIC-7* is a slowly voltage-gated 2Cl(-)/1H(+)-exchanger and requires *Ostm1* for transport activity. *EMBO J* 2011;30:2140–52.
- De Stefano S, Pusch M, Zifarelli G. A single point mutation reveals gating of the human *CIC-5* Cl(-)/H(+) antiporter. *J Physiol*. 2013;591:5879–93.
- Friedrich T, Breiderhoff T, Jentsch TJ. Mutational analysis demonstrates that *CIC-4* and *CIC-5* directly mediate plasma membrane currents. *J Biol Chem*. 1999;274:896–902.
- Piccolo A, Pusch M. Chloride/proton antiporter activity of mammalian CLC proteins *CIC-4* and *CIC-5*. *Nature* 2005;436:420–3.
- Zhou P, He N, Zhang JW, Lin ZJ, Wang J, Yan LM, et al. Novel mutations and phenotypes of epilepsy-associated genes in epileptic encephalopathies. *Genes Brain Behav*. 2018;17:e12456.
- Accardi A, Miller C. Secondary active transport mediated by a prokaryotic homologue of *CIC* Cl(-) channels. *Nature* 2004;427:803–7.
- Klee EW, Cousin MA, Pinto E, Vairo F, Morales-Rosado JA, Macke EL, et al. Impact of integrated translational research on clinical exome sequencing. *Genet Med*. 2021;23:498–507.
- Palmer EE. Dissecting the clinical outcome and cause of abnormalities of the corpus callosum. *Dev Med Child Neurol*. 2016;58:430–1.
- Palmer EE, Mowat D. Agenesis of the corpus callosum: A clinical approach to diagnosis. *Am J Med Genet Part C: Semin Med Genet*. 2014;166:184–97.
- Drossman DA, Hasler WL. Rome IV-Functional GI Disorders: Disorders of gut-brain interaction. *Gastroenterology* 2016;150:1257–61.

37. Ludwig CF, Ullrich F, Leisle L, Stauber T, Jentsch TJ. Common gating of both CLC transporter subunits underlies voltage-dependent activation of the 2Cl⁻/1H⁺-exchanger ClC-7/Ostm1. *J Biol Chem*. 2013;288:28611–9.
38. Polovitskaya MM, Barbini C, Martinelli D, Harms FL, Cole FS, Calligari P, et al. A recurrent Gain-of-Function Mutation in CLCN6, Encoding the ClC-6 Cl. *Am J Hum Genet*. 2020;107:1062–77.
39. Posey JE, Harel T, Liu P, Rosenfeld JA, James RA, Coban Akdemir ZH, et al. Resolution of disease phenotypes resulting from multilocus genomic variation. *N Engl J Med*. 2017;376:21–31.
40. Richards S, Aziz N, Bale S, Bick D, Das S, Gastier-Foster J, et al. Standards and guidelines for the interpretation of sequence variants: a joint consensus recommendation of the American College of Medical Genetics and Genomics and the Association for Molecular Pathology. *Genet Med*. 2015;17:405–24.
41. Menke C, Nagaraj CB, Dawson B, He H, Tawde S, Wakefield EG. Understanding and interpretation of a variant of uncertain significance (VUS) genetic test result by pediatric providers who do not specialize in genetics. *J Genet Couns*. 2021;30:1559–69.
42. Palmer S, Perry J, Ashworth A. A contravention of Ohno's law in mice. *Nat Genet*. 1995;10:472–6.
43. Mossink B, Verboven AHA, van Hugte EJH, Klein Gunnewiek TM, Parodi G, Linda K, et al. Human neuronal networks on micro-electrode arrays are a highly robust tool to study disease-specific genotype-phenotype correlations in vitro. *Stem Cell Rep*. 2021;16:2182–96.
44. Banerjee-Basu S, Packer A. SFARI Gene: an evolving database for the autism research community. *Dis Model Mech*. 2010;3:133–5.

ACKNOWLEDGEMENTS

The authors would like to thank the individuals and their families who participated in this study. We thank the families and members of the Australia Disorders of the Corpus Callosum (AusDoCC) for their support and time in being involved in this research. We thank T. Jentsch for providing plasmids. For one of the families this study was made possible through access to the data and findings generated by the 100,000 Genomes Project. The 100,000 Genomes Project is managed by Genomics England Limited (a wholly owned company of the Department of Health). The 100,000 Genomes Project is funded by the National Institute for Health Research and NHS England. The Wellcome Trust, Cancer Research UK, and the Medical Research Council have also funded research infrastructure. The 100,000 Genomes Project uses data provided by patients and collected by the National Health Service as part of their care and support. This study makes use of data generated by the DECIPHER community. A full list of centres who contributed to the generation of the data is available from <https://deciphergenomics.org/about/stats> and via email from contact@deciphergenomics.org. Funding for the DECIPHER project was provided by Wellcome.

AUTHOR CONTRIBUTIONS

CF, MHN, AM, MN, BC, CB, FSA, AC, MOH, HS, SW, AV, BC, SR, KN, SA, MR., CSM, CW-M, KJ, MM, DB, ND, MG, TBH, EC, AMc, DH, ST, MW, LJR, CS, GC, LD, RM-L, TD-B, JB, CS, EF, SEC, M-AS, AP, BG, M-TAW, GR, CM, SD, SB, CA, JMB, TTS, GNW, EJS, LM, DL, RS, RM, OM, FC, MC, LR, MHW, CWO, RP, SDK, MF, FERL, AMF, ARS, VM, SN, SG, DDW, LMB, JF, VC, SJ, LP, PMC, MB, EKB, JAR, CB, ZP, KMcW, TB, ET, MMA, SSM, and RA were responsible for compilation of genetic and clinical information and critical review and approval of manuscript. AP, VS, JG, AH, LS, and DK, were responsible for performing experimental work and data analysis, and approval of manuscript. EP, MP, and VMK were responsible for conceiving the idea of the study, performing experimental work and or data analysis, drafting and finalizing, and approval of the manuscript.

FUNDING

MP received funding from the Fondazione AIRC per la Ricerca sul Cancro (grant # IG 21558) and from the Italian Ministry for University and Research (MIUR) (PRIN 20174TB8KW). SW received support from FWO (1861419N), and the Queen Elisabeth Medical Foundation. TBH was supported by the Deutsche Forschungsgemeinschaft (DFG, German Research Foundation)—418081722, 433158657. SSM received support from CPA grant PG01217. LJR was supported by an NHMRC Principal Research Fellowship (GNT1120615) and received support from BICARE, Australia. This work was funded in part by King Salman Center for Disability Research through Research Group no RG-2022-010 and RG-2022-011. This work is partly funded by the NIHR GOSH BRC. The views expressed are those of the author(s) and not necessarily those of the NHS, the NIHR or the Department of Health. AMc is funded by MRC (MR/T007087/1), GOSH Charity (VS0122), and Rosetrees Trust (CF2/100018). EP is supported by a NHMRC Investigator Grant (GNT20081). Open Access funding enabled and organized by Projekt DEAL.

COMPETING INTERESTS

SW received consultancy fees from UCB, Biocodex, Xenon Pharmaceuticals, Zogenix, Lundbeck, Knopp Biosciences, and Uncod Therapeutics. KMc, TB, and ET are employees of GeneDx. ZP is an employee of Ambyr. CB is an employee of Centogene, GmbH. The Department of Molecular and Human Genetics at Baylor College of Medicine receives revenue from clinical genetic testing completed at Baylor Genetics Laboratories. All other authors declare no competing interests.

ADDITIONAL INFORMATION

Supplementary information The online version contains supplementary material available at <https://doi.org/10.1038/s41380-022-01852-9>.

Correspondence and requests for materials should be addressed to Elizabeth E. Palmer, Michael Pusch or Vera M. Kalscheuer.

Reprints and permission information is available at <http://www.nature.com/reprints>

Publisher's note Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.



Open Access This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons licence, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons licence, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons licence and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this licence, visit <http://creativecommons.org/licenses/by/4.0/>.

© The Author(s) 2022

¹Centre for Clinical Genetics, Sydney Children's Hospital Network, Randwick, NSW, Australia. ²Discipline of Paediatrics and Child Health, Faculty of Medicine and Health, University of New South Wales, Randwick, NSW, Australia. ³Istituto di Biofisica, CNR, Genova, Italy. ⁴Department of Clinical Genetics, Liverpool Hospital, Liverpool, NSW, Australia. ⁵Max Planck Institute for Molecular Genetics, Group Development and Disease, Berlin, Germany. ⁶Department of Biomedical and Clinical Sciences, Linköping University, Linköping 581 83, Sweden. ⁷Department of Human Genetics, Gilbert and Rose-Marie Chagoury School of Medicine, Lebanese American University, Byblos, Lebanon. ⁸Institut Jerome Lejeune, Paris, France. ⁹Service de Génétique Médicale, CHU de Nantes, Nantes Université, Nantes, France. ¹⁰Nantes Université, CNRS, INSERM, l'Institut du Thorax, Nantes, France. ¹¹Department of Translational Genomics, Center for Genomic Medicine, King Faisal Specialist Hospital and Research Center, Riyadh, Saudi Arabia. ¹²Department of Neurosciences, King Faisal Specialist Hospital and Research Center, Riyadh, Saudi Arabia. ¹³Applied and Translational Neurogenomics Group, VIB Center for Molecular Neurology, VIB, Antwerp, Belgium. ¹⁴Neurology Department, Antwerp University Hospital, Antwerp, Belgium. ¹⁵Translational Neurosciences, Faculty of Medicine and Health Science, University of Antwerp, Antwerp, Belgium. ¹⁶Department of Child Neurology & Metabolism, Ghent University Hospital, Ghent, Belgium. ¹⁷Department of Pediatric Neurology, Antwerp University Hospital, University of Antwerp, Antwerp, Belgium. ¹⁸Children's Hospital at Westmead, Sydney Children's Hospitals Network, Sydney, Australia. ¹⁹Service de Génétique Clinique, Centre Hospitalier Régional Universitaire de Tours, Tours, France. ²⁰Genetic Center, Akron Children's Hospital, Akron, OH, USA. ²¹Center of Medical Genetics, University Hospital Antwerp/University of Antwerp, Edegem, Belgium. ²²Department of Pediatric Neurology, University Hospital Antwerp/University of Antwerp, Edegem, Belgium. ²³Institute of Human Genetics, Heidelberg University, Heidelberg, Germany. ²⁴Institute of Medical Genetics and Applied Genomics, University of Tuebingen, Tuebingen, Germany. ²⁵Department of Clinical Genetics, Great Ormond Street Hospital for Children, London, UK. ²⁶Developmental Neurosciences, UCL Great Ormond Street Institute of Child Health, London, UK. ²⁷Department of Neurology, Great Ormond Street Hospital, London, UK. ²⁸Wessex Clinical Genetics Service, Princess Anne Hospital, Southampton, UK. ²⁹Genetic Services of WA, King Edward Memorial Hospital, Subiaco, WA, Australia.

³⁰Department of Neuroscience, Washington University in St Louis School of Medicine, St Louis, MI, USA. ³¹The University of Queensland, Queensland Brain Institute, St Lucia, QLD, Australia. ³²Centre for Population Genomics, Murdoch Children's Research Institute, Melbourne, Australia. ³³Garvan Institute of Medical Research, UNSW, Sydney, NSW, Australia. ³⁴Division of Clinical and Metabolic Genetics, The Hospital for Sick Children, Toronto, ON, Canada. ³⁵Genetics of Learning Disability Service, Newcastle, NSW, Australia. ³⁶University of Newcastle Grow Up Well Priority Research Centre, Newcastle, NSW, Australia. ³⁷Department of Pathology and Laboratory Medicine, Children's Mercy Hospital and Clinics, MI, Kansas City, USA. ³⁸Kansas City School of Medicine, University of Missouri, Kansas City, MI, USA. ³⁹Division of Clinical Genetics, Children's Mercy Hospital and Clinics, Kansas City, MI, USA. ⁴⁰Service de Génétique Médicale, Institut de Génétique Médicale d'Alsace (IGMA), Hôpitaux Universitaires de Strasbourg, Strasbourg, France. ⁴¹Laboratoire de Génétique Médicale, UMRS_1112, Institut de Génétique Médicale d'Alsace (IGMA), Université de Strasbourg et INSERM, Strasbourg, France. ⁴²Service de Pédiatrie, Hôpital de Hautepierre, Hôpitaux Universitaires de Strasbourg, Strasbourg, France. ⁴³Laboratoires de Diagnostic Génétique, Institut de Génétique Médicale d'Alsace (IGMA), Hôpitaux Universitaires de Strasbourg, Nouvel Hôpital Civil, Strasbourg, France. ⁴⁴Pediatric Neurology Department, CHU de Strasbourg, Strasbourg, France. ⁴⁵Northern Ireland Regional Genetics Service, Belfast, Northern Ireland. ⁴⁶Department of Medical Genetics, Haukeland University Hospital, Bergen, Norway. ⁴⁷Division of Evolution, Infection and Genomics, School of Biological Sciences, Faculty of Biology, Medicine and Health, University of Manchester, Manchester, UK. ⁴⁸Manchester Centre for Genomic Medicine, Saint Mary's Hospital, Manchester University NHS Foundation Trust, Manchester, UK. ⁴⁹Department of Neurology, Division of Child Neurology, Columbia University Irving Medical Center, New York, USA. ⁵⁰Texas Tech Health Sciences Center Lubbock and KinderGenome Medical Genetics, Dallas, TX, USA. ⁵¹Texas Sports Psychiatry and Integrative Health, Austin, TX, USA. ⁵²Hillcrest Internal Medicine, Waco, TX, USA. ⁵³Centre de Génétique Humaine, Institut de Pathologie et de Génétique ASBL, Gosselies, Belgium. ⁵⁴Department of Medical Genomics/Clinical Genetics, Royal Prince Alfred Hospital, Camperdown, Sydney, NSW, Australia. ⁵⁵Department of Genetics, Children's Hospital of Eastern Ontario, Ottawa, ON, Canada. ⁵⁶Division of Medical Genetics, Department of Pediatrics, San Antonio Military Medical Center, San Antonio, TX, USA. ⁵⁷Department of Pediatrics, Long School of Medicine-UT Health San Antonio, San Antonio, TX, USA. ⁵⁸Department of Human Genetics, Radboud University Medical Center, Nijmegen, The Netherlands. ⁵⁹Pluryn, Residential Care Setting, Groesbeek, The Netherlands. ⁶⁰Centro de Genética Médica Jacinto Magalhães, Centro Hospitalar Universitário do Porto, Porto, Portugal. ⁶¹Unit for Multidisciplinary Research in Biomedicine, School of Medicine and Biomedical Sciences, Porto University, Porto, Portugal. ⁶²Service de Génétique Médicale, CHU Bordeaux, Bordeaux, France. ⁶³INSERM U1211, Laboratoire Maladies Rares: Génétique et Métabolisme, Bordeaux, Univ., Bordeaux, France. ⁶⁴Child Neurology and Neurodevelopmental Medicine Thompson Autism Center, CHOC Hospital, Orange County, CA, USA. ⁶⁵Indiana University School of Medicine, Indianapolis, USA. ⁶⁶University of California, San Diego, Rady Children's Hospital San Diego, San Diego, CA, USA. ⁶⁷North West Thames Regional Genetics Service, London North West University Healthcare NHS Trust, Harrow, London, UK. ⁶⁸Imperial College London, London, UK. ⁶⁹West of Scotland Centre for Genomic Medicine, Queen Elizabeth University Hospital, Glasgow, UK. ⁷⁰Centrum Medische Genetica, Universitair Ziekenhuis Brussel, Vrije Universiteit Brussel (VUB), Brussels, Belgium. ⁷¹CHU Sainte-Justine Research Center, University of Montreal, Montreal, QC, Canada. ⁷²Division Clinical Genetics Texas A&M University Health Science Center, College Station, TX, USA. ⁷³Department of Clinical Genetics, Leiden University Medical Centre, Leiden, The Netherlands. ⁷⁴Molecular and Human Genetics, Baylor College of Medicine, Houston, TX, USA. ⁷⁵Baylor Genetics Laboratories, Houston, TX, USA. ⁷⁶Centogene GmbH, Rostock, Germany. ⁷⁷Clinical Genomics, Ambry Genetics, Aliso Viejo, CA, USA. ⁷⁸GeneDx LLC, Gaithersburg, MA, USA. ⁷⁹Neuropediatric Unit, CHU UCL-Namur, Namur, Belgium. ⁸⁰Children's Hospital at Westmead, Sydney Children's Hospitals Network, Sydney, NSW, Australia. ⁸¹East Anglian Medical Genetics Service, Clinical Genetics, Addenbrooke's Treatment Centre, Addenbrooke's Hospital, Cambridge, UK. ⁸²These authors contributed equally: Elizabeth E. Palmer, Michael Pusch. [✉]email: elizabeth.palmer@unsw.edu.au; michael.pusch@ibf.cnr.it; kalscheu@molgen.mpg.de