

UC Merced

UC Merced Previously Published Works

Title

Complete Genome Sequence of Methanolinea tarda NOBI-1T, a Hydrogenotrophic Methanogen Isolated from Methanogenic Digester Sludge

Permalink

<https://escholarship.org/uc/item/606440zg>

Journal

Microbiology Resource Announcements, 2(5)

ISSN

2576-098X

Authors

Yamamoto, Kyosuke

Tamaki, Hideyuki

Cadillo-Quiroz, Hinsby

et al.

Publication Date

2014-10-30

DOI

10.1128/genomea.00876-14

Peer reviewed

Complete Genome Sequence of *Methanolinea tarda* NOBI-1^T, a Hydrogenotrophic Methanogen Isolated from Methanogenic Digester Sludge

Kyosuke Yamamoto,^a Hideyuki Tamaki,^{a,b} Hinsby Cadillo-Quiroz,^{c,d} Hiroyuki Imachi,^e Nikos Kyrpides,^{f,g} Tanja Woyke,^f Lynne Goodwin,^f Stephen H. Zinder,^h Yoichi Kamagata,^a Wen-Tso Liu^b

Bioproduction Research Institute, National Institute of Advanced Industrial Science and Technology (AIST), Higashi, Tsukuba, Ibaraki, Japan^a; Department of Civil and Environmental Engineering, University of Illinois at Urbana-Champaign, Urbana, Illinois, USA^b; School of Life Sciences, Arizona State University, Tempe, Arizona, USA^c; Swette Center for Environmental Biotechnology at the Biodesign Institute, Arizona State University, Tempe, Arizona, USA^d; Department of Subsurface Geobiology Analysis and Research (D-SUGAR), Japan Agency for Marine-Earth Science & Technology (JAMSTEC), Yokosuka, Kanagawa, Japan^e; DOE Joint Genome Institute, Walnut Creek, California, USA^f; Department of Biological Sciences, Faculty of Science, King Abdulaziz University, Jeddah, Saudi Arabia^g; Department of Microbiology, Cornell University, Ithaca, New York, USA^h

Here, we report a 2.0-Mb complete genome sequence of *Methanolinea tarda* NOBI-1^T, a methanogenic archaeon isolated from an anaerobic digested sludge. This is the first genome report of the genus *Methanolinea* isolate belonging to the family *Methanoregulaceae*, a recently proposed novel family within the order *Methanomicrobiales*.

Received 5 August 2014 Accepted 8 August 2014 Published 4 September 2014

Citation Yamamoto K, Tamaki H, Cadillo-Quiroz H, Imachi H, Kyrpides N, Woyke T, Goodwin L, Zinder SH, Kamagata Y, Liu W-T. 2014. Complete genome sequence of *Methanolinea tarda* NOBI-1^T, a hydrogenotrophic methanogen isolated from methanogenic digester sludge. *Genome Announc.* 2(5):e00876-14. doi:10.1128/genomeA.00876-14.

Copyright © 2014 Yamamoto et al. This is an open-access article distributed under the terms of the [Creative Commons Attribution 3.0 Unported license](https://creativecommons.org/licenses/by/3.0/).

Address correspondence to Hideyuki Tamaki, tamaki-hideyuki@aist.go.jp, or Wen-Tso Liu, wtliu@illinois.edu.

Methanolinea tarda NOBI-1^T, an H₂/CO₂-using methanogen, was isolated from mesophilic methanogenic sludge digesting municipal sewage in Nagaoka, Niigata Prefecture, Japan and described as a novel species of a novel genus within the order *Methanomicrobiales* (1). Based on 16S rRNA gene-based phylogeny, *M. tarda* NOBI-1^T belongs to the family level clade E1/E2, which is the outer group of the previously known three families. Recently, several strains belonging to the E1/E2 clade have been isolated and characterized, and a novel family *Methanoregulaceae* was proposed for the E1/E2 group as the fourth family within the order *Methanomicrobiales* (2). The family *Methanoregulaceae* is comprised of five isolated strains: *M. tarda* NOBI-1^T, *Methanolinea mesophila* TNR^T (2), *Methanoregula boonei* 6A8^T (3), *Methanoregula formicica* SMSP^T (4), and *Methanosphaerula palustris* E1-9c^T (5). Since the natural habitats and physiological features of these *Methanoregulaceae* members vary by species, the taxonomic identification of the species relies largely on molecular phylogeny, and the characteristic genetic and physiological properties of this group distinguishable from other families remain largely unclear. The whole-genome sequence of *M. tarda* NOBI-1^T provides the first genomic information of the species belonging to the genus *Methanolinea* and will contribute to an improved understanding of the unique features of the family *Methanoregulaceae*.

The whole-genome shotgun sequencing was performed using a combined Illumina and Roche GS-FLX Titanium approach. Sequence assembly was carried out using the GS De Novo assembler Newbler (version 2.3). Manual finishing efforts raised the quality of the assembly to that of a finished genome. Genes were identified using Prodigal (6) as part of the JGI genome annotation pipeline (7), followed by a round of manual curation using the JGI GenePRIMP pipeline (8). Additional gene functional an-

notation and comparative analysis were performed within the Integrated Microbial Genomes (IMG-ER) platform (9).

The complete genome sequence length was 2,052,856 bp with a G+C content of 56.5%. The genome contains 2,057 protein-coding sequences, 55 pseudo genes, 46 tRNA genes, and an rRNA operon including one 23S large-subunit gene and one 16S small-subunit gene. A total of 77.5% of open reading frames (1,634) are protein-coding genes with function prediction.

Gene classification by the NCBI Clusters of Orthologous Groups (COG) categories (10) reveals that the genome harbors all of the genes involved in hydrogenotrophic methanogenesis pathway and genes for formate dehydrogenase, which are essential to utilize H₂/CO₂ and formate for growth and methane production. The genome contains one clustered regularly interspaced short palindromic repeat (CRISPR) locus and 9 CRISPR-associated genes, and 16 transposases and inactivated derivatives, suggesting the contribution of viral predation and gene transfer events by mobile genetic elements to the evolution of *M. tarda* NOBI-1^T. Further comparative analyses with the genomes of other species belonging to the *Methanoregulaceae* and/or species within other taxa will provide insights into the unique genetic and physiological characteristics of the species within the *Methanoregulaceae* lineage.

Nucleotide sequence accession numbers. This whole-genome shotgun project has been deposited in DDBJ/EMBL/GenBank under the accession no. [AGIY000000000](https://www.ncbi.nlm.nih.gov/nuccore/AGIY000000000). The version described in this paper is AGIY000000002.

ACKNOWLEDGMENTS

This work was performed under the auspices of the U.S. Department of Energy's Office of Science, Biological and Environmental Research Pro-

gram, and by the University of California, Lawrence Berkeley National Laboratory under contract no. DE-AC02-05CH11231. Additionally, this study was financially supported in part by JSPS KAKENHI Grant Numbers 23657069, 23681044, and 24687011. The JGI project no. is 402816.

REFERENCES

1. Imachi H, Sakai S, Sekiguchi Y, Hanada S, Kamagata Y, Ohashi A, Harada H. 2008. *Methanolinea tarda* gen. nov., sp. nov., a methane-producing archaeon isolated from a methanogenic digester sludge. *Int. J. Syst. Evol. Microbiol.* 58:294–301. <http://dx.doi.org/10.1099/ijs.0.65394-0>.
2. Sakai S, Ehara M, Tseng IC, Yamaguchi T, Bräuer SL, Cadillo-Quiroz H, Zinder SH, Imachi H. 2012. *Methanolinea mesophila* sp. nov., a hydrogenotrophic methanogen isolated from rice field soil, and proposal of the archaeal family *Methanoregulaceae* fam. nov. within the order *Methanomicrobiales*. *Int. J. Syst. Evol. Microbiol.* 62:1389–1395. <http://dx.doi.org/10.1099/ijs.0.035048-0>.
3. Bräuer SL, Cadillo-Quiroz H, Ward RJ, Yavitt JB, Zinder SH. 2011. *Methanoregula boonei* gen. nov., sp. nov., an acidiphilic methanogen isolated from an acidic peat bog. *Int. J. Syst. Evol. Microbiol.* 61:45–52. <http://dx.doi.org/10.1099/ijs.0.021782-0>.
4. Yashiro Y, Sakai S, Ehara M, Miyazaki M, Yamaguchi T, Imachi H. 2011. *Methanoregula formicica* sp. nov., a methane-producing archaeon isolated from methanogenic sludge. *Int. J. Syst. Evol. Microbiol.* 61:53–59. <http://dx.doi.org/10.1099/ijs.0.014811-0>.
5. Cadillo-Quiroz H, Yavitt JB, Zinder SH. 2009. *Methanosphaerula palustris* gen. nov., sp. nov., a hydrogenotrophic methanogen isolated from a minerotrophic fen peatland. *Int. J. Syst. Evol. Microbiol.* 59:928–935. <http://dx.doi.org/10.1099/ijs.0.006890-0>.
6. Hyatt D, Chen G-L, Locascio PF, Land ML, Larimer FW, Hauser LJ. 2010. Prodigal: prokaryotic gene recognition and translation initiation site identification. *BMC Bioinformatics* 11:119. <http://dx.doi.org/10.1186/1471-2105-11-119>.
7. Mavromatis K, Ivanova NN, Chen I-MA, Szeto E, Markowitz VM, Kyrpides NC. 2009. The DOE-JGI standard operating procedure for the annotations of microbial genomes. *Stand. Genomics Sci.* 1:63–67. <http://dx.doi.org/10.4056/sigs.632>.
8. Pati A, Ivanova NN, Mikhailova N, Ovchinnikova G, Hooper SD, Lykidis A, Kyrpides NC. 2010. GenePRIMP: a gene prediction improvement pipeline for prokaryotic genomes. *Nat. Methods* 7:455–457. <http://dx.doi.org/10.1038/nmeth.1457>.
9. Markowitz VM, Mavromatis K, Ivanova NN, Chen IM, Chu K, Kyrpides NC. 2009. IMG ER: a system for microbial genome annotation expert review and curation. *Bioinformatics* 25:2271–2278. <http://dx.doi.org/10.1093/bioinformatics/btp393>.
10. Tatusov RL, Natale DA, Garkavtsev IV, Tatusova TA, Shankavaram UT, Rao BS, Kiryutin B, Galperin MY, Fedorova ND, Koonin EV. 2001. The COG database: new developments in phylogenetic classification of proteins from complete genomes. *Nucleic Acids Res.* 29:22–28. <http://dx.doi.org/10.1093/nar/29.1.22>.