

UC Irvine

UC Irvine Previously Published Works

Title

Curl pattern classification: A potential tool for communication and risk stratification

Permalink

<https://escholarship.org/uc/item/6111t25m>

Journal

International Journal of Women's Dermatology, 8(2)

ISSN

2352-6475

Authors

Krueger, Loren

Aguh, Crystal

Peterson, Erik

et al.

Publication Date

2022-06-01

DOI

10.1097/jw9.0000000000000015

Copyright Information

This work is made available under the terms of a Creative Commons Attribution License, available at <https://creativecommons.org/licenses/by/4.0/>

Peer reviewed

Curl pattern classification: A potential tool for communication and risk stratification

Loren Krueger, MD^{a,b,*}, Crystal Aguh, MD^c, Erik Peterson, MD^a, Jamael Thomas, MD MPH^c, JaBreia James, BS^d, Kristen Lo Sicco, MD^a, Jerry Shapiro, MD^a

Keywords: Alopecia, classification scheme, curl pattern, hair loss, hair texture, skin of color

Hair and hair loss disorders lack adequate tools for quantitative assessment, impacting the quality of our care. Even though alopecia is among the top 10 conditions for which Black patients seek dermatologic care, many dermatologists are less familiar or confident with evaluation of hair loss in ethnic hair. For example, we do not utilize a widely accepted measure for hair texture, yet we do consider hair texture when evaluating our hair loss patients as it is relevant to hair fragility, shaft shape, and styling practices. This gap in objectivity likely lowers dermatologists' confidence and accuracy in addressing hair disorders in these patients.

The lay public has developed and widely adopted classification schemes for hair texture, often termed "curl pattern," enabling communication regarding optimal hair products and styling. These schemes have garnered significant publicity with frequent features in lifestyle and beauty magazines as well as social media. Additionally, hair care products reference curl patterns in their marketing.^{1,2} As this is already widely accepted in curly hair communities, it is prudent to become familiar to augment patient-physician communication. With this pilot study, we aim to demonstrate that this classification scheme may also have utility as a risk-stratification tool in alopecia. Adoption of this tool could guide assessment, treatment recommendations, and patient compliance, yet remains underutilized in clinical practice.

We performed an institutional review board-approved, pilot survey study at New York University and Johns Hopkins University regarding female patients with a clinical and/or histopathologic diagnosis of androgenetic alopecia (AGA), traction alopecia, frontal fibrosing alopecia, central centrifugal cicatricial alopecia (CCCA), alopecia areata, or telogen effluvium. During a clinical encounter, consent was obtained and the provider determined the participant's curl pattern. Curl patterns were divided into pattern 1 (straight), pattern 2 (wavy), pattern 3 (curly), and pattern 4 (coily). Of note, the publicly

adopted curl pattern classification subdivides each of these categories further (Fig. 1).

A total of 74 participants were enrolled with 61 reporting ethnicity (43/61 = Caucasian, 9 = Hispanic, 8 = Black, 1 = Asian). Inclusion criteria required that the patients have clear clinical or histologically proven subtype of alopecia for a diagnosis to eliminate any discrepancies. Regarding alopecia subtypes of the 74 participants, 30/74 = frontal fibrosing alopecia, 25 = AGA, 7 = CCCA, 6 = telogen effluvium, 5 = alopecia areata, 1 = traction alopecia. Overall, the subtypes of alopecia are significantly different amongst the 4 curl patterns ($P = 0.011$) (Table 1), mostly between patterns 1 and 4 ($P = 0.0009$). Although AGA occurs with similar frequency across the 4 curl patterns, the risk of CCCA compared to AGA increases with increasing curl pattern ($P = 0.011$) (Table 1). Likewise, the study found that the distributions of alopecia types and curly hair patterns are significantly different between European or Caucasian patients and African or African American patients ($P = 0.0025$) (Tables 2 and 3). Curl pattern appears to be more significant than ethnicity in determining risk of associated alopecia subtype.

A limitation is the small sample size. However, our cohort does include all curl patterns and the proportions of Black and Hispanic participants enrolled are relatively reflective of the US population. A larger and more ethnically diverse cohort would enable better understanding of the relationship between curl pattern and ethnicity on alopecia. For example, we now acknowledge the genetic drivers in CCCA but have yet to parse out independent contributions of race/ethnicity versus hair texture.

What is known about this subject in regard to women and their families?

- Dermatologists currently lack adequate tools for objective assessment in hair and hair loss disorders, which impacts women.
- Although we consider hair texture when evaluating hair loss patients as it relates to styling practices, hair health and treatment recommendations, we do not have a tool for categorization of hair texture.
- The public has adopted and widely utilizes "curl pattern" classifications for communication and marketing of products.

What is new from this article as messages for women and their families?

- We suggest dermatologists become familiar with curl pattern classification to optimize communication with our patients.
- Curl pattern classification may have predictive value in hair loss disorders and more work is needed to move towards its validation.

^a The Ronald O. Perleman Department of Dermatology, New York University, New York, New York

^b Department of Dermatology, Emory University, Atlanta, Georgia

^c Department of Dermatology, Johns Hopkins University, Baltimore, Maryland

^d Medical College of Georgia at Augusta University, Augusta, Georgia

K.L. and J.S. have contributed equally to this work.

*Corresponding author. E-mail address: loren.d.krueger@emory.edu (L. Krueger).

Copyright © 2022 The Authors. Published by Wolters Kluwer Health, Inc. on behalf of Women's Dermatologic Society. This is an open-access article distributed under the terms of the Creative Commons Attribution-Non Commercial-No Derivatives License 4.0 (CCBY-NC-ND), where it is permissible to download and share the work provided it is properly cited. The work cannot be changed in any way or used commercially without permission from the journal.

International Journal of Women's Dermatology (2022) 8:e015

Received: 24 August 2021; Accepted 21 February 2022

Published online 5 April 2022

DOI: 10.1097/JW9.0000000000000015



Fig. 1. Proposed curl pattern classification scheme.

Table 1.

Alopecia type and curl pattern

Curl pattern	(AGA), n (%)	(CCCA), n (%)	(FFA), n (%)	(TA), n (%)	(AA), n (%)	(TE), n (%)	Fisher exact test <i>P</i>
Pattern 1 (n = 17)	6 (35.3)	0 (0)	8 (47.0)	0 (0)	3 (17.7)	0 (0)	0.011
Pattern 2 (n = 37)	12 (32.4)	2 (5.4)	17 (46.0)	0 (0)	2 (5.4)	4 (10.8)	
Pattern 3 (n = 12)	4 (33.3)	1 (8.3)	5 (41.7)	0 (0)	0 (0)	2 (16.7)	
Pattern 4 (n = 8)	3 (37.5)	4 (50.0)	0 (0)	1 (12.5)	0 (0)	0 (0)	

AA, alopecia areata; AGA, androgenetic alopecia; CCCA, central centrifugal cicatricial alopecia; FFA, frontal fibrosing alopecia; TA, traction alopecia; TE, telogen effluvium.

Table 2.

Race and curl pattern

Race/ethnicity	Pattern 1 (straight), n (%)	Pattern 2 (wavy), n (%)	Pattern 3 (curly), n (%)	Pattern 4 (coiled), n (%)	Fisher exact test <i>P</i>
Hispanic or Latino (n = 9)	1 (11.1)	7 (77.7)	1 (11.1)	0 (0)	<0.0001
Asian or Pacific Islander (n = 1)	1 (100)	0 (0)	0 (0)	0 (0)	
European or Caucasian (n = 43)	14 (32.6)	22 (51.2)	6 (14.0)	1 (2.3)	
African, African American (n = 8)	0 (0)	1 (12.5)	1 (12.5)	6 (75.0)	

Table 3.

Race and alopecia types

Race/ethnicity	(AGA), n (%)	(CCCA), n (%)	(FFA), n (%)	(TA), n (%)	(AA), n (%)	(TE), n (%)	Fisher exact test <i>P</i>
Hispanic or Latino (n = 9)	3 (33.3)	1 (11.1)	4 (44.4)	0 (0)	0 (0)	1 (11.1)	0.043
Asian or Pacific Islander (n = 1)	0 (0)	0 (0)	1 (100)	0 (0)	0 (0)	0 (0)	
European or Caucasian (n = 43)	12 (27.9)	1 (2.3)	21 (48.8)	1 (2.3)	4 (9.3)	4 (9.3)	
African, African American (n = 8)	5 (62.5)	3 (37.5)	0 (0)	0 (0)	0 (0)	0 (0)	

AA, alopecia areata; AGA, androgenetic alopecia; CCCA, central centrifugal cicatricial alopecia; FFA, frontal fibrosing alopecia; TA, traction alopecia; TE, telogen effluvium.

Dermatologists lack a widely adopted, objective classification scheme of curl pattern, while the lay public successfully navigates hair care practices using similar tools. This gap limits our objective assessment in the already challenging realm of hair loss disorders. The authors aim to introduce this visual tool

to facilitate quality communication with patients. Additionally, the results of our pilot study suggest this classification may also have utility in risk stratification, as subtypes of alopecia, including CCCA, are statistically different amongst the 4 curl patterns. Ultimately, this tool has the potential to guide assessment and

facilitate patient-centered education and treatment, and we are hopeful future studies will move towards its validation.

Author contributions

LK: Study conception and design, draft manuscript preparation, data collection, analysis and interpretation of results; CA: Data collection, manuscript review; EP: Data collection; JT: Data collection; JBJ: Draft manuscript preparation; KLS: Study conception and design, analysis and interpretation of results; JS: Data collection

Conflicts of interest

None.

Funding

This was funded by the Rudin Research Foundation Grant at the Department of Dermatology, New York University, to

support resident research. This funding was used to support the development of medical imagery. We thank Dr. Michael Natter for his valuable contributions.

Study approval

The author(s) confirm that any aspect of the work covered in this manuscript that has involved human patients has been conducted with the ethical approval of all relevant bodies.

References

1. Here K, Growth H, Products H, et al. The Only Hair Typing System Article You'll Ever Need. [online] Curl Centric® | Rewrite the Rules of Natural Hair Care. 2018. Available from: <https://www.curlcentric.com/hair-typing-system/>. Accessed November 9, 2021.
2. Andre Walker Hair (n.d.). Hair Products for Natural and Black Hair. Retrieved November 22, 2021. Available from: <https://andrewalkerhair.com/#andre-walker>. Accessed January 5, 2018.