

# Lawrence Berkeley National Laboratory

## LBL Publications

### Title

Improved Draft Genome Sequence of *Microbacterium* sp. Strain LKL04, a Bacterial Endophyte Associated with Switchgrass Plants

### Permalink

<https://escholarship.org/uc/item/616716pb>

### Journal

Microbiology Resource Announcements, 8(45)

### ISSN

2576-098X

### Authors

Sahib, Mohammad Radhi

Yang, Piao

Bokros, Norbert

et al.

### Publication Date

2019-11-07

### DOI

10.1128/mra.00927-19

Peer reviewed



# Improved Draft Genome Sequence of *Microbacterium* sp. Strain LKL04, a Bacterial Endophyte Associated with Switchgrass Plants

Mohammad Radhi Sahib,<sup>a,b</sup> Piao Yang,<sup>c</sup> Norbert Bokros,<sup>a</sup> Nicole Shapiro,<sup>d</sup> Tanja Woyke,<sup>d</sup> Nikos C. Kyrpides,<sup>d</sup> Ye Xia,<sup>c</sup> Seth DeBolt<sup>a</sup>

<sup>a</sup>Department of Horticulture, University of Kentucky, Lexington, Kentucky, USA

<sup>b</sup>Department of Horticulture, Al-Qasim Green University, Babylon, Iraq

<sup>c</sup>Department of Plant Pathology, The Ohio State University, Columbus, Ohio, USA

<sup>d</sup>Department of Energy Joint Genome Institute, Walnut Creek, California, USA

**ABSTRACT** We report here the genome assembly and analysis of *Microbacterium* strain sp. LKL04, a Gram-positive bacterial endophyte isolated from switchgrass plants (*Panicum virgatum*) grown on a reclaimed coal-mining site. The 2.9-Mbp genome of this bacterium was assembled into a single contig encoding 2,806 protein coding genes.

Members of the genus *Microbacterium* have previously been isolated from a wide range of environments, including soils, marine ecosystems, air, and sewage, and from plants and insects (1–5). We report here information about the sequenced and assembled genome of the bacterial endophyte *Microbacterium* sp. strain LKL04, a Gram-positive actinobacterium, isolated from leaves of switchgrass plants grown on a reclaimed coal-mining site in western Kentucky (6).

Switchgrass samples were collected from the coal-mining site in July 2010. Leaf samples were cut into 1- to 1.5-cm-long segments, surface sterilized with a 20% bleach solution, and rinsed 5 times with autoclaved tap water. The surface-sterilized segments were incubated on tryptic soy agar (TSA) plates for 3 to 5 days at 26°C before the individual colonies were isolated and restreaked at least three times on new TSA plates (6). Single purified colonies were then isolated and grown at room temperature for 1 to 2 days in tryptic soy broth (TSB). A modified cetyltrimethylammonium bromide (CTAB) bacterial DNA isolation protocol (7; <https://1ofdmq2n8tc36m6i46scovo2e-wpengine.netdna-ssl.com/wp-content/uploads/2014/02/JGI-Bacterial-DNA-isolation-CTAB-Protocol-2012.pdf>) was followed to isolate the bacterial DNA for sequencing.

The genome of *Microbacterium* sp. strain LKL04 was sequenced at 212× coverage using Pacific Biosciences (PacBio) sequencing technology (8). A PacBio SMRTbell library was constructed and sequenced with the PacBio RS platform, generating 198,113 filtered subreads with an average read length of 3,930 bp ± 2,621 bp, totaling 778.5 Mbp. Reads were trimmed and assembled using Hierarchical Genome Assembly Process (HGAP) v.2.3.0 (9). The final genome assembly contains a single contig spanning the complete 2.922-Mbp length of the bacterial genome, with a GC content of 69.7%, which is characteristic of actinobacteria. The genome is predicted to be circular.

Genes were identified using Prodigal v.2.5, followed by a round of manual curation using GenePRIMP, resulting in a total of 2,862 predicted genes (10, 11). From these, 2,806 predicted protein coding genes were translated and used to search the National Center for Biotechnology Information (NCBI) nonredundant, UniProt, TIGRFam, Pfam, Kyoto Encyclopedia of Genes and Genomes (KEGG), Clusters of Orthologous Genes (COG), PANTHER, and InterPro databases (12–18). For the remaining 56 genes, the tRNAScan-SE tool was used to further identify 45 tRNA genes, 6 rRNA genes, and 5

**Citation** Sahib MR, Yang P, Bokros N, Shapiro N, Woyke T, Kyrpides NC, Xia Y, DeBolt S. 2019. Improved draft genome sequence of *Microbacterium* sp. strain LKL04, a bacterial endophyte associated with switchgrass plants. *Microbiol Resour Announc* 8:e00927-19. <https://doi.org/10.1128/MRA.00927-19>.

**Editor** Julie C. Dunning Hotopp, University of Maryland School of Medicine


**Copyright** © 2019 Sahib et al. This is an open-access article distributed under the terms of the [Creative Commons Attribution 4.0 International license](https://creativecommons.org/licenses/by/4.0/).

Address correspondence to Ye Xia, [xia.374@osu.edu](mailto:xia.374@osu.edu), or Seth DeBolt, [sdebo2@uky.edu](mailto:sdebo2@uky.edu).

**Received** 14 August 2019

**Accepted** 15 October 2019

**Published** 7 November 2019



**FIG 1** Circular representation of the LKL04 genome using Circos (25). The circles, from outside to inside, denote protein coding genes colored by size (A), RNA genes (B), transmembrane helix regions (C), GC content along a 1-kb window, with red lines indicating regions above the 69.7% genome average and black lines indicating regions below the genome average (D) GC skew, with red lines indicating a skew greater than zero and black lines indicating a skew less than zero (E), and genes annotated into distinct PANTHER protein classes (F). The repository for storage of scripts used to construct the figure can be found at [https://github.com/nbo245/LKL04/tree/master/circos\\_plot](https://github.com/nbo245/LKL04/tree/master/circos_plot).

noncoding RNAs. For the noncoding RNAs, the RNA components of the protein secretion complex and RNase P were identified by searching the genome for the corresponding Rfam profiles using Infernal (19, 20). CheckM v.1.0.8, hosted on KBase, was used to estimate the completeness of the LKL04 genome (21, 22). Overall, the LKL04 genome returned a completeness score of 99.5% and a contamination level of only 0.67%. Using the PANTHER hidden Markov model (HMM) scoring tool pantherScore v.2.1, the protein sequences were further mapped against the PANTHER HMM database v.14.1 to functionally annotate the LKL04 genes and query for significantly overrepresented genes (23). Default parameters were used for each software program, unless otherwise specified. Selected annotations and genome characteristics are shown in Fig. 1. Additional gene prediction analysis and manual functional annotation were performed within the Integrated Microbial Genomes (IMG) platform developed by the Joint Genome Institute (Walnut Creek, CA) (24).

**Data availability.** The whole-genome sequence has been deposited in DDBJ/EMBL/GenBank under the accession no. [PRJNA322991](https://www.ncbi.nlm.nih.gov/nuccore/PRJNA322991). Original forward and reverse sequencing reads can be retrieved from NCBI under SRA accession no. [SRR4232145](https://www.ncbi.nlm.nih.gov/sra/SRR4232145) and [SRR4232146](https://www.ncbi.nlm.nih.gov/sra/SRR4232146). The associated sequence data can also be found at the Joint Genome Institute (JGI) portal with the IMG taxon identifier (ID) 2667527218 (<https://genome.jgi.doe.gov/portal/MicspLKL04/MicspLKL04.info.html>) or at <https://www.ncbi.nlm.nih.gov/Taxonomy/Browser/wwwtax.cgi?id=912630>. Scripts used to construct Fig. 1 can be found at [https://github.com/nbo245/LKL04/tree/master/circos\\_plot](https://github.com/nbo245/LKL04/tree/master/circos_plot).

## ACKNOWLEDGMENTS

Genome sequencing and data annotation were carried out in the U.S. Department of Energy (DOE) Joint Genome Institute (JGI), a DOE Office of Science User Facility

supported under contract DE-AC02-05CH11231. This work was also partially supported by the Hatch Project from USDA-NIFA-OHO01392 and by OARD Seed Award OHOA-1615 at The Ohio State University.

## REFERENCES

- Alves A, Correia A, Igual JM, Trujillo ME. 2014. *Microbacterium endophyticum* sp. nov. and *Microbacterium halimionae* sp. nov., endophytes isolated from the salt-marsh plant *Halimione portulacoides* and emended description of the genus *Microbacterium*. *Syst Appl Microbiol* 37: 474–479. <https://doi.org/10.1016/j.syapm.2014.08.004>.
- Kim KK, Lee KC, Oh H-M, Lee J-S. 2008. *Microbacterium aquimaris* sp. nov., isolated from seawater. *Int J Syst Evol Microbiol* 58:1616–1620. <https://doi.org/10.1099/ijs.0.65763-0>.
- Yoon J-H, Schumann P, Kang S-J, Lee C-S, Lee S-Y, Oh T-K. 2009. *Microbacterium insulae* sp. nov., isolated from soil. *Int J Syst Evol Microbiol* 59:1738–1742. <https://doi.org/10.1099/ijs.0.007591-0>.
- Mawlankar RR, Mual P, Sonalkar VV, Thorat MN, Verma A, Srinivasan K, Dastager SG. 2015. *Microbacterium enclense* sp. nov., isolated from sediment sample. *Int J Syst Evol Microbiol* 65:2064–2070. <https://doi.org/10.1099/ijs.0.000221>.
- Kim DY, Shin D-H, Jung S, Kim H, Lee JS, Cho H-Y, Bae KS, Sung C-K, Rhee YH, Son K-H, Park H-Y. 2014. Novel alkali-tolerant GH10 endo- $\beta$ -1,4-xylanase with broad substrate specificity from *Microbacterium trichothecenolyticum* HY-17, a gut bacterium of the mole cricket *Gryllotalpa orientalis*. *J Microbiol Biotechnol* 24:943–953. <https://doi.org/10.4014/jmb.1405.05032>.
- Xia Y, Greissworth E, Mucci C, Williams MA, De Bolt S. 2013. Characterization of culturable bacterial endophytes of switchgrass (*Panicum virgatum* L.) and their capacity to influence plant growth. *Glob Change Biol Bioenergy* 5:674–682. <https://doi.org/10.1111/j.1757-1707.2012.01208.x>.
- Wilson K. 2001. Preparation of genomic DNA from bacteria. *Curr Protoc Mol Bio* 56:241–245. <https://doi.org/10.1002/0471142727.mb0204s56>.
- Rhoads A, Au KF. 2015. PacBio sequencing and its applications. *Genomics Proteomics Bioinformatics* 13:278–289. <https://doi.org/10.1016/j.gpb.2015.08.002>.
- Chin C-S, Alexander DH, Marks P, Klammer AA, Drake J, Heiner C, Clum A, Copeland A, Huddleston J, Eichler EE, Turner SW, Korlach J. 2013. Nonhybrid, finished microbial genome assemblies from long-read SMRT sequencing data. *Nat Methods* 10:563–569. <https://doi.org/10.1038/nmeth.2474>.
- Hyatt D, Chen G-L, LoCascio PF, Land ML, Larimer FW, Hauser LJ. 2010. Prodigal: prokaryotic gene recognition and translation initiation site identification. *BMC Bioinformatics* 11:119. <https://doi.org/10.1186/1471-2105-11-119>.
- Pati A, Ivanova NN, Mikhailova N, Ovchinnikova G, Hooper SD, Lykidis A, Kyrpides NC. 2010. GenePRIMP: a gene prediction improvement pipeline for prokaryotic genomes. *Nat Methods* 7:455–457. <https://doi.org/10.1038/nmeth.1457>.
- Pruitt KD, Tatusova T, Maglott DR. 2007. NCBI reference sequences (RefSeq): a curated non-redundant sequence database of genomes, transcripts and proteins. *Nucleic Acids Res* 35:D61–D65. <https://doi.org/10.1093/nar/gkl842>.
- UniProt Consortium. 2019. UniProt: a worldwide hub of protein knowledge. *Nucleic Acids Res* 47:D506–D515. <https://doi.org/10.1093/nar/gky1049>.
- Haft DH, Selengut JD, White O. 2003. The TIGRFAMs database of protein families. *Nucleic Acids Res* 31:371–373. <https://doi.org/10.1093/nar/gkg128>.
- Finn RD, Bateman A, Clements J, Coggill P, Eberhardt RY, Eddy SR, Heger A, Hetherington K, Holm L, Mistry J, Sonnhammer ELL, Tate J, Punta M. 2014. Pfam: the protein families database. *Nucleic Acids Res* 42: D222–D230. <https://doi.org/10.1093/nar/gkt1223>.
- Kanehisa M, Goto S. 2000. KEGG: Kyoto Encyclopedia of Genes and Genomes. *Nucleic Acids Res* 28:27–30. <https://doi.org/10.1093/nar/28.1.27>.
- Tatusov RL, Fedorova ND, Jackson JD, Jacobs AR, Kiryutin B, Koonin EV, Krylov DM, Mazumder R, Mekhedov SL, Nikolskaya AN, Rao BS, Smirnov S, Sverdlov AV, Vasudevan S, Wolf YI, Yin JJ, Natale DA. 2003. The COG database: an updated version includes eukaryotes. *BMC Bioinformatics* 4:41. <https://doi.org/10.1186/1471-2105-4-41>.
- Thomas PD, Campbell MJ, Kejariwal A, Mi H, Karlak B, Daverman R, Diemer K, Muruganujan A, Narechania A. 2003. PANTHER: a library of protein families and subfamilies indexed by function. *Genome Res* 13:2129–2141. <https://doi.org/10.1101/gr.772403>.
- Quast C, Pruesse E, Yilmaz P, Gerken J, Schweer T, Yarza P, Peplies J, Glöckner FO. 2013. The SILVA ribosomal RNA gene database project: improved data processing and Web-based tools. *Nucleic Acids Res* 41:D590–D596. <https://doi.org/10.1093/nar/gks1219>.
- Nawrocki EP, Eddy SR. 2013. Infernal 1.1: 100-fold faster RNA homology searches. *Bioinformatics* 29:2933–2935. <https://doi.org/10.1093/bioinformatics/btt509>.
- Parks DH, Imelfort M, Skennerton CT, Hugenholtz P, Tyson GW. 2015. CheckM: assessing the quality of microbial genomes recovered from isolates, single cells, and metagenomes. *Genome Res* 25:1043–1055. <https://doi.org/10.1101/gr.186072.114>.
- Arkin AP, Cottingham RW, Henry CS, Harris NL, Stevens RL, Maslov S, Dehal P, Ware D, Perez F, Canon S, Sneddon MW, Henderson ML, Riehl WJ, Murphy-Olson D, Chan SY, Kamimura RT, Kumari S, Drake MM, Brettin TS, Glass EM, Chivian D, Gunter D, Weston DJ, Allen BH, Baumohl J, Best AA, Bowen B, Brenner SE, Bun CC, Chandonia J-M, Chia J-M, Colasanti R, Conrad N, Davis JJ, Davison BH, DeJongh M, Devoid S, Dietrich E, Dubchak I, Edirisinghe JN, Fang G, Faria JP, Frybarger PM, Gerlach W, Gerstein M, Greiner A, Gurtowski J, Haun HL, He F, Jain R, Joachimiak MP, Keegan KP, Kondo S, Kumar V, Land ML, Meyer F, Mills M, Novichkov PS, Oh T, Olsen GJ, Olson R, Parrello B, Pasternak S, Pearson E, Poon SS, Price GA, Ramakrishnan S, Ranjan P, Ronald PC, Schatz MC, Seaver SMD, Shukla M, Sutormin RA, Syed MH, Thomason J, Tintle NL, Wang D, Xia F, Yoo H, Yoo S, Yu D. 2018. KBase: the United States Department of Energy Systems Biology Knowledgebase. *Nat Biotechnol* 36:566–569. <https://doi.org/10.1038/nbt.4163>.
- Mi H, Muruganujan A, Huang X, Ebert D, Mills C, Guo X, Thomas PD. 2019. Protocol update for large-scale genome and gene function analysis with the PANTHER classification system (v.14.0). *Nat Protoc* 14:703. <https://doi.org/10.1038/s41596-019-0128-8>.
- Markowitz VM, Chen I-MA, Palaniappan K, Chu K, Szeto E, Grechkin Y, Ratner A, Jacob B, Huang J, Williams P, Huntemann M, Anderson I, Mavromatis K, Ivanova NN, Kyrpides NC. 2012. IMG: the integrated microbial genomes database and comparative analysis system. *Nucleic Acids Res* 40:D115–D122. <https://doi.org/10.1093/nar/gkr1044>.
- Krzywinski M, Schein H, Birol I, Connors J, Gascoyne R, Horsman D, Jones SJ, Marra MA. 2009. Circos: an information aesthetic for comparative genomics. *Genome Res* 19:1639–1645. <https://doi.org/10.1101/gr.092759.109>.