

UC Irvine

UC Irvine Previously Published Works

Title

Measurement of the parameters of the $\psi(3770)$ resonance

Permalink

<https://escholarship.org/uc/item/61d3b670>

Journal

Physical Review D, 21(9)

ISSN

2470-0010

Authors

Schindler, RH
Siegrist, JL
Alam, MS
et al.

Publication Date

1980-05-01

DOI

10.1103/physrevd.21.2716

Copyright Information

This work is made available under the terms of a Creative Commons Attribution License, available at <https://creativecommons.org/licenses/by/4.0/>

Peer reviewed

Measurement of the parameters of the $\psi(3770)$ resonance

R. H. Schindler, J. L. Siegrist, M. S. Alam, A. M. Boyarski, M. Breidenbach, D. L. Burke, J. Dorenbosch, J. M. Dorfan, G. J. Feldman, M. E. B. Franklin, G. Hanson, K. G. Hayes, T. Himel, D. G. Hitlin,* R. J. Hollebeek, W. R. Innes, J. A. Jaros, P. Jenni, R. R. Larsen, V. Lüth, M. L. Perl, B. Richter, D. L. Scharre, R. F. Schwitters,[†] H. Taureg, M. Tonutti,[‡] R. A. Vidal, J. M. Weiss, and H. Zaccane[§]
Stanford Linear Accelerator Center, Stanford University, Stanford, California 94305

G. S. Abrams, C. A. Blocker, W. C. Carithers, W. Chinowsky, M. W. Coles, S. Cooper, W. E. Dieterle, J. B. Dillon, M. W. Eaton, G. Gidal, G. Goldhaber, A. D. Johnson, J. A. Kadyk, A. J. Lankford, R. E. Millikan, M. E. Nelson, C. Y. Pang, J. F. Patrick, J. Strait, G. H. Trilling, E. N. Vella, and I. Videau

Lawrence Berkeley Laboratory and Department of Physics, University of California, Berkeley, California 94720

(Received 26 November 1979)

We present a measurement of the cross section for hadron production by e^+e^- annihilation in the vicinity of the previously observed resonance near 3.77 GeV. The data are used to determine the parameters of the $\psi(3770)$ resonance. The values found are: mass, 3764 ± 5 MeV/ c^2 , total width, 23.5 ± 5 MeV, and partial width to electron pairs, 276 ± 50 eV.

The $\psi(3770)$ resonance observed in e^+e^- annihilation to hadrons lies just above the threshold for pair production of D mesons.^{1,2} Because of the simple threshold kinematics and relatively large pair-production rates for charged and neutral D mesons, the $\psi(3770)$ represents a convenient source of charmed D mesons for studies of their decay properties. Precise knowledge of the parameters of this resonance is important both for comparison with the charmonium model of the ψ family³ as well as assisting in the determination of absolute branching ratios of D -meson decay. The two previous measurements of the leptonic partial width differed significantly.^{1,2}

We present here an analysis of cross-section data obtained during a contiguous period of time at 23 center-of-mass energies ($E_{c.m.}$) over the range 3.670 to 3.870 GeV. The data were taken with the SLAC-LBL MARK II magnetic detector at the e^+e^- storage ring SPEAR. The cross-section results were fitted to determine parameters of the $\psi(3770)$ resonance.

A detailed description of the MARK II detector may be found in Ref. 4. Briefly, it is a 3-m-diameter solenoidal magnet with a field of 0.4 T. Charged particles are tracked by drift chambers over $0.85 \times 4\pi$ sr. Their identities may be established over restricted kinematic ranges by time-of-flight counters, liquid-argon electromagnetic calorimeters, and range counters (for π - μ separation). Photons are detected in the liquid-argon calorimeter.

The detector trigger required that at least two charged tracks be within the active volume of the drift chambers, with at least one within the central $0.65 \times 4\pi$ sr with a transverse momentum

greater than 100 MeV/ c . Event selection for determination of the total hadronic cross section was essentially the same as that used in earlier SPEAR experiments.⁵ Aside from time-of-flight criteria for rejecting cosmic rays, only the charged-particle tracking information was used. Hadronic events were required to have three or more charged tracks with transverse momenta greater than 100 MeV/ c , or two charged prongs of momenta greater than 300 MeV/ c that were acoplanar with the incident e^+e^- beam direction by at least 20° .

Contamination of the hadron sample from two-photon processes⁶ and τ lepton pair production⁷ were estimated by Monte Carlo calculations and were subtracted from the number of hadron candidates. The estimated two-photon contribution in the two-prong events was typically 6%; the overall two-photon event subtraction was typically 1%. Similarly, the τ contribution to the two-prong events was approximately 20%, while only 8% of the total number of hadronic events are expected to come from $\tau^+\tau^-$ production.

Because the liquid-argon calorimeter was unavailable during this running, corrections had to be made for contamination of the hadronic event sample by Bhabha scattering events that radiate energetic photons and subsequently produce e^+e^- pairs. These corrections were estimated by studying samples of data taken near the peak of the $\psi(3770)$ when the shower system was operational. A subtraction of 1% of the total number of hadron candidate events was made to account for this multiprong Bhabha contamination.

The average hadronic-event detection efficiency was estimated from the observed charged multi-

plicity distributions and Monte Carlo simulations of this experiment using a phase-space model and a D -production model. The estimates are both consistent with a smooth variation in the detection efficiency from 0.66 at $E_{c.m.} = 3.67$ GeV to 0.69 at 3.87 GeV.

Luminosity was measured by detection of small-angle Bhabha scattering events in counter telescopes centered at 22 mrad from the incident beam direction. The system has been calibrated at other energies by comparison with the yield of Bhabha events in the central $0.65 \times 4\pi$ sr of the detector.

During data taking, the storage-ring energy was monitored with a precision flip coil in a reference bend magnet connected in series with the SPEAR bend magnets. The relative E_{beam} were held to




FIG. 1. (a) The observed cross section $R (= \sigma_{hadronic} / \sigma_{\mu\mu})$ with continuum radiative correction near the $\psi(3770)$. The curve is the fit to the data. (b) The cross section in units of R after radiative corrections for $\psi(3095)$, $\psi(3684)$, and $\psi(3770)$. The curve is the fitted Breit-Wigner form.

within ± 0.2 MeV of the desired central value for data points in the range 1.835 to 1.858 GeV and to within ± 0.4 MeV for the remaining data points presented. The systematic error on the energy scale ($E_{c.m.}$) of SPEAR is $\pm 0.13\%$.

The observed cross-section results are presented in Fig. 1(a) in terms of R , the ratio of the total hadronic cross section to the theoretical pointlike cross section for muon pair production $\sigma_{\mu\mu} [\sigma_{\mu\mu} = 86.9/E_{c.m.}^2, \text{ with } \sigma \text{ in nb and } E \text{ in GeV}]$. The most prominent features of the data presented in Fig. 1(a) are the $\psi(3770)$ resonance and the radiative tail of the $\psi(3684)$. The error bars are statistical only; the overall systematic uncertainty is estimated to be $\pm 10\%$ with contributions predominantly from event selection (5%), efficiency estimates (5%), and luminosity monitoring (7%).

The determination of the resonance parameters (mass $M_{\psi'}$, total width Γ_{tot} , and leptonic width Γ_{ee}) was accomplished by fitting the observed data to a function that described the resonance shape and the dominant nonresonant hadronic backgrounds, and included radiative corrections for the resonance itself, the nearby $\psi(3684)$ and $\psi(3095)$, and the continuum.^{8,9} The resonance shape was taken to be a nonrelativistic p -wave Breit-Wigner shape with an energy-dependent total width $[\Gamma_{tot}(E_{c.m.})]$:

$$R(E_{c.m.}) = \frac{1}{\sigma_{\mu\mu}} \frac{3\pi}{M_{\psi'}^2} \times \frac{\Gamma_{ee}\Gamma_{tot}(E_{c.m.})}{(E_{c.m.} - M_{\psi'})^2 + \Gamma_{tot}^2(E_{c.m.})/4}. \quad (1)$$

The form of Γ_{tot} was chosen assuming that $\psi(3770)$ is a resonance of unique isospin (0 or 1) which decays predominantly to D^+D^- or $D^0\bar{D}^0$ through the strong interaction. These assumptions are justified by its previously measured large total width (~ 25 MeV), proximity to the substantially narrower $\psi(3684)$, and its association with the onset of D -meson production.¹ We therefore chose the energy dependence of the total width to account for both threshold dependence and the otherwise equal production of $D^0\bar{D}^0$ or D^+D^- :¹⁰

$$\Gamma_{tot}(E_{c.m.}) \propto \frac{p_*^3}{1 + (rp_*)^2} + \frac{p_0^3}{1 + (rp_0)^2}. \quad (2)$$

Here p_* , p_0 refer to the momentum of the pair-produced D^+ , D^0 , respectively, and are calculated using the respective masses of 1863.3 and 1868.3 MeV.¹¹ The width $\Gamma_{tot}(E_{c.m.})$ in (2) is normalized to Γ_{tot} (the parameter of the fit) at the peak of the resonance ($E_{c.m.} = M_{\psi'}$). The parameter r is an interaction radius taken to be 2.5 F for the

analysis. While the resonance parameters were found to be insensitive to the exact value of r , it does affect the charged-to-neutral decay fraction determined by (2). This variation is less than 5% and is negligible in comparison to the uncertainties in total and leptonic widths.

The background shape includes a flat term and terms proportional to $(p_+)^3$ and $(p_0)^3$. The latter account for the onset of nonresonant D production. Because of the proximity to threshold, these p^3 contributions are found to be negligibly small.

The radiative correction to the continuum amounts to $\sim 9\%$ (a constant $R = 2.5$ was assumed) and is subtracted before fitting the data. The radiative tails of the $\psi(3095)$ and $\psi(3684)$ have line shapes that depend only on their masses and leptonic widths.⁸ The fit is not sensitive to the nearly flat contribution (~ 0.2 units of R) of the $\psi(3095)$ and therefore the previously measured values are used.¹ The fit is consistent with the known parameters of the $\psi(3684)$.¹ Finally, the self-radiative corrections for the $\psi(3770)$ were performed numerically in the fit.⁸ At the peak, the correction amounts to an enhancement of $\sim 25\%$ of resonance height.

Figure 1(b) shows the radiatively corrected data and fit. The results for this fit are

$$\begin{aligned} M_{\psi''} &= 3764 \pm 5 \text{ MeV}, \\ \Gamma_{ee} &= 276 \pm 50 \text{ eV}, \\ \Gamma_{\text{tot}} &= 23.5 \pm 5.0 \text{ MeV}, \\ \chi^2 &= 12.4 \text{ for 17 degrees of freedom.} \end{aligned} \quad (3)$$

All errors in (3) reflect statistical as well as systematic uncertainties based on our cross-section measurements and assumptions about signal and background shapes. The error in $M_{\psi''}$ includes the additional uncertainty (0.13%) in the absolute energy calibration of SPEAR. The error in $M_{\psi''}$ associated only with the fit is 1.9 MeV, reflecting an insensitivity to background shapes and normalization of R .

Figure 2 shows the observed cross section with




FIG. 2. The observed cross section for $D\bar{D}$ production.

$\psi(3095)$ and $\psi(3684)$ radiative tails and continuum background removed. The resonance is not corrected for self-radiative effects. It is therefore the measured cross section we attribute to D -meson pair production. The value at the peak ($E_{\text{c.m.}} = 3764$ MeV) is found to be 9.3 ± 1.4 nb.

In Table I, we compare these measurements with those of the two previous experiments.^{1,2}

The result we obtain for the partial leptonic width lies between the two earlier measurements. In the framework of the standard charmonium model¹³ the magnitude of Γ_{ee} suggests the need for mixing of the $\psi(3684)$ and $\psi(3770)$ by continuum states, in addition to the much smaller tensor-force mixing of bound states.³ We obtain a mixing angle of $20.3 \pm 2.8^\circ$ for the two levels. The total width we obtain is in good agreement with both experiments. We find, however, a mass between 6 and 8 MeV lower than those experiments. The systematic error associated with the storage-ring calibration can be removed by measuring masses relative to the $\psi(3684)$. This

TABLE I. Comparison of $\psi(3770)$ parameters. $M_{\psi''}$ is the mass of the ψ'' , Γ_{tot} the total width at the peak, and Γ_{ee} the partial leptonic width. ΔM is the mass difference of the $\psi(3684)$ and the $\psi(3770)$.

Experiment	$M_{\psi''}$ (MeV)	Γ_{tot} (MeV)	Γ_{ee} (eV)	ΔM (MeV)
LGW ^a	3772 ± 6	28 ± 5	345 ± 85	88 ± 3
DELCO ^b	3770 ± 6	24 ± 5	180 ± 60	86 ± 2^c
This experiment	3764 ± 5	24 ± 5	276 ± 50	80 ± 2

^a See Ref. 1.

^b See Ref. 2.

^c See Ref. 12.

mass difference (ΔM) is presented in Table I. Finally, we note that the peak cross section appears to be in reasonable agreement (about 10% lower) with the value 10.3 ± 2.5 nb reported earlier.¹

This work was supported primarily by the Department of Energy under Contract Nos. DE-AC03-76SF00515 and W7405-ENG-48. The work of M. T. was supported by Universitat Bonn. The work of H. Z. was supported by CEN-Saclay.

*Present address: California Institute of Technology, Pasadena, California 91125.

†Present address: Harvard University, Cambridge, Massachusetts 02138.

‡Present address: Universitat Bonn, Federal Republic of Germany.

§Present address: CEN-Saclay, France.

¹P. Rapidis *et al.*, Phys. Rev. Lett. **39**, 526 (1977); I. Peruzzi *et al.*, Phys. Rev. Lett. **39**, 1301 (1977).

²W. Bacino *et al.*, Phys. Rev. Lett. **40**, 671 (1978).

³E. Eichten *et al.*, Phys. Rev. D **17**, 3090 (1978); T. Appelquist *et al.*, Annu. Rev. Nucl. Part. Sci. **29**, 387 (1978).

⁴W. Davies-White *et al.*, Nucl. Instrum. Methods **160**, 227 (1979); G. S. Abrams *et al.*, Phys. Rev. Lett. **43**, 477 (1979); G. S. Abrams *et al.*, *ibid.* **43**, 481 (1979).

⁵J. -E. Augustin *et al.*, Phys. Rev. Lett. **34**, 764 (1975).

⁶S. J. Brodsky *et al.*, Phys. Rev. D **4**, 1532 (1971).

⁷G. Feldman, in *Proceedings of the 19th International Conference on High Energy Physics, Tokyo, 1978*, edited by S. Homma, M. Kawaguchi, and H. Miyazawa (Phys. Soc. of Japan, Tokyo, 1979).

⁸J. D. Jackson and D. L. Scharre, Nucl. Instrum. Methods **128**, 13 (1975); G. Bonneau and F. Martin, Nucl. Phys. **B27**, 381 (1971).

⁹A. M. Boyarski *et al.*, Phys. Rev. Lett. **34**, 1357 (1975); V. Luth *et al.*, *ibid.* **35**, 1124 (1975).

¹⁰A. Barbaro-Galtieri, in *Advances in Particle Physics*, edited by R. L. Cool and R. L. Marshak (Wiley, New York, 1968), Vol. II, p. 193.

¹¹Particle Data Group, Phys. Lett. **75B** (1978).

¹²W. Bacino, UCLA (private communication).

¹³Here $\psi(3684)$ is assigned the 2^3S_1 and $\psi(3770)$ is assigned the 1^3D_1 level.