

# UCLA

## UCLA Previously Published Works

### Title

Correction: Perfluorooctane Sulfonate Disturbs Nanog Expression through miR-490-3p in Mouse Embryonic Stem Cells.

### Permalink

<https://escholarship.org/uc/item/623345tt>

### Journal

PLoS One, 14(8)

### Authors

Xu, Bo

Chen, Xiaojiao

Mao, Zhilei

et al.

### Publication Date

2019

### DOI

10.1371/journal.pone.0221486

### Copyright Information

This work is made available under the terms of a Creative Commons Attribution License, available at <https://creativecommons.org/licenses/by/4.0/>


Peer reviewed

## CORRECTION

# Correction: Perfluorooctane Sulfonate Disturbs Nanog Expression through miR-490-3p in Mouse Embryonic Stem Cells

Bo Xu, Xiaojiao Chen, Zhilei Mao, Minjian Chen, Xiumei Han, Guizhen Du, Xiaoli Ji, Chunxin Chang, Virender K. Rehan, Xinru Wang, Yankai Xia

In [Fig 2b](#), the image appearing in the GADPH row is incorrect. Please see the complete, correct [Fig 2](#) here.




## OPEN ACCESS

**Citation:** Xu B, Chen X, Mao Z, Chen M, Han X, Du G, et al. (2019) Correction: Perfluorooctane Sulfonate Disturbs Nanog Expression through miR-490-3p in Mouse Embryonic Stem Cells. PLoS ONE 14(8): e0221486. <https://doi.org/10.1371/journal.pone.0221486>

**Published:** August 16, 2019

**Copyright:** © 2019 Xu et al. This is an open access article distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.



**Fig 2. Effects of PFOS on pluripotency and expressions of *miR-145*, *miR-490-3p* in mESCs.** Cells were cultured with various concentrations of PFOS (0.2  $\mu$ M, 2  $\mu$ M, 20  $\mu$ M, and 200  $\mu$ M) or DMSO as control for 24 h. (A) Oct-4/Sox-2/Nanog mRNA levels were determined by quantitative real-time PCR using a housekeeping gene GAPDH as an internal control. (B) The protein levels of Oct-4/Sox-2/Nanog were determined by Western blot analysis using GAPDH as an internal control. (C) miRNA levels (*miR-145*, *miR-490-3p*) were determined by quantitative real-time PCR and were normalized to U6 as an internal control. Each data point was normalized to the control (DMSO) and represented the means  $\pm$  S.E. from three independent experiments. (D) Relative protein levels of Oct4, Sox2 and Nanog. \* indicates significant difference when the values were compared to that of the control ( $p < 0.05$ ).

<https://doi.org/10.1371/journal.pone.0221486.g001>

## Reference

- Xu B, Chen X, Mao Z, Chen M, Han X, Du G, et al. (2013) Perfluorooctane Sulfonate Disturbs Nanog Expression through miR-490-3p in Mouse Embryonic Stem Cells. PLoS ONE 8(10): e74968. <https://doi.org/10.1371/journal.pone.0074968> PMID: 24098361