UC Davis

UC Davis Previously Published Works

Title

Transcervical sterilization: population sterilization rates overestimated.

Permalink

<https://escholarship.org/uc/item/6241v42h>

Journal

European journal of obstetrics, gynecology, and reproductive biology, 180(1)

ISSN

0301-2115

Author

Creinin, Mitchell D

Publication Date

2014-09-01

DOI

10.1016/j.ejogrb.2014.05.034

Peer reviewed

Hiromi Nakamura

Department of Obstetrics and Gynecology, Gifu Prefectural Tajimi Hospital, Tajimi, Gifu, Japan

* Corresponding author at: Department of Obstetrics and Gynecology, Gifu Prefectural Tajimi Hospital, Maebata-cho 5-161, Tajimi, Gifu 507-8522, Japan. Tel.: +81 572225311; ?fax: +81 572251246.
E-mail address: gyendoscopy@gmail.com (A. Takeda).

Received 16 April 2014
Accepted 22 April 2014

<http://dx.doi.org/10.1016/j.ejogrb.2014.04.028>

Conflict of interest

None.

Authors' contributions

I conceived and wrote the entire piece.

Authors' information

Dr Kieran Walsh is clinical director of BMJ Learning and Quality.

Acknowledgment

None.

Reference

- [1] Scheele F, Novak Z, Vetter K, Caccia N, Goverde A. Obstetrics and gynaecology training in Europe needs a next step. *Eur J Obstet Gynecol Reprod Biol* 2014 (May) pii: S0301-2115(14)00223-1.

Reforms needed to the system of obstetrics and gynaecology training in Europe

Dear Editor,

Scheele et al. have offered an interesting overview of the reforms needed to the system of obstetrics and gynaecology training in Europe [1].

Their suggestions are likely to be what is required – however it is sobering to review the number of reforms suggested. The suggestions include e-learning programmes, simulation, team training, professionalism development, reflective practice, continuous formative assessment and feedback, summative assessment for independent practice, faculty development, patient feedback, assessing quality of care, and assessing the quality of training. How will the average department of postgraduate training in obstetrics and gynaecology deliver all these reforms (eleven in total) and at the same time continue to provide clinical care and conduct research?

The answer might lie in packaging a number of these reforms together and seeking efficiencies as a result. There are a number of examples that come to mind almost immediately. Firstly simulation and team training need not be twin track ideas – they can be integrated. Simulation works best when it is a team based learning activity and as a corollary team learning works best when teams role play how they would perform in realistic situations. Secondly there is considerable overlap between the ideas of

e-learning programmes and simulation. With modern technology e-learning programmes can be interactive and multimedia and can simulate the real world with a virtual one. Equally simulation scenarios can be filmed and resultant media put online. Thirdly the paper mentions formative assessment and summative assessment – however there is similarly potential for integration here. The concept of progress testing allows the assessor to take the best of formative and summative assessment and develop a solution that works well for the examiners and examinees. Lastly and perhaps most importantly there should not be an artificial divide between assessing quality of care and assessing the quality of training. Training should only happen in a high quality care environment – in this circumstance the trainees will learn from experience and from their senior role models.

Scheele et al. have rightly presented the challenges that modern curriculum designers face – the next step will be strategic implementation. Resources might be limited and so we will have to think how to design curricula that are both effective and efficient.

Kieran Walsh*

BMJ Learning, BMJ Group, BMA House, Tavistock Square, London WC1H 9JR, United Kingdom

* Tel.: +44 7985 755 333; fax: +44 207 383 6242.
E-mail address: kmwalsh@bmjgroup.com (K. Walsh).

Received 10 June 2014
Accepted 14 June 2014

<http://dx.doi.org/10.1016/j.ejogrb.2014.06.014>

Transcervical sterilization: Population sterilization rates overestimated

Dear Editor,

I read with interest the study by Fernandez et al. [1] reporting the pregnancy rates in France between women who had sterilization by laparoscopic or hysteroscopic methods. Unfortunately, the authors fail to recognize a significant limitation of the dataset and the results, accordingly, are misleading to both patients and providers. Fernandez et al. [1] provide pregnancy rates from a national database in women who successfully had each procedure. For the patient, she enters a procedure or operating room to have a sterilization procedure performed and, with transcervical sterilization, she is significantly more likely to not have that procedure completed. In fact, the likelihood that a woman will be successfully sterilized within one year of the initial procedure is significantly lower with transcervical than laparoscopic sterilization [2]. All women attempting sterilization, not just those who successfully have the procedure, need to be considered when evaluating the public health benefits of a procedure. Because women who desire transcervical sterilization are significantly less likely to achieve sterilization, the pregnancy rate in the total population (not just those that successfully have the procedure) is significantly greater over 10 years in women desiring transcervical sterilization as compared to laparoscopic sterilization [3]. The conclusion of

Fernandez et al. [1] that the “lower pregnancy rate for sterilization by tubal microinserts justifies the development of this technique and its preference by the health authorities” is not supported by the totality of the available information in the literature.

Acknowledgement

Research: Merck, Medicines 360; Consulting: Merck, Danco.
Funding: None.

References

- [1] Fernandez H, Legendre G, Blein C, Lamarsalle L, Panel P. Tubal sterilization: pregnancy rates after hysteroscopic versus laparoscopic sterilization in France. *Eur J Obstet Gynecol Reprod Biol* 2006–2010 [published ahead of print 14 May 2014].
- [2] Gariepy AM, Creinin MD, Schwarz EB, Smith KJ. Reliability of laparoscopic compared with hysteroscopic sterilization at 1 year: a decision analysis. *Obstet Gynecol* 2011;118:273–9.
- [3] Gariepy AM, Creinin MD, Smith KJ, Xu X. Probability of pregnancy after sterilization: a comparison of hysteroscopic versus laparoscopic sterilization. *Contraception* 2014 [published ahead of print 28 April 2014].

Mitchell D. Creinin*

*Department of Obstetrics and Gynecology University of California,
Davis, Sacramento, CA 95817, USA*

* Corresponding author at: University of California, Davis 4860 Y
Street, Suite 2500, Sacramento, CA, USA. Tel.: + 916 784 6670.
E-mail address: mitchell.creinin@ucdmc.ucdavis.edu (M. Creinin).

Received 19 May 2014

Accepted 22 May 2014

<http://dx.doi.org/10.1016/j.ejogrb.2014.05.034>