

UCLA

UCLA Previously Published Works

Title

The association between coffee consumption and bladder cancer in the bladder cancer epidemiology and nutritional determinants (BLEND) international pooled study

Permalink

<https://escholarship.org/uc/item/62b2k3hc>

Journal

Cancer Causes & Control, 30(8)

ISSN

0957-5243

Authors

Yu, Evan Yi-Wen
Wesselius, Anke
van Osch, Frits
[et al.](#)

Publication Date

2019-08-01

DOI

10.1007/s10552-019-01191-1

Copyright Information

This work is made available under the terms of a Creative Commons Attribution License, available at <https://creativecommons.org/licenses/by/4.0/>

Peer reviewed



Published in final edited form as:

Cancer Causes Control. 2019 August ; 30(8): 859–870. doi:10.1007/s10552-019-01191-1.

The association between coffee consumption and bladder cancer in the bladder cancer epidemiology and nutritional determinants (BLEND) international pooled study

Evan Yi-Wen Yu^{1,2}, Anke Wesselius¹, Frits van Osch^{1,3}, Mariana Carla Stern⁴, Xuejuan Jiang⁴, Eliane Kellen⁵, Chih-Ming Lu⁶, Hermann Pohlabein⁷, Gunnar Steineck⁸, James Marshall⁹, Mohamed Farouk Allam¹⁰, Carlo La Vecchia¹¹, Kenneth C. Johnson¹², Simone Benhamou¹³, Zuo-Feng Zhang¹⁴, Cristina Bosetti¹⁵, Jack A. Taylor¹⁶, Maurice P. Zeegers^{1,2}

¹NUTRIM School for Nutrition and Translational Research in Metabolism, University of Maastricht, Universiteitssingel 40 (Room C5.564), 6229 ER Maastricht, The Netherlands

²CAPHRI School for Public Health and Primary Care, University of Maastricht, Maastricht, The Netherlands

³Institute of Cancer and Genomic Sciences, University of Birmingham, Birmingham, UK

⁴Department of Preventive Medicine, University of Southern California, Los Angeles, CA, USA

⁵Leuven University Centre for Cancer Prevention (LUCC), Louvain, Belgium

⁶Department of Urology, Buddhist Dalin Tzu Chi General Hospital, Dalin Township, Chiayi County, Taiwan

⁷Leibniz Institute for Prevention Research and Epidemiology-BIPS, Bremen, Germany

⁸Clinical Cancer Epidemiology, Department of Oncology, Clinical Sciences, Sahlgrenska Academy, University of Gothenburg, Gothenburg, Sweden

⁹Department of Cancer Prevention and Control, Roswell Park Cancer Institute, Buffalo, NY, USA

¹⁰Department of Medical and Surgical Specialties, Radiological Sciences and Public Health, Section of Public Health and Human Sciences, University of Brescia, Brescia, Italy

¹¹Department of Clinical Medicine and Community Health, University of Milan, Milan, Italy

¹²Department of Epidemiology and Community Medicine, University of Ottawa, Ottawa, ON, Canada

Open Access This article is distributed under the terms of the Creative Commons Attribution 4.0 International License (<http://creativecommons.org/licenses/by/4.0/>), which permits unrestricted use, distribution, and reproduction in any medium, provided you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons license, and indicate if changes were made.

✉Evan Yi-Wen Yu, evan.yu@maastrichtuniversity.nl.

Author contributions Study conception and design: AW and MPZ; Analyses and interpretation of data: EYY and FVO; Drafting of the manuscript: EYY; Revised the manuscript: AW, FVO, and MPZ; Provided the data: MCS, XJ, EK, CML, HP, GS, JM, MFA, CLV, KCJ, SB, ZFZ, CB, JAT, and MPZ; Approved the manuscript: all authors.

Electronic supplementary material The online version of this article (<https://doi.org/10.1007/s10552-019-01191-1>) contains supplementary material, which is available to authorized users.

Conflict of interest All the authors declare that they have no conflicts of interest.

Ethics approval Each participating study has been approved by the local ethic committee.

¹³INSERM U946, Variabilite Genetique et Maladies Humaines, Fondation Jean Dausset/CEPH, Paris, France

¹⁴Departments of Epidemiology, UCLA Center for Environmental Genomics, Fielding School of Public Health, University of California, Los Angeles (UCLA), Los Angeles, CA, USA

¹⁵Laboratory of Oncology, Istituto di Ricerche Farmacologiche “Mario Negri” IRCCS, Milan, Italy

¹⁶Epidemiology Branch and Epigenetic and Stem Cell Biology Laboratory, National Institute of Environmental Health Sciences, NIH, Research Triangle Park, NC, USA

Abstract

Background—Inconsistent results for coffee consumption and bladder cancer (BC) risk have been shown in epidemiological studies. This research aims to increase the understanding of the association between coffee consumption and BC risk by bringing together worldwide case–control studies on this topic.

Methods—Data were collected from 13 case–control comprising of 5,911 cases and 16,172 controls. Pooled multivariate odds ratios (ORs), with corresponding 95% confidence intervals (CIs), were obtained using multilevel logistic regression models. Furthermore, linear dose–response relationships were examined using fractional polynomial models.

Results—No association of BC risk was observed with coffee consumption among smokers. However, after adjustment for age, gender, and smoking, the risk was significantly increased for never smokers (ever vs. never coffee consumers: OR_{model2} 1.30, 95% CI 1.06–1.59; heavy (> 4 cups/day) coffee consumers vs. never coffee consumers: OR_{model2} 1.52, 95% CI 1.18–1.97, p trend = 0.23). In addition, dose–response analyses, in both the overall population and among never smokers, also showed a significant increased BC risk for coffee consumption of more than four cups per day. Among smokers, a significant increased BC risk was shown only after consumption of more than six cups per day.

Conclusion—This research suggests that positive associations between coffee consumption and BC among never smokers but not smokers.

Keywords

Bladder cancer; Coffee consumption; Smoking; Dose–response analyses; Population-attributable risk

Introduction

Bladder cancer (BC) is the most common malignancy of the urinary tract and the seventh cause of death from cancer (2.8% of all cancer deaths), with nearly 430,000 new diagnoses and 165,000 deaths per year worldwide [1, 2]. Three-quarters of all BC cases occur in men [3], and most BC cases occur in the United States, Canada, and the European Union [4–7]. As with many solid tumors, BC incidence increases with age and it rarely occurs before the age of 40–50 years [8]. Given that this cancer is easy to relapse, BC is reported to be the most expensive life-time treatment of all cancers ranging from €80,000 to €160,000 per patient [9]. The strongest risk factors for BC occurrence, such as tobacco smoking and

harmful chemicals [10], have long been identified. However, as the bladder is an excretory organ, the role of fluid consumption in the development of BC could also be important.

Coffee is one of the most consumed beverages in the world. Since early 1970s, the possible relationship between coffee consumption and BC has been of considerable interest, when Cole et al. [11] suggested for the first time that coffee was a potential risk factor for BC. The International Agency for Research on Cancer (IARC) Monographs Programme in 1991 stated that there was limited evidence on the effect of coffee consumption on the BC risk and subsequently classified coffee as “possibly carcinogenic” (*group 2B; Monographs Volume 51*) [12]. Since then, several epidemiological studies focused on the relationship between coffee consumption and BC, however, results remained inconclusive [13–27]. In May 2016, a subsequent IARC Working Group of 23 scientists from 10 countries met to evaluate the carcinogenicity of drinking coffee and concluded that: “no consistent evidence of an association with drinking coffee, or of an exposure–response gradient”. This conclusion was based on evidence from 10 cohort studies and several population-based case–control studies conducted in Europe, the United States, and Japan [28]. An explanation for this inconsistency may be that previous studies on the relation between coffee consumption and BC risk lacked sufficient sample size to identify a significant association. In addition, since heavy coffee consumption is shown to be strongly associated with tobacco smoking [29], positive associations reported in some studies could possibly have been due to inadequate control for tobacco smoking. Moreover, several studies showed smoking to be interactive with caffeine in coffee [30–35], and, thereby, lead to misleading results in un-stratified analysis on the relation between coffee consumption and BC risk.

The present study aims to update the understanding and find more conclusive answers on the influence of coffee consumption on the BC risk by bringing together available case–control studies on the topic including almost 6,000 BC cases.

Methods

Study population

Data were derived from the BLadder cancer Epidemiology and Nutritional Determinants study (BLEND). BLEND is a large international epidemiology consortium aimed to pool data from available epidemiological studies on diet and BC. For the present study, 13 case–control studies (including 5,911 cases/16,172 controls), originated from nine different countries in three continents (i.e., Europe, North America, and Asia) had sufficient information on coffee consumption to be eligible for inclusion. BC cases were diagnosed and histologically confirmed through each study center of the included individual studies, with International Classification of Diseases (ICD) nine or ten. Most of the BC cases were identified in 1990s.

Data collection and coding

Details on the methodology of the BLEND consortium have been described elsewhere [36]. Taking into account the local context of the included studies, different dietary assessment methods were adopted: (1) self-administrated food frequency questionnaire (FFQ) were used

in Germany-1 [37], USA-2 [38], Canada-1 [39], France-1 [40], USA-3 [41], USA-4 [42]; (2) FFQ administered by a trained interviewer were used in USA-1 [43], Belgium-1 [44], China-1 [45], Sweden-1 [46], Spain-1 [47], Italy-1 [48], Italy-2 [49]. Coffee consumption was categorized using the hierarchical Eurocode 2 food coding system developed by the European Union [50]. To obtain unified consumption across studies, weekly, monthly, or yearly coffee cups were converted to daily cups of coffee consumption.

In addition to information on coffee and other dietary intake data, the BLEND dataset also included data on: study characteristics (design, method of dietary assessment, recall time of dietary consumption, and geographical region), participant demographics (age and gender), smoking status, and smoking pack-years.

Statistical analyses

To assess the influence of coffee consumption on the BC risk, multilevel logistic regression analyses were used to estimate the pooled odds ratios (ORs) and 95% confidence intervals (CIs). Coffee consumption was expressed as: (1) ever (individuals who drank coffee > 0 cup/day) or never consumption; (2) based on the available data, coffee consumption was divided into six categories: never, 0–1 cup/day, 1–2 cups/day, 2–3 cups/day, 3–4 cups/day, and more than four cups/day; (3) caffeinated or decaffeinated coffee consumption. In addition, standardized analysis on coffee cup size was performed. For this we transformed a United States (U.S.) cup size to a 237 ml coffee cup size according to U.S. Food and Drug Administration [51] and an Asian cup size to a 500 ml cup size according to the questionnaire used in the study of Lu et al. [45].

The logistic regression models used never coffee consumers as the reference group and were computed as “crude model” (model 1), adjusted for age, gender, smoking (model 2), or fully adjustment (model 3) additionally adjusted other fluid consumption (i.e., water, liquid milk, tea, alcohol, carbonated drink, and juice). Smoking was defined as: 0 (never smokers); 1 [current light smokers (i.e., smoking less than 20 pack-years)]; 2 (current heavy smokers (i.e., smoking more than 20 pack-years)); 3 (former light smokers (i.e., smokers who ceased smoking over 1 year and smoked less than 20 pack-years)); 4 [former heavy smokers (i.e., smokers who ceased smoking over 1 year and smoked more than 20 pack-years)]. In addition, the effect of ever versus never coffee consumption was also assessed using a meta-analysis approach; for this, pooled ORs (PORs) and 95% CIs were calculated by using a random-effect model stratified by geographical regions (i.e., Europe, North America and Asia) and study designs (i.e., hospital-based case–control studies and population-based case–control studies). Due to the lack of data, the influence of caffeine on BC risk was only assessed by comparing ever (caffeinated vs. decaffeinated) versus never coffee consumers based on multilevel logistic regression (model 2). To understand the relevance of the effect modification, the interaction terms between coffee consumption and age, gender, smoking status with pack-years were added to the model 2. $P_{\text{interaction}} < 0.10$ was considered statistically significant where upon analyses were stratified for the covariate of interest to understand the relevance of the effect modification.

In our secondary analysis, a potential dose–response relationship between coffee consumption and BC was assessed by using fractional polynomial regression, in which

the best fitting second order fractional polynomial regression model was defined as the model with the lowest deviance [52, 53]. A likelihood ratio test was used to assess the difference between the nonlinear and linear models to test for nonlinearity [54]. The results of the dose–response analyses were presented for each one coffee cup/day increment up to ten cups/day with stratification by smoking status (i.e., ever smokers and never smokers) and (i.e., hospital-based case–control studies and population-based case–control studies). Adjustments (model 2) were made for age, gender, and smoking (in overall population).

Finally, the population-attributable risk (PAR) of heavy coffee consumption (i.e., > 4 cups/day) on BC risk was estimated for Europe and North America, using the pooled risk estimates and the proportion of BC incidence in the population of interest.

All statistical analyses were performed with *STATA* version 14 SE (Stata Corporation, Texas, USA). *P* values below 0.05 were considered statistically significant.

Results

Baseline characteristics

The baseline characteristics of the study population are shown in Table 1. Of more than 22,000 participants, 5,911 cases of BC (4,639 men, 1,272 women) were identified. The median age at baseline was 61.4 years for cases and 57.2 years for controls, respectively. Approximately 35% of participants reported drinking coffee more than four cups per day, with an average consumption of four cups/day overall. At baseline, a higher coffee consumption was observed among smokers (five cups/day) compared to never smokers (three cups/day). In addition, coffee consumption showed strong interaction with smoking status as well as pack-years (*p* interaction: 0.001 and 0.042, respectively), while not interaction was found with age (*p* interaction: 0.17) and gender (*p* interaction: 0.16).

Associations between coffee consumption and BC risk

Ever versus never coffee consumption—The results comparing ever versus never coffee consumers are shown in Table 2 and Fig. 1. Overall, after adjustment for possible confounders, no statistically significant difference in BC risk could be observed between coffee consumers versus never coffee consumers (OR_{model2} 1.11, 95% CI 0.98–1.26; OR_{model3} 1.09, 95% CI 0.94–1.25). Among never smokers, a statistically significant association between coffee consumption and the risk of BC was found after further adjustment (OR_{model2} 1.30, 95% CI 1.06–1.59; OR_{model3} 1.31, 95% CI 1.03–1.66). For smokers, no significant association was observed comparing ever versus never coffee consumers. However, the estimates for former light smokers showed to be slightly higher than the estimates for other smokers. In addition, the meta-analysis stratified by geographical regions presented similar PORs based on model 2 (POR_{Overall} 1.11, 95% CI 0.96–1.25; POR_{Europe} 1.13, 95% CI 0.88–1.37; POR_{North America} 1.11, 95% CI 0.88–1.34); with low heterogeneity ($I^2 = 0.0\%$; *p* = 0.57).

Categories of coffee consumption with BC risk—The results of multilevel logistic regressions for subsequent categories of coffee consumption are shown in Table 2. Overall, coffee consumption of more than four cups/day results in an increased BC risk of 1.27 (95%

CI 1.11–1.46, p trend = 0.05, model 2) compared to never coffee consumers. A similar increased risk was observed among never smokers when comparing high (> 4 cups/day) coffee consumption to never coffee consumers (OR_{model2} 1.52, 95% CI 1.19–1.94, p trend = 0.23). Among smokers no significant association could be observed; however, former light smokers showed again slightly higher and borderline significant results compared to other type of smokers (OR_{model2} 1.41, 95% CI 0.99–2.02, p trend = 0.06). The coffee cup size standardized analysis also showed significantly increased BC risk estimates with more than four cups/day coffee consumption (Supplementary Table 3).

Results for the comparison of caffeinated and decaffeinated coffee are shown in Supplementary Tables 1 and 2. For the analysis on caffeinated coffee consumption, only 1 (France-1) out of the 13 included case–control studies could be included (including 187 cases/296 controls). For the analysis on decaffeinated coffee, two studies (Italy-1 and USA-4) had sufficient data to be included in our analyses (including 1,048 cases/1,487 controls). Consumers of caffeinated coffee showed significant increased risks (compared to decaffeinated coffee consumers: OR_{model2} 1.88, 95% CI 1.42–2.48; compared to never coffee consumers: OR_{model2} 1.52, 95% CI 1.06–2.21), whereas, decaffeinated coffee consumers showed a null association with BC compared to never coffee consumers.

Dose–response analyses

Dose–response relationships between coffee consumption and the risk of BC are displayed in Fig. 2. The tests for nonlinearity were not statistically significant; hence, linear models were applied in the dose–response analyses. The curves for the overall population showed a slightly increased BC risks with the increment of coffee consumption. Similar results were found among ever and never smokers; however, among never smokers the increased BC risk was significant for consuming over four cups/day, while for ever smokers for consumption over six cups/day. Adjusted ORs and 95% CIs for one cup/day increment were 1.14 (95% CI 1.05–1.24) in overall study population, 1.05 (95% CI 1.02–1.15) in ever smokers, 1.16 (95% CI 1.04–1.41) in never smokers. In addition, both curves for hospital-based studies and population-based studies showed increased BC risks with the increment of cups of coffee consumption per day. However, in hospital-based studies a statistically significant increase BC risk was observed for consuming over two cups/day, while for population-based studies a significant increase was observed for consuming over five cups/day (Supplementary Fig. 2).

Population-attributable risks

Assuming an incidence of heavy coffee drinking of 28% and 17%, in Europe and North America respectively, a PAR of 7.94% for Europe and 4.45% for North America was observed.

Discussion

This large multi centric study found an overall increased risk of BC with high (> 4 cups/day) coffee consumption. In addition, we showed that this increased risk was only observed among never smokers, but not found among smokers.

The interaction between coffee and smoking has already been studied with some detail, and experimental studies showed that smokers eliminate caffeine faster, suggesting that the effect of coffee consumption on BC risk is lower among smokers [33, 55]. In fact, faster metabolism of caffeine in smokers would allow them to consume higher levels before experiencing symptoms of caffeine toxicity [56–58]. Moreover, this hypothesis is strengthened by experimental studies reporting that the cytochrome P450 1A2 (CYP1A2) metabolic pathway is upregulated by both caffeine and compounds in tobacco smoke, including nicotine and polycyclic aromatic hydrocarbons (PAHs) [59–61], so that the effect of caffeine is potentially weaker among smokers than among never smokers. These experimental studies are in line with our result showing an increased BC risk among never smokers only. Several epidemiological studies (both case–control and cohort) also suggested a higher BC risks for coffee consumers among never smokers only or a null association among smokers [40, 62–66]. A meta-analysis of epidemiological studies also suggested that the increased BC risk of coffee consumption was higher among never smokers than among smokers [31]. However, a more recent prospective cohort study (2017) [67] conducted in the United States, which showed that high coffee consumption (> 4 cups/day) was positively associated with BC risk, could not observe an increased BC risk among never smokers only. This discrepancy may be due to the limited number of cases in their smoking-stratified analysis.

Coffee contains high content of caffeine, which has shown mutagenic effects in human cells [68], a proven influence on suppressing the activation of the protein kinase ataxia-telangiectasia mutated (ATM) and the phosphorylation of the kinase Chk2, both important for the activation of the tumor suppressor gene P53 [69, 70]. It could, therefore, be suggested that the increased BC risk is due to the caffeine content of coffee. Previous epidemiological studies already confirmed this hypothesis by showing a null association between BC risk and decaffeinated coffee [71] and an increased BC risk for caffeinated coffee [72, 73]. The present study also showed an increased BC risk when comparing caffeinated coffee consumers versus decaffeinated coffee consumers. However, due to lack of data, we were unable to perform further analyses, i.e., the association between the intensity of caffeinated coffee consumption and BC.

Besides caffeine, coffee also included other compounds, such as several phenolic compounds (i.e., chlorogenic, caffeic acid, ferulic, and coumaric acids), melanoidins, and diterpenes (i.e., cafestol and kahweol), with anti-carcinogenic properties [74–80]. This might explain why the consumption of less than four coffee cups/day showed a null association with BC risk. In addition, experimental research on the mechanisms of action of coffee compounds on P53 suppression showed this effect to be concentration dependent [69, 81–84].

The present study found a four cups/day threshold for an increased BC risk in both the overall study population and among never smokers. This finding is in line with previous reported meta-analyses based on case–control studies, also showing an increased BC risk over four cups/day (respectively OR 1.29, 95% CI 1.12–1.48; OR 1.20, 95% CI 1.12–1.24) [31, 85]. Among smokers, however, a threshold of six cups/day was observed; this again

shows that the effect of coffee on BC risk might be influenced by the faster caffeine metabolism among smokers.

An issue that may arise in evaluating the influence of coffee consumption on BC risk in hospital-based case–control studies is that the controls includes patients suffering from a disease that may influence coffee intake (i.e., heart issues), resulting in inflated ORs. In the present study, therefore, stratified analyses by study design (i.e., hospital-based controls or population-based controls) were performed, and the association with coffee consumption was higher, though not significantly, in hospital-based studies than it was in population-based studies, where similar results were found in both ever versus never coffee consumption and dose–response analyses (Supplementary Figs. 1 and 2).

For the present study, the estimates of PAR showed that in Europe 7.94% and in North America 4.45% of incidence of BC cases could be attributed to heavy coffee consumption (i.e., > 4 cups/day). Unfortunately, the exact prevalence of heavy coffee drinking in the Eastern Asian population is yet unknown. However, since anecdotal evidence suggests that coffee consuming is on the rise in Eastern Asia, the PAR is expected to increase Eastern Asian countries over the next years.

Among the strengths of the BLEND study there is the large sample size, allowing to perform detailed analyses with enough statistical power to detect smaller effects, the study has also some limitations. First, it is known that the size of standard coffee cups is varied around the world, and the effect of a cup of coffee on BC might, therefore, differ between different countries. However, our per-center analyses as well our standardized analysis, in which we transformed a United States (U.S.) cup size to a 237 ml coffee cup size according to U.S. Food and Drug Administration [51] and an Asian cup size to a 500 ml cup size according to the questionnaire used in the study of Lu et al. [45], showed similar results (Supplementary Table 3: $OR_{\text{model2}} 1.25$, 95% CI 1.06–1.47, p trend = 0.18). Due to large heterogeneity among coffee cup size among European countries, Europe was not included in these standardized analyses. In addition, it is suggested that the strength of coffee brew may compensate for the different serving size between countries [86].

Second, it is often suggested that case–control studies are limited in showing causal relation, due to the potential recall bias of case–control studies. This might have led to a lower reliability of the results compared to those of cohort studies. However, although this issue has been addressed and analyzed for its consequences in many epidemiological/methodological papers [87–90], no clear answer on the magnitude of the effect of this specific type of bias could be draw.

Thirdly, limited information was available on possible risk factors, other than age, gender and smoking, for the development of BC, such as body mass index (BMI), physical activity, socioeconomic status (SES), disinfection byproducts, arsenic in the drinking water, and occupational exposures to potentially carcinogenic chemicals. Although, adjustments for these factors could have influenced the results, current literature shows that only a small proportion of BC cases can be attributed to these factors [72]. Last, although status as well as duration and intensity of smoking were taken into account in our analysis, the

adjustment for smoking might still be imperfect due to differences in smoking practices (e.g., depth of inhalation or amount of inhalation), or differences in types of smoke exposure [67]. In addition, since smoking is perceived as a socially undesirable behavior, the use of self-reported questionnaires for smoking status, duration, and intensity might have led to underreporting of the actual smoking habits.

Conclusion

In summary, the present study, with more than 5,900 cases, observed an increased risk between high (> 4 cups/day) coffee consumption and BC among never smokers, while no association with BC risk was observed with coffee consumption among smokers. Additionally, it indicates that around 7.94% of BC cases for Europe and 4.45% of BC cases for North America in the population might be attributable to heavy coffee consumption (> 4 cups/day).

Supplementary Material

Refer to Web version on PubMed Central for supplementary material.

Acknowledgments

We gratefully acknowledge all principal investigators for their willingness to participate in this jointed project. The author E.Y.W. YU gives thanks to the financial support from China Scholarship Council (No. 201706310135).

Funding

This work was partly funded by the World Cancer Research Fund International (WCRF 2012/590) and European Commission. The Hessen Case-control study on bladder cancer was supported by the Bundesanstalt für Arbeitsschutz (No. F 1287). The Kaohsiung study was supported by grant NSC 85-2332-B-037-066 from the National Scientific Council of the Republic of China. The Stockholm Case-control study was supported by grant from the Swedish National Cancer Society and from the Swedish Work Environment Fund. The New England bladder cancer study was funded in part by Grant Nos. 5 P42 ES007373 from the National Institute of Environmental Health Sciences, NIH and CA57494 from the National Cancer Institute, NIH. The Italian Case-control study on bladder cancer was conducted within the framework of the CNR (Italian National Research Council) Applied Project “Clinical Application of Oncological Research” (contracts: 94.01321.PF39 and 94.01119.PF39), and with the contributions of the Italian Association for Cancer Research, the Italian League against Tumours, Milan, and Mrs. Angela Marchegiano Borgomainerio. The Brescia bladder cancer study was partly supported by the International Agency for Research on Cancer. The French INSERM study was supported by a grant from the Direction Générale de la Santé, Ministère des Affaires Sociales, France. The Molecular Epidemiology of Bladder Cancer and Prostate Cancer was supported in part by grants ES06718 (to Z.-F.Z.), U01 CA96116 (to A.B.), and CA09142 from the NIH National Institute of Environmental Health Sciences, the National Cancer Institute, the Department of Health and Human Services, and by the Ann Fitzpatrick Alper Program in Environmental Genomics at the Jonsson Comprehensive Cancer Centre, UCLA.

Abbreviations

BC	Bladder cancer
IARC	International Agency for Research on Cancer
BLEND	Bladder cancer epidemiology and nutritional determinants
FFQ	Food frequency questionnaires
OR	Odds ratio

CI	Confidence interval
POR	Pooled odds ratio
CYP1A2	Cytochrome P450 1A2
PAHs	Polycyclic aromatic hydrocarbons
ATM	Ataxia-telangiectasia mutated
PAR	Population-attributable risk
BMI	Body mass index
SES	Socioeconomic status

References

1. Ferlay J, Soerjomataram I, Dikshit R et al. (2015) Cancer incidence and mortality worldwide: sources, methods and major patterns in GLOBOCAN 2012. *Int J Cancer* 136(5):E359–E386 [PubMed: 25220842]
2. Siegel RL, Miller KD, Jemal A (2017) Cancer statistics, 2017. *CA Cancer J Clin* 67(1):7–30 [PubMed: 28055103]
3. Antoni S, Ferlay J, Soerjomataram I et al. (2017) Bladder cancer incidence and mortality: a global overview and recent trends. *Eur Urol* 71(1):96–108 [PubMed: 27370177]
4. Jemal A, Bray F, Center MM et al. (2011) Global cancer statistics. *CA Cancer J Clin* 61(2):69–90 [PubMed: 21296855]
5. Ploeg M, Aben KK, Kiemeny LA (2009) The present and future burden of urinary bladder cancer in the world. *World J Urol* 27(3):289–293 [PubMed: 19219610]
6. Marugame T, Mizuno S (2005) Comparison of prostate cancer mortality in five countries: France, Italy, Japan, UK and USA from the WHO mortality database (1960–2000). *Jpn J Clin Oncol* 35(11):690–691 [PubMed: 16306486]
7. Stewart B, Wild CP (2017) World cancer report 2014. International Agency for Research on Cancer, World Health Organization. WHO Press, Geneva
8. Letašiová S, Medveová A, Šovčíková A et al. (2012) Bladder cancer, a review of the environmental risk factors. *Environ Health* 11(Suppl 1):S11 [PubMed: 22759493]
9. Botteman MF, Pashos CL, Redaelli A, Laskin B, Hauser R (2003) The health economics of bladder cancer. *Pharmacoeconomics* 21(18):1315–1330 [PubMed: 14750899]
10. Johansson SL, Cohen SM (1997) Epidemiology and etiology of bladder cancer. In: *Seminars in surgical oncology*. Wiley, New York
11. Cole P (1971) Coffee-drinking and cancer of the lower urinary tract. *Lancet* 1(7713):1335–1337 [PubMed: 4103399]
12. IARC Working Group on the Evaluation of Carcinogenic Risks to Humans (1991) Coffee, tea, mate, methylxanthines and methylglyoxal: World Health Organization. International Agency for Research on Cancer, vol 51
13. Pannelli F, La Rosa F, Saltalamacchia G et al. (1989) Tobacco smoking, coffee, cocoa and tea consumption in relation to mortality from urinary bladder cancer in Italy. *Eur J Epidemiol* 5(3):392–397 [PubMed: 2792314]
14. Risch HA, Burch JD, Miller AB et al. (1988) Dietary factors and the incidence of cancer of the urinary bladder. *Am J Epidemiol* 127(6):1179–1191 [PubMed: 3369417]
15. Ciccone G, Vineis P (1988) Coffee drinking and bladder cancer. *Cancer Lett* 41(1):45–52 [PubMed: 3390802]

16. Jensen OM, Wahrendorf J, Knudsen JB, Sorensen BL (1986) The Copenhagen case-control study of bladder cancer. II. Effect of coffee and other beverages. *Int J Cancer* 37(5):651–657 [PubMed: 3699928]
17. Bravo P, del Rey J, Sanchez J, Conde M (1986) Coffee and analgesics as risk factors for cancer of the bladder. *Arch Esp Urol* 39(5):337–341 [PubMed: 3753045]
18. Rebelakos A, Trichopoulos D, Tzonou A et al. (1985) Tobacco smoking, coffee drinking, and occupation as risk factors for bladder cancer in Greece. *J Natl Cancer Inst* 75(3):455–461 [PubMed: 3861898]
19. Gonzalez CA, Lopez-Abente G, Errezola M et al. (1985) Occupation, tobacco use, coffee, and bladder cancer in the county of Mataro (Spain). *Cancer* 55(9):2031–2034 [PubMed: 3978583]
20. Hopkins J (1984) Coffee drinking and bladder cancer. *Food Chem Toxicol* 22(6):481–483 [PubMed: 6539739]
21. Marrett LD, Walter SD, Meigs JW (1983) Coffee drinking and bladder cancer in Connecticut. *Am J Epidemiol* 117(2):113–127 [PubMed: 6829543]
22. Hartge P, Hoover R, West DW, Lyon JL (1983) Coffee drinking and risk of bladder cancer. *J Natl Cancer Inst* 70(6):1021–1026 [PubMed: 6574270]
23. Howe GR, Burch JD, Miller AB et al. (1980) Tobacco use, occupation, coffee, various nutrients, and bladder cancer. *J Natl Cancer Inst* 64(4):701–713 [PubMed: 6928984]
24. Abou-Daoud KT (1980) Cancer of the bladder and cigarette smoking, coffee and alcohol drinking in Lebanon. *J Med Liban* 31(3):251–257 [PubMed: 7218288]
25. Bross ID, Tidings J (1973) Another look at coffee drinking and cancer of the urinary bladder. *Prev Med* 2(3):445–451 [PubMed: 4774263]
26. Zeitlin BR (1972) Coffee and bladder cancer. *Lancet* 1(7759):1066
27. Fraumeni JF Jr, Scotto J, Dunham LJ (1971) Coffee-drinking and bladder cancer. *Lancet* 2(7735):1204
28. Loomis D, Guyton KZ, Grosse Y et al. (2016) Carcinogenicity of drinking coffee, mate, and very hot beverages. *Lancet Oncol* 17(7):877–878 [PubMed: 27318851]
29. Johan HB, Nordestgaard AT, Taylor AE et al. (2017) Heavier smoking increases coffee consumption: findings from a Mendelian randomization analysis. *Int J Epidemiol* 46(6):1958–1967 [PubMed: 29025033]
30. Swanson JA, Lee JW, Hopp JW (1994) Caffeine and nicotine: a review of their joint use and possible interactive effects in tobacco withdrawal. *Addict Behav* 19(3):229–256 [PubMed: 7942243]
31. Zhou Y, Tian C, Jia C (2012) A dose-response meta-analysis of coffee consumption and bladder cancer. *Prev Med* 55(1):14–22 [PubMed: 22564775]
32. Benowitz NL, Peng M, Jacob P (2003) Effects of cigarette smoking and carbon monoxide on chlorzoxazone and caffeine metabolism. *Clin Pharmacol Ther* 74(5):468–474 [PubMed: 14586387]
33. Zevin S, Benowitz NL (1999) Drug interactions with tobacco smoking. *Clin Pharmacokinet* 36(6):425–438 [PubMed: 10427467]
34. De Leon J, Diaz FJ, Rogers T et al. (2003) A pilot study of plasma caffeine concentrations in a US sample of smoker and nonsmoker volunteers. *Prog Neuropsychopharmacol Biol Psychiatry* 27(1):165–171 [PubMed: 12551740]
35. Plowchalk DR, Yeo KR (2012) Prediction of drug clearance in a smoking population: modeling the impact of variable cigarette consumption on the induction of CYP1A2. *Eur J Clin Pharmacol* 68(6):951–960 [PubMed: 22258279]
36. Goossens ME, Isa F, Brinkman M et al. (2016) International pooled study on diet and bladder cancer: the bladder cancer, epidemiology and nutritional determinants (BLEND) study: design and baseline characteristics. *Arch Public Health* 74(1):30–39 [PubMed: 27386115]
37. Pohlbeln H, Jockel KH, Bolm-Audorff U (1999) Non-occupational risk factors for cancer of the lower urinary tract in Germany. *Eur J Epidemiol* 15(5):411–419 [PubMed: 10442466]
38. Mettlin C, Graham S (1979) Dietary risk factors in human bladder cancer. *Am J Epidemiol* 110(3):255–263 [PubMed: 582494]

39. Johnson K, Mao Y, Argo J et al. (1998) The National Enhanced Cancer Surveillance System: a case-control approach to environment-related cancer surveillance in Canada. *Environmetrics* 9(5):495–504
40. Clavel J, Cordier S (1991) Coffee consumption and bladder cancer risk. *Int J Cancer* 47(2):207–212 [PubMed: 1988365]
41. Cao W, Cai L, Rao JY et al. (2005) Tobacco smoking, GSTP1 polymorphism, and bladder carcinoma. *Cancer* 104(11):2400–2408 [PubMed: 16240451]
42. Taylor JA, Umbach DM, Stephens E et al. (1998) The role of N-acetylation polymorphisms in smoking-associated bladder cancer: evidence of a gene-gene-exposure three-way interaction. *Cancer Res* 58(16):3603–3610 [PubMed: 9721868]
43. Jiang X, Castela JE, Groshen S et al. (2007) Alcohol consumption and risk of bladder cancer in Los Angeles County. *Int J Cancer* 121(4):839–845 [PubMed: 17440923]
44. Kellen E, Zeegers M, Paulussen A, Van Dongen M, Buntinx F (2006) Fruit consumption reduces the effect of smoking on bladder cancer risk. The Belgian case control study on bladder cancer. *Int J Cancer* 118(10):2572–2578 [PubMed: 16380991]
45. Lu CM, Lan SJ, Lee YH et al. (1999) Tea consumption: fluid intake and bladder cancer risk in Southern Taiwan. *Urology*. 54(5):823–828 [PubMed: 10565741]
46. Steineck G, Hagman U, Gerhardsson M, Norell SE (1990) Vitamin A supplements, fried foods, fat and urothelial cancer A case-referent study in Stockholm in 1985–87. *Int J Cancer* 45(6):1006–1011 [PubMed: 2351481]
47. Baena AV, Allam MF, Del Castillo AS et al. (2006) Urinary bladder cancer risk factors in men: a Spanish case-control study. *Eur J Cancer Prev* 15(6):498–503 [PubMed: 17106329]
48. Shen M, Hung RJ, Brennan P et al. (2003) Polymorphisms of the DNA repair genes XRCC1, XRCC3, XPD, interaction with environmental exposures, and bladder cancer risk in a case-control study in northern Italy. *Cancer Epidemiol Biomark Prev* 12(11 Pt 1):1234–1240
49. D'Avanzo B, La Vecchia C, Negri E, Decarli A, Benichou J (1995) Attributable risks for bladder cancer in northern Italy. *Ann Epidemiol* 5(6):427–431 [PubMed: 8680604]
50. Poortvliet E, Klensin J, Kohlmeier L (1992) Rationale document for the Eurocode 2 food coding system (version 91/2). *Eur J Clin Nutr* 46(Suppl 5):S9–S24
51. US Food Drug Administration (2012) Guidance for industry: drug interaction studies—study design, data analysis, implications for dosing, and labeling recommendations. US Food and Drug Administration, Rockville, pp 1–75
52. Royston P, Altman DG (1997) Approximating statistical functions by using fractional polynomial regression. *J R Stat Soc* 46(3):411–422
53. Jones BL, Nagin DS (2012) A Stata plugin for estimating group-based trajectory models. *Sociol Methods Res* 42(4):608–613
54. Bagnardi V, Zambon A, Quatto P, Corrao G (2004) Flexible meta-regression functions for modeling aggregate dose-response data, with an application to alcohol and mortality. *Am J Epidemiol* 159(11):1077–1086 [PubMed: 15155292]
55. Mihi YTK, Takahiko K, Koji M (1998) Effects of lifestyle and genetic polymorphisms on consumption of coffee or black tea and urinary caffeine levels. *Biomarkers* 3(4–5):367–377 [PubMed: 23899363]
56. Joeres R, Klinker H, Heusler H et al. (1988) Influence of smoking on caffeine elimination in healthy volunteers and in patients with alcoholic liver cirrhosis. *Hepatology* 8(3):575–579 [PubMed: 3371873]
57. Langmann P, Bienert A, Zilly M et al. (2000) Influence of smoking on cotinine and caffeine plasma levels in patients with alcoholic liver cirrhosis. *Eur J Med Res* 5(5):217–221 [PubMed: 10806124]
58. Ossip DJ, Epstein LH (1981) Relative effects of nicotine and coffee on cigarette smoking. *Addict Behav* 6(1):35–39 [PubMed: 7257914]
59. Landi M, Sinha R, Lang N, Kadlubar F (1999) Human cytochrome P4501A2. *IARC Sci Publ* 148:173–195
60. Kalow W, Tang BK (1991) Use of caffeine metabolite ratios to explore CYP1A2 and xanthine oxidase activities. *Clin Pharmacol Ther* 50(5–1):508–519 [PubMed: 1934864]

61. Gunes A, Dahl M-L (2008) Variation in CYP1A2 activity and its clinical implications: influence of environmental factors and genetic polymorphisms. *Pharmacogenomics* 9(5):625–637 [PubMed: 18466106]
62. Donato F, Boffetta P, Fazioli R et al. (1997) Bladder cancer, tobacco smoking, coffee and alcohol drinking in Brescia, northern Italy. *Eur J Epidemiol* 13(7):795–800 [PubMed: 9384269]
63. Vena JE, Freudenheim JO, Graham S et al. (1993) Coffee, cigarette smoking, and bladder cancer in western New York. *Ann Epidemiol* 3(6):586–591 [PubMed: 7921304]
64. Escolar Pujolar A, Gonzalez CA, Lopez-Abente G et al. (1993) Bladder cancer and coffee consumption in smokers and non-smokers in Spain. *Int J Epidemiol* 22(1):38–44 [PubMed: 8449645]
65. D'Avanzo B, La Vecchia C, Franceschi S et al. (1992) Coffee consumption and bladder cancer risk. *Eur J Cancer* 28(8–9):1480–1484
66. Hashemian M, Sinha R, Murphy G et al. (2019) Coffee and tea drinking and risk of cancer of the urinary tract in male smokers. *Ann Epidemiol*. 10.1016/j.annepidem.2019.03.014
67. Loftfield E, Freedman ND, Inoue-Choi M, Graubard BI, Sinha R (2017) A prospective investigation of coffee drinking and bladder cancer incidence in the United States. *Epidemiology* 28(5):685–693 [PubMed: 28768299]
68. Kuhlmann W, Fromme H-G, Heege E-M, Ostertag W (1968) The mutagenic action of caffeine in higher organisms. *Cancer Res* 28(11):2375–2389 [PubMed: 4881507]
69. Zhang Z-W, Xiao J, Luo W, Wang B-H, Chen J-M (2015) Caffeine suppresses apoptosis of bladder cancer RT4 cells in response to ionizing radiation by inhibiting ataxia telangiectasia mutated-Chk2-p53 axis. *Chin Med J* 128(21):2938–2945 [PubMed: 26521794]
70. Sarkaria JN, Busby EC, Tibbetts RS et al. (1999) Inhibition of ATM and ATR kinase activities by the radiosensitizing agent, caffeine. *Cancer Res* 59(17):4375–4382 [PubMed: 10485486]
71. Turati F, Bosetti C, Polesel J et al. (2015) Coffee, tea, cola, and bladder cancer risk: dose and time relationships. *Urology* 86(6):1179–1184 [PubMed: 26416008]
72. Sugiyama K, Sugawara Y, Tomata Y, Nishino Y, Fukao A, Tsuji I (2017) The association between coffee consumption and bladder cancer incidence in a pooled analysis of the Miyagi Cohort Study and Ohsaki Cohort Study. *Eur J Cancer Prev* 26(2):125–130 [PubMed: 26882077]
73. Slattery ML, West DW, Robison LM (1988) Fluid intake and bladder cancer in Utah. *Int J Cancer* 42(1):17–22 [PubMed: 3391705]
74. Cavin C, Holzhaeuser D, Scharf G et al. (2002) Cafestol and kahweol, two coffee specific diterpenes with anticarcinogenic activity. *Food Chem Toxicol* 40(8):1155–1163 [PubMed: 12067578]
75. Huber WW, Scharf G, Nagel G et al. (2003) Coffee and its chemo-preventive components Kahweol and Cafestol increase the activity of O 6-methylguanine-DNA methyltransferase in rat liver-comparison with phase II xenobiotic metabolism. *Mutat Res* 522(1):57–68 [PubMed: 12517412]
76. Bravi F, Bosetti C, Tavani A et al. (2007) Coffee drinking and hepatocellular carcinoma risk: a meta-analysis. *Hepatology* 46(2):430–435 [PubMed: 17580359]
77. Bravi F, Scotti L, Bosetti C et al. (2009) Coffee drinking and endometrial cancer risk: a metaanalysis of observational studies. *Am J Obstet Gynecol* 200(2):130–135 [PubMed: 19110217]
78. Je Y, Liu W, Giovannucci E (2009) Coffee consumption and risk of colorectal cancer: a systematic review and meta-analysis of prospective cohort studies. *Int J Cancer* 124(7):1662–1668 [PubMed: 19115212]
79. Galeone C, Tavani A, Pelucchi C et al. (2010) Coffee and tea intake and risk of head and neck cancer: pooled analysis in the international head and neck cancer epidemiology consortium. *Cancer Epidemiol Biomark Prev* 19(7):1723–1736
80. Turati F, Galeone C, La Vecchia C, Garavello W, Tavani A (2010) Coffee and cancers of the upper digestive and respiratory tracts: meta-analyses of observational studies. *Ann Oncol* 22(3):536–544 [PubMed: 20943597]
81. Ames BN, Gold LS (1997) The causes and prevention of cancer: gaining perspective. *Environ Health Perspect* 105(Suppl 4):865–873 [PubMed: 9255573]

82. Asaad NA, Zeng Z-C, Guan J, Thacker J, Iliakis G (2000) Homologous recombination as a potential target for caffeine radiosensitization in mammalian cells: reduced caffeine radiosensitization in XRCC2 and XRCC3 mutants. *Oncogene* 19(50):5788–5800 [PubMed: 11126366]
83. Saiki S, Sasazawa Y, Imamichi Y et al. (2011) Caffeine induces apoptosis by enhancement of autophagy via PI3 K/Akt/mTOR/p70S6 K inhibition. *Autophagy* 7(2):176–187 [PubMed: 21081844]
84. Cortez D (2003) Caffeine inhibits checkpoint responses without inhibiting the ataxia-telangiectasia-mutated (ATM) and ATM-and Rad3-related (ATR) protein kinases. *J Biol Chem* 278(39):37139–37145 [PubMed: 12847089]
85. Wu W, Tong Y, Zhao Q et al. (2015) Coffee consumption and bladder cancer: a meta-analysis of observational studies. *Sci Rep* 5:1–9
86. Bracken MB, Triche E, Grosso L et al. (2002) Heterogeneity in assessing self-reports of caffeine exposure: implications for studies of health effects. *Epidemiology* 13(2):165–171 [PubMed: 11880757]
87. Barry D (1996) Differential recall bias and spurious associations in case/control studies. *Stat Med* 15(23):2603–2616 [PubMed: 8961466]
88. Barry D, Livingstone V (2006) The investigation and correction of recall bias for an ordinal response in a case-control study. *Stat Med* 25(6):965–975 [PubMed: 15977299]
89. Chouinard E, Walter S (1995) Recall bias in case-control studies: an empirical analysis and theoretical framework. *J Clin Epidemiol* 48(2):245–254 [PubMed: 7869070]
90. Gefeller O (2009) Invited commentary: recall bias in melanoma—much ado about almost nothing? *Am J Epidemiol* 169(3):267–270 discussion 71–2 [PubMed: 19011114]




Fig. 1. Forest Plot of Meta-Analysis with ORs and 95% CIs for Ever and Never Coffee Consumption with Bladder Cancer Risk Adjusted for Age, Gender and Smoking by Geographical Regions. *OR* odds ratio, *CI* confidence interval. Circle dots denote the odds ratios (ORs); Horizontal lines represent the 95% confidence intervals (CIs); Weights are from random-effect model. *Europe* pooled OR of studies from Europe; *North America* pooled OR of studies from North America; *Overall* pooled OR of all studies




Fig. 2. Dose-response Relationships between Coffee Consumption and the Risk of Bladder Cancer among I) Overall study population; II) Ever Smokers; III) Never Smokers. The solid lines represent the odds ratios (ORs). The dashed lines represent the 95% confidence intervals (CIs) for the trend. The ORs were adjusted for age, gender and smoking (in the overall study population) (model 2). *OR* odds ratio; *CI* confidence interval. P test < 0.05 were considered statistically significant; p increase < 0.05 were considered statistically significant

Table 1

Characteristics of the study population (5,911 cases and 16,172 controls) and coffee consumption category according to age, gender and smoking status

Characteristics	Coffee consumption					Mean (\pm SD) cups/day	p value ^a	p interaction			
	Total	Never	Ever	1 cups/day	1-2 cups/day				2-3 cups/day	3-4 cups/day	> 4 cups/day
n (%)	22,083 (100)	2,372 (10.74)	19,711 (89.26)	2,752 (12.46)	3,616 (16.37)	2,962 (13.41)	2,750 (12.45)	7,631 (34.56)	4.00 (\pm 3.97)	< 0.001	
Case (%)	5,911 (100)	499 (8.44)	5,412 (91.56)	659 (11.15)	1,077 (18.22)	716 (12.11)	801 (13.55)	2,159 (36.53)	4.58 (\pm 4.84)		
Control (%)	16,172 (100)	1,873 (11.58)	14,299 (88.42)	2,093 (12.94)	2,539 (15.70)	2,246 (13.89)	1,949 (12.05)	5,472 (33.84)	3.78 (\pm 3.58)		
Gender											0.16
Male (%)											
Case	4,639 (100)	369 (7.95)	4,270 (92.05)	488 (10.52)	865 (18.65)	560 (12.07)	646 (13.93)	1,711 (36.88)	4.70 (\pm 4.99)	< 0.001	
Control	8,983 (100)	1,002 (11.15)	7,981 (88.85)	1,240 (13.80)	1,468 (16.34)	1,240 (13.80)	1,114 (12.40)	2,919 (32.49)	3.84 (\pm 3.98)		
Female (%)											
Case	1,272 (100)	130 (10.22)	1,142 (89.78)	171 (13.44)	212 (16.67)	156 (12.26)	155 (12.19)	448 (35.22)	4.13 (\pm 4.21)	0.06	
Control	7,189 (100)	871 (12.12)	6,318 (87.88)	853 (11.87)	1,071 (14.90)	1,006 (13.99)	835 (11.61)	2,553 (35.51)	3.71 (\pm 3.01)		
Age (Mean \pm SD)											
Case	5,911 (100)	60.48 (\pm 12.70)	5,412 (91.56)	65.38 (\pm 9.88)	64.36 (\pm 10.36)	64.34 (\pm 8.65)	61.61 (\pm 9.98)	59.88 (\pm 9.21)	4.52 (\pm 4.82)	< 0.001	0.17
Control	16,172 (100)	59.92 (\pm 8.49)	14,299 (88.42)	61.18 (\pm 8.52)	61.94 (\pm 8.80)	61.57 (\pm 8.15)	60.52 (\pm 9.32)	58.92 (\pm 10.13)	3.80 (\pm 3.59)		
Smoking status											
Current smokers (%)											0.001
Case	2,317 (100)	140 (6.04)	2,177 (93.96)	176 (7.60)	417 (18.00)	258 (11.14)	359 (15.49)	967 (41.74)	4.95 (\pm 4.60)	< 0.001	
Control	5,350 (100)	357 (6.67)	4,993 (93.33)	324 (6.06)	782 (14.62)	478 (8.93)	751 (14.04)	2,658 (49.68)	4.92 (\pm 3.75)		
Former smokers (%)											
Case	2,491 (100)	187 (7.51)	2,304 (92.49)	310 (12.44)	412 (16.54)	356 (14.29)	298 (11.96)	928 (37.25)	4.88 (\pm 5.50)	< 0.001	
Control	4,636 (100)	464 (10.01)	4,172 (89.99)	845 (18.23)	637 (13.74)	946 (20.41)	415 (8.95)	1,329 (28.67)	3.49 (\pm 3.92)		
Never smokers (%)											
Case	1,103 (100)	172 (15.59)	931 (84.41)	173 (15.68)	248 (22.48)	102 (9.25)	144 (13.06)	264 (23.93)	3.11 (\pm 3.14)	< 0.001	
Control	6,186 (100)	1,052 (17.01)	5,134 (82.99)	924 (14.94)	1,120 (18.11)	822 (13.29)	783 (12.66)	1,485 (24.01)	3.01 (\pm 2.84)		

n number, SD standard deviation

^a Calculated by χ^2 test for categorical variables and t test for continuous variables

p values < 0.05 were considered statistically significant
p interaction < 0.10 were considered statistically significant

Author Manuscript

Author Manuscript

Author Manuscript

Author Manuscript

Table 2

Adjusted odds ratios and 95% confidence intervals of bladder cancer according to coffee consumption level stratified by smoking

Study sub-groups	Model adjustments	Coffee consumption (ORs and 95% CI)					p trend		
		Never	Ever	1 cups/day	1-2 cups/day	2-3 cups/day		3-4 cups/day	> 4 cups/day
Overall (n = 22,083)	Model 1 ^a	Reference	1.42 (1.26-1.59)	1.05 (0.91-1.22)	1.35 (1.17-1.55)	1.23 (1.06-1.42)	1.51 (1.31-1.76)	1.82 (1.60-2.07)	0.04
	Model 2 ^b	Reference	1.11 (0.98-1.26)	0.94 (0.80-1.10)	1.12 (0.96-1.29)	1.01 (0.86-1.17)	1.16 (0.99-1.35)	1.27 (1.11-1.46)	0.05
	Model 3 ^c	Reference	1.09 (0.94-1.25)	0.92 (0.77-1.10)	1.04 (0.88-1.24)	0.97 (0.81-1.15)	1.15 (0.96-1.37)	1.24 (1.06-1.45)	0.11
Current light smokers (n = 3,548)	Model 1 ^a	Reference	1.01 (0.68-1.47)	0.75 (0.43-1.30)	0.99 (0.64-1.53)	0.71 (0.42-1.19)	1.31 (0.84-2.04)	1.09 (0.72-1.63)	0.34
	Model 2 ^b	Reference	0.97 (0.65-1.43)	0.63 (0.36-1.11)	0.92 (0.59-1.44)	0.71 (0.42-1.21)	1.24 (0.79-1.94)	1.09 (0.72-1.66)	0.23
	Model 3 ^c	Reference	1.16 (0.71-1.90)	0.71 (0.37-1.35)	1.09 (0.63-1.88)	0.80 (0.43-1.49)	1.46 (0.85-2.48)	1.25 (0.78-2.07)	0.14
Current heavy smokers (n = 3,458)	Model 1 ^a	Reference	1.01 (0.71-1.41)	0.91 (0.60-1.39)	1.08 (0.74-1.60)	0.92 (0.63-1.35)	0.97 (0.65-1.44)	1.06 (0.74-1.51)	0.66
	Model 2 ^b	Reference	1.01 (0.71-1.42)	0.83 (0.54-1.28)	0.99 (0.67-1.47)	0.91 (0.62-1.35)	0.97 (0.65-1.46)	1.15 (0.80-1.66)	0.57
	Model 3 ^c	Reference	0.92 (0.59-1.41)	0.75 (0.45-1.24)	0.78 (0.48-1.29)	0.82 (0.51-1.32)	0.89 (0.55-1.45)	1.03 (0.66-1.61)	0.34
Former light smokers (n = 3,314)	Model 1 ^a	Reference	1.24 (0.91-1.70)	1.03 (0.71-1.51)	1.20 (0.82-1.76)	1.36 (0.95-1.97)	1.37 (0.90-2.08)	1.29 (0.91-1.83)	0.07
	Model 2 ^b	Reference	1.28 (0.93-1.75)	1.01 (0.68-1.48)	1.19 (0.81-1.75)	1.44 (0.99-2.09)	1.36 (0.89-2.08)	1.41 (0.99-2.02)	0.06
	Model 3 ^c	Reference	1.30 (0.90-1.88)	1.05 (0.68-1.62)	1.14 (0.74-1.77)	1.50 (0.99-2.27)	1.39 (0.87-2.21)	1.38 (0.93-2.07)	0.09
Former heavy smokers (n = 3,202)	Model 1 ^a	Reference	1.02 (0.76-1.37)	0.83 (0.58-1.17)	0.96 (0.67-1.37)	0.90 (0.64-1.26)	1.03 (0.71-1.49)	1.28 (0.93-1.76)	0.19
	Model 2 ^b	Reference	1.01 (0.75-1.35)	0.80 (0.57-1.13)	0.93 (0.65-1.33)	0.89 (0.63-1.24)	1.03 (0.71-1.49)	1.28 (0.93-1.76)	0.18
	Model 3 ^c	Reference	0.92 (0.66-1.27)	0.72 (0.49-1.05)	0.79 (0.53-1.18)	0.78 (0.54-1.13)	0.96 (0.64-1.44)	1.20 (0.85-1.70)	0.07
Never smokers (n = 7,289)	Model 1 ^a	Reference	1.36 (1.12-1.67)	1.31 (1.01-1.70)	1.53 (1.20-1.96)	1.04 (0.78-1.39)	1.31 (0.99-1.74)	1.52 (1.19-1.94)	0.23

Study sub-groups	Model adjustments	Coffee consumption (ORs and 95% CI)					p trend	
		Never	Ever	1 cups/day	1-2 cups/day	2-3 cups/day		3-4 cups/day
Model 2 ^b	Reference	1.30 (1.06-1.59)	1.18 (0.90-1.55)	1.36 (1.05-1.76)	0.96 (0.71-1.30)	1.22 (0.91-1.64)	1.52 (1.18-1.97)	0.23
	Reference	1.31 (1.03-1.66)	1.28 (0.95-1.73)	1.42 (1.06-1.89)	0.97 (0.70-1.34)	1.24 (0.91-1.70)	1.51 (1.15-1.99)	0.24
Model 3 ^c								

Referent group was never coffee consumers

OR odds ratio, CI confidence interval

^aModel 1: Crude model

^bModel 2: Adjusted for age, gender and smoking (in the overall analyses)

^cModel 3: Additionally adjusted for water, liquid milk, alcohol, carbonated drinks, tea and juice

p trend < 0.05 were considered statistically significant