

UC San Diego

UC San Diego Previously Published Works

Title

Clinical and evoked pain, personality traits, and emotional states: Can familial confounding explain the associations?

Permalink

<https://escholarship.org/uc/item/62c6q56s>

Journal

Journal of Psychosomatic Research, 78(1)

ISSN

0022-3999

Authors

Strachan, Eric
Poeschla, Brian
Dansie, Elizabeth
[et al.](#)

Publication Date

2015

DOI

10.1016/j.jpsychores.2014.09.019

Peer reviewed

Published in final edited form as:

J Psychosom Res. 2015 January ; 78(1): 58–63. doi:10.1016/j.jpsychores.2014.09.019.

Clinical and Evoked Pain, Personality Traits, and Emotional States: Can Familial Confounding Explain the Associations?

Eric Strachan, PhD^{#1,2}, Brian Poeschla, MD^{#1}, Elizabeth Dansie, PhD³, Annemarie Succop, BA^{2,4}, Laura Chopko, BA^{1,2}, and Niloofar Afari, PhD^{5,2}

¹ Department of Psychiatry & Behavioral Sciences, University of Washington, Seattle, WA

² University of Washington Twin Registry, University of Washington, Seattle, WA

³ Department of Anesthesiology, University of Washington, Seattle, WA

⁴ Department of Medicine, University of Washington, Seattle, WA

⁵ VA Center of Excellence for Stress and Mental Health, VA San Diego Healthcare System, and Department of Psychiatry, University of California, San Diego, CA

These authors contributed equally to this work.

Abstract

Objectives—Pain is a complex phenomenon influenced by context and person-specific factors. Affective dimensions of pain involve both enduring personality traits and fleeting emotional states. We examined how personality traits and emotional states are linked with clinical and evoked pain in a twin sample.

Methods—99 female twin pairs were evaluated for clinical and evoked pain using the McGill Pain Questionnaire (MPQ) and dolorimetry, and completed the 120-item International Personality Item Pool (IPIP), the Positive and Negative Affect Scale (PANAS), and ratings of stress and mood. Using a co-twin control design we examined a) the relationship of personality traits and emotional states with clinical and evoked pain, and b) whether genetics and common environment (i.e. familial factors) may account for the associations.

Results—Neuroticism was associated with the sensory component of the MPQ; this relationship was not confounded by familial factors. None of the emotional state measures was associated with the MPQ. PANAS Negative Affect was associated with lower evoked pressure pain threshold and tolerance; these associations were confounded by familial factors. There were no associations between IPIP traits and evoked pain.

© 2014 Elsevier Inc. All rights reserved.

Correspondence to: Eric Strachan, PhD, Psychiatry and Behavioral Sciences, (Box 357236), University of Washington School of Medicine, Seattle, WA 98104, USA. Tel.: +1 206 543 2833; fax: +1 206 543-3830, erstrach@u.washington.edu..

Publisher's Disclaimer: This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final citable form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

CONFLICTS OF INTEREST

None of the authors have any conflicts of interest to declare.

Conclusions—A relationship exists between neuroticism and clinical pain that is not confounded by familial factors. There is no similar relationship between negative emotional states and clinical pain. In contrast, the relationship between negative emotional states and evoked pain is strong while the relationship with enduring personality traits is weak. The relationship between negative emotional states and evoked pain appears to be non-causal and due to familial factors.

Keywords

clinical pain; evoked pain; personality; extraversion; neuroticism; emotions; twins

INTRODUCTION

The sensation of physical pain, familiar to all human beings, is strongly influenced by a variety of corresponding and sometimes competing variables. Demographic characteristics, genetic factors (1), and personality dispositions (2) all influence pain responses and individual differences in responses to chronic pain tend to be large (1, 3). However, situational variables, temporary emotional states, and stress also play a role (4, 5). For example, acute negative emotions are often associated with increased pain sensitivity (6, 7).

Evoked pain sensitivity is defined as the acute response to potentially pain-inducing stimuli such as pressure or very cold water. Clinical pain, which can be defined as the subjective perception of spontaneous pain, is often chronic pain. Chronic clinical pain syndromes, such as fibromyalgia, are defined by the presence of both spontaneous and evoked pain, and fibromyalgia patients have been distinguished clinically from other patients with pain by finding evoked pain, that is pain evoked by 4 kg manual pressure, in at least 11 of 18 defined tender points. Fibromyalgia patients have increased sensitivity to several types of evoked pain, including pressure at non-tender-point sites (25) and heat and cold pain (26). Sensitivity to evoked pain via biological and affective mechanisms may thus be a risk factor for chronic clinical pain syndromes.

Typical models of pain recognize interactions between intensity, unpleasantness, and affective dimensions but the complex biological, psychological, and environmental factors are as yet poorly understood (8). The affective dimensions alone are complex, involving both enduring personality traits and fleeting emotional states at the time the pain is experienced. Personality traits may exert their influence by means of cognitive processes related to the ways in which individuals constitute the meanings and implications of pain (9, 10). For example, positive personality traits such as optimism, hope, and self-efficacy, are protective against enhanced pain perception, and this protective effect may be mediated by a lower tendency to pain catastrophizing in these individuals (11). Trait neuroticism—the tendency to experience strong negative emotions—also has been associated with chronic pain (12-14) and with chronic fatigue syndrome, which includes muscle and multi-joint pain in its diagnostic criteria (15-23). In long term prospective twin studies neuroticism has been linked with the likelihood of developing a range of painful physical conditions including chronic fatigue, and these studies identified familial influences suggesting genetic pathways between neuroticism and these pain experiences (22, 24).

There also is a trait component to the tendency to experience certain affective states and those affective states can, in turn, affect pain sensitivity (6, 7). For example, the personality trait of neuroticism is correlated with both trait and state anxiety (25) as well as a number of clinical pain types in a variety of settings (9, 22, 26-29). A profitable way of exploring pain's multiple manifestations, therefore, is to examine both evoked and clinical forms of pain and their associations with affective traits and states. This, however, can be complicated because personality traits, the tendency to experience certain emotional states, and pain responding all have heritable components (i.e. variability in those traits is partially due to genetics) and all may be influenced by developmental experiences (e.g. parental modelling of pain responding during childhood). Twin studies of clinical pain syndromes (30-32) and of experimental pain sensitivity (33-36) have shown that genetics play a moderate to large role in both clinical and evoked pain. There is also evidence that chronic pain symptoms and symptoms of anxiety and depression are mediated by shared genetic factors (37).

Twins are ideal for studying complex phenotypes, such as pain responding, for which appropriate comparison groups are often ill-defined. Monozygotic (MZ) twins who are reared together share 100% of their genes and 100% of their common developmental environment, and dizygotic (DZ) twins share an average of 50% of their genes and 100% of their common developmental environment. Because of this, co-twin control studies can examine whether shared genes and common developmental environment—referred to jointly as “familial factors” in this paper—account for the relationship between two traits. Thus, co-twin studies can help to distinguish between potentially causal relationships among clinical phenotypes, or alternatively, between shared genetic or environmental linkages among phenotypes (39). In cases such as the present study, where randomization to conditions is impossible, co-twin studies provide the best available alternative to understanding the direct, potentially causal, links among variables of interest.

The aim of this study was to examine how both personality traits and emotional states are linked with clinical and evoked pain, and to accomplish this in a twin sample that allowed us to examine confounding by familial factors. We hypothesized that personality traits would show a stronger association with clinical pain than evoked pain; and that conversely, fleeting emotional states would show a stronger association with evoked pain than clinical pain. Based on prior research on personality and chronic fatigue (22) we hypothesized that any association between personality traits and clinical pain would be at least partially confounded by shared familial factors common to both personality traits and clinical pain. We had no a priori hypothesis regarding familial mediation of emotional state and evoked pain.

METHODS

Participants

All twin pairs in this study ($n = 99$ pairs) were female-female members of the University of Washington Twin Registry (UWTR), a community-based registry of twins identified through the Washington State Department of Licensing. The general characteristics of the UWTR have been reported elsewhere (40, 41). Pairs were recruited between 2006 and 2010 to participate in a twin study of risk factors for chronic pain, investigating the psychological,

behavioral, and physiological risk factors for and correlates of medically unexplained pain in women. Women were targeted because of the greater incidence and burden of medically unexplained pain in women compared to men. Pairs were either discordant for medically unexplained pain or were both pain free (i.e. healthy control pairs). As reported previously (42, 43), in order to participate, both twins were required to be free from severe or chronic medical conditions that could potentially be the source of the unexplained pain (e.g. autoimmune disorders, cancer, etc.), or other medical conditions that could interfere with the study measurements, including cardiopulmonary disorders, uncontrolled endocrine conditions, and uncontrolled allergic or sleep disorders. Potential participants also were excluded if they were current smokers, tested positive for drugs of abuse during the study visit, had body mass index of < 20 or > 30 kg/m², were pregnant or planning to become pregnant during the course of the study, had an amputated limb, or were completely blind or deaf.

Potential participants who were either pain-free or had pain not related to a known disease or organic etiology and did not meet the above exclusionary criteria were eligible to participate. Participants were asked to discontinue all medications that affect sleep, hypothalamic-pituitary-adrenal axis function, or autonomic nervous system functioning for two weeks prior to and during the study evaluation. In addition, all pain medications were discontinued for the duration of the study. The study protocol allowed participants to take short-acting over-the-counter pain relievers (e.g., Tylenol) as needed. However, none of the participants needed to take pain relievers during the 2-day laboratory visit. Alcohol (2 drinks per week) and caffeine (1 cup of coffee per day) were restricted for the duration of the study.

Assignment of Zygosity

We employed a commonly-used self-report measure to assign zygosity. This measure contains multiple questions about childhood similarity in twin pairs. Twin pairs whose responses were concordant for the question, “As children, were you and your twin as alike as two peas in a pod, or of only ordinary family resemblance?” were classified as either MZ (both twins agreed on “alike as two peas in a pod”) or DZ (both twins agreed on “ordinary family resemblance”). In cases where responses were discordant, zygosity was assigned by using an algorithm based on several additional questions assessing childhood similarity. The use of this method to assign zygosity has been demonstrated to be 95%-98% accurate compared to biological indicators (44, 45). Pairs with indeterminate zygosity were excluded from the analyses.

General Procedures

Once enrolled, twin pairs completed a 7-day at-home protocol, kept a daily diary of sleep, stress, mood, and pain levels, and completed a comprehensive battery of questionnaires which included the measures of personality and clinical pain. Immediately after the at-home protocol, the twin pairs presented for a 2-day visit at the University of Washington General Clinical Research Center, where they participated in a number of study procedures including the evoked pain testing that is presented here. Twins came to the Research Center together but were evaluated separately at each step. Written informed consent was obtained from all

participants and the study was approved by the Institutional Review Boards at the University of Washington and University of California, San Diego.

Measures

Personality Traits—Personality traits were measured using the IPIP-NEO short form (46) (available at <http://www.personal.psu.edu/faculty/j/5/j5j/IPIP/>; see also <http://ipip.ori.org/>). The IPIP-NEO short form is a public-domain 120-item version of the IPIP-NEO that was developed to represent the commercially-available NEO PI-R (47), a widely used and well-validated measure of five factor personality traits. (For a review of reliability and validity estimates of IPIP scales see <http://ipip.ori.org/newBroadbandText.htm>. The five personality domains of extraversion, neuroticism (emotional instability), agreeableness, conscientiousness, and openness to experience were assessed. (For a review of these domains and their hierarchically structured sub-facets see Digman, 1990 (48).)

Emotional State Measures—A visual analog scale (VAS) rating of current stress and mood was administered each day during the at-home protocol. For this study, we used the VAS ratings completed on the same day as the Short Form McGill Pain Questionnaire (MPQ) (49) which was our measure of clinical pain. The Positive and Negative Affect Scale (PANAS) (50) was administered on the same day as, and in a sequence prior to, the dolorimetry test which was our measure of evoked pain. The PANAS is a well-validated measure of mood states and demonstrated good reliability in the present sample (positive affect $\alpha = .91$; negative affect $\alpha = .87$). The instructions for the PANAS were focused on the way each person was feeling “right now, that is, at the present moment.” The VAS and PANAS measures were added after the study began and 12 pairs do not have data.

Clinical Pain—Clinical pain was measured using the Short Form McGill Pain Questionnaire (MPQ) (49). The Questionnaire is a widely used instrument designed to assess multiple dimensions of the pain experience (51). Higher scores indicate greater sensory or affective components to pain severity. This instrument has shown good reliability and validity for a wide variety of acute and chronic pain populations (52). The MPQ was completed approximately one week prior to tests of evoked pain and asked about pain over the past four weeks. In order to identify twins with existing chronic/persistent pain, participants also completed the London Fibromyalgia Study Screening Questionnaire (LFSSQ), a validated screening tool that assesses for continuous pain in muscles, bones, or joints, above and below the waist, and in the head, neck, spine, or back in the past 3 months (53).

Evoked Pain—Evoked pain was measured as pressure pain threshold and tolerance using dolorimetry. A dolorimeter is a calibrated pressure algometer with a 3.14 cm² footplate which reads the amount of pressure exerted on an object in kilograms. Using the same sites as assessed for the fibromyalgia manual tender point examination, trained research personnel used the dolorimeter to apply pressure at a rate of 1 kg/second up to 12 kg. Both pain threshold, defined as the point at which the stimulus is experienced as painful, and pain tolerance, the point at which the twin can no longer endure the stimulus, were recorded. For

data analysis, we used the average pressure across all tender points for both threshold and tolerance.

Statistical Analyses

Descriptive statistics were computed as means and standard deviations for continuous measures and percentages for categorical measures. We also calculated basic correlation coefficients to show the associations among the clinical and evoked measures of pain. Prior to conducting regression analyses, bivariate associations of each personality and emotional state measure with each of the pain indices were estimated by using generalized estimating equations (GEE) analyses which correct for interdependency within twin pairs. Only those trait and state measures that were associated with the pain indices at a probability level of 0.15 (corrected for interdependency among observations) were included in the subsequent linear regression models (54). We then used linear regression to estimate the associations of those personality and emotional state indices that were identified through bivariate associations, with clinical and evoked pain. The regression models were fit using GEE with robust variance estimation to control for the correlation within twin pairs (55). In total, four separate regression models were estimated using each of the four pain measures as outcomes. To estimate the most parsimonious models, a sequential backward elimination approach using a more liberal 0.15 probability level was used to eliminate personality and emotional state predictors not contributing significantly to the prediction equations for these models.

Our analytic strategy was to first estimate the overall association between personality traits and emotional state with each pain outcome in all twins as described above, to determine if a phenotypic association existed. Then, we used GEE regression to estimate the within-twin pair effects of personality and emotional state on pain. These within-pair analyses adjust for familial influences by estimating the extent to which differences in personality or emotional states within twin pairs predict differences in pain outcomes. Only those predictors that remained in the overall models were included in the subsequent within-pair analyses. Because twin pairs share a similar family environment and a portion of their genes (100% in monozygotic and on average 50% in dizygotic pairs), within-twin pair estimates are adjusted for familial and some genetic influences. If within-twin pair associations are attenuated and rendered non-significant compared to the overall effect obtained in our initial regression analyses, we can conclude these associations were confounded by familial factors (i.e. familial factors contribute to these associations). Alternately, a within-twin pair association that remains robust compared to the overall effect provides evidence that familial factors do not play a prime role in the association between personality and pain, and instead a causal effect of exposure on the outcome is possible given that other conditions of causation are met (39). Models were adjusted for age and chronic pain status as measured by the LFSSQ. Significance level was set at $p < 0.05$. Data were analyzed using Predictive Analytics Software (PASW v. 18.0, SPSS Inc.).

RESULTS

Twin Participants

Of the 198 twins who participated in the study, 4 participants (2%) were excluded due to an indeterminate zygosity. The participants were all female, and 75% were identified as monozygotic. The average age of the sample was 29 years ($SD = 10$); 89% identified themselves as White and 92% had some college education or greater. In terms of pairwise concordance on the LFSSQ, 52 pairs were concordant for no pain, 11 had one member with no pain and one member with only transient pain (i.e. lasting for less than 1 week during the prior three months), 22 had one member with no pain and one member with persistent pain (i.e. lasting for more than one week), and the remaining 14 pairs had members with widespread persistent pain.

Bivariate Associations

Table 1 displays the bivariate correlations between all of the clinical (MPQ) and evoked pain (dolorimetry) measures. As expected, clinical and evoked pain measures were significantly correlated (r 's ranging from an absolute value of 0.15 - 0.84). Greater sensory and affective clinical pain ratings were significantly associated with lower evoked pressure pain threshold and tolerance.

Table 2 provides the bivariate associations of personality trait and emotional state measures with clinical and evoked pain measures. Neuroticism was the personality trait most strongly correlated with sensory and affective clinical pain indices, with higher neuroticism related to greater sensory and affective clinical pain. As expected, neither of the positive and negative PANAS sub-scales was significantly correlated with either of the clinical pain measures because the PANAS data were not collected on the same day as the MPQ.

However, the VAS stress and mood ratings, which were collected on the same day as the MPQ, were significantly correlated with the clinical pain indices, with higher VAS ratings related to greater sensory and affective clinical pain. There were no significant associations between the personality traits and evoked pressure pain threshold, but the personality traits of agreeableness and conscientiousness were either significant or met our 0.15 probability cut-off to be included in the overall multivariate models for evoked pain tolerance. In addition, the PANAS negative affect sub-scale was strongly negatively correlated with both evoked pain measures.

Adjusted Associations

Clinical Pain—The overall multivariate regression analyses for both clinical pain indices included neuroticism, and VAS stress and mood ratings as predictors. The top portion of Table 3 displays the overall and within-twin pair regression estimates for both indices of clinical pain. After adjusting for age and chronic pain status in the overall model, only neuroticism remained as a significant predictor for sensory pain. After adjusting for familial factors in the within-twin pair analyses, the association between neuroticism and sensory pain ratings remained significant and the magnitude of the association was the same as in the overall association. The regression estimates for the associations of neuroticism and VAS

stress ratings with affective pain ratings were not significant in the overall model and were not significant in the within-twin pair analyses.

Evoked Pain—The overall multivariate regression analyses for evoked pain threshold included the PANAS negative affect sub-scale as a predictor; PANAS negative affect, agreeableness, and conscientiousness were predictors in the model for evoked pain tolerance. The bottom portion of Table 3 displays the overall and within-twin pair regression estimates for both indices of evoked pain. After adjusting for age and chronic pain status in the overall models, only PANAS negative affect remained as a significant predictor for both evoked pain threshold and tolerance. Unlike the clinical pain measures, the relationships between negative affect and both evoked pain measures were attenuated and rendered non-significant after adjusting for familial factors in the within-twin pair analyses.

DISCUSSION

To our knowledge, this is the first study to examine the differential relationship of personality and emotional states with clinical and evoked pain in the same sample. We found that while measures of clinical and evoked pain were related to each other in the expected directions, the pattern of associations with personality trait and emotional state differed between clinical and evoked pain. In multivariate analyses, we found a significant positive association of neuroticism with the sensory component but not the affective component of pain. In contrast, negative emotional state as measured by the PANAS was negatively associated with both evoked pressure pain threshold and tolerance. Further, the link between the trait of neuroticism and sensory pain was not confounded by familial factors, whereas the associations between negative affect and evoked pain threshold and tolerance were confounded by familial factors, indicating that the links between emotional state and evoked pain may be due to the shared familial and genetic mechanisms common to both negative affect and evoked pain.

In their broadest outline, our results suggest that the relationship between the enduring *trait* of neuroticism and clinical pain is stronger than the relationship between transient *states* of negative affect (stress and mood) and clinical pain; and that the relationship of the trait of neuroticism to clinical pain is not due to the influences of familial factors including shared genetics and common environment. That is, our results suggest that the more emotionally unstable an individual is temperamentally, the more likely that individual is to complain of pain and that this relationship between temperament and pain is robust and relatively uninfluenced by transient mood and stress states. These results provide evidence that neuroticism as a temperamental trait is involved in the expression of clinical pain. In contrast, for evoked pain, the relationship with negative emotional states is strong while the relationship with enduring personality traits is weak or nonexistent. Further, we found that the association between emotional states and evoked pain appears to be confounded by familial factors. Therefore, negative emotional states and evoked pain may share a common familial origin. Because this was observed in a model controlling for neuroticism, the trait aspects of neuroticism do not appear to be involved in this potential mediation by familial factors.

It is possible that the potentially shared genetic factors that underlie the negative emotional states and lower evoked pain threshold and tolerance are a product of underlying neural circuitry, perhaps a broader “neural alarm system” involved in responding to noxious stimuli or threats (56). The anterior cingulate cortex has long been implicated in regulating the felt unpleasantness and distressing affective components of physical pain. Individuals who are dispositionally pain-sensitive show more anterior cingulate cortex activity and report greater levels of perceived unpleasantness to painful stimulation (56-58). Patients who have undergone cingulotomy for intractable pain have reported that they are able to still feel the pain but that it no longer distresses them (56). Regions involved in pain response networks include the dorsal subdivision of the anterior cingulate cortex and its links to the insula, somatosensory cortex, thalamus, and periaqueductal gray area (58). There is mounting evidence from the animal lesion and human neuroimaging studies that both physical and social pain overlap in their underlying neural circuitry and computational processes, and that the anterior cingulate cortex plays a key role in this overlap (56). It may be that shared genetic factors underlying evoked pain sensitivity and emotional state intensity influence the feedback loops within this neural alarm and response system.

Neural circuits are obviously influenced by genetics but social learning through the common family environment may also play a role in the development of neural alarms and responses. Several studies support that observational learning processes play an important role in acquisition of pain-related information and response trajectories (59, 60). Thus individuals appear to gain input from their environment regarding the potential danger of specific pain stimuli and appropriate coping responses. Additionally, the impact of social modeling is supported by investigations of parents and children as well as healthy college students (60-62).

Our differing results for clinical versus evoked pain with respect to the influence of personality traits and affective states have several clinical implications. The finding of a positive and potentially direct association between the trait of neuroticism and the sensory component of clinical pain—that is, an association that is not influenced by familial factors—provides further evidence for the hypothesis that clinical pain and its associated syndromes in general, whatever their precise etiology from case to case, are inherently multifactorial in the way they arise and are sustained, and involve important psychological trait factors in addition to physiological factors. This hypothesis can be extended to specific syndromes involving pain complaints such as chronic widespread pain or fibromyalgia. The case for the multifactorial nature of these syndromes, and the necessity of employing both psychosocial and neurobiological explanations to understand them, has been made by Wessely and others (63-67). Interventions that can address enduring personality trait and behavioral difficulties, such as cognitive behavioral therapy and graded exercise therapy may be of significant benefit in managing chronic pain syndromes (68-70). Alternately, the weak relationship we found between enduring traits and evoked pain and the strong relationship between transient negative emotional states and evoked pain suggest that a patient's personality characteristics might contribute less to the variation in how patients will experience pain stimuli in a medical setting, for example when undergoing a painful procedure. Stress reduction and interventions to improve mood may be of considerable benefit in managing pain in those settings.

This study has several limitations. First, the sample was all female, predominantly White, and well educated. Although this matches the demographics of both the Puget Sound region and people who present to physicians with chronic pain complaints, the generalizability of the findings is limited and additional research should be conducted on males and more heterogeneous samples. Second, the sample consisted primarily of individuals who reported minimal long-lasting chronic pain. Therefore, these findings should be replicated in samples of patients with substantial amounts of clinical pain. Third, the evoked pain assessment focused on pressure pain which may not be representative of other pain modalities. Additionally, our measure of evoked pain is commonly employed to diagnose clinical pain in fibromyalgia and does not have established validity as a measure of acute pain. Future research should further validate our findings and use a variety of evoked pain protocols to examine the potential differential association of personality traits and mood states with clinical and evoked pain. Fourth, because of the limited sample of DZ pairs, we were not able to separately examine the confounding effects of shared genetics versus common environmental influences. Finally, despite the power of twin studies to disentangle familial confounding in bivariate associations, a longitudinal study—particularly one in which personality and evoked pain are measured in a large inception cohort prior to onset of chronic pain—could provide additional evidence regarding personality traits, clinical pain, and evoked pain sensitivity.

In conclusion, we found that the trait of neuroticism, but not negative emotional state, was associated with clinical pain. This association was not influenced by familial factors, suggesting that high neuroticism may be directly linked to clinical pain as a risk factor. In contrast, negative emotional state but not the trait of neuroticism, was associated with evoked pain indices. These associations were confounded by shared familial mechanisms suggesting shared genetic and/or common environmental vulnerabilities to both negative emotions and evoked pain. This study highlights the need to investigate cognitive or coping styles as potential risk factors for increased clinical pain and the potential for strategies to change cognitive and emotional coping that reduce the impact on clinical pain. The link between emotional states and evoked pain especially at the level of neural circuits also warrants further investigation.

ACKNOWLEDGEMENTS

This research was supported by National Institutes of Health award R01AR051524 (Dr. Afari). Portions of this research were conducted at the University of Washington Institute of Translational Health Sciences supported by National Institutes of Health awards UL1RR025014, KL2RR025015, and TL1RR025016. Dr. Afari also is supported by the VA Center of Excellence for Stress and Mental Health. We wish to thank the twins for taking part in the University of Washington Twin Registry and for their time and enthusiasm for this project.

REFERENCES

1. Young EE, Lariviere WR, Belfer I. Genetic basis of pain variability: recent advances. *J Med Genet.* 2012; 49:1–9. [PubMed: 22058430]
2. Riley, JL., III; Wade, JB. Psychological and demographic factors that modulate the different stages and dimensions of pain.. In: Price, DD.; Bushnell, MC., editors. *Psychological methods of pain control: basic science and clinical perspectives.* IASP; Seattle, WA: 2004. p. 19-41.
3. Nielsen CS, Price DD, Vassend O, Stubhaug A, Harris JR. Characterizing individual differences in heat-pain sensitivity. *PAIN.* 2005; 119:65–74. [PubMed: 16298065]

4. Rainville, P. Pain and emotions.. In: Price, DD.; Bushnell, MC., editors. Psychological methods of pain control: basic science and clinical perspectives. IASP; Seattle, WA: 2004. p. 117-41.
5. Vassend O, Roysamb E, Nielsen CS. Five-factor personality traits and pain sensitivity: a twin study. *PAIN*. 2013; 154:722–8. [PubMed: 23473786]
6. Sibille KT, Kindler LL, Glover TL, Staud R, Riley JL 3rd, Fillingim RB. Affect balance style, experimental pain sensitivity, and pain-related responses. *Clin J Pain*. Feb 23.2012 PubMed PMID: 22367502. Eng.
7. Zautra A, Smith B, Affleck G, Tennen H. Examinations of chronic pain and affect relationships: applications of a dynamic model of affect. *J Consult Clin Psychol*. Oct; 2001 69(5):786–95. PubMed PMID: 11680555. eng. [PubMed: 11680555]
8. Price DD. Psychological and neural mechanisms of the affective dimension of pain. *Science*. Jun 9; 2000 288(5472):1769–72. PubMed PMID: 10846154. eng. [PubMed: 10846154]
9. Harkins SW, Price DD, Braith J. Effects of extraversion and neuroticism on experimental pain, clinical pain, and illness behavior. *PAIN*. Feb; 1989 36(2):209–18. PubMed PMID: 2919101. eng. [PubMed: 2919101]
10. Wade JB, Dougherty LM, Hart RP, Raffi A, Price DD. A canonical correlation analysis of the influence of neuroticism and extraversion on chronic pain, suffering, and pain behavior. *PAIN*. Oct; 1992 51(1):67–73. PubMed PMID: 1454407. eng. [PubMed: 1454407]
11. Pulvers K, Hood A. The role of positive traits and pain catastrophizing in pain perception. *Curr Pain Headache Rep*. 2013; 17:330. [PubMed: 23512722]
12. Malin K, Littlejohn GO. Neuroticism in young women with fibromyalgia links to key clinical features. *Pain Res Treat*. 2012:730741. PubMed PMID: 22454770. eng. [PubMed: 22454770]
13. Malt EA, Olafsson S, Lund A, Ursin H. Factors explaining variance in perceived pain in women with fibromyalgia. *BMC Musculoskelet Disord*. Apr 25.2002 3:12. PubMed PMID: 12019032. eng. [PubMed: 12019032]
14. Netter P, Hennig J. The fibromyalgia syndrome as a manifestation of neuroticism? *Z Rheumatol*. 1998; 57(Suppl 2):105–8. PubMed PMID: 10025096. eng. [PubMed: 10025096]
15. Blakely AA, Howard RC, Sosich RM, Murdoch JC, Menkes DB, Spears GF. Psychiatric symptoms, personality and ways of coping in chronic fatigue syndrome. *Psychol Med*. May; 1991 21(2):347–62. PubMed PMID: 1876640. eng. [PubMed: 1876640]
16. Buckley L, MacHale SM, Cavanagh JT, Sharpe M, Deary IJ, Lawrie SM. Personality dimensions in chronic fatigue syndrome and depression. *J Psychosom Res*. Apr; 1999 46(4):395–400. PubMed PMID: 10340240. eng. [PubMed: 10340240]
17. Christodoulou C, DeLuca J, Johnson SK, Lange G, Gaudino EA, Natelson BH. Examination of Cloninger's basic dimensions of personality in fatiguing illness: chronic fatigue syndrome and multiple sclerosis. *J Psychosom Res*. Dec; 1999 47(6):597–607. PubMed PMID: 10661606. eng. [PubMed: 10661606]
18. Fiedler N, Lange G, Tiersky L, DeLuca J, Policastro T, Kelly-McNeil K, et al. Stressors, personality traits, and coping of Gulf War veterans with chronic fatigue. *J Psychosom Res*. Jun; 2000 48(6):525–35. PubMed PMID: 11033371. eng. [PubMed: 11033371]
19. Johnson SK, DeLuca J, Natelson BH. Personality dimensions in the chronic fatigue syndrome: a comparison with multiple sclerosis and depression. *J Psychiatr Res*. Jan-Feb;1996 30(1):9–20. PubMed PMID: 8736462. eng. [PubMed: 8736462]
20. Kersh BC, Bradley LA, Alarcon GS, Alberts KR, Sotolongo A, Martin MY, et al. Psychosocial and health status variables independently predict health care seeking in fibromyalgia. *Arthritis Rheum*. Aug; 2001 45(4):362–71. PubMed PMID: 11501724. eng. [PubMed: 11501724]
21. Masuda A, Munemoto T, Yamanaka T, Takei M, Tei C. Psychosocial characteristics and immunological functions in patients with postinfectious chronic fatigue syndrome and noninfectious chronic fatigue syndrome. *J Behav Med*. Oct; 2002 25(5):477–85. PubMed PMID: 12442562. eng. [PubMed: 12442562]
22. Kato K, Sullivan PF, Evengard B, Pedersen NL. Premorbid predictors of chronic fatigue. *Arch Gen Psychiatry*. Nov; 2006 63(11):1267–72. PubMed PMID: 17088507. eng. [PubMed: 17088507]
23. Poeschla B, Strachan E, Dansie E, Buchwald D, Afari N. Chronic fatigue and personality: a twin study of causal pathways and shared liabilities. *Ann Behav Med*. 2012 in press.

24. Charles ST, Gatz M, Kato K, Pedersen NL. Physical health 25 years later: the predictive ability of neuroticism. *Health Psychol.* May; 2008 27(3):369–78. 2008. [PubMed: 18624602]
25. Flett G, Hewitt P, Dyck D. Self-oriented perfectionism, neuroticism and anxiety. *Pers Individ Differ.* 1989; 10:731–5.
26. Costa P, Fleg J, McCrae R, Lakatta E. Neuroticism, coronary artery disease, and chest pain complaints: cross-sectional and longitudinal studies. *Exp Aging Res.* 1982; 8:37–44.
27. Geertzen JH, de Bruijn-Kofman AT, de Bruijn HP, van de Wiel HB, Dijkstra PU. Stressful life events and psychological dysfunction in Complex Regional Pain Syndrome type I. *Clin J Pain.* Jun; 1998 14(2):143–7. PubMed PMID: 9647456. eng. [PubMed: 9647456]
28. Himmelfarb S, Murrell SA. The prevalence and correlates of anxiety symptoms in older adults. *J Psychol.* Mar; 1984 116(2d Half):159–67. PubMed PMID: 6609232. eng. [PubMed: 6609232]
29. Vervoort T, Goubert L, Eccleston C, Bijttebier P, Crombez G. Catastrophic thinking about pain is independently associated with pain severity, disability, and somatic complaints in school children and children with chronic pain. *J Pediatr Psychol.* Aug; 2006 31(7):674–83. PubMed PMID: 16093515. eng. [PubMed: 16093515]
30. Battie MC, Videman T, Levalathi E, Gill K, Kaprio J. Heritability of low back pain and the role of disc degeneration. *PAIN.* 2007; 131:272–80. [PubMed: 17335977]
31. Fillingim RB, Wallace MR, Herbstman DM, Ribeiro-Dasilva M, Staud R. Genetic contribution to pain: a review of findings in humans. *Oral Dis.* 2008; 14:673–83. [PubMed: 19193196]
32. Macgregor AJ, Andrew T, Sambrook PN, Spector TD. Structural, psychological, and genetic influences on low back and neck pain: a study of adult female twins. *Arthritis Rheum.* 2004; 51:160–7. [PubMed: 15077255]
33. Angst MS, Phillips NG, Drover DR, Tingle M, Ray A, Swan GE, et al. Pain sensitivity and opioid analgesia: a pharmacogenomic twin study. *PAIN.* 2012; 153:1397–409. [PubMed: 22444188]
34. Macgregor AJ, Griffiths GO, Baker J, Spector TD. Determinants of pressure pain threshold in adult twins: evidence that shared environmental influences predominate. *PAIN.* 1997; 73:253–7. [PubMed: 9415513]
35. Nielsen CS, Stubhaug A, Price DD, Vassend O, Czajkowski N, Harris JR. Individual differences in pain sensitivity: genetic and environmental contributions. *PAIN.* 2008; 136:21–9. [PubMed: 17692462]
36. Norbury TA, MacGregor AJ, Urwin J, Spector TD, McMahon SB. Heritability of responses to painful stimuli in women: a classical twin study. *Brain.* 2007; 130:3041–9. [PubMed: 17932101]
37. Reichborn-Kjennerud T, Stoltenberg C, Tambs K, Roysamb E, Kringlen E, Torgersen S, et al. Back-neck pain and symptoms of anxiety and depression: a population-based twin study. *Psychol Med.* 2002; 32:1009–20. [PubMed: 12214782]
38. Gustin SM, Peck CC, Macey PM, Murray GM, Henderson LA. Unraveling the effects of plasticity and pain on personality. *J Pain.* 2013; 14:1642–52. [PubMed: 24290444]
39. McGue M, Osler M, Christensen K. Causal inference and observational research: The utility of twins. *Perspect Psychol Sci.* Oct 11; 2010 5(5):546–56. PubMed PMID: 21593989. Pubmed Central PMCID: 3094752. Epub 2011/05/20. Eng. [PubMed: 21593989]
40. Afari N, Noonan C, Goldberg J, Edwards K, Gadepalli K, Osterman B, et al. University of Washington Twin Registry: construction and characteristics of a community-based twin registry. *Twin Res Hum Genet.* Dec; 2006 9(6):1023–9. PubMed PMID: 17254446. Pubmed Central PMCID: 2953369. Epub 2007/01/27. eng. [PubMed: 17254446]
41. Strachan E, Hunt C, Afari N, Duncan G, Noonan C, Schur E, et al. The University of Washington Twin Registry: Poised for the next generation of twin research. *Twin Res Hum Genet.* Dec.2012 7:1–8. PubMed PMID: 23218177.
42. Afari N, Mostoufi S, Noonan C, Poeschla B, Succop A, Chopko L, et al. C-reactive protein and pain sensitivity: findings from female twins. *Ann Behav Med.* Oct; 2011 42(2):277–83. PubMed PMID: 21785898. eng. [PubMed: 21785898]
43. Godfrey K, Strachan E, Dansie E, Crofford L, Buchwald D, Goldberg J, et al. Salivary cortisol and cold pain sensitivity in female twins. *Ann Behav Med.* 2014; 47(2):180–8. [PubMed: 23955075]

44. Eisen S, Neuman R, Goldberg J, Rice J, True W. Determining zygosity in the Vietnam Era Twin Registry: an approach using questionnaires. *Clin Genet.* Jun; 1989 35(6):423–32. PubMed PMID: 2736790. eng. [PubMed: 2736790]
45. Torgersen S. The determination of twin zygosity by means of a mailed questionnaire. *Acta Genet Med Gemellol (Roma).* 1979; 28(3):225–36. PubMed PMID: 297423. eng. [PubMed: 297423]
46. Goldberg L, Johnson J, Eber H, Hogan R, Ashton M, Cloninger C, et al. The International Personality Item Pool and the future of public-domain personality measures. *J Res Pers.* 2006; 40:84–96.
47. Costa, P.; McCrae, R. Revised NEO Personality Inventory (NEO-PI-R) and NEO Five-Factor Inventory (NEO-FFI) professional manual. Psychological Assessment Resources, Inc.; Odessa, FL: 1992.
48. Digman J. Personality structure: emergence of the five-factor model. *Ann Rev Psychol.* 1990; 41:417–40.
49. Melzack R. The short-form McGill Pain Questionnaire. *PAIN.* Aug; 1987 30(2):191–7. PubMed PMID: 3670870. eng. [PubMed: 3670870]
50. Watson D, Clark LA, Tellegen A. Development and validation of brief measures of positive and negative affect: the PANAS scales. *J Pers Soc Psychol.* Jun; 1988 54(6):1063–70. PubMed PMID: 3397865. eng. [PubMed: 3397865]
51. Melzack R. The McGill Pain Questionnaire: major properties and scoring methods. *Pain.* Sep; 1975 1(3):277–99. PubMed PMID: 1235985. eng. [PubMed: 1235985]
52. Melzack, R.; Katz, J. The McGill Pain Questionnaire: appraisal and current status.. In: Turk, D.; Melzack, R., editors. *Handbook of Pain Assessment.* Guildford; New York: 1992. p. 152-68.
53. White KP, Harth M, Speechley M, Ostbye T. Testing an instrument to screen for fibromyalgia syndrome in general population studies: the London Fibromyalgia Epidemiology Study Screening Questionnaire. *J Rheumatol.* Apr; 1999 26(4):880–4. PubMed PMID: 10229410. eng. [PubMed: 10229410]
54. Tabachnick, G.; Fidell, L. *Using Multivariate Statistics.* 5th ed.. Allyn and Bacon; Boston: 2007.
55. Diggle, P.; Heagerty, P.; Liang, K.; Zeger, S. *Analysis of longitudinal data.* 2nd ed.. Oxford University Press; Oxford, England: 2002.
56. Eisenberger NI, Lieberman MD. Why rejection hurts: a common neural alarm system for physical and social pain. *Trends Cogn Sci.* Jul; 2004 8(7):294–300. PubMed PMID: 15242688. eng. [PubMed: 15242688]
57. Eisenberger, NI.; Lieberman, MD. Why it hurts to be left out: the neurocognitive overlap between physical and social pain. In: Williams, KD.; Forgas, JP.; von Hippel, W., editors. *The social outcast: ostracism, social exclusion, rejection, and bullying.* Cambridge University Press; New York: 2005. p. 109-27.
58. Lieberman MD, Eisenberger NI. Neuroscience. Pains and pleasures of social life. *Science.* Feb 13; 2009 323(5916):890–1. PubMed PMID: 19213907. eng. [PubMed: 19213907]
59. Goubert L, Vlaeyen JW, Crombez G, Craig KD. Learning about pain from others: an observational learning account. *J Pain.* Feb; 2011 12(2):167–74. PubMed PMID: 21111682. [PubMed: 21111682]
60. Trost Z, France CR, Vervoort T, Lange JM, Goubert L. Learning about pain through observation: the role of pain-related fear. *J Behav Med.* Apr; 2014 37(2):257–65. PubMed PMID: WOS: 000332689300008. English. [PubMed: 23264004]
61. Hermann, C. Modeling, Social Learning in Pain.. In: Willis, RSW., editor. *The Encyclopedia of Pain.* Springer Publishing; Heidelberg: 2007. p. 491-3.
62. Craig, KD. Social Modeling Influences on Pain.. In: Sternbach, R., editor. *The Psychology of Pain.* 2nd ed.. Raven Press; New York, NY: 1986. p. 67-96.
63. Wessely S. Old wine in new bottles: neurasthenia and 'ME'. *Psychol Med.* Feb; 1990 20(1):35–53. PubMed PMID: 2181519. eng. [PubMed: 2181519]
64. Harvey SB, Wessely S. Chronic fatigue syndrome: identifying zebras amongst the horses. *BMC Med.* 2009; 7:58. PubMed PMID: 19818158. eng. [PubMed: 19818158]
65. Kouyanou K, Pither CE, Rabe-Hesketh S, Wessely S. A comparative study of iatrogenesis, medication abuse, and psychiatric morbidity in chronic pain patients with and without medically

- explained symptoms. *PAIN*. Jun; 1998 76(3):417–26. PubMed PMID: 9718260. eng. [PubMed: 9718260]
66. Gatchel RJ, Peng YB, Peters ML, Fuchs PN, Turk DC. The biopsychosocial approach to chronic pain: scientific advances and future directions. *Psychol Bull*. Jul; 2007 133(4):581–624. PubMed PMID: 17592957. eng. [PubMed: 17592957]
67. Turk DC, Okifuji A. Psychological factors in chronic pain: evolution and revolution. *J Consult Clin Psychol*. Jun; 2002 70(3):678–90. PubMed PMID: 12090376. eng. [PubMed: 12090376]
68. Bagby R, Joffe R, Parker J, Kalemka B, Harkness K. Major depression and the five-factor model of personality. *J Personal Disord*. 1995; 9:224–34.
69. Bagby RM, Quilty LC, Segal ZV, McBride CC, Kennedy SH, Costa PT. Personality and differential treatment response in major depression: a randomized controlled trial comparing cognitive-behavioural therapy and pharmacotherapy. *Can J Psychiat*. Jun; 2008 53(6):361–70. PubMed PMID: 18616856. eng.
70. White PD, Goldsmith KA, Johnson AL, Potts L, Walwyn R, DeCesare JC, et al. Comparison of adaptive pacing therapy, cognitive behaviour therapy, graded exercise therapy, and specialist medical care for chronic fatigue syndrome (PACE): a randomised trial. *Lancet*. Mar 5; 2011 377(9768):823–36. PubMed PMID: 21334061. eng. [PubMed: 21334061]

HIGHLIGHTS

- We examined how personality traits and emotional states link with clinical and evoked pain in twins
- Personality traits were associated with clinical pain controlling for familial factors
- Emotional states were not associated with clinical pain
- Emotional states were associated with evoked pain; this was confounded by familial factors
- Personality traits were not associated with evoked pain

Table 1Correlations for relationships between clinical and evoked pain indices ($n = 192 - 194$).

	McGill Sensory	McGill Affective	Dolorimetry Threshold
McGill Affective	0.56 ***		
Dolorimetry Threshold	-0.20 **	-0.19 **	
Dolorimetry Tolerance	-0.15 *	-0.15 *	0.84 ***

Note. n = number of twins.

* $p < 0.05$

** $p < 0.01$

*** $p < 0.001$.

Table 2

Bivariate associations between personality trait or emotional state measures and pain indices.

Pain Indices	IPIP-NEO Personality Trait					Emotional State			
	N	E	O	A	C	PANAS Positive	PANAS Negative	VAS Stress	VAS Mood
<i>Clinical Pain</i>									
McGill Sensory	0.23* (0.005)	0.04 (0.59)	0.002 (0.98)	-0.06 (0.42)	-0.01 (0.92)	-0.02 (0.86)	0.02 (0.70)	0.19* (0.04)	0.20* (0.03)
McGill Affective	0.28* (0.01)	-0.12 (0.06)	-0.07 (0.18)	-0.16* (0.05)	-0.13* (0.09)	0.02 (0.73)	-0.03 (0.61)	0.27* (0.02)	0.23* (0.04)
<i>Evoked Pain</i>									
Dolorimetry Threshold	-0.04 (0.77)	0.05 (0.56)	-0.06 (0.40)	0.06 (0.34)	0.10 (0.16)	-0.03 (0.68)	-0.14* (0.07)	NA	NA
Dolorimetry Tolerance	-0.09 (0.39)	0.08 (0.33)	0.03 (0.71)	0.17* (0.01)	0.12* (0.14)	0.04 (0.59)	-0.64* (0.002)	NA	NA

Note. N = neuroticism, E = extraversion, O = openness, A = agreeableness, C = conscientiousness, Positive = PANAS positive affect, Negative = PANAS negative affect, Stress = VAS stress rating at time of McGill completion, Mood = VAS mood rating at time of McGill completion, NA = not applicable. All *p* values in parentheses are corrected for interdependency among observations using GEE analyses.

* All bivariate relationships with a *p* value < 0.15 were included in subsequent adjusted overall individual-level GEE analyses.

Table 3

Adjusted associations of personality and emotional state with clinical and evoked pain.

Outcome	Predictor	Overall			Within-Pair*	
		<i>n</i>	B	<i>p</i>	B	<i>p</i>
<i>Clinical Pain</i>						
McGill Sensory	N	192	0.048	0.002	0.047	0.04
	Stress			<i>NS</i>	--	--
	Mood			<i>NS</i>	--	--
McGill Affective	N	174	0.01	0.09	0.02	0.07
	Stress		0.01	0.06	0.01	0.22
	Mood			<i>NS</i>	--	--
<i>Evoked Pain</i>						
Threshold	Negative	174	-0.03	0.02	-0.003	0.89
Tolerance	Negative	174	-0.05	0.003	-0.03	0.16
	A			<i>NS</i>	--	--
	C			<i>NS</i>	--	--

Note.

N = neuroticism, Stress = VAS stress rating at time of McGill completion, Mood = VAS mood rating at time of McGill completion, Negative = PANAS negative affect, A = agreeableness, C = conscientiousness. All analyses were adjusted for age, zygosity, and pain status. Sample size = 174 when the VAS or PANAS measures were included in the models.

* Adjusted for familial contributions.