

# UC San Francisco

## UC San Francisco Previously Published Works

### Title

Mechanisms, Pathophysiology, and Management of Obesity

### Permalink

<https://escholarship.org/uc/item/63p9z787>

### Journal

NEW ENGLAND JOURNAL OF MEDICINE, 376(15)

### ISSN

0028-4793

### Authors

Stigler, FL

Lustig, RH

Ma, JI

### Publication Date

2017-04-13

Peer reviewed

Hugh Calkins, M.D.

Johns Hopkins Medical Institutions  
Baltimore, MD

Since publication of their article, the authors report no further potential conflict of interest.

1. Rigato I, Bauce B, Rampazzo A, et al. Compound and digenic heterozygosity predicts lifetime arrhythmic outcome and sudden cardiac death in desmosomal gene-related arrhythmogenic right ventricular cardiomyopathy. *Circ Cardiovasc Genet* 2013;6:533-42.
2. Corrado D, Wichter T, Link MS, et al. Treatment of arrhythmo-

genic right ventricular cardiomyopathy/dysplasia: an International Task Force consensus statement. *Circulation* 2015;132:441-53.

3. Hodgkinson KA, Howes AJ, Boland P, et al. Long-term clinical outcome of arrhythmogenic right ventricular cardiomyopathy in individuals with a p.S358L mutation in TMEM43 following implantable cardioverter defibrillator therapy. *Circ Arrhythm Electrophysiol* 2016;9(9):e003589.

4. Haywood AF, Merner ND, Hodgkinson KA, et al. Recurrent missense mutations in TMEM43 (ARVD5) due to founder effects cause arrhythmogenic cardiomyopathies in the UK and Canada. *Eur Heart J* 2013;34:1002-11.

DOI: 10.1056/NEJMc1701400

## Mechanisms, Pathophysiology, and Management of Obesity

**TO THE EDITOR:** The review article by Heymsfield and Wadden (Jan. 19 issue)<sup>1</sup> is valuable with respect to the clinical management of obesity, but information about the contribution of mitochondrial genes to obesity is not included. Mitochondrial dysfunction is associated with an accumulation of fat that can occur during aging and in patients with obesity, the metabolic syndrome, or diabetes mellitus.<sup>2</sup>

Zheng et al.<sup>3</sup> found that obese participants with a high metabolic syndrome score have increased DNA methylation in the mitochondrial genes *MT-CO1* and *MT-ND6* and in the mitochondrion-related nuclear gene *PPARGC1A*. Flaquer et al.<sup>4</sup> conducted a study using samples obtained from 6528 participants in the KORA (Cooperative Health Research in the Region of Augsburg) studies and found that two mitochondrial single-nucleotide polymorphisms (SNPs) located in the cytochrome *c* oxidase subunit genes (*MT-CO1* and *MT-CO3*) and three mitochondrial SNPs located in the NADH dehydrogenase subunit genes (*MT-ND1*, *MT-ND2*, and *MT-ND4L*) were significantly associated with a higher body-mass index (BMI). Latorre-Pellicer et al.<sup>5</sup> systematically characterized conplastic mice (mice in which the nuclear genome of one mouse is backcrossed into the cytoplasm of another, so that the nuclear genes and mitochondrial genes are from different parents) throughout their lifespan. They found that the mitochondrial DNA haplotype profoundly influences mitochondrial proteostasis and generation of reactive oxygen species, insulin signaling, telomere shortening, the development of obesity, and mitochondrial dysfunction. These findings highlight the importance of the contribution of mitochondrial genetic variants to the risk of a high BMI.

Nai-Wei Sheu, M.D.

Yu-Chih Lin, M.D.

Kaohsiung Medical University Hospital  
Kaohsiung, Taiwan

Chung-Jen Chen, M.D.

Kaohsiung Medical University College of Medicine  
Kaohsiung, Taiwan  
cjchen@kmu.edu.tw

No potential conflict of interest relevant to this letter was reported.

1. Heymsfield SB, Wadden TA. Mechanisms, pathophysiology, and management of obesity. *N Engl J Med* 2017;376:254-66.
2. López-Lluch G. Mitochondrial activity and dynamics changes regarding metabolism in ageing and obesity. *Mech Ageing Dev* 2016 December 16 (Epub ahead of print).
3. Zheng LD, Brooke J, Smith C, Almeida FA, Cheng Z. Mitochondrial epigenetic changes and progression from metabolically healthy obesity to metabolically unhealthy obesity: a cross-sectional study. *Lancet Diabetes Endocrinol* 2016;4:Suppl 1:S16.
4. Flaquer A, Baumbach C, Kriebel J, et al. Mitochondrial genetic variants identified to be associated with BMI in adults. *PLoS One* 2014;9(8):e105116.
5. Latorre-Pellicer A, Moreno-Loshuertos R, Lechuga-Vieco AV, et al. Mitochondrial and nuclear DNA matching shapes metabolism and healthy ageing. *Nature* 2016;535:561-5.

DOI: 10.1056/NEJMc1701944

**TO THE EDITOR:** Heymsfield and Wadden identify environmental and genetic factors as well as energy-balance dysregulation as the leading mechanisms of obesity, and they describe therapeutic lifestyle changes, adjunctive pharmacotherapy, and bariatric surgery as the main treatment strategies for this condition. We would like to point out that emotional factors can influence overeating that results in overweight and obesity.<sup>1</sup>

Emerging data suggest that addictive overeating is a common experience of obese persons.<sup>1</sup> In one randomized trial, mothers had significant and clinically important reductions in weight when their school-age children were taught about nutri-

tion and influenced the family's dietary and physical activity habits.<sup>2</sup> We think that further research concerning these issues would be important and timely. Future findings regarding emotional factors, in general, and binge eating, in particular, may lead to important clinical and public health measures to combat the increasing pandemic of obesity that is a major contributor to the emergence of cardiovascular disease as the leading cause of death worldwide.<sup>3,4</sup>

Felicita Andreotti, M.D., Ph.D.

Filippo Crea, M.D.

Catholic University Hospital  
Rome, Italy  
felicita.andreotti@iol.it

Charles H. Hennekens, M.D., Dr.P.H.

Charles E. Schmidt College of Medicine  
Boca Raton, FL

Dr. Andreotti reports receiving consulting fees from Amgen, Bayer, Boehringer Ingelheim, and Daiichi Sankyo and speaking fees from Bayer, Bristol-Myers Squibb, Daiichi Sankyo, the Menarini International Foundation, and Pfizer; and Dr. Hennekens, receiving fees from Amgen, AstraZeneca, Bayer, the British Heart Foundation, Cadila Pharmaceuticals, DalCor Pharmaceuticals, Lilly, and Regeneron Pharmaceuticals for serving as a chairperson or member of data and safety monitoring boards and advisory fees from Aralez Pharmaceuticals, Pfizer and its legal counsel, and the Food and Drug Administration (FDA), and being a co-inventor on patents for inflammatory markers and cardiovascular disease that are held by Brigham and Women's Hospital. No other potential conflict of interest relevant to this letter was reported.

- Schroder R, Sellman D, Elmslie J. Addictive overeating: lessons learned from medical students' perceptions of Overeaters Anonymous. *N Z Med J* 2010;123:15-21.
- Gunawardena N, Kurotani K, Indrawansa S, Nonaka D, Mizoue T, Samarasinghe D. School-based intervention to enable school children to act as change agents on weight, physical activity and diet of their mothers: a cluster randomized controlled trial. *Int J Behav Nutr Phys Act* 2016;13:45-55.
- Hennekens CH, Andreotti F. Leading avoidable cause of premature deaths worldwide: case for obesity. *Am J Med* 2013;126:97-8.
- Andreotti F, Rio T, Lavorgna A. Body fat and cardiovascular risk: understanding the obesity paradox. *Eur Heart J* 2009;30:752-4.

DOI: 10.1056/NEJMc1701944

**TO THE EDITOR:** We challenge the recommendation by Heymsfield and Wadden that people who are overweight or obese should lose weight using a reduced-calorie diet. This recommendation assumes that obesity is the cause of metabolic dysfunction and that weight loss is the cure for metabolic dysfunction. We argue rather that obesity is a marker for metabolic dysfunction, and "real food"<sup>1</sup> is its cure.

If recommendations to lose weight with the use of a calorie-restricted diet were a drug, they

probably would not receive FDA approval. The studies presented by the authors indicate that even high-intensity weight-loss interventions are ineffective in the long term and adverse effects such as stigmatization of and bias against obese persons are common.<sup>2</sup> Not surprisingly, many physicians and patients are frustrated by this approach.

The recommendation to improve metabolic health unrelated to weight loss can improve clinical end points (e.g., a Mediterranean diet reduces mortality from cardiovascular disease<sup>3</sup> and is more sustainable than other approaches). Finally, we physicians need to use our societal standing to advocate for change from obesogenic environments loaded with high-sugar, low-fiber foods<sup>1</sup> toward environments that make the healthy choice the easy choice.<sup>4</sup>

Florian L. Stigler, M.D., M.P.H.

Medical University of Graz  
Graz, Austria  
florian.stigler@lshtm.ac.uk

Robert H. Lustig, M.D.

Janet I. Ma, M.D.  
University of California, San Diego  
San Diego, CA

No potential conflict of interest relevant to this letter was reported.

- Lustig RH. Processed food — an experiment that failed. *JAMA Pediatr* 2017;171:212-4.
- Phelan SM, Burgess DJ, Yeazel MW, Hellerstedt WL, Griffin JM, van Ryn M. Impact of weight bias and stigma on quality of care and outcomes for patients with obesity. *Obes Rev* 2015;16:319-26.
- Estruch R, Ros E, Salas-Salvadó J, et al. Primary prevention of cardiovascular disease with a Mediterranean diet. *N Engl J Med* 2013;368:1279-90.
- Lustig RH, Schmidt LA, Brindis CD. Public health: the toxic truth about sugar. *Nature* 2012;482:27-9.

DOI: 10.1056/NEJMc1701944

**TO THE EDITOR:** Although they discuss the mechanisms and pathophysiological features of obesity, Heymsfield and Wadden do not emphasize the role of insulin, a metabolically important hormone. High levels of insulin inhibit lipolysis and suppress hepatic glucose production.<sup>1</sup> As a consequence, levels of plasma fatty acids and glucose are reduced. The deficiency of fuel in the blood induces intense hunger and physiological weakness, which in turn probably induce overfeeding and physical inactivity, leading to positive energy balance and eventual obesity.

Basal hyperinsulinemia is induced by a carbo-

hydrate-rich diet.<sup>2</sup> This is why a carbohydrate-rich diet is generally much more obesogenic than a fat-rich diet, which is more likely to induce basal hypoinsulinemia.<sup>2</sup> Diet is as important as environmental or genetic factors in inducing obesity.

Song J. Lee, Ph.D.

Chungnam National University  
Daejeon, South Korea

Sang W. Shin, M.D., Ph.D.

Korea University  
Seoul, South Korea  
shinsw@kumc.or.kr

No potential conflict of interest relevant to this letter was reported.

1. Morimoto C, Tsujita T, Okuda H. Antilipolytic actions of insulin on basal and hormone-induced lipolysis in rat adipocytes. *J Lipid Res* 1998;39:957-62.
2. Farquhar JW, Frank A, Gross RC, Reaven GM, Brown EP. Glucose, insulin, and triglyceride responses to high and low carbohydrate diets in man. *J Clin Invest* 1966;45:1648-56.

DOI: 10.1056/NEJMc1701944

**THE AUTHORS REPLY:** We thank Sheu and colleagues for expanding our discussion to the role played by mitochondria in the pathogenesis and management of obesity. Mitochondrial mechanisms are directly involved in cellular energy production and are increasingly recognized as playing a role in the control of energy balance. A study of mitochondrial DNA has also identified genetic variants linked with excess adiposity and metabolic disturbances.<sup>1</sup>

We agree with Andreotti et al. that emotional factors play an important role in the genesis of overweight and obesity and that these effects should be the focus of additional research. In support of this viewpoint, longitudinal studies show that depression, anxiety, and life stress increase the odds of weight gain with the development of overweight and obesity through multiple as yet incompletely established mechanisms.<sup>2</sup> These effects are bidirectional, with obesity increasing the risk of depression and anxiety. Binge eating disorder, which is characterized by consumption of an objectively large amount of food and loss of control over eating, occurs in only a small percentage of obese persons who may also be at an increased risk for depression and anxiety. Studies are currently in progress to examine whether the concept of food addiction contributes to our understanding of and treat-

ment approaches for disturbed eating behaviors and binge eating.<sup>3</sup>

We support Stigler and colleagues' call on physicians to serve as advocates for changing the nation's obesogenic food environment in favor of one that makes vegetables and fresh fruits available to all citizens. We also think that data are lacking on the possible health benefits, independent of weight loss, of diets with various macronutrient compositions aligned with the nutritional needs of patients with weight-related chronic diseases. Unlike Stigler et al., we think the data show that high-intensity lifestyle interventions, which include calorie restriction, reduce the short-term and long-term risks of type 2 diabetes and other health complications, even when lost weight is partially regained.<sup>4</sup>

Lee and Shin emphasize the important roles played by insulin in the pathogenesis of obesity and carbohydrate-rich diets in the management of this condition. South Korea, the home country of Lee and Shin, has a long history of a high carbohydrate intake from vegetables and fresh fruits, even though obesity rates are low.<sup>5</sup> We agree that excess intake of "added" sugars, particularly those in sweetened beverages, contributes to the pathogenesis of obesity and that reducing the intake of these "free" carbohydrates should be a component of efforts to control weight.

Steven B. Heymsfield, M.D.

Pennington Biomedical Research Center  
Baton Rouge, LA  
steven.heymsfield@pbrc.edu

Thomas A. Wadden, Ph.D.

Perelman School of Medicine  
Philadelphia, PA

Since publication of their article, the authors report no further potential conflict of interest.

1. Flaquer A, Baumbach C, Kriebel J, et al. Mitochondrial genetic variants identified to be associated with BMI in adults. *PLoS One* 2014;9(8):e105116.
2. Berkowitz RI, Fabricatore AN. Obesity, psychiatric status, and psychiatric medications. *Psychiatr Clin North Am* 2011;34:747-64.
3. Ziauddeen H, Fletcher PC. Is food addiction a valid and useful concept? *Obes Rev* 2013;14:19-28.
4. Knowler WC, Fowler SE, Hamman RF, et al. 10-Year follow-up of diabetes incidence and weight loss in the Diabetes Prevention Program Outcomes Study. *Lancet* 2009;374:1677-86.
5. Lee MJ, Popkin BM, Kim S. The unique aspects of the nutrition transition in South Korea: the retention of healthful elements in their traditional diet. *Public Health Nutr* 2002;5:197-203.

DOI: 10.1056/NEJMc1701944

Correspondence Copyright © 2017 Massachusetts Medical Society.