

UC Berkeley

UC Berkeley Previously Published Works

Title

The Association Between BMI and QCT-Derived Proximal Hip Structure and Strength in Older Men: A Cross-Sectional Study

Permalink

<https://escholarship.org/uc/item/6417p576>

Journal

Journal of Bone and Mineral Research, 30(7)

ISSN

0884-0431

Authors

Shen, Jian
Nielson, Carrie M
Marshall, Lynn M
[et al.](#)

Publication Date

2015-07-01

DOI

10.1002/jbmr.2450

Peer reviewed



Published in final edited form as:

J Bone Miner Res. 2015 July ; 30(7): 1301–1308. doi:10.1002/jbmr.2450.

The association between BMI and QCT-derived proximal hip structure and strength in older men: a cross-sectional study

Jian Shen¹, Carrie M. Nielson¹, Lynn M. Marshall², David C. Lee³, Tony M. Keaveny^{3,4}, and Eric S. Orwoll¹ for the Osteoporotic Fractures in Men (MrOS) Research Group

¹Department of Medicine, Bone and Mineral Unit, Oregon Health and Science University, Portland, OR

²Departments of Orthopaedics and Rehabilitation, Oregon Health and Science University, Portland, OR

³O.N. Diagnostics, Berkeley, CA

⁴Department of Mechanical Engineering, University of California, Berkeley, Berkeley, CA

Abstract

Although higher body mass index (BMI) is associated with higher bone mineral density, recent evidence indicates that increased BMI may not be consistently associated with reduced hip fracture risk. Moreover, substantial proportions of hip fractures occur among overweight and obese men and women. The role of increased BMI and obesity on bone density, structure, and strength at the hip is not well understood.

We conducted cross-sectional analyses between BMI and various density and structure measures derived from quantitative computed tomography (QCT)-scans of the proximal femur, in 3067 men (mean age: 73 y) from the Osteoporotic Fractures in Men Study (MrOS). Finite element (FE) analysis of hip QCT scans was performed for a subcohort of 672 men to provide a measure of femoral strength for a simulated sideways fall. The impact force was estimated using patient-specific weight and height information. Multivariable general linear models were used to examine the associations between BMI and hip QCT measures.

The relationship of BMI with hip QCT measures was significantly different between men categorized as non-obese and obese (P for interaction = 0.014). For non-obese men (BMI < 30), increasing BMI was associated with higher integral, cortical and trabecular vBMD, integral volume, cross-sectional area, and percent cortical volume (all $p < 0.001$). For obese men (BMI ≥ 30), increasing BMI was not associated with any of those parameters. In addition, compared to non-obese men, obese men had a higher hip strength, but also a higher ratio of impact force to strength ($P < 0.0001$), in theory increasing their risk of hip fracture despite their increased strength. These results provide a better understanding of hip fracture risk in obese men.

To whom correspondence may be addressed: Eric S Orwoll, Bone and Mineral Unit, Department of Medicine, Oregon Health and Science University, Portland, OR, Phone: 503-494-0225, orwoll@ohsu.edu, Fax: 503-494-4816.

Supplemental data is included in the submission

The authors' responsibilities were as follows— study concept and design: JS, CMN, ESO; Data analysis: JS; Data interpretation: JS, CMN, LMM, TMK and ESO; drafting of the manuscript: JS; critical revision of the manuscript: JS, CMN, LMM, DCL, TMK and ESO.

Keywords

men; obesity; hip fracture risk; hip structure; hip strength

Introduction

While obesity is a well-established risk factor for many chronic diseases, it is generally believed that individuals with higher body mass index (BMI) have a low risk of osteoporosis, in part due to their increased bone mineral density (BMD)^{1,2}. However, fracture in obese individuals remains a public health concern³. The relationship between obesity and fracture is complex and not well understood. While there are some evidence showing that increases in BMI is associated with reduced risk of fracture^{4,5}, more recent studies suggest obesity is not protective and may increase risk for fracture. A study of postmenopausal women with low-trauma fracture reported a high prevalence of obesity despite the fact that BMD was normal in most of the fracture cases⁶. Likewise, among middle aged men, higher waist circumference has been linked to increased risk of hip and wrist fracture⁷. In a large cohort of elderly men, we observed that a substantial proportion of hip and non-spine fractures occur among overweight or obese men⁸. The recent meta-analysis of the association of fracture risk and BMI that was comprised of 398,610 women from 25 countries demonstrated that when adjusted for BMD, high BMI was a risk factor for all osteoporotic fractures⁹. In view of this, understanding the relationships of skeletal strength and the mechanisms of fracture in obese individuals should provide insights for improving preventive and therapeutic approaches for obese populations.

Bone structure and the relative amount of bone in the cortical and trabecular compartments contribute to femoral strength¹⁰. Evaluations of bone structure^{11–13} and strength^{14–17} provide mechanistic insight into the etiology of hip fracture. Measures of femoral structure from hip structure analysis (HSA) using DXA scans and measures of distal radius/tibia strength from finite element analysis using peripheral QCT scans are reduced relative to body weight in overweight and obese older women^{18,19}. However, these associations have not been studied in men, and hip structural analysis from DXA is not as refined for analysis of femoral strength as is finite element analysis of CT scans²⁰. The purpose of this study was to investigate the associations between BMI and QCT-derived hip bone density and structural measurements and to further evaluate the association of BMI with finite element-estimated hip strength and load to strength ratio in older men participating in the Osteoporotic Fractures in Men Study (MrOS). We hypothesized that obesity may have deleterious effects on bone structure or volumetric BMD, and that in the obese bone strength may not be sufficient to protect against the increased impact force exerted on bone during a fall.

Materials and methods

Subjects

MrOs is a prospective cohort study designed to determine risk factors for fracture in men aged 65 years and older. The design and recruitment have been previously described^{21,22}.

Briefly, 5994 community-dwelling, ambulatory men were recruited from six US academic medical centers (Birmingham, AL, Minneapolis, MN, Palo Alto, CA, Pittsburgh, PA, Portland, OR, and San Diego, CA) from March 2000 through April 2002. Eligible participants were at least 65 years old, able to walk without assistance from another person, and had not had bilateral hip replacement surgery. The institutional review board at each study site approved the study and written informed consent was obtained from all participants.

Analytic Sample

A total of 3785 participants (63% of MrOS) were referred for QCT scans as previously reported²³. Of these, 122 were not eligible for hip scan because of hip replacement and 103 hip scans were not available for processing because they were lost or corrupted during data transfer, leaving 3560 hip scans available for processing. Of the initial 3560 hip scans available for processing, 133 failed image processing for various reasons²³. A total of 3427 with complete data for femoral neck integral BMD was included in these analyses. We excluded men reporting the use of osteoporosis medications, inhaled or oral corticosteroids and hormone therapy (n=354) or having missing BMI or BMI < 18.5 (n=6). Thus, a sample of 3067 men formed the analytic sample for the association analyses with QCT measures.

Finite element (FE) analysis was performed in QCT scans selected with a stratified sampling plan. Baseline QCT scans was selected at random for 384 men. In addition, FE was performed in baseline QCT for all remaining 378 men from Portland and Birmingham sites whose scans had not been selected for the random sample. Eighty-eight men were excluded for reporting the use of osteoporosis medications, corticosteroid, or androgens, and two men were excluded for having missing BMI measurements or low BMI < 18.5, leaving a total of 672 men for the analyses of FE measures.

Baseline characteristics

Height was measured on Harpenden stadiometers and weight on standard balance beam or digital scales, with participants wearing light clothing without shoes. Body mass index (BMI) was calculated as weight (kilograms) per square height (meter). Subjects were classified as normal weight (BMI: 18.5 to < 25 kg/m²), overweight (25 to < 30 kg/m²), obese (>= 30 kg/m²), using World Health Organization categories of BMI. Demographic, lifestyle factors including smoking status and alcohol intake, medical history were obtained from standardized questionnaires. Physical activity level was assessed with the Physical Activity Scale for the Elderly (PASE)²⁴. Dietary calcium and vitamin D were assessed using a modified Block Food Frequency Questionnaire²⁵. A computerized dictionary, based on the original Established Populations for Epidemiologic Studies of the Elderly (EPESI) coding system²⁶ was used to categorize medications. All prescription medications validated by the clinics were recorded in an electronic medication inventory database (San Francisco Coordinating Center, San Francisco, CA). Each medication was matched to its ingredients based on the Iowa Drug Information Service (IDIS) Drug Vocabulary (College of Pharmacy, University of Iowa, Iowa City, IA).

DXA and QCT scanning

DXA BMD (g/cm^2) was measured in using fan-beam DXA (QDR 4500W, Hologic Inc., Waltham, MA, USA) and using standardized procedures and centralized quality control procedures²². Hip QCT scans were obtained from the femoral head to 3.5 cm below the lesser trochanter at settings of 80 kVp, 280 mA, 3-mm slice thickness, and 512×512 matrix in spiral reconstruction mode. Scanner models used at the sites were described²³. Calibration standards with known hydroxyapatite concentrations (150, 75, and $0 \text{ mg}/\text{cm}^3$; Image Analysis) were included in each scan²³.

QCT-derived femoral measures

Image processing was described previously²³. Briefly, regions of interest (ROIs) in the left proximal femur were identified in the QCT images reformatted along the centroidal axis of the femoral neck. The femoral neck ROI was defined as at the portion of the neck extending from the slice with minimum cross-sectional area (medial boundary) to a point 25% of the proximal distance of the neck axis length toward the maximal cross-sectional area. Within this femoral neck ROI, the following measures were obtained. The cross-sectional area (cm^2) was computed as the area within the periosteal boundary at the minimum cross-section. Integral volume (cm^3) was computed as the total volume within the periosteal boundary. The cortical volume was defined by applying a threshold of $0.35 \text{ g}/\text{cm}^3$ to all voxels between the periosteal boundary and the internal trabecular bone. Medullary volume was computed by subtracting the cortical volume from the integral volume. The percent cortical volume was computed as cortical volume divided by integral volume times 100. Volumetric BMD was computed over all voxels in the integral, trabecular, and cortical volumes.

Estimation of femoral strength and the load-to-strength ratio (Φ)

FE analyses were performed as described previously¹⁷¹⁶. Briefly, FE models were created to simulate a fall to the side of the hip, the diaphysis angled at 15° with respect to the ground with 15° of internal rotation. This represents a severe, unprotected, fall to the side of the hip, which is known to be associated with a high risk of fracture. Strength was calculated from the resulting nonlinear force–deformation curve as the force at 4% deformation of the femoral head with respect to the greater trochanter. The load-to-strength ratio (Φ) was calculated for a simulated sideways fall orientation and impact directly on the greater trochanter. The *in vivo* impact force (the “load”) on the side of the trochanter was estimated for each subject from biomechanical theory using patient-specific weight and height information. A uniform value of trochanteric soft-tissue thickness was assumed for all men. The resulting expression for the load-to-strength ratio was directly proportional to patient mass and the square root of patient height and was inversely and non-linearly proportional to the FE-derived strength²⁷. Theoretically, if the *in vivo* impact force is correctly calculated for each patient, when $\Phi \geq 1$, a fracture is predicted to occur, whereas no fracture is predicted when $\Phi < 1$ ²⁷ regardless, higher values of Φ denote a greater risk of fracture.

Statistical Analyses

We compared the baseline characteristics of men across BMI category using ANOVA tests for continuous variables or chi-square tests for categorical variables. To examine the association between BMI and hip QCT measures, we used multivariable general linear models (GLM) to estimate least square (LS) means of each QCT outcome variable by BMI categories and performed linear trend tests. To test our hypothesis of differential associations of BMI with QCT parameters between non-obese and obese individuals, we specifically evaluated the interactions between obesity and BMI, and stratified obese status to assess the relationships between BMI and QCT parameters expressed as SD change per unit increase of BMI in each stratum. We generated Loess curves to further visualize the nonlinearity in the associations between BMI and QCT measurements. A logistic regression model was used to evaluate the OR and 95% CI of having a theoretical biomechanical fracture threshold (defined as a load-to-strength ratio >1.0) according to obese status. Covariates for all analyses included age, race, enrollment site and physical activity. Other potential confounders included calcium and vitamin D intake, baseline history of diabetes and fracture after age 50 and were assessed by their change in the associations between BMI categories and each QCT and FE measures by 10% or more. A two-tailed alpha of 0.05 was used.

Because of two different sampling strategies involved in FE samples, we first evaluated potential sampling effect on the associations. We examined the associations of BMI with FE strength and load-to-strength ratio in the random sample and the sample from Portland and Birmingham sites separately (Supplemental table 1a and 1b), and then compared the point estimates from two samples. We found no significant difference of point estimates for both strength and load-to strength ratio between two samples (P for interaction > 0.05) (Supplemental table 1c). We further evaluated the associations between BMI and FE-derived biomechanical measurements stratified by obese status in the random sample and the sample from Portland and Birmingham sites separately (Supplemental table 2a and 2b). The point estimates were compatible between the two samples. Thus the two samples were combined for the association analyses, and results from the combined samples were presented.

Results

Baseline characteristics of the 3067 men included in these analyses are presented in Table 1. Compared to men in lower BMI categories, those in higher BMI categories were younger and physically less active and had lower intakes of calcium and vitamin D. A greater proportion of obese men reported poor health status in general and had a history of chronic diseases than did men in lower BMI categories. There was no difference in race, smoking status and alcohol consumption across BMI categories.

All bone density and geometry parameters at the femoral neck were significantly different across BMI categories adjusted for age, race and clinical center (Figure 1). Compared with normal weight men, those who were overweight or obese had greater volumetric and areal BMD, greater total bone size (minimal cross-sectional area and integral volume) and cortical thickness (percent cortical volume).

The associations between BMI and QCT measures (except for medullary volume) were nonlinear as shown in Supplemental figure 1 (Loess curve of femoral neck integral vBMD and percent cortical volume are shown as examples), and were significantly different between non-obese and obese men (Table 2) (P for interaction ≤ 0.014). For men with BMI < 30 , there was a linear relationship between BMI and most QCT measures. In those men, increasing BMI was associated with higher integral, cortical and trabecular vBMD, integral volume, cross-sectional area, and percent cortical volume (SD increase per unit of BMI: 0.05 for integral vBMD, 0.042 for cortical vBMD and 0.002 for trabecular vBMD, 0.027 for integral volume, 0.03 for cross-sectional area and 0.042 for percent cortical volume, all $P < 0.001$). However, among men with BMI ≥ 30 , increasing BMI was not associated with further increases in those parameters. Although increasing BMI was associated with increases in areal femoral neck DXA BMD for both non-obese and obese men, the increment in femoral neck BMD per unit increase in BMI was less for obese men than for non-obese men (SD increase per unit of BMI : 0.035 for obese men vs 0.105 for non-obese men, P for interaction < 0.0001). Inclusion of potential confounders had no substantial effect on these associations.

FE analyses suggested there was a statistically significant linear trend for increasing hip strength from the normal to the obese categories (Figure 2A). Compared to normal-weight men, obese men had higher strength (mean: 6130N vs 5300N, $P < 0.0001$). However, there was also a higher load-to-strength ratio in obese men (mean: 0.79 for obese men vs 0.66 for the normal weight, $P < 0.0001$) (Figure 2B). The increment in femoral strength per unit increase in BMI for obese men was not significantly different from that for non-obese men (SD increase per unit BMI: -0.03 vs 0.06 , P for interaction $= 0.36$) (Table 3). Similarly, there was no significant difference of the increment in load-to-strength ratio between obese and non-obese men (SD increase per unit BMI: 0.14 vs 0.1, P for interaction $= 0.99$) (Table 3). Further, using the load-to-strength ratio = 1.0 as a theoretical value of the fracture threshold, 24% in the obese vs 14% in the non-obese had a load-to-strength ratio greater than the threshold. Obese men were 4 times more likely to have a load-to-strength ratio > 1.0 compared to normal-weight men (OR: 4.66; 95% CI: 2.16–10.05; $P < 0.0001$).

Discussion

In this cross-sectional analysis of BMI and QCT-derived structural and densitometric measures of proximal femur in older men, the relationship between BMI and hip QCT measures was significantly different between men categorized as non-obese and obese. Whereas for non-obese subjects, BMI was associated with a linear increase in femoral neck volumetric BMD, percent cortical volume, and minimal cross-sectional area, for obese men there was no further increase in these measures as BMI increased. Furthermore, compared to normal-weight men, obese men had higher hip strength but also higher load-to-strength ratio, suggesting that in obese men increments in strength with increased BMI might not be sufficient to counter the higher forces involved in a fall.

In theory, individuals with high BMI may accrue more bone and greater bone strength to support a greater body mass^{1,2}. In fact, in this study men meeting the criteria for obesity based on their BMI did have greater femoral neck volumetric BMD, bone size and cortical

thickness compared to the overweight and normal weight men. However, the relationship of increasing BMI with progressive increases in those QCT measures appeared to reach a plateau at BMIs of 30 kg/m², with no additional increment in those QCT parameters with further increases in BMI. Previous studies of older women also found that relative to their body weight, heavier individuals have lower BMD and weaker radial and tibial geometry compared to those with normal weight¹⁸. Together, these findings provide evidence that increments in hip BMD and structural elements in the older obese men are not commensurate with their higher weight. Because these QCT-derived structural and densitometric parameters have been shown to significantly affect bone strength^{10,28–31} and are associated with hip fracture risk independent of areal BMD¹¹, our findings may have some important implications for understanding the pathophysiology of hip fracture in subjects with high BMI. In addition, our finding could be of clinical importance regarding fracture prediction. The FRAX algorithm uses a series of clinical risk factors, including BMI as a continuous variable, to calculate fracture risk³². It has been noted that FRAX may underestimate fracture probability in obese individuals and thus the US-based FRAX-based intervention thresholds (eg, >20% probability of a major osteoporotic fracture or >3% probability of hip fracture in the following 10 years) may be too high for obese subjects³³. Therefore, incorporating the non-linear association of BMI with bone measures in FRAX model may improve the identification of individual at risk for fracture in the obese populations.

In our study, increasing BMI was associated with progressive increase in QCT measures in non-obese men, but that positive association of BMI appears to reach plateau in the men categorized as obese, in whom there was not a continuing improvement in bone density and structural parameters with increasing BMI. Although we didn't observe significant deleterious effect of obesity on bone structure or volumetric BMD, our finding provides support that adiposity in obese individual may not be beneficial to bone health. The beneficial effect of fat mass on bone is to some extent mediated through its mechanical loading and hormones secreted from either the pancreatic beta cell or the adipocyte^{1,2}. However, both experimental and human studies have suggested the inverse relationship between fat, in particular visceral fat, and BMD^{34,35} and that adipocyte-derived adipokine, chemokines and cytokines may impair bone remodeling processes³⁶. A recent study in obese men reported that those with high visceral adipose tissue had weak QCT-derived microarchitecture and mechanical properties compared with those with low visceral adipose tissue³⁷. Another study reported that when the mechanical loading effect of body weight is statistically removed, higher body fat mass is correlated with lower bone mass³⁸. Moreover, although weight loss overall may lead to greater bone loss^{39,40}, there is evidence that the loss of fat mass in particular may increase BMD⁴¹ and bone strength⁴², supporting the negative impact of fat mass on bone. Our findings, together with evidence from these other reports, suggest that adverse effects of excess fat in the obese may outweigh the benefit of increased mechanic loading on bone density and structure³⁴. On the other hand, lean mass is associated with high BMD and bone strength^{43,44}. Because lean mass is progressively smaller fraction of total mass as BMI increases¹⁸, reducing lean mass relative to total body mass in the obese may also explain no additional benefit on QCT measures with increasing

BMI among obese men in our study. Finally, obese men have more medical conditions, and more limited activity, that may impair the potential to respond to mechanical stimuli.

FE analysis of hip QCT scans showed that both hip strength and load-to-strength ratio increased with increasing BMI. That trend seemed to be similar in men categorized as obese and non-obese, although the power to detect differences in this relationship between obese and non-obese men may have been limited due to the relatively fewer men with FE analyses than in the full cohort. However, it was clear that the load-to-strength ratio was higher in obese men. In fact, compared to non-obese men, a high proportion of obese men had load-to-strength ratios greater than the theoretical biomechanical fracture threshold $^{27}(\Phi= 1.0)$ (24% in the obese vs 14% in the non-obese). Those findings suggest that in obese men the increase in strength with increasing BMI might not be sufficient to compensate for fall forces that increase in direct proportion to body weight. As a result, obese individuals may be more likely to suffer fracture with a sideways fall.

Our study has several strengths. It is the first evaluation of volumetric and structural properties derived from hip QCT scans across BMI categories in a large population of older men. Analysis of QCT scans provides a more accurate assessment of hip structural characteristics than does HSA analysis of DXA images ⁴⁵. Moreover, biomechanical analysis using FE provides an estimate of femoral strength in a clinically relevant type of fall ¹⁷, the load-to-strength ratio accounts for subject-specific estimates of force encountered by the hip in a fall, and FE-derived femoral strength and load-to-strength ratio also predict hip fracture^{16,17}.

Our study also has several limitations. It has been estimated that alteration of body composition can result in artifacts in the assessment of BMD by both DXA and QCT ⁴⁶, but the impact of changes in body composition on QCT seems to be small ⁴⁷. A recent study using simulated increases in body fat suggested that whereas fat layering introduces error and decreases the reproducibility of both DXA and QCT BMD measurements, the error in QCT is smaller and more uniform than with DXA ⁴⁷. However, the extent to which excess fat tissue in obese individuals influence the quality of QCT image remains unclear. In addition, proximal femur QCT scans in our study were obtained at a relatively low energy dose which may increase the possibility of image artifacts in participants with large body mass. Our estimates of the hip impact force did not account for any patient-specific variations in cushioning effects of trochanteric fat thickness, and instead assumed a constant value of trochanteric fat thickness for all subjects, which may have led to a relative overestimation of the load-to-strength ratio in overweight and obese men, who presumably have higher values of trochanteric fat tissue. While increased fat thickness should reduce the impact force to some degree, however, it is not clear if very large values of fat thickness add any further cushioning or protective effect^{48,49}. Future studies are needed to incorporate real fat thickness measurements, and the associated biomechanics, into these analyses. Different sampling strategies involved in FE sub-cohort may bias the association of FE measures with BMI. Also, although BMI is widely used and accepted method to define obesity, it is not a precise measure of excess body fat and does not reflect fat distribution, particularly among older population. Future study using more precise assessment of body fat, such as QCT measures, may better characterize the relationship between obesity and hip structures. In

addition, the cross-sectional design of the study makes it impossible to assess causality in the observed associations. Finally, our participants were older men, and most were of European ancestry. The generalizability of our findings to older women, other races /ethnic groups or younger populations is unknown.

In conclusion, the relationships between BMI and femoral BMD and structure appear to be nonlinear and are different between men categorized as non-obese and obese. In non-obese men, increasing BMI is associated with increasingly greater bone size, cortical thickness and cortical and trabecular BMDs. In individuals with BMI levels in the obese range, there is no further increase in those volumetric and structure parameters with additional increases in BMI. FE analyses suggest that the increase in strength is not proportionate with increased impact force in obese men. Our data provide some insight into the relationship between skeletal structure and strength and hip fracture risk in obese men.

Supplementary Material

Refer to Web version on PubMed Central for supplementary material.

Acknowledgments

The Osteoporotic Fractures in Men (MrOS) Study is supported by National Institutes of Health funding. The following institutes provide support: the National Institute of Arthritis and Musculoskeletal and Skin Diseases (NIAMS), the National Institute on Aging (NIA), the National Center for Research Resources (NCRR), and NIH Roadmap for Medical Research under the following grant numbers: U01 AR45580, U01 AR45614, U01 AR45632, U01 AR45647, U01 AR45654, U01 AR45583, U01 AG18197, U01-AG027810, and UL1 RR024140. Additional support was provided to Dr. Nielson by the Office of Research on Women's Health and the National Institute of Child Health and Human Development, Building Interdisciplinary Research Careers in Women's Health (BIRCWH) grant number HD043488-08

References

1. Albala C, Yanez M, Devoto E, Sostin C, Zeballos L, Santos JL. Obesity as a protective factor for postmenopausal osteoporosis. *Int J Obes Relat Metab Disord*. Nov; 1996 20(11):1027–1032. [PubMed: 8923160]
2. Klein KO, Larmore KA, de Lancey E, Brown JM, Considine RV, Hassink SG. Effect of obesity on estradiol level, and its relationship to leptin, bone maturation, and bone mineral density in children. *J Clin Endocrinol Metab*. Oct; 1998 83(10):3469–3475. [PubMed: 9768648]
3. Kanis JA, Bianchi G, Bilezikian JP, et al. Towards a diagnostic and therapeutic consensus in male osteoporosis. *Osteoporos Int*. Nov; 2011 22(11):2789–2798. [PubMed: 21509585]
4. Kanis J, Johnell O, Gullberg B, et al. Risk factors for hip fracture in men from southern Europe: the MEDOS study. *Mediterranean Osteoporosis Study*. *Osteoporos Int*. 1999; 9(1):45–54. [PubMed: 10367029]
5. Premaor MO, Compston JE, Fina Aviles F, et al. The association between fracture site and obesity in men: a population-based cohort study. *J Bone Miner Res*. Aug; 2013 28(8):1771–1777. [PubMed: 23371035]
6. Premaor MO, Pilbrow L, Tonkin C, Parker RA, Compston J. Obesity and fractures in postmenopausal women. *J Bone Miner Res*. Feb; 2010 25(2):292–297. [PubMed: 19821769]
7. Owusu W, Willett W, Ascherio A, et al. Body anthropometry and the risk of hip and wrist fractures in men: results from a prospective study. *Obes Res*. Jan; 1998 6(1):12–19. [PubMed: 9526965]
8. Nielson CM, Marshall LM, Adams AL, et al. BMI and fracture risk in older men: the osteoporotic fractures in men study (MrOS). *J Bone Miner Res*. Mar; 2011 26(3):496–502. [PubMed: 20814955]

9. Johansson H, Kanis JA, Oden A, et al. A meta-analysis of the association of fracture risk and body mass index in women. *J Bone Miner Res.* Jan; 2014 29(1):223–233. [PubMed: 23775829]
10. Nawathe S, Akhlaghpour H, Bouxsein ML, Keaveny TM. Microstructural failure mechanisms in the human proximal femur for sideways fall loading. *J Bone Miner Res.* Feb; 2014 29(2):507–515. [PubMed: 23832419]
11. Black DM, Bouxsein ML, Marshall LM, et al. Proximal femoral structure and the prediction of hip fracture in men: a large prospective study using QCT. *J Bone Miner Res.* Aug; 2008 23(8):1326–1333. [PubMed: 18348697]
12. Carballido-Gamio J, Harnish R, Saeed I, et al. Proximal femoral density distribution and structure in relation to age and hip fracture risk in women. *J Bone Miner Res.* Mar; 2013 28(3):537–546. [PubMed: 23109068]
13. Carballido-Gamio J, Harnish R, Saeed I, et al. Structural patterns of the proximal femur in relation to age and hip fracture risk in women. *Bone.* Nov; 2013 57(1):290–299. [PubMed: 23981658]
14. Amin S, Kopperdahl DL, Melton LJ 3rd, et al. Association of hip strength estimates by finite-element analysis with fractures in women and men. *J Bone Miner Res.* Jul; 2011 26(7):1593–1600. [PubMed: 21305605]
15. Keaveny TM, Kopperdahl DL, Melton LJ 3rd, et al. Age-dependence of femoral strength in white women and men. *J Bone Miner Res.* May; 2010 25(5):994–1001. [PubMed: 19874201]
16. Kopperdahl DL, Aspelund T, Hoffmann PF, et al. Assessment of incident spine and hip fractures in women and men using finite element analysis of CT scans. *J Bone Miner Res.* Mar; 2014 29(3):570–580. [PubMed: 23956027]
17. Orwoll ES, Marshall LM, Nielson CM, et al. Finite element analysis of the proximal femur and hip fracture risk in older men. *J Bone Miner Res.* Mar; 2009 24(3):475–483. [PubMed: 19049327]
18. Beck TJ, Petit MA, Wu G, LeBoff MS, Cauley JA, Chen Z. Does obesity really make the femur stronger? BMD, geometry, and fracture incidence in the women's health initiative-observational study. *J Bone Miner Res.* Aug; 2009 24(8):1369–1379. [PubMed: 19292617]
19. Sornay-Rendu E, Boutroy S, Vilayphiou N, Claustrat B, Chapurlat RD. In Obese Postmenopausal Women, Bone Microarchitecture and Strength Are Not Commensurate to Greater Body Weight: The Os des Femmes de Lyon (OFELY) Study. *J Bone Miner Res.* Jul; 2013 28(7):1679–1687. [PubMed: 23371055]
20. Danielson ME, Beck TJ, Karlamangla AS, et al. A comparison of DXA and CT based methods for estimating the strength of the femoral neck in post-menopausal women. *Osteoporos Int.* Apr; 2013 24(4):1379–1388. [PubMed: 22810918]
21. Blank JB, Cawthon PM, Carrion-Petersen ML, et al. Overview of recruitment for the osteoporotic fractures in men study (MrOS). *Contemp Clin Trials.* Oct; 2005 26(5):557–568. [PubMed: 16085466]
22. Orwoll E, Blank JB, Barrett-Connor E, et al. Design and baseline characteristics of the osteoporotic fractures in men (MrOS) study--a large observational study of the determinants of fracture in older men. *Contemp Clin Trials.* Oct; 2005 26(5):569–585. [PubMed: 16084776]
23. Marshall LM, Lang TF, Lambert LC, Zmuda JM, Ensrud KE, Orwoll ES. Dimensions and volumetric BMD of the proximal femur and their relation to age among older U.S. men. *J Bone Miner Res.* Aug; 2006 21(8):1197–1206. [PubMed: 16869717]
24. Washburn RA, Smith KW, Jette AM, Janney CA. The Physical Activity Scale for the Elderly (PASE): development and evaluation. *J Clin Epidemiol.* Feb; 1993 46(2):153–162. [PubMed: 8437031]
25. Block G, Woods M, Potosky A, Clifford C. Validation of a self-administered diet history questionnaire using multiple diet records. *J Clin Epidemiol.* 1990; 43(12):1327–1335. [PubMed: 2254769]
26. Pahor M, Chrischilles EA, Guralnik JM, Brown SL, Wallace RB, Carbonin P. Drug data coding and analysis in epidemiologic studies. *Eur J Epidemiol.* Aug; 1994 10(4):405–411. [PubMed: 7843344]
27. Keaveny TM, Bouxsein ML. Theoretical implications of the biomechanical fracture threshold. *J Bone Miner Res.* Oct; 2008 23(10):1541–1547. [PubMed: 18410232]

28. Bauer JS, Kohlmann S, Eckstein F, Mueller D, Lochmuller EM, Link TM. Structural analysis of trabecular bone of the proximal femur using multislice computed tomography: a comparison with dual X-ray absorptiometry for predicting biomechanical strength in vitro. *Calcif Tissue Int.* Feb; 2006 78(2):78–89. [PubMed: 16467973]
29. Bousson V, Le Bras A, Roqueplan F, et al. Volumetric quantitative computed tomography of the proximal femur: relationships linking geometric and densitometric variables to bone strength. Role for compact bone. *Osteoporos Int.* 2006; 17(6):855–864. [PubMed: 16547689]
30. Lang TF, Keyak JH, Heitz MW, et al. Volumetric quantitative computed tomography of the proximal femur: precision and relation to bone strength. *Bone.* Jul; 1997 21(1):101–108. [PubMed: 9213015]
31. Orwoll ES. Toward an expanded understanding of the role of the periosteum in skeletal health. *J Bone Miner Res.* Jun; 2003 18(6):949–954. [PubMed: 12817746]
32. Kanis JA, Johnell O, Oden A, Johansson H, McCloskey E. FRAX and the assessment of fracture probability in men and women from the UK. *Osteoporos Int.* Apr; 2008 19(4):385–397. [PubMed: 18292978]
33. Caffarelli C, Alessi C, Nuti R, Gonnelli S. Divergent effects of obesity on fragility fractures. *Clinical interventions in aging.* 2014; 9:1629–1636. [PubMed: 25284996]
34. Gilsanz V, Chalfant J, Mo AO, Lee DC, Dorey FJ, Mittelman SD. Reciprocal relations of subcutaneous and visceral fat to bone structure and strength. *J Clin Endocrinol Metab.* Sep; 2009 94(9):3387–3393. [PubMed: 19531595]
35. Reid IR. Relationships between fat and bone. *Osteoporos Int.* May; 2008 19(5):595–606. [PubMed: 17965817]
36. Rosen CJ, Bouxsein ML. Mechanisms of disease: is osteoporosis the obesity of bone? *Nat Clin Pract Rheumatol.* Jan; 2006 2(1):35–43. [PubMed: 16932650]
37. Bredella MA, Lin E, Gerweck AV, et al. Determinants of bone microarchitecture and mechanical properties in obese men. *J Clin Endocrinol Metab.* Nov; 2012 97(11):4115–4122. [PubMed: 22933540]
38. Zhao LJ, Liu YJ, Liu PY, Hamilton J, Recker RR, Deng HW. Relationship of obesity with osteoporosis. *J Clin Endocrinol Metab.* May; 2007 92(5):1640–1646. [PubMed: 17299077]
39. Schwartz AV, Johnson KC, Kahn SE, et al. Effect of 1 year of an intentional weight loss intervention on bone mineral density in type 2 diabetes: results from the Look AHEAD randomized trial. *J Bone Miner Res.* Mar; 2012 27(3):619–627. [PubMed: 22354851]
40. Villareal DT, Shah K, Banks MR, Sinacore DR, Klein S. Effect of weight loss and exercise therapy on bone metabolism and mass in obese older adults: a one-year randomized controlled trial. *J Clin Endocrinol Metab.* Jun; 2008 93(6):2181–2187. [PubMed: 18364384]
41. Christensen P, Bartels EM, Riecke BF, et al. Improved nutritional status and bone health after diet-induced weight loss in sedentary osteoarthritis patients: a prospective cohort study. *Eur J Clin Nutr.* Apr; 2012 66(4):504–509. [PubMed: 22190136]
42. Scibora LM, Buchwald H, Petit MA, Hughes J, Ikramuddin S. Bone Strength Is Preserved Following Bariatric Surgery. *Obes Surg.* Jun 28, 2014
43. Travison TG, Araujo AB, Esche GR, Beck TJ, McKinlay JB. Lean mass and not fat mass is associated with male proximal femur strength. *J Bone Miner Res.* Feb; 2008 23(2):189–198. [PubMed: 17922610]
44. Wang MC, Bachrach LK, Van Loan M, Hudes M, Flegal KM, Crawford PB. The relative contributions of lean tissue mass and fat mass to bone density in young women. *Bone.* Oct; 2005 37(4):474–481. [PubMed: 16040285]
45. Beck TJ. Extending DXA beyond bone mineral density: understanding hip structure analysis. *Curr Osteoporos Rep.* Jun; 2007 5(2):49–55. [PubMed: 17521505]
46. Bolotin HH, Sievanen H, Grashuis JL. Patient-specific DXA bone mineral density inaccuracies: quantitative effects of nonuniform extraosseous fat distributions. *J Bone Miner Res.* Jun; 2003 18(6):1020–1027. [PubMed: 12817754]
47. Yu EW, Thomas BJ, Brown JK, Finkelstein JS. Simulated increases in body fat and errors in bone mineral density measurements by DXA and QCT. *J Bone Miner Res.* Sep 13, 2011

48. Buxsein ML, Szulc P, Munoz F, Thrall E, Sornay-Rendu E, Delmas PD. Contribution of trochanteric soft tissues to fall force estimates, the factor of risk, and prediction of hip fracture risk. *J Bone Miner Res.* Jun; 2007 22(6):825–831. [PubMed: 17352651]
49. Robinovitch SN, McMahon TA, Hayes WC. Force attenuation in trochanteric soft tissues during impact from a fall. *J Orthop Res.* Nov; 1995 13(6):956–962. [PubMed: 8544034]




Figure 1. Least squares means for femoral neck QCT measurements by BMI category adjusted for age, site, race and physical activity in men of the MrOS study. The error bars indicate standard errors.




Figure 2.

BMI and FE-derived biomechanical measurements (A) Strength (B) Load-to-strength ratio, adjusted for age, site, race and physical activity in the men of the MrOS study. P is for trend test. § P < 0.05 and * P < 0.0001 compared with normal-weight men.

Sample size: N for normal weight=202; N for overweight=380; N for obese=90

The length of the box represents the interquartile range (the distance between the 25th and 75th percentiles). The symbol in the box interior represents the group mean. The horizontal line in the box interior represents the group median. The whiskers on the boxes depict the extreme values.

Table 1

Characteristics of men in MrOS study by BMI category (N = 3067)

	Normal (18.5–24.9 kg/m ²)	Overweight (25–29.9 kg/m ²)	Obese (30–50.7 kg/m ²)	P
N (%)	829(27.0)	1568(51.1)	670 (21.9)	
Age (yr)	74.5(6.3)	73.4(5.8)	72.2(5.2)	<0.0001
Caucasian, n (%)	716(86.4)	1386(88.4)	574(85.7)	0.14
Current smoker, n (%)	32(3.9)	64(4.1)	15(2.2)	0.09
Alcohol (drinks/wk)	4.0(5.8)	4.3(7.0)	4.0(6.7)	0.39
Physical activity score	152(70)	149(65)	144(67)	0.08
Self-reported good health, n (%)	740(89.3)	1376(87.8)	536(80)	<0.0001
Calcium intake (mg)	1,206(620)	1,136(589)	1066(557)	<0.0001
Vitamin D intake (IU)	408(243)	392(246)	367(233)	0.005
Self-reported medical history				
Osteoporosis, n (%)	12(1.5)	30(1.9)	13(1.9)	0.68
Diabetes, n (%)	55(6.6)	152(9.7)	137(20.5)	<0.0001
Arthritis, n (%)	319(38.5)	717(45.7)	367(54.8)	<0.0001
Fracture after age 50, n (%)	404(48.7)	851(54.3)	379(56.6)	0.005

Values are means (SD) unless otherwise noted. P values for categorical variables are from chi-square tests (Fisher's exact test for small expected frequencies); P values for continuous variables were calculated using one-way ANOVA.

Table 2

The association of BMI with proximal femoral QCT measurements stratified by obesity status in men of the MrOS study

	BMI < 30 (n=2397)			BMI ≥ 30 (n=670)		
	SD increase per unit of BMI	P	SD increase per unit of BMI	P	P for Interaction *	
Femoral neck						
BMD (g/cm ³)						
Integral	0.05 ± 0.008	<0.0001	-0.012 ± 0.014	0.39	0.0003	
Cortical	0.042 ± 0.008	<0.0001	-0.008 ± 0.013	0.54	0.002	
Trabecular	0.002 ± 0.0004	<0.0001	-0.0008 ± 0.0006	0.18	0.0005	
Volume (cm ³)						
Integral	0.027 ± 0.008	0.001	-0.015 ± 0.014	0.28	0.014	
Medullary	0.006 ± 0.009	0.47	-0.011 ± 0.014	0.45	0.33	
Percent cortical volume	0.042 ± 0.008	<0.0001	-0.009 ± 0.013	0.49	0.002	
Cross-sectional area (cm ²)	0.03 ± 0.009	0.0005	-0.02 ± 0.014	0.14	0.003	
DXA BMD	0.105 ± 0.008	<0.0001	0.035 ± 0.014	0.011	<0.0001	

P-value was adjusted for age, site, race and physical activity.

* The interaction was to test the difference of point estimates between the non-obese and the obese.

The association of BMI with FE-derived measures stratified by obesity status in men of the MrOS study

Table 3

	BMI < 30 (n=582)		BMI ≥ 30 (n=90)		P for Interaction*
	SD increase per unit of BMI	P	SD increase per unit of BMI	P	
Strength	0.056 ± 0.014	0.0002	-0.03 ± 0.064	0.69	0.36
Load-to-strength ratio	0.094 ± 0.015	<0.0001	0.14 ± 0.07	0.06	0.99

P-value was adjusted for age, site, race and physical activity.

*The interaction was to test the difference of point estimates between the non-obese and the obese.