



RESEARCH ARTICLE

1-Octen-3-ol – the attractant that repels [version 1; referees: 4 approved]

Pingxi Xu, Fen Zhu, Garrison K. Buss, Walter S. Leal

Department of Molecular and Cellular Biology, University of California-Davis, Davis, CA, 95616, USA

v1 First published: 18 Jun 2015, 4:156 (doi: [10.12688/f1000research.6646.1](https://doi.org/10.12688/f1000research.6646.1))
 Latest published: 18 Jun 2015, 4:156 (doi: [10.12688/f1000research.6646.1](https://doi.org/10.12688/f1000research.6646.1))

Abstract

Since the discovery in the early 1980s that 1-octen-3-ol, isolated from oxen breath, attracts tsetse fly, there has been growing interest in exploring the use of this semiochemical as a possible generic lure for trapping host-seeking mosquitoes. Intriguingly, traps baited with 1-octen-3-ol captured significantly more females of the malaria mosquito, *Anopheles gambiae*, and the yellow fever mosquito, *Aedes aegypti*, than control traps, but failed to attract the southern house mosquito, *Culex quinquefasciatus*. Additionally, it has been demonstrated that this attractant is detected with enantioselective odorant receptors (ORs) expressed only in maxillary palps. On the basis of indoor behavioral assays it has even been suggested that 1-octen-3-ol might be a repellent to the southern house mosquito. Our approach was two-prong, i.e., to isolate 1-octen-3-ol-sensitive ORs expressed in maxillary palps and antennae of southern house female mosquito, and test the hypothesis that this semiochemical is a repellent. An OR with high transcript levels in maxillary palps, CquiOR118b, showed remarkable selectivity towards (*R*)-1-octen-3-ol, whereas an OR expressed in antennae, CquiOR114b, showed higher preference for (*S*)-1-octen-3-ol than its antipode. Repellency by a surface landing and feeding assay showed that not only racemic, but enantiopure (*R*)- and (*S*)-1-octen-3-ol are repellents at 1% dose thus suggesting the occurrence of other (*S*)-1-octen-3-ol-sensitive OR(s). Female mosquitoes with ablated maxillary palps were repelled by 1-octen-3-ol, which implies that in addition to OR(s) in the maxillary palps, antennal OR(s) are essential for repellency activity.

Open Peer Review

Referee Status:

	Invited Referees			
	1	2	3	4
version 1 published 18 Jun 2015				
	report	report	report	report

- Joseph Dickens**, United States
Department of Agriculture USA
- Jeffery K. Tomberlin**, Texas A&M University USA
- Wynand van der Goes van Naters**, Cardiff University UK
- Kenneth F. Haynes**, University of Kentucky USA

Discuss this article

Comments (0)

Corresponding author: Walter S. Leal (wsleal2010@gmail.com)

How to cite this article: Xu P, Zhu F, Buss GK and Leal WS. 1-Octen-3-ol – the attractant that repels [version 1; referees: 4 approved] *F1000Research* 2015, 4:156 (doi: [10.12688/f1000research.6646.1](https://doi.org/10.12688/f1000research.6646.1))

Copyright: © 2015 Xu P *et al.* This is an open access article distributed under the terms of the [Creative Commons Attribution Licence](https://creativecommons.org/licenses/by/4.0/), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited. Data associated with the article are available under the terms of the [Creative Commons Zero "No rights reserved" data waiver](https://creativecommons.org/licenses/by/4.0/) (CC0 1.0 Public domain dedication).

Grant information: This work was supported in part by the National Institute of Allergy and Infectious Diseases of the National Institutes of Health under award R01AI095514. FZ sabbatical leave at UC Davis was supported in part by the Chinese Scholarship Council. *The funders had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript.*

Competing interests: No competing interests were disclosed.

First published: 18 Jun 2015, 4:156 (doi: [10.12688/f1000research.6646.1](https://doi.org/10.12688/f1000research.6646.1))

Introduction

1-Octen-3-ol (Figure 1) is a natural product derived from linoleic acid, which was first isolated from the matsutake pine mushroom¹ and thereafter from plants and other fungi. It is approved by US Food and Drug Administration (ASP 1154, Regnum 172.515) as a food additive and also considered a wine fault – an unpleasant characteristic of wine. Since it was discovered as an emanation from oxen breath that attracts tsetse fly², there has been growing interest in using 1-octen-3-ol as an insect attractant. Indeed, it was demonstrated earlier on that 1-octen-3-ol synergizes with CO₂ and thus increase mosquito trapping efficacy³. Intriguingly, field experiments demonstrated that the effect of 1-octen-3-ol on mosquito captures is species specific⁴. Of notice, 1-octen-3-ol seem to have little or no effect on trapping of the southern house mosquito, *Culex quinquefasciatus*, although being undoubtedly an attractant (kairomone) for *Anopheles* and *Aedes* mosquitoes⁵.




Figure 1. Chemical structures of tested compounds. (S) and (R)-1-octen-3-ol, (S) and (R)-1-octyn-3-ol, (S) and (R)-3-octanol.

Club-shaped olfactory basiconic sensilla (peg sensilla) in the maxillary palps of the southern house mosquito harbor three types of olfactory receptor neurons (ORNs) identifiable by their spike amplitudes⁵. The second largest neurons, ORN-B, responded to 1-octen-3-ol with very high sensitivity. Cell B also showed a remarkable selectivity between the two enantiomers of 1-octen-3-ol, with the (R)-(-)-isomer eliciting robust responses at 10 ng dose (256.6 ± 12 spikes/s), whereas the (S)-(+)-antipode eliciting only 115.5 ± 23 spikes/s even when challenged with 100x higher doses, i.e., 1 μg ⁵. It was also demonstrated that neuron-B in the maxillary palps of the malaria mosquito, *Anopheles gambiae*, responds to 1-octen-3-ol⁶, and chiral discrimination was also observed with electrophysiological recordings from the neuron B in the maxillary palps of the yellow fever mosquito, *Aedes aegypti*^{7,8}. Additionally, odorant receptors housed in the maxillary palps of the malaria mosquito⁶ and yellow fever mosquito⁹, AgamOR8 and AegOR8, respectively, showed significant preference for the (R)-enantiomer when co-expressed in *Xenopus* oocytes along with the obligatory co-receptor Orco.

In-door behavioral studies demonstrated that at two doses (R)-(-)-1-octen-3-ol caused an increase in activation for *Cx. quinquefasciatus*, and at seven of the doses tested (R:S)-1-octen-3-ol mixture (84:16) caused significantly more mosquitoes to sustain their flight and reach the capture chambers in a two-choice, Y-tube olfactometer thus suggesting that the isomeric mixture has an excitatory effect⁷. Additionally, they observed that at the highest concentration, mosquitoes that reached the capture chambers moved towards the control chamber rather than the arm containing (R)-(-)-1-octen-3-ol per se or in mixtures, i.e., a reduced attraction response mediated by the (R)-enantiomer⁷. Since the ability of the olfactory system to detect the two enantiomers at this close ratio (approximately 5:1) was not observed in our electrophysiological recordings from peg sensilla⁵, we aimed at testing chiral discrimination at the receptor level. Using cDNA template from the maxillary palps, we cloned the *Culex* ortholog of AgamOR8 and AegOR8, co-expressed it along with CquiOrco in *Xenopus* oocytes, and observed an ability to discriminate enantiomers that reflects our previous findings with single sensillum recordings. Additionally, we cloned a paralogous odorant receptor (OR) from antennae, which responded to both enantiomers of 1-octen-3-ol. These findings provide evidence that peripheral reception of 1-octen-3-ol is enantioselective at the maxillary palps, but random (racemic) at the antennae.

Materials and methods

cDNA preparation

Culex quinquefasciatus mosquitoes used in this study were from a laboratory colony¹⁰, maintained for the last 5 years at $27 \pm 1^\circ\text{C}$ under a photoperiod of 12:12 h (light:dark). Our Davis colony was derived from mosquitoes collected in Merced, California, in the 1950s and maintained by Dr. Anthon Cornel in the Kearney Agricultural Center, University of California. Twenty pairs of antennae and maxillary palps of 9-day old gravid female adults were dissected on ice under a light microscope. Total RNA was extracted using RNeasy Micro Kit (QIAGEN, Valencia, CA). Before synthesizing first-strand cDNA, RNA concentrations from antennae and maxillary palps extracts were adjusted (normalized). First-strand

cDNA was synthesized with oligo (dT) primer (BioRad, Hercules, CA) and GoScript Reverse Transcriptase (Promega, Madison, WI) following the manufacturer's protocol.

Gene cloning

Full-length sequences of *CquiOR114b* and *CquiOR118b* were amplified from female antennae and maxillary palps cDNAs, respectively. The In-Fusion cloning strategy was taken by using In-Fusion® HD Cloning Kit (Clontech® Laboratories, Mountain View, CA). Briefly, the PCR primers were designed with 16 overlapped nucleotides at 5'-end homologous to the linearized ends of the destination vector (pGEMHE), which was double digested by *XbaI* and *XmaI*. The primers for *CquiOR114b* were: forward, **AGATCAATTCCCCGGGacc**ATGGCTACGAAGAAGGTTGCATTC; and two reverse primers, reverse-1: **TCAAGCTTGCTCTAGAT**TACGATCCTTCATAAACCCTT and reverse-2: **TCAAGCTTGCTCTAGAT**TACAACCTCAAGGAACTCTGCTAACTCC. Low case "acc" stands for Kozak sequence.

For *CquiOR118b* two forward and one reverse primers were used; forward-1: **AGATCAATTCCCCGGGATGAACGAC**CTGGTGCAGTTCGAG and forward-2: **AGATCAATTC**CCCCGGGATGCATGTGGGCAACTCCAAGATTTCCG; reverse, **TCAAGCTTGCTCTAGAT**TATTTCTCGCTGGGATCATAAATAGTTTTTCAGCAG. Underline denotes homologous sequence for In-Fusion reaction. PfuUltra II Fusion HS DNA Polymerase (Agilent Technologies, Santa Clara, CA) was used for PCR. PCR products were directly cloned into pGEMHE by using In-Fusion® HD Cloning Kit, following the manufacturer's protocol. In brief, a mix of PCR product, In-Fusion HD Enzyme Premix, pGEMHE vector was incubated at 50°C for 15 min. One microliter of the reaction was added to Stellar™ competent cells for transformation. Plasmids were purified by plasmid mini prep columns SpinSmart (Denville Scientific, South Plainfield, NJ) and sequenced by Davis Sequencing Inc. (Davis, CA).

Quantitative RT-PCR (qPCR)

SsoAdvanced™ SYBR® Green from BioRad was used for qPCR. The reactions were carried out in a BioRad C1000 thermal cycler with CFX96 detection module. The detection primers for *CquiOr114b* were: forward, TTAGCGGGA GAAAACATGGG; reverse, ACTGACTTTGGTACAC GTGG. For *CquiOr118b*, they were: forward, GTCGTTGCTTTTCCTGATGG; reverse, CACGGCATT CTCATATTTTACT. The following primers were used for a reference gene, *CquiOrco*: forward, GCCGGATACGTTTTCTCCTTC; reverse, GCGCATAATTCCTTCAGATG. The reaction system (total volume, 20 µl) included SsoAdvanced SYBR green mix (2x) 10 µl, cDNA 100 ng, paired primer mix 350 nM, and double distilled H₂O. The qPCR program was 95°C for 30 s, 95°C for 5 s, 62°C for 30 s, 72°C for 30 s, and 40 cycles. The melt curves were made from 65°C to 95°C with increment of 5°C, 5 s.

Electrophysiology

Two-Electrode Voltage Clamp Records

The two-electrode voltage-clamp (TEVC) technique was used to measure odorant-induced currents in *Xenopus* oocytes at a holding potential of -80 to -70 mV. Oocytes on stage V or VI were purchased from Ecocyte Bioscience (Austin, TX). Signals were

amplified with an OC-725C amplifier (Warner Instruments), low-pass-filtered at 50 Hz, and digitized at 1 kHz. Data acquisition and analysis were conducted with Digidata 1440A and software pCLAMP10 (Molecular Devices). Traces were collected from same batches and same age of oocytes to make data consistent. Data were analyzed with GraphPad Prism 6 (La Jolla, CA). The following chiral compounds were gifts from Bedoukian Research Inc.: (*R*)-(-)-1-octen-3-ol (CAS# 3687-48-7), (*S*)-(+)-1-octen-3-ol (CAS# 24587-53-9); (*R*)-(-)-1-octyn-3-ol (CAS#32556-70-0), (*S*)-(+)-1-octyn-3-ol (CAS#322556-71-1); (*R*)-(-)-3-octanol (CAS#70492-66-9). Racemic 1-octen-3-ol (CAS # 3391-86-4) and (*S*)-(+)-3-octanol (CAS# 22658-92-0) were acquired from Fluka and Aldrich, respectively (Sigma-Aldrich, Milwaukee, WI).

Behavioral assays

Repellence was measured by using a previously described surface-landing and feeding assay¹⁰. In short, 3–5 days-old female mosquitoes (30–40 mosquitoes per assay) were placed on a two-choice arena designed to attract host-seeking mosquitoes. For physical stimuli, water at 37°C was circulating inside of Dudley tubes, which were painted black in the internal surfaces. Chemical stimuli were provided by stream of CO₂ at 50 ml/min and dental cotton rolls impregnated with defibrinated sheep blood, which were placed on the top of the Dudley tubes. For each test, filter paper rings freshly treated at the outer perimeter with 200 µl of hexane only or 200 µl of a tested compound in hexane were placed to surround each Dudley tube. Mosquito activity was observed and recorded for 5 min with a camcorder equipped with Super NightShot Plus infrared system (Sony Digital Hanycam, DCR-DVD 810). Control and treatment sides were rotated between trials. For all experiments except the concentration screen (used twice), the rings were prepared fresh for each assay. The number of mosquitoes responding to control (hexane only) and treatment were counted in real time and the information also retrieved from video recordings. When testing repellency by racemic and enantiomers, experiments were carried out by using all three compounds, R, S, and racemic in a single set of assays. Each compound was tested interactively, such that R was followed by S and S was followed by racemic, giving rise to a "block" of 3 trials. Three blocks (n=9 for each compound) were conducted per assay. In about half of the trials (n=9 repetitions per compound) test compounds were placed in one of the two sides of the arena. Data were arcsin-transformed before paired two-tailed Student t test comparisons.

Results and discussion

Dataset 1. Raw data

<http://dx.doi.org/10.5256/f1000research.6646.d49878>

Dataset 1: qPCR data obtained for *CquiOR114b*, *CquiOR118b*, and *CquiOrco* (reference gene) with cDNAs from antennae and maxillary palps.

Dataset 2: Concentration-response data generated with *CquiOR118b* and enantiomers of 1-octen-3-ol, 1-octyn-3-ol, and 3-octanol.

Dataset 3: Concentration-response data generated with *CquiOR114b* and enantiomers of 1-octen-3-ol, 1-octyn-3-ol, and 3-octanol.

Dataset 4: Data for repellency activity elicited by 1-octen-3-ol and its enantiomers on female *Culex quinquefasciatus* in a surface-landing and feeding assays.

Cloning and tissue expression

We aimed at cloning *CquiOR118*, the *Cx. quinquefasciatus* ortholog of *AgamOR8* and *AaegOR8*. Despite several attempts, we were unable to clone the full length cDNA (VectorBase, CPIJ013954). On the basis of our RNA-Seq findings suggesting a shorter N-terminal amino acid sequence¹¹, we designed a new forward primer considering the starting codon as the next ATG. Indeed, this led to the full length sequence, which was cloned and confirmed by DNA sequencing. We named the shorter version of this gene *CquiOR118b*, which encodes a protein with 391 amino acid residues and is predicted to have seven transmembrane topology (OCTOPUS, <http://octopus.cbr.su.se/>). While this manuscript was in preparation it has been reported² that a longer version of *CquiOR118*, as predicted in VectorBase, was cloned from another strain of *Cx. quinquefasciatus*. Thus, both *CquiOR118b* from the Merced strain (see below) and *CquiOR118* from the Thai strain are functional². Out of 5 clones we sequenced, we obtained two isoforms of *CquiOR118b*, which differed only in the residue 163 predicted to be in the external cellular loop-2: Ile vs. Val.

Our previous differential expression analysis¹¹ suggested that transcript levels of two genes from the same clade, *CquiOR114* and *CquiOR117*, are significantly higher in antennae than control tissues (legs). Because of the predicted longer C-terminus amino acid sequences encoded by these genes¹¹, we designed primers that would allow us to clone the short and longer versions of these genes. No PCR product was generated with primers for the short sequences, but we cloned and sequenced cDNAs (*CquiOR114b*) encoding proteins with 405 amino acid residues (longer C-terminus) and predicted seven transmembrane topology. Out of 5 cloned sequenced, we found two isoforms, which differed in 3 amino acid residues. We named them *CquiOR114b-1* (Leu-63, Gly-122, and Asp-129) and *CquiOR114b-2* (Trp-63, Glu-122, and Asn-129). These residues are predicted to be part of the first transmembrane segment formed by residues 60 to 80, and the internal cellular loop-2 formed by residues 112–150.

Quantitative PCR analysis showed that indeed *CquiOR114b* was expressed in antennae but not in the maxillary palps, whereas *CquiOR118b* was expressed in the maxillary palps but not in antennae (Figure 2). Next, we used the *Xenopus* oocyte recording system to compare the responses of the newly cloned ORs and their isoforms.

Chiral discrimination by *CquiOR118b*

Initially, the responses of oocytes co-expressing one of the isoform of *CquiOR118* and the obligatory co-receptor *CquiOrco* were compared. Since there were no significant difference in the responses elicited by Ile-163-*CquiOR118b* and Val-163-*CquiOR118b*, we used only colony 1 (GenBank, KT022418) in subsequent analysis. Next, we challenged *CquiOR118b*-*CquiOrco*-expressing oocytes with enantiomers of 1-octen-3-ol and C8 analogs, namely, 1-octyn-3-ol and 3-octanol. While robust responses were elicited by (*R*)-1-octen-3-ol in a dose-dependent manner, currents generated by its antipode, (*S*)-1-octen-3-ol, were relatively very small (Figure 3).




Figure 2. qPCR analysis of newly cloned receptors. Data show high transcription levels of *CquiOR114b* and *CquiOR118b* in antennae and maxillary palps, respectively.




Figure 3. Concentration-response relationships for *CquiOR118b*. *CquiOR118b*-*CquiOrco*-expressing oocytes were challenged with enantiomers of 1-octen-3-ol, 1-octyn-3-ol, and 3-octanol at 0.01, 0.1, and 1 μ M doses. (N = 3)

The remarkable ability of CquiOR118b expressed in a heterologous system to discriminate enantiomers of 1-octen-3-ol is in line with the observations with the intact olfactory system⁵. Likewise, CquiOR118b showed dramatic enantioselectivity towards (*R*)- as compared to (*S*)-1-octyn-3-ol. The receptor showed reduced selectivity towards the saturated analog, 3-octanol. Although at higher doses it preferred (*S*)-3-octanol, the responses to (*R*)-3-octanol were relatively high. It is worth mentioning that the hydroxyl group in the (*S*)-enantiomer of the saturated analog has the orientation as in the (*R*)-isomers of the unsaturated counterparts (Figure 1), their nomenclature differing (*S* vs. *R*) because of the IUPAC rules, not the orientation of the polar moiety expected to fit in the binding cavity of the receptor. Our findings suggest that with CquiOR118b per se the mosquito olfactory system is unlikely to be able to detect the behaviorally relevant ratio of the isomers of 1-octen-3-ol, i.e., R/S, 84:16⁷. It is, therefore, likely that 1-octen-3-ol is also detected by other receptor(s).

Random reception by CquiOR114b

We first compared the two isoforms of CquiOR114b by challenging with 1-octen-3-ol, 1-octyn-3-ol, and 3-octanol oocytes expressing each isoform, CquiOR114b-1 (GenBank, KT022419) or CquiOR114b-2 (GenBank, KT022420) along with CquiOrco (Figure 4). Traces comparing these ligands at three different doses were almost indistinguishable, with the responses recorded from CquiOR114b-1-CquiOrco-expressing oocytes being slightly higher than those from CquiOR114b-2. We, therefore, used CquiOR114b-1 to obtain dose-dependent curves.

CquiOR114b-CquiOrco-expressing oocytes gave robust responses to 3-octanol, with responses to the (*R*)- and (*S*)-stereoisomers being almost indistinguishable (Figure 5). Likewise CquiOR114b responded to the unsaturated compounds, with a slightly preference for (*S*)-isoforms. Of notice, currents elicited by (*R*)-1-octen-3-ol were significantly lower than those obtained with its antipode, (*S*)-1-octen-3-ol, particularly at 0.1 mM (Figure 5). It is, therefore, likely that this antennal receptor contributes to the overall reception of (*S*)-enantiomers of unsaturated C8 alcohols.




Figure 4. Traces obtained with oocytes co-expressing CquiOR114b-1 and CquiOrco (top trace) and CquiOR114b-2 and CquiOrco (lower trace). Compounds were delivered in the following order: 1-octyn-3-ol, 1-octen-3-ol, and 3-octanol from 1 to 10 μ M (left to right).




Figure 5. Concentration-response relationships for CquiOR114b. CquiOR118b-CquiOrco-expressing oocytes were challenged with enantiomers of 1-octen-3-ol, 1-octyn-3-ol, and 3-octanol at 1, 10, and 100 μ M doses. (N = 3)

Behavioral responses

Previously, Cook and collaborators observed an intriguing reduced relative attraction response elicited by (*R*)-1-octen-3-ol in Y-tube olfactometer, but they were unable to conclude if the effect was true repellency as the design of their arena did not allow repellency measurement. With a recently designed surface landing and feeding assay¹⁰, we tested the hypothesis that 1-octen-3-ol is a repellent. Although at very low concentrations of racemic 1-octen-3-ol (0.01 and 0.1%) (Figure 6) mosquitoes were attracted to both sides of the arena, at higher doses (1 and 10%) they were repelled by 1-octen-3-ol. Next, we compared repellency elicited by enantiomers and racemic 1-octen-3-ol. Surprisingly, both (*R*)- and (*S*)-1-octen-3-ol were repellent at the 1% dose (Figure 7). We then surmised on the basis of dose dependence curves obtained with CquiOR118b (Figure 3) that other odorant receptor(s) must mediate repellency elicited by (*S*)-1-octen-3-ol, possible candidates being CquiOR114b, which we identified from antennae and the recently reported CquiOR113 from maxillary palps¹².

We then attempted to combine surgery with behavioral measurement to determine if the maxillary palps are the only olfactory tissues involved in reception of this repellent. Mosquitoes with ablated antennae show little or no flight activity. It might be that impairing a significant component of the olfactory system, and possibly hygroscopic and thermal detectors, may render mosquitoes completely inactive. By contrast, ablating one or two of the maxillary palps had little effect on mosquito activity. Interestingly, mosquitoes with single or double ablated maxillary palps were still repelled by 1-octen-3-ol (Figure 8). We, therefore, concluded that the maxillary palps are not sufficient for repellency by 1-octen-3-ol. Other appendages, most likely antennae, are involved in the reception of this repellent.




Figure 6. Female *Cx. quinquefasciatus* are repelled by 1-octen-3-ol. In the surface-landing and feeding assay, females of the southern house mosquito were significantly repelled by racemic 1-octen-3-ol at 1 and 10% doses, but not at lower doses. Filled bars represent control.




Figure 7. The southern house mosquito is repelled by both enantiomers of 1-octen-3-ol. Female mosquitoes were repelled not only by racemic but also enantiopure isomers, (*R*)- and (*S*)-1-octen-3-ol at 1% dose. Filled bars represent control.




Figure 8. Mosquito with ablated maxillary palps are repelled by 1-octen-3-ol. The effect of surgery on response of female *Culex* mosquitoes to 1-octen-3-ol was minimal given that mosquitoes with one or two maxillary palps ablated were repelled by 1-octen-3-ol. Filled bars represent control.

Conclusion

We have isolated and cloned two odorant receptors from the southern house mosquito sensitive to 1-octen-3-ol and related compounds. CquiOR118b, which is expressed in the maxillary palps, showed remarkable selective and sensitivity towards (*R*)-1-octen-3-ol and the related alkyne, (*R*)-1-octynyl-3-ol. To a much lower extent, CquiOR118b-CquiOrco-expressing oocytes discriminated enantiomers of 3-octanol. By contrast, antennal CquiOR114b responded equally to enantiomers of 3-octanol and showed preference for (*S*)-isomers of 1-octen-3-ol and 1-octyn-3-ol. Repellency assays showed that both isomers of 1-octen-3-ol, a known attractant for *Anopheles* and *Aedes* mosquitoes, were indeed repellents to *Cx. quinquefasciatus*. However, the maxillary palps alone are not enough for detection of this repellent.

Data availability

F1000Research: Dataset 1. Raw data, [10.5256/f1000research.6646.d49878](https://doi.org/10.5256/f1000research.6646.d49878)¹³

Author contributions

PX and WSL designed the experiments. PX, FZ, and GKB carried out the research. PX, FZ, GKB, and WSL analyzed the data.

WL wrote the manuscript. All authors have agreed to the final content of the manuscript.

Competing interests

No competing interests were disclosed.

Grant information

This work was supported in part by the National Institute of Allergy and Infectious Diseases of the National Institutes of Health under award R01AI095514. FZ sabbatical leave at UC Davis was supported in part by the Chinese Scholarship Council.

I confirm that the funders had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript.

Acknowledgements

We thank Dr. Anthon Cornel (University of California, Department of Entomology & Nematology) for providing mosquitoes that allowed us to duplicate his colony at the Davis campus, Dr. Nik Nikbakht, for maintaining the Davis colony, supplying mosquitoes for behavioral assays, and commenting on an earlier version of the manuscript, Dr. Robert Bedoukian (Bedoukian Research Inc.) for providing chemicals used in this research.

References

- Murahashi S: **About the fragrances of matsutake.** *Sci Pap Inst Phys Chem Res.* 1936; **30**: 263–71.
- Hall DR, Beevor PS, Cork A, *et al.*: **1-Octen-3-ol. A potent olfactory stimulant and attractant for tsetse isolated from cattle odours.** *Insect Sci Applic.* 1984; **5**(5): 335–9. [PubMed Abstract](#) [Publisher Full Text](#)
- Takken W, Kline DL: **Carbon dioxide and 1-octen-3-ol as mosquito attractants.** *J Am Mosq Control Assoc.* 1989; **5**(3): 311–6. [PubMed Abstract](#)
- Kline DL, Allan SA, Bernier UR, *et al.*: **Evaluation of the enantiomers of 1-octen-3-ol and 1-octyn-3-ol as attractants for mosquitoes associated with a freshwater swamp in Florida, U.S.A.** *Med Vet Entomol.* 2007; **21**(4): 323–31. [PubMed Abstract](#) [Publisher Full Text](#)
- Syed Z, Leal WS: **Maxillary palps are broad spectrum odorant detectors in *Culex quinquefasciatus*.** *Chem Senses.* 2007; **32**(8): 727–38. [PubMed Abstract](#) [Publisher Full Text](#)
- Lu T, Qiu YT, Wang G, *et al.*: **Odor coding in the maxillary palp of the malaria vector mosquito *Anopheles gambiae*.** *Curr Biol.* 2007; **17**(18): 1533–44. [PubMed Abstract](#) [Publisher Full Text](#) [Free Full Text](#)
- Cook JI, Majeed S, Ignell R, *et al.*: **Enantiomeric selectivity in behavioural and electrophysiological responses of *Aedes aegypti* and *Culex quinquefasciatus* mosquitoes.** *Bull Entomol Res.* 2011; **101**(5): 541–50. [PubMed Abstract](#) [Publisher Full Text](#)
- Grant AJ, Dickens JC: **Functional characterization of the octenol receptor neuron on the maxillary palps of the yellow fever mosquito, *Aedes aegypti*.** *PLoS One.* 2011; **6**(6): e21785. [PubMed Abstract](#) [Publisher Full Text](#) [Free Full Text](#)
- Bohbot JD, Dickens JC: **Characterization of an enantioselective odorant receptor in the yellow fever mosquito *Aedes aegypti*.** *PLoS One.* 2009; **4**(9): e7032. [PubMed Abstract](#) [Publisher Full Text](#) [Free Full Text](#)
- Xu P, Choo YM, De La Rosa A, *et al.*: **Mosquito odorant receptor for DEET and methyl jasmonate.** *Proc Natl Acad Sci U S A.* 2014; **111**(46): 16592–7. [PubMed Abstract](#) [Publisher Full Text](#) [Free Full Text](#)
- Leal WS, Choo YM, Xu P, *et al.*: **Differential expression of olfactory genes in the southern house mosquito and insights into unique odorant receptor gene isoforms.** *Proc Natl Acad Sci U S A.* 2013; **110**(46): 18704–9. [PubMed Abstract](#) [Publisher Full Text](#) [Free Full Text](#)
- Hill SR, Majeed S, Ignell R: **Molecular basis for odorant receptor tuning: a short C-terminal sequence is necessary and sufficient for selectivity of mosquito Or8.** *Insect Mol Biol.* 2015. [PubMed Abstract](#) [Publisher Full Text](#)
- Leal W, Xu P, Zhu F, *et al.*: **Dataset 1 in: 1-Octen-3-ol – the attractant that repels.** *F1000Research.* 2015. [Data Source](#)

Open Peer Review

Current Referee Status:



Version 1

Referee Report 02 July 2015

doi:10.5256/f1000research.7140.r9111



Kenneth F. Haynes

Department of Entomology, University of Kentucky, Lexington, KY, USA

This is a first report that connects olfactory receptors to 1-octen-3-ol in a *Culex* mosquito to behavioral response. In addition, both enantiomers were shown to be repellents. Interestingly the response in the *Culex quinquefasciatus* contrasts with that seen in *Anopheles* and *Aedes* mosquitoes. The adaptive explanation for the contrast between species remains to be explained. This paper reports many important steps towards understanding the mechanisms underlying the behavioral response.

I have read this submission. I believe that I have an appropriate level of expertise to confirm that it is of an acceptable scientific standard.

Competing Interests: No competing interests were disclosed.

Referee Report 02 July 2015

doi:10.5256/f1000research.7140.r9110



Wynand van der Goes van Naters

School of Biosciences, Cardiff University, Cardiff, UK

Xu *et al.* report experiments on the reception and behavioral response in the southern house mosquito *Culex quinquefasciatus* to enantiomers and analogs of 1-octen-3-ol. Specifically, the authors find that the maxillary palp receptor Or118b is especially sensitive to the R(-) enantiomer while the antennal receptor Or114b mediates stronger responses to the S(+) than to the R(-) enantiomer. The authors show that both enantiomers repel female *Cx. quinquefasciatus*, as does a racemic mixture. Females in which the maxillary palps have been ablated are similarly repelled by racemic 1-octen-3-ol, proving that the maxillary palps are not solely responsible for the response to this chemical. The paper makes an important contribution within the context of the literature on insect reception of 1-octen-3-ol, which is an attractive kairomone for several haematophagous insects. Experiments are compelling and the conclusions follow from the data. I have only a few minor suggestions:

1. Could the authors please provide more detail on the stimulus method in the oocyte recordings? How long was each stimulus pulse? Black lines above traces sometimes indicate the timing of the stimulus pulse, but this is not the case in Figure 4.

2. Legend of Figure 4: please check the concentration range (to 100 micromolar).
3. Legend of Figure 5, below the title of the legend, starts with "CquiOR118..." instead of "CquiOR114..."
4. Please indicate the meaning of the error bars in the figures (standard deviation or SEM or CI?).
5. While the reader can infer that Or118 was identified by BLAST as an ortholog of Or8 from *An. gambiae*, it is not obvious how Or114 was identified. Could the authors perhaps include a paragraph on the bioinformatics that underlies this work?
6. It is interesting that 1-octen-3-ol repels this mosquito species, but is attractive to several other species. Is there a possibility that 1-octen-3-ol could be attractive for *Culex quinquefasciatus* when combined with other chemicals?

I have read this submission. I believe that I have an appropriate level of expertise to confirm that it is of an acceptable scientific standard.

Competing Interests: No competing interests were disclosed.

Referee Report 01 July 2015

doi:10.5256/f1000research.7140.r9109



Jeffery K. Tomberlin

Department of Entomology, Texas A&M University, College Station, TX, USA

I very much enjoyed reading this article. And, I believe it makes a significant positive contribution to our understanding of mosquito responses to volatiles associated with hosts. I believe the title and abstract are appropriate for the publication.

With regards to the experiment design, were the trials conducted on the same day from the same population of mosquitoes (could there be a generation effect)? Would the authors consider using logistic regression to analyze the data? Such an approach would determine if there is an interaction between dose, response, and trial?

Furthermore, was there much movement between the treatments (i.e., were mosquitoes flying from one treatment to the next)? If so, how did these responses vary across concentrations? These data would determine if there was increased activity as a response to the treatment.

Another interesting aspect of the study would be to look at the raw data in conjunction with percent response as these data would also lend towards appreciating the compound "exciting" the mosquitoes.

The discussion and conclusions are well developed. One aspect that would be important to consider is the relationship between the volatile and its source (most likely a fungus) as this compound operates similarly to quorum sensing molecules (concentration of the compound dictates "behavior" of the microbe). By building a bridge between the role of the compound with its source, one would be able to tie together the ecology of the mosquito with the source (i.e., microbe) and host health. Such an approach could provide greater explanation as to why certain doses are attractant while others are repellent.

I have read this submission. I believe that I have an appropriate level of expertise to confirm that it is of an acceptable scientific standard.

Competing Interests: No competing interests were disclosed.

Referee Report 29 June 2015

doi:[10.5256/f1000research.7140.r9112](https://doi.org/10.5256/f1000research.7140.r9112)



Joseph Dickens

Invasive Insect Biocontrol and Behaviour Laboratory, United States Department of Agriculture, Beltsville, MD, USA

The authors detail studies aimed at understanding the olfactory detection of 1-octen-3-ol in the southern house mosquito, *Culex quinquefasciatus*. Behavioral studies showed mosquitoes to be repelled by high concentrations of 1-octen-3-ol. An odorant receptor (OR) identified in the maxillary palps responded selectively to (*R*)-1-octen-3-ol in a manner similar to ORs in other mosquitoes. However, when the palps were ablated, the repellent effects of octenol at high concentrations remained. The authors then demonstrated a second OR expressed primarily in antennae that responded to high concentrations of (*S*)-1-octen-3-ol. The involvement of this antennal OR in the repellency of octenol at high concentrations is postulated. This is an intriguing paper that expands our knowledge octenol reception in the southern house mosquito. Since the antennal OR postulated for octenol reception is activated only at relatively high concentrations (10^{-5} M and above), it would be interesting to discover its natural ligand.

I have read this submission. I believe that I have an appropriate level of expertise to confirm that it is of an acceptable scientific standard.

Competing Interests: No competing interests were disclosed.
