

UCLA

Department of Statistics Papers

Title

Consistent Parametric Estimation of the Intensity of a Spatial-temporal Point Process

Permalink

<https://escholarship.org/uc/item/6584c641>

Author

Schoenberg, Frederic P.

Publication Date

2003

**Consistent parametric estimation of the intensity
of a spatial-temporal point process.**

Frederic Paik Schoenberg

Department of Statistics, University of California, Los Angeles, CA 90095–1554, USA.

phone: 310-794-5193

fax: 310-472-3984

email: frederic@stat.ucla.edu

Postal address: UCLA Dept. of Statistics

8142 Math-Science Building

Los Angeles, CA 90095–1554, USA.

Abstract

We consider conditions under which parametric estimates of the intensity of a spatial-temporal point process are consistent. Although the actual point process being estimated may not be Poisson, an estimate involving maximizing a function that corresponds exactly to the log-likelihood if the process is Poisson is consistent under certain simple conditions. A second estimate based on weighted least squares is also shown to be consistent under quite similar assumptions. The conditions for consistency are simple and easily verified, and examples are provided to illustrate the extent to which consistent estimation may be achieved. An important special case is when the point processes being estimated are in fact Poisson, though other important examples are explored as well.

Key words: maximum likelihood estimation, weighted least squares estimation, consistency, Poisson process, conditional intensity.

1 Introduction.

Maximum likelihood estimates (MLEs) have been extensively used in point process inference for decades, at least partly because of their known asymptotic properties. The consistency and asymptotic normality of the maximum likelihood estimate of the intensity of a stationary point process on the line are elementary (see e.g. Cox and Lewis 1966), and for the parameters governing the conditional intensity of an arbitrary stationary point process on the line, the consistency, asymptotic normality and efficiency of the MLE were proven by Ogata (1978).

There have since been a host of similar proofs, generalizing the important results in

Ogata (1978) to more general point processes under various conditions, of which we name a few. The case of non-stationary Poisson processes on the line was investigated by Kutoyants (1984) and more recently by Helmers and Zitikis (1999), and nice summaries of results for general non-stationary point processes on the line were given by Karr (1986) and Andersen et al. (1993). Regarding higher-dimensional point processes, conditions for the consistency and asymptotic normality of the MLE were derived by Brillinger (1975) for stationary multivariate Poisson processes, by Rathbun and Cressie (1994) for the case of non-stationary Poisson processes in \mathbf{R}^d , by Krickeberg (1982) for such processes in locally compact Hausdorff spaces, by Nishayama (1995) for a class of sequential marked point processes, and by Rathbun (1996a) for non-stationary spatial-temporal point processes. Jensen (1993) derived the asymptotic normality of the MLE for spatial Gibbs point processes under conditions similar to those in Rathbun (1996a), and Rathbun (1996b) established conditions for consistent estimation in the case of a spatial modulated Poisson process with partially observed covariates. Using simulations, Huang and Ogata (1999) assessed the relative efficiency of the MLE, the maximum pseudo-likelihood estimator and an approximate maximum likelihood estimator for spatial processes with strong interactions.

The results above are very important since point processes are commonly modeled via their conditional intensities, with parameters estimated by maximum likelihood. However, in some cases one may wish to estimate the unconditional intensity or mean measure, i.e. the expected value of the conditional intensity, assuming it exists. (Hereafter we refer to the unconditional intensity simply as the *intensity*). The present paper explores the problem of parametric estimation of the intensity of an arbitrary simple spatial-temporal point process,

keeping assumptions about the point process and its conditional intensity to a minimum.

Estimating point process intensities may be important in applications, for several reasons. First, whereas the conditional intensity uniquely characterizes the finite-dimensional distributions of any simple point process (see e.g. Daley and Vere-Jones, 1988), the intensity uniquely determines the mean number of points such a process has in any measurable subset of its domain. Hence accurate estimation of the intensity is critical in cases where the mean behavior of a point process is of interest. Second, in many cases the parametric form of the intensity may be more readily suggested than that of the conditional intensity. For example, a functional form for the intensity may be inferred by examining nonparametric intensity estimates, such as those produced by smoothing the point process using kernels, splines, or wavelets (see e.g. Vere-Jones 1992, Brillinger 1998). Third, the assumptions required for the proofs listed above of the consistency of the MLE for the conditional intensity of a point process are unfortunately quite stringent, involving multiple restrictions on the derivatives of the conditional intensity. These conditions can be extremely difficult to verify in applications. For the simpler case of estimating the intensity some results are obtainable under more general and much simpler assumptions which can readily be verified in applications. Under such conditions the consistency of the Poisson maximum likelihood estimate (PMLE), defined as the MLE of the intensity if the process were Poisson, can be demonstrated. Only a slight variant of these conditions is needed to establish the consistency of the weighted least squares estimator (WLSE) as well. The simplicity of these assumptions may greatly facilitate an analysis of when the PMLE and WLSE are consistent, and equally importantly, when they are not. Note that in the case where the point process being estimated is Poisson,

the intensity and conditional intensity are the same, as are the PMLE and MLE; hence for this situation our results represent a proof of the consistency of the MLE under conditions that are easily verifiable, without restrictions on the derivatives of the intensity function.

The structure of this paper is as follows. After formally introducing the PMLE in Section 2, Section 3 summarizes previous results on the MLE and then gives simpler conditions and a simple proof of the consistency of the PMLE. Section 4 provides similar conditions for the consistency of the WLSE. Several examples and counterexamples are given in Section 5 to demonstrate the need for the conditions in the previous Sections and to clarify under what conditions consistent estimation of the intensity is achievable, and Section 6 summarizes the results as well as directions for future research.

2 Preliminaries

Following Brémaud (1981), we consider a spatial-temporal point process to be a measurable mapping from a filtered probability space $(\Omega, \mathcal{F}_t, P)$ onto Φ , the collection of all boundedly finite counting measures on the spatial-temporal domain $\mathcal{S} \times [0, \infty)$. The filtration \mathcal{F}_t is assumed to be increasing and right continuous, and the spatial domain \mathcal{S} any measurable space equipped with measure $\mu_{\mathcal{S}}$ defined on the Borel subsets of \mathcal{S} . Let \mathcal{B} denote the Borel subsets of space-time $\mathcal{S} \times [0, \infty)$. For any spatial-temporal subset $B \in \mathcal{B}$ the random variable $N(B)$ represents the number of points in B .

Suppose N has first moment measure M with $M(B) := E[N(B)] < \infty$ for any $B \in \mathcal{B}$. Then the \mathcal{F} -compensator A of N may be defined as the unique \mathcal{F} -predictable process such that $N - A$ is a local \mathcal{F} -martingale. The existence and uniqueness of the compensator were

shown by Jacod (1975). If A is deterministic (and hence equal to M), then N is a Poisson process.

Let $\mu_{\mathbf{R}}$ denote Lebesgue measure on the real (time-) line, and let $\mu_{\mathcal{B}}$ denote the product measure $\mu_{\mathcal{S}} \times \mu_{\mathbf{R}}$ on space-time. If there exists an integrable, non-negative, real-valued, \mathcal{F} -predictable process λ such that, with probability 1, for all $B \in \mathcal{B}$,

$$\int_B \lambda^*(s, t) d\mu_{\mathcal{B}}(s, t) = A(B),$$

then λ^* is called an \mathcal{F} -conditional intensity of N . Let the intensity $\lambda(s, t)$ denote the expectation with respect to P of $\lambda^*(s, t)$, provided it exists. Hence λ^* is the Radon-Nikodym derivative of the compensator A , and λ the derivative of the first moment measure M .

In what follows we consider sequences of point processes, $\{N_T\}$, $T = 1, 2, \dots$, where only the points of N_T occurring from time 0 to time T , over all of \mathcal{S} , may be observed. We assume throughout that each process N_T has a conditional intensity λ_T^* whose expectation λ_T exists and is known up to a fixed parameter vector θ , within a complete separable metric space Θ of possibilities.

The (partial) log-likelihood function for N_T is conventionally expressed in terms of the conditional intensity λ_T^* as:

$$\int_{\mathcal{S}} \int_0^T \log \lambda_T^*(s, t) dN_T(s, t) - \int_{\mathcal{S}} \int_0^T \lambda_T^*(s, t) d\mu_{\mathcal{B}}(s, t).$$

When a functional form for λ_T^* is known, the parameters governing λ_T^* are typically estimated using the MLE, i.e. the value of the parameters maximizing the log-likelihood function above.

If N_T is a Poisson process, then λ_T^* and λ_T are identical, so in this case the log-likelihood

$L_T(\theta)$ may be written

$$\int_S \int_0^T \log \lambda_T(s, t; \theta) dN_T(s, t) - \int_S \int_0^T \lambda_T(s, t, \theta) d\mu_{\mathcal{B}}(s, t). \quad (1)$$

Hence the estimator $\hat{\theta}$ maximizing (1) may be called the Poisson maximum likelihood estimator (PMLE) of θ . In the next section we examine the case where $\hat{\theta}$ is used to estimate θ even though N may not be Poisson.

3 Asymptotic Properties of the PMLE

As mentioned in the introduction, several authors have proven the consistency and asymptotic normality of the MLE for the parameter vector governing the conditional intensity of a point process. These proofs typically proceed in standard fashion by writing a Taylor expansion of $dL_T(\theta)/d\theta_j$, where $L_T(\theta)$ is the log-likelihood function (1), yielding the approximation

$$\partial L_T(\hat{\theta})/\partial\theta_j \approx \partial L_T(\theta^*)/\partial\theta_j + \sum_{k=1}^K (\theta_k^* - \hat{\theta}_k) I_T^{jk}(\theta^*),$$

where θ_j is the j th coordinate of θ , θ^* is the true parameter vector being estimated, and $I_T^{jk}(\theta) = \partial^2 L_T(\theta)/\partial\theta_j\partial\theta_k$ is the jk element of the Fisher information matrix. The asymptotic results then follow by observing that $\partial L_T(\theta^*)/\partial\theta_j$ are local square integrable martingales and invoking the martingale central limit theorem. (For details see e.g. Theorems VI.1.1 and VI.1.2 of Andersen et al., 1993.) Conditions are required, however, to ensure that the remainder terms in the Taylor approximation are negligible and that the assumptions for the martingale central limit theorem are met. For instance, consider the following conditions

of Rathbun (1996a), for a sequence $\{N_T\}$ of spatial-temporal point process with λ_T and λ_T^* as defined in Section 2, and where the functional form of λ_T^* is known up to a fixed parameter $\theta \in \Theta$:

(A1) Θ is a compact subset of \mathbf{R}^K .

(A2) $\lambda_T^*(u; \theta)$ is continuous as a function of θ , non-negative almost surely almost everywhere in \mathcal{S} , for all $\theta \in \Theta$.

(A3) For each $T = 1, 2, \dots$, the compensator satisfies

$$A_T(\mathcal{S} \times [0, T]; \theta) := \int_{\mathcal{S}} \int_0^T \lambda_T^*(s, t; \theta) d\mu_{\mathcal{B}}(s, t) < \infty$$

with probability one.

(A4) λ_T^* has derivatives $\partial \lambda_T^*(s, t; \theta) / \partial \theta_i$ and $\partial^2 \lambda_T^*(s, t; \theta) / \partial \theta_i \partial \theta_j$ that are continuous functions of θ for all $\theta \in \Theta$, for all $i, j = 1, \dots, K$, and almost all $s \in \mathcal{S}$ and $t \in [0, T]$.

(A5) For all $i, j = 1, \dots, K$,

$$\sup_{\theta \in \Theta} \sup_{t \in [0, T]} E \left[\int_{\mathcal{S}} \frac{(\partial^2 \lambda_T^*(s, t; \theta) / \partial \theta_i \partial \theta_j)^2}{\lambda_T^*(s, t; \theta)} d\mu_{\mathcal{S}}(s) \right] < \infty,$$

and

$$\sup_{\theta \in \Theta} \sup_{t \in [0, T]} E \left[\int_{\mathcal{S}} \frac{(\partial \lambda_T^*(s, t; \theta) / \partial \theta_i \partial \lambda_T^*(s, t; \theta) / \partial \theta_j)^2}{(\lambda_T^*(s, t; \theta))^3} d\mu_{\mathcal{S}}(s) \right] < \infty.$$

(A6) For all $i, j = 1, \dots, K$,

$$\frac{1}{T} \int_{\mathcal{S}} \int_0^T \frac{\partial \lambda_T^*(s, t; \theta) / \partial \theta_i \partial \lambda_T^*(s, t; \theta) / \partial \theta_j}{\lambda_T^*(s, t; \theta)} d\mu_{\mathcal{B}}(s, t) \xrightarrow[T \rightarrow \infty]{p} \sigma_{ij}(\theta)$$

uniformly in θ , and the matrix $\Sigma(\theta)$ whose (i, j) element is $\sigma_{ij}(\theta)$ is positive definite and deterministic.

(A7) For all $i, j = 1, \dots, K$, and all $c > 0$,

$$\sup_{\theta, \theta' \in \Theta} \left\{ \frac{1}{T} \int_{\mathcal{S}} \int_0^T \left| \frac{\partial \lambda_T^*(s, t; \theta)}{\partial \theta_i} \frac{\partial \lambda_T^*(s, t; \theta)}{\partial \theta_j} - \frac{\partial \lambda_T^*(s, t; \theta')}{\partial \theta_i} \frac{\partial \lambda_T^*(s, t; \theta')}{\partial \theta_j} \right| d\mu_{\mathcal{B}}(s, t); \sqrt{T} |\theta' - \theta| \leq c \right\}$$

converges in probability to 0 as $T \rightarrow \infty$, uniformly in θ .

Theorem 3.1. Under conditions (A1-A7), the MLE is consistent and asymptotically normal.

Theorem 3.1 was proven by Rathbun (1996a), using a result of Sweeting (1980). Andersen et al. (1993) consider conditions slightly stronger than those of Rathbun (1996a), including conditions on the third partial derivatives of the conditional intensity. However, even Rathbun's conditions (A1-A7) can be difficult to check in applications, as noted on page 62 of Rathbun (1996a).

The proofs of Rathbun (1996a) and Andersen et al. (1993) for the consistency and asymptotic normality of the MLE for the parameters governing λ^* extend readily to the use of the PMLE as an estimate of the parameters governing λ . However, for this simpler case of estimating the intensity λ , fewer conditions are needed and much simpler machinery is required. Below we present assumptions for the consistency of the PMLE which we believe are considerably simpler and easier to verify than (A1-A7).

Let θ^* denote the true value of the parameter vector being estimated, and let $\hat{\theta}_T$ denote the PMLE of θ^* . Given any value of θ^* , we assume that there exists a function $\phi(t)$ and subsets $\Theta_T \subseteq \Theta$ such that:

$$(B1) P(\hat{\theta}_T \notin \Theta_T) \xrightarrow{T \rightarrow \infty} 0.$$

(B2) Θ_T contains θ^* and admits a finite partition of compact subsets $\Theta_T^1, \dots, \Theta_T^J$ such that

$\lambda_T(s, t; \theta)$ is continuous as a function of θ within each subset Θ_T^j .

(B3) For all θ in Θ_T , $V \left[\int_{\mathcal{S}} \int_0^T \log \lambda_T(s, t; \theta) dN_T(s, t) \right] = o(\phi(T)^2)$.

(B4) Given any neighborhood U of θ^* , there exists $\gamma_1 > 0$ so that for all sufficiently large T , there is a subset of $\mathcal{S} \times [0, T]$ of $\mu_{\mathcal{B}}$ -measure at least $\gamma_1 \phi(T)$ on which $\lambda_T(s, t; \theta^*)$ and $|\log \lambda_T(s, t; \theta^*) - \log \lambda_T(s, t; \theta)|$ are uniformly bounded away from zero, for $\theta \in \Theta_T \setminus U$.

Assumptions (B1) and (B2) control the regularity of Θ . We note in passing that for the case where the parameter space Θ contains only countably many elements, the continuity requirement (B2) is not required for Theorem 3.2 below. Assumption (B3) controls the variance of the process N . Assumption (B4) ensures that any neighborhood U of θ^* provides sufficient restriction on $\lambda_T(s, t; \theta)$. It is demonstrated in Section 5 why these assumptions are needed to establish the consistency of the PMLE.

Theorem 3.2. Under assumptions B1-B4, the PMLE $\hat{\theta}$ is consistent.

Proof.

Consider the value θ^* fixed. We seek to show that $\forall \epsilon > 0$, for any neighborhood U of θ^* , for all sufficiently large T ,

$$P(\hat{\theta}_T \notin U) < \epsilon. \quad (2)$$

Fix $\epsilon > 0$ and U . We first show that there exists $\delta > 0$ such that for sufficiently large T ,

$$EL_T(\theta^*)/\phi(T) - \sup_{\theta \in \Theta_T \setminus U} EL_T(\theta)/\phi(T) \geq \delta, \quad (3)$$

where now $L_T(\theta)$ is defined by (1).

Observe that

$$EL_T(\theta^*) - \sup_{\theta \in \Theta_T \setminus U} EL_T(\theta)$$

$$\begin{aligned}
&= E \int_S \int_0^T \log \lambda_T(s, t; \theta^*) dN_T(s, t) - \int_S \int_0^T \lambda_T(s, t; \theta^*) d\mu_S(s) dt \\
&\quad - \sup_{\theta \in \Theta_T \setminus U} \left\{ E \int_S \int_0^T \log \lambda_T(s, t; \theta) dN_T(s, t) - \int_S \int_0^T \lambda_T(s, t; \theta) d\mu_S(s) dt \right\} \\
&= \int_S \int_0^T \log \lambda_T(s, t; \theta^*) \lambda_T(s, t; \theta^*) d\mu_S(s) dt - \int_S \int_0^T \lambda_T(s, t; \theta^*) d\mu_S(s) dt \\
&\quad - \sup_{\theta \in \Theta_T \setminus U} \left\{ \int_S \int_0^T \log \lambda_T(s, t; \theta) \lambda_T(s, t; \theta^*) d\mu_S(s) dt - \int_S \int_0^T \lambda_T(s, t; \theta) d\mu_S(s) dt \right\} \\
&= \inf_{\theta \in \Theta_T \setminus U} \left\{ \int_S \int_0^T \lambda_T(s, t; \theta^*) \left[\log \lambda_T(s, t; \theta^*) - \log \lambda_T(s, t; \theta) - 1 + \frac{\lambda_T(s, t; \theta)}{\lambda_T(s, t; \theta^*)} \right] d\mu_S(s) dt \right\} \\
&= \inf_{\theta \in \Theta_T \setminus U} \left\{ \int_S \int_0^T \lambda_T(s, t; \theta^*) [\exp(\psi(s, t, \theta)) - \psi(s, t, \theta) - 1] d\mu_S(s) dt \right\},
\end{aligned}$$

where $\psi(s, t, \theta) = \log \lambda_T(s, t; \theta) - \log \lambda_T(s, t; \theta^*)$.

Assumption (B4) ensures that there exists some positive constants $\gamma_1, \gamma_2, \gamma_3$ such that for sufficiently large T , for all t in a subset of $[0, T]$ with measure at least $\gamma_1 \phi(T)$, $\lambda_T(s, t; \theta^*) > \gamma_2$ and $|\psi(s, t, \theta)| > \gamma_3$. Let $\gamma_4 = \min\{\exp(\gamma_3) - \gamma_3 - 1, \exp(\gamma_3 + \gamma_3) - 1\}$. Recalling that $\gamma_3 > 0$ and that the inequality $\exp(x) \geq x + 1$ has equality iff. $x = 0$, (see e.g. Abramowitz 1964), it follows that $\gamma_4 > 0$.

Hence, for sufficiently large T , $EL_T(\theta^*) - \sup_{\theta \in \Theta_T \setminus U} EL_T(\theta) \geq \gamma_1 \gamma_2 \gamma_4 \phi(T)$, which establishes (3) for $\delta = \gamma_1 \gamma_2 \gamma_4$.

With Θ_T defined as in assumption (B1) and $\Theta_T^1, \dots, \Theta_T^J$ as in assumption (B2), fix m elements $\theta^1 \in \Theta_T^1, \dots, \theta^J \in \Theta_T^J$. By assumption (B3), for each such value θ^j ,

$$V \left[\frac{L_T(\theta^j)}{\phi(T)} \right] = V \left[\frac{1}{\phi(T)} \int_S \int_0^T \log \lambda_T(s, t; \theta^j) dN_T \right] \rightarrow 0.$$

Hence, for each θ^j , since each of the following terms obviously has mean zero and has variance

converging to zero,

$$\frac{L_T(\theta^j) - EL_T(\theta^j)}{\phi(T)} \xrightarrow[T \rightarrow \infty]{p} 0 \quad (4)$$

by Chebyshev's inequality.

Since by assumption (B2) the function $\lambda_T(s, t; \theta)$ is continuous with respect to θ on Θ_T^j , so is the function

$$\frac{L_T(\theta) - EL_T(\theta)}{\phi(T)} = \frac{\int_S^T \int_0^T \log \lambda_T(s, t; \theta) dN_T - \int_S^T \int_0^T \log \lambda_T(s, t; \theta) \lambda_T(s, t; \theta^*) d\mu_S(s) dt}{\phi(T)}.$$

Thus the compactness of Θ_T^j implies that $[L_T(\theta) - EL_T(\theta)]/\phi(T) \xrightarrow[T \rightarrow \infty]{p} 0$ uniformly on Θ_T^j . Since $\Theta_T = \bigcup_{j=1}^J \Theta_T^j$, $[L_T(\theta) - EL_T(\theta)]/\phi(T) \xrightarrow[T \rightarrow \infty]{p} 0$ uniformly on all of Θ_T .

Hence there is a $\delta > 0$ such that for sufficiently large T ,

$$P \left(\sup_{\theta \in \Theta_T} [L_T(\theta) - EL_T(\theta)]/\phi(T) \geq \delta/2 \right) < \epsilon/3 \quad (5)$$

and

$$P(\hat{\theta} \notin \Theta_T) < \epsilon/3. \quad (6)$$

Let $\check{\theta}_T$ denote a (possibly non-unique) value of θ maximizing $L_T(\theta)$ among $\check{\theta}_T \in \Theta_T \setminus U$, i.e. $L_T(\check{\theta}_T) \geq L_T(\theta), \forall \theta \in \Theta_T \setminus U$. Putting together (3), (5), and (6) yields, for sufficiently large T ,

$$\begin{aligned} P(\hat{\theta}_T \notin U) &\leq P(\hat{\theta}_T \notin \Theta_T) + P(L_T(\check{\theta}_T) \geq L_T(\theta^*)) \\ &\leq P(\hat{\theta}_T \notin \Theta_T) + P(L_T(\check{\theta}_T) - EL_T(\check{\theta}_T) \geq \delta\phi(T)/2) \\ &\quad + P(EL_T(\check{\theta}_T) - EL_T(\theta^*) > -\delta\phi(T)) + P(EL_T(\theta^*) - L_T(\theta^*) \geq \delta\phi(T)/2) \\ &< \epsilon/3 + \epsilon/3 + 0 + \epsilon/3, \end{aligned}$$

establishing (2).

Assumptions (B1-B4) are by no means minimal, but they are quite straightforward to verify, in contrast to the conditions in previous results regarding maximum likelihood estimation. In particular, no conditions on the derivatives of λ are required. In addition, the replacement of assumption (A1) by (B1) is quite simple and scarcely affects the proof, yet this feature of Theorem 3.2 may be quite attractive since typically in applications the domain for each estimated parameter is, a priori, the whole real line \mathbf{R} or in some cases the half-line \mathbf{R}^+ , rather than some compact subset thereof. The assumption (A1) of compactness of the parameter space is nevertheless commonly assumed in the results of previous authors on the asymptotic properties of the MLE for point processes. Note that while this assumption is stated explicitly by Ogata (1978) and by Rathbun and Cressie (1994), in other works (e.g. Rathbun 1996a) this condition is not explicitly listed in the assumptions but instead appears in the statement of the theorem or in the definition of consistency.

Remark 3.3. Note that assumption (B4) is quite a bit stronger than what is minimally necessary for Theorem 3.2. (B4) is only used in the proof of relation (2) and hence may be discarded for processes where this inequality can be proven directly.

4 Weighted Least Squares Estimates

The parameters θ^* governing the intensity of a spatial-temporal point process can alternatively be estimated by weighted least squares (WLS). Here the estimator $\tilde{\theta}_T$ is chosen to

minimize the quadratic variation:

$$Q_T(\theta) = \sum_{i=1}^{I_T} w_i^T \left[N_T(B_i^T) - E\{N_T(B_i^T); \theta\} \right]^2, \quad (7)$$

where for given T , the sets $\{B_1^T, \dots, B_{I_T}^T\}$ form a partition of the product space $\mathcal{S} \times [0, T]$, and the weights w_i^T are non-negative constants. Here $E\{N_T(B_i^T); \theta\} = \int_{B_i^T} \lambda(s, t; \theta) d\mu_{\mathcal{S}}(s) dt$; with this notation, $EN_T(B_i^T) = E\{N_T(B_i^T); \theta^*\}$. For simplicity, assume that for each T , the number of bins I_T in the partition is finite.

We consider the following replacements for assumptions (B3-B4):

(C3) For all θ in Θ_T , $\max_i V \left[\int_{B_i^T} dN_T \right] = o(\phi(T)^2)$.

(C4) Given any neighborhood U of θ^* , there exist constants $\nu_1, \nu_2, \nu_3 > 0$ so that for sufficiently large T , a fraction of at least $\nu_1 \phi(T)$ of the bins B_i^T have product measure $\mu_{\mathcal{B}}(B_i^T)$ at least $\nu_2 / \sqrt{w_i^T}$ and the property that either $\lambda_T(s, t; \theta) - \lambda_T(s, t; \theta^*) > \nu_3$ or $\lambda_T(s, t; \theta) - \lambda_T(s, t; \theta^*) < -\nu_3$ for all $s, t \in B_i^T$ and all $\theta \in \Theta_T \setminus U$.

Assumption (C3) guarantees that the process N is not too volatile. Like (B4), assumption (C4) ensures that outside neighborhoods U of θ^* , λ_T is uniformly bounded away from its value at θ^* within a sufficient fraction of adequately-sized (and adequately-weighted) bins. As with assumptions (B3-B4), these assumptions are relatively easy to verify.

Theorem 4.1.

Under assumptions (B1-B2) and (C3-C4), the WLSE $\tilde{\theta}_T$ is consistent.

Proof.

$$\begin{aligned}
Q_T(\theta) &= \sum_i w_i^T \left[N_T(B_i^T) - E\{N_T(B_i^T); \theta\} \right]^2 \\
&= \sum_i w_i^T \left[N_T(B_i^T)^2 - 2N_T(B_i^T)E\{N_T(B_i^T); \theta\} + (E\{N_T(B_i^T); \theta\})^2 \right].
\end{aligned}$$

Taking expectations yields

$$EQ_T(\theta) = \sum_i w_i^T \left[EN_T(B_i^T)^2 - 2EN_T(B_i^T)E\{N_T(B_i^T); \theta\} + (E\{N_T(B_i^T); \theta\})^2 \right]. \quad (8)$$

Fix θ^* and a neighborhood U around it. Letting $\delta = \nu_1\nu_2^2\nu_3^2 > 0$, from (8) and (C4) one obtains, for sufficiently large T ,

$$\begin{aligned}
&\inf_{\theta \in \Theta_T \setminus U} \frac{1}{I_T\phi(T)} [EQ_T(\theta) - EQ_T(\theta^*)] \\
&= \inf_{\theta \in \Theta_T \setminus U} \frac{1}{I_T\phi(T)} \sum_{i_T} w_{i_T}^T \left[(E\{N_T(B_{i_T}^T); \theta\})^2 - 2EN_T(B_{i_T}^T)E\{N_T(B_{i_T}^T); \theta\} + (EN_T(B_{i_T}^T))^2 \right] \\
&= \inf_{\theta \in \Theta_T \setminus U} \frac{1}{I_T\phi(T)} \sum_{i_T} w_{i_T}^T \left[E\{N_T(B_{i_T}^T); \theta\} - EN_T(B_{i_T}^T) \right]^2 \\
&= \inf_{\theta \in \Theta_T \setminus U} \frac{1}{I_T\phi(T)} \sum_{i_T} w_{i_T}^T \left[\int_{t \in B_{i_T}} (\lambda_T(s, t; \theta) - \lambda_T(s, t; \theta^*)) d\mu_S(s) dt \right]^2 \\
&\geq \frac{1}{I_T\phi(T)} \nu_1\phi(T) I_T[\nu_2\nu_3]^2 \\
&= \delta.
\end{aligned} \quad (9)$$

Note that assumption (C3) implies that $V[Q_T(\theta)] = o(I_T^2\phi(T)^2)$ by the Cauchy-Schwarz inequality. Thus $[Q_T(\theta) - EQ_T(\theta)]/I_T\phi(T) \xrightarrow{p} 0$ for all θ in Θ_T , and assumption (B2) ensures that this convergence is uniform over all θ in Θ_T , just as in Theorem 3.2.

Hence if $\hat{\theta}_T$ denotes the WLSE of θ among $\hat{\theta}_T \in \Theta_T \setminus U$, then for sufficiently large T ,

$$\begin{aligned}
P(\tilde{\theta}_T \notin U) &\leq P(\tilde{\theta}_T \notin \Theta_T) + P(Q_T(\hat{\theta}_T) \geq Q_T(\theta^*)) \\
&\leq P(\tilde{\theta}_T \notin \Theta_T) + P(Q_T(\hat{\theta}_T) - EQ_T(\hat{\theta}_T) \leq -\delta T\phi(T)/2)
\end{aligned}$$

$$\begin{aligned}
& +P\left(EQ_T(\hat{\theta}_T) - EQ_T(\theta^*) < \delta T\phi(T)\right) + P\left(EQ_T(\theta^*) - Q_T(\theta^*) \leq \delta T\phi(T)/2\right) \\
& < \epsilon/3 + \epsilon/3 + 0 + \epsilon/3,
\end{aligned}$$

which completes the proof.

5 Examples and Counterexamples

Some examples may help to clarify when the conditions for consistent estimation are satisfied.

Our first two examples consider the Poisson case, where $\lambda = \lambda^*$ and the PMLE and MLE are equivalent.

Example 5.1. Suppose N_T is a sequence of spatial-temporal versions of the cyclic Poisson process, studied for example by Helmers and Zitikis (1999) and Helmers et al. (2003). That is, suppose N_T is Poisson with separable intensity function

$$\lambda_T(s, t; \theta) = f(s; \theta)g(t; \theta), \quad (10)$$

where $f, g > 0$, $\theta \in \mathbf{R}^K$ for some positive integer K , and g is any integrable cyclic function with (possibly unknown) period τ , i.e. $g(t; \theta) = g(t + j\tau; \theta)$, for all t and any integer j . Let f and g be continuous in θ with $|\log fg|$ bounded for each θ by some constant B_θ , and suppose $\alpha := \int_{\mathcal{S}} f(s; \theta^*) d\mu_{\mathcal{S}}(s) < \infty$ and $\beta := \int_0^\tau g(t) dt < \infty$. Finally, suppose that condition (B4) holds with $\phi(T) = T$; note for example that one only needs $f(s; \theta^*) > c_1 > 0$ for t in some non-null subset of $[0, \tau)$, and $g(t; \theta^*) > c_2 > 0$ for s in some non-null subset of \mathcal{S} , in order to ensure that $\lambda_T(s, t; \theta^*)$ is uniformly bounded away from zero on a subset of $\mu_{\mathcal{B}}$ -measure at least $\gamma_1 T$.

Assumptions (B1) and (B2) are satisfied with $\Theta_T = [-T, T]^K$, and since N_T is Poisson,

$$\begin{aligned}
V_T(\theta) &:= \text{Var} \left[\int_{\mathcal{S}} \int_0^T \log \lambda_T(s, t; \theta) dN_T(s, t) \right] & (11) \\
&= E \left[\int_{\mathcal{S}} \int_0^T \log \lambda_T(s, t; \theta) dN_T(s, t) \right]^2 - \left[E \int_{\mathcal{S}} \int_0^T \log \lambda_T(s, t; \theta) dN_T(s, t) \right]^2 \\
&= \int_{\mathcal{S}} \int_0^T [\log \lambda_T(s, t; \theta)]^2 \lambda_T(s, t; \theta^*) d\mu_{\mathcal{S}}(s) dt + \left[\int_{\mathcal{S}} \int_0^T \log \lambda_T(s, t; \theta) \lambda_T(s, t; \theta^*) d\mu_{\mathcal{S}}(s) dt \right]^2 \\
&\quad - \left[\int_{\mathcal{S}} \int_0^T \log \lambda_T(s, t; \theta) \lambda_T(s, t; \theta^*) d\mu_{\mathcal{S}}(s) dt \right]^2 \\
&= \int_{\mathcal{S}} \int_0^T [\log \lambda_T(s, t; \theta)]^2 \lambda_T(s, t; \theta^*) d\mu_{\mathcal{S}}(s) dt \\
&\leq B_{\theta}^2 \alpha \beta (1 + T/\tau) \\
&= o(T^2),
\end{aligned}$$

so condition (B3) is satisfied with $\phi(T) = T$.

Example 5.2. Let N_T be a sequence of spatial-temporal Poisson processes, but not necessarily cyclical or separable. Suppose $\log \lambda_T(s, t; \theta)$ is continuous in $\theta \in \mathbf{R}^K$ and bounded in absolute value by some constant $B_{\theta} < \infty$, and that the space \mathcal{S} has finite, positive measure $\mu_{\mathcal{S}}(\mathcal{S})$. Finally, suppose that λ_T is parameterized such that for θ outside any neighborhood U of θ^* , $|\log \lambda_T(s, t; \theta^*) - \log \lambda_T(s, t; \theta)|$ is uniformly bounded away from zero.

Then

$$\begin{aligned}
V_T(\theta) &\leq B_{\theta}^2 V \left[\int_{\mathcal{S}} \int_0^T dN_T(s, t) \right] \\
&= B_{\theta}^2 E N_T(\mathcal{S} \times [0, T]) \\
&\leq B_{\theta}^2 \exp(B_{\theta}) T \mu_{\mathcal{S}}(\mathcal{S}),
\end{aligned}$$

so requirement (B3) is fulfilled with $\phi(T) = T$. (B1) and (B2) are similarly satisfied by $\Theta_T = [-T, T]^K$ as in example 5.1, and assumption (B4) is satisfied since $\lambda_T(s, t; \theta^*) > \exp(-B_\theta^*) > 0$ on $\mathcal{S} \times [0, T]$ which has measure $T\mu_{\mathcal{S}}(\mathcal{S})$.

Example 5.3. Suppose N_T are spatial-temporal versions of the Isham and Westcott (1979) self-correcting point process, as described by Rathbun (1996a). Such processes have conditional intensity

$$\lambda_T^*(s, t; \theta) = \exp[\theta_1 + \theta_2(t - \theta_3 N_T \{b(s, r) \times [0, t]\})],$$

where $b(s, r)$ is a ball of radius r around location s , and θ_2 and θ_3 are positive. Such processes are called self-correcting since the further $\theta_3 N_T \{b(s, r) \times [0, t]\}$ happens to deviate from t , the more the conditional intensity λ_T^* adjusts until $\theta_3 N_T \{b(s, r) \times [0, t]\}$ comes back to t . Hence the variance of $N_T \{b(s, r) \times [0, t]\}$ is actually smaller than that of the Poisson process with equivalent intensity, and (B3) and (B4) are easily satisfied with $\phi(T) = T$ as in the previous example, provided $0 < \mu_{\mathcal{S}}(\mathcal{S}) < \infty$.

Example 5.4. For point processes with rapidly increasing intensities, assumption (B4) will typically not be satisfied, but in such cases one may appeal to Remark 2.3. For example, if λ_T increases exponentially, i.e. if λ_T is separable with $g(t) \propto \exp(\theta_K t)$, then (B3) and (2) are satisfied with $\phi(T) = \exp(\theta_K^* T)$.

The next two examples illustrate limitations on the possibilities for consistent estimation of the intensity.

Example 5.5. Suppose N_T are observations on $\mathcal{S} \times [0, T]$ of a *finite* point process,

N , i.e. a process that, with positive probability, may contain only finitely many points on $\mathcal{S} \times [0, \infty)$. In this case consistent estimation of θ is unachievable, as is well known (see for instance Example 6.3 of Rathbun and Cressie 1994). An example is when N is a spatial-temporal Poisson process with separable intensity as in (10), with $g(t; \theta) \propto \exp(\theta_i t)$, where $\theta_i < 0$. One may inquire which of the conditions (B1-B4) are not met in this case. Since the integral $\lim_{T \rightarrow \infty} \int_{\mathcal{S}} \int_0^T \lambda_T(s, t; \theta^*) < \infty$, the set on which λ_T is uniformly bounded away from zero must have finite product measure; hence $\phi(T) = O(1)$ in condition (B4). However, since N is a Poisson process $V_T(\theta)$ is non-decreasing as a function of T , so assumption (B3) is violated: $V_T(\theta)/\phi(T)^2 \not\rightarrow 0$ as $T \rightarrow \infty$.

Example 5.6. If λ_T has a change-point governed by a parameter in θ , then in general consistent estimation of θ is unachievable. For a simple example let N_T be Poisson with

$$\lambda_T(s, t; \theta) = \theta_1 \mathbf{1}_{\{t \leq \theta_2\}} + \theta_3 \mathbf{1}_{\{t > \theta_2\}},$$

and where $0 < \mu_{\mathcal{S}}(\mathcal{S}) < \infty$. Assumption (B4) is violated, since for any neighborhood $U = (\theta_1^* - \epsilon_1, \theta_1^* + \epsilon_1) \times (\theta_2^* - \epsilon_2, \theta_2^* + \epsilon_2) \times (\theta_3^* - \epsilon_3, \theta_3^* + \epsilon_3)$ of θ^* , $\lambda_T(s, t; (\theta_1^*, \theta_2^*, \theta_3^*)) = \lambda_T(s, t; (\theta_1^*, \theta_2^* - \epsilon_2, \theta_3^*))$ for $t > \theta_2^*$. Thus the set of (s, t) on which $|\log \lambda_T(s, t; \theta^*) - \log \lambda_T(s, t; \theta)|$ are uniformly bounded away from zero for $\theta \in \Theta_T \setminus U$ has $\mu_{\mathcal{B}}$ -measure less than $\theta_2^* \mu_{\mathcal{S}}(\mathcal{S}) = O(1)$, which violates (B4) since $V_T(\theta) = O(T)$. In general if λ_T is governed by a parameter θ_i that does not affect $\lambda_T(s, t; \theta)$ on a set of (s, t) with infinite measure, then consistent estimation of θ_i is unachievable, so assumption (B4) is needed to ensure such cases are excluded.

6 Discussion

Although maximum likelihood estimation for point processes is very common, examples of applications where the assumptions necessary to establish the consistency of the MLE are verified are rather elusive. The general impression among applied researchers appears to be that asymptotic properties of the MLE such as consistency and asymptotic normality apply quite generally, and that verification of these properties for particular point process models is difficult and unnecessary. The aim of the current paper is to provide conditions for consistent estimation of point process intensities that may readily be verified in practice. These results are stated for the estimation of the intensity using the PMLE and NLSE; a special case includes estimation of Poisson intensities by MLE.

Our results involve conditions on the rate of increase of the variance of the point process. Since variances are rather easy to check and are easily interpretable, it is possible that these conditions may be checked in applications without explicit assumption of a parametric form for the conditional intensity of the point process, but rather by examining the variance directly, or by appealing to subject matter information or knowledge of the mechanism driving the point process. By contrast, it is difficult to see how the applied researcher could justify assumptions involving the second (and higher-order) partial derivatives of the conditional intensity of the point process without specifying the conditional intensity in detail.

Although our assumptions may be easily verifiable, they are by no means minimal, nor are our results optimally strong. In particular, only consistent estimation is investigated here. Similar conditions under which estimates may be shown to be asymptotically normal

and/or efficient are important subjects for further research.

Acknowledgements

This material is based upon work supported by the National Science Foundation under Grant No. 9978318. Any opinions, findings, and conclusions or recommendations expressed in this material are those of the author and do not necessarily reflect the views of the National Science Foundation.

References

- Abramowitz, M., 1964. Handbook of Mathematical Functions with Formulas, Graphs, and Mathematical Tables. M. Abramowitz and I. Stegun (Eds.), U.S. Government Printing Office, Washington D.C.
- Andersen, P.K., Borgan, O., Gill, R.D., Keiding, N., 1993. Statistical Models Based on Counting Processes. Springer, NY.
- Brémaud, P., 1981. Point Processes and Queues. Springer, NY.
- Brillinger, D.R., 1998. Some wavelet analyses of point process data. In: 31st Asilomar Conference on Signals, Systems and Computers, IEEE Computer Society, Los Alamitos, California, 1087–1091.
- Cox, D.R., Lewis, P.A.W., 1966. The Statistical Analysis of Series of Events. Methuen, London.
- Daley, D., Vere-Jones, D., 1988. An Introduction to the Theory of Point Processes. Springer, NY.
- Helmers, R., Mangku, W., Zitikis, R., 2003. Consistent estimation of cyclic Poisson intensity functions. *J. Multiv. Anal.* 84, 19–39.
- Helmers, R., Zitikis, R., 1999. On estimation of Poisson intensity functions. *Ann. Inst. Statist. Math.* 51, 265–280.
- Huang, F., Ogata, Y., 1999. Improvements of the maximum pseudo-likelihood estimators in various spatial statistical models. *J. Comp. Graph. Statist.* 8, 510–530.

- Isham, V., Westcott, M., 1979. A self-correcting point process. *Stoch. Proc. Appl.* 8, 335–347.
- Jensen, J.L., 1993. Asymptotic normality of estimates in spatial point processes. *Scand. J. Statist.* 20, 97–109.
- Karr, A.F., 1986. *Point Processes and their Statistical Inference*. Marcel Dekker, NY.
- Krickeberg, K., 1982. Processus ponctuels en statistique. In: Hennequin, P.L. (Ed.), *Ecole d'Eté de Probabilités de Saint-Flour X*, International Cooperative Publishing House, Fairland, MD, 205–313.
- Kutoyants, Y.A., 1984. *Parameter Estimation for Stochastic Processes*. Heldermann Verlag, Berlin.
- Nishiyama, Y., 1995. Local asymptotic normality of a sequential model for market point processes and its applications. *Ann. Inst. Statist. Math.* 47, 195–209.
- Ogata, Y., 1978. The asymptotic behaviour of maximum likelihood estimators for stationary point processes. *Ann. Inst. Statist. Math.* 30, 243–261.
- Ogata, Y., 1988, Statistical models for earthquake occurrences and residual analysis for point processes. *J. Amer. Statist. Assoc.* 83, 9–27.
- Ogata, Y., 1998, Space-time point process models for earthquake occurrences. *Ann. Inst. Statist. Math.* 50, 379–402.
- Rathbun, S.L., 1996a. Asymptotic properties of the maximum likelihood estimator for spatio-temporal point processes. *J. Statist. Plan. Inf.* 51, 55–74.

- Rathbun, S.L., 1996b. Estimation of Poisson intensity using partially observed concomitant variables. *Biometrics* 52, 226–242.
- Rathbun, S.L., Cressie, N. 1994. Asymptotic properties of estimators for the parameters of spatial inhomogeneous Poisson point processes. *Adv. Appl. Probab.* 26, 122–154.
- Sweeting, T.J., 1980. Uniform asymptotic normality of the maximum likelihood estimator. *Ann. Statist.* 8, 1375–1381.
- Vere-Jones, D., 1992. Statistical methods for the description and display of earthquake catalogs. In: Walden, A., Guttorp, P. (Eds.), *Statistics in the Environmental and Earth Sciences*, Edward Arnold, London, 220–236.