

UC Irvine

UC Irvine Previously Published Works

Title

Ligation of CD40 in Human Müller Cells Induces P2X7 Receptor-Dependent Death of Retinal Endothelial Cells
CD40, Endothelial and Müller Cells

Permalink

<https://escholarship.org/uc/item/65d369mn>

Journal

Investigative Ophthalmology & Visual Science, 57(14)

ISSN

0146-0404

Authors

Portillo, Jose-Andres C
Corcino, Yalitza Lopez
Dubyak, George R
[et al.](#)

Publication Date

2016-11-17

DOI

10.1167/iovs.16-20301

Peer reviewed

Ligation of CD40 in Human Müller Cells Induces P2X₇ Receptor–Dependent Death of Retinal Endothelial Cells

Jose-Andres C. Portillo,¹ Yalitz Lopez Corcino,^{1,2} George R. Dubyak,³ Timothy S. Kern,^{4–6} Shigemi Matsuyama,^{7,8} and Carlos S. Subauste^{1,2,5}

¹Division of Infectious Diseases and HIV Medicine, Department of Medicine, Case Western Reserve University, Cleveland, Ohio, United States

²Department of Pathology, Case Western Reserve University, Cleveland, Ohio, United States

³Department of Biophysics, Case Western Reserve University, Cleveland, Ohio, United States

⁴Division of Molecular Endocrinology, Department of Medicine, Case Western Reserve University, Cleveland, Ohio, United States

⁵Department of Ophthalmology and Visual Sciences, Case Western Reserve University, Cleveland, Ohio, United States

⁶Veterans Administration Medical Center, Research Service 151, Cleveland, Ohio, United States

⁷Department of Medicine, Case Western Reserve University, Cleveland, Ohio, United States

⁸Case Comprehensive Cancer Center, Case Western Reserve University, Cleveland, Ohio, United States

Correspondence: Carlos S. Subauste, Division of Infectious Diseases and HIV Medicine, Department of Medicine, Case Western Reserve University, Cleveland, OH 44106, USA; carlos.subauste@case.edu.

Submitted: July 10, 2016

Accepted: October 12, 2016

Citation: Portillo J-AC, Lopez Corcino Y, Dubyak GR, Kern TS, Matsuyama S, Subauste CS. Ligation of CD40 in human Müller cells induces P2X₇ receptor–dependent death of retinal endothelial cells. *Invest Ophthalmol Vis Sci.* 2016;57:6278–6286. DOI: 10.1167/iops.16-20301

PURPOSE. Cluster of differentiation 40 (CD40) is required for retinal capillary degeneration in diabetic mice, a process mediated by the retinal endothelial cells (REC) death. However, CD40 activates prosurvival signals in endothelial cells. The purpose of this study was to identify a mechanism by which CD40 triggers programmed cell death (PCD) of RECs and address this paradox.

METHODS. Human RECs and Müller cells were incubated with CD154 and L-N6-(1-Iminoethyl)lysine (L-Nil, nitric oxide synthase 2 inhibitor), α -lipoic acid (inhibitor of oxidative stress), anti-Fas ligand antibody, or A-438079 (P2X₇ adenosine triphosphate [ATP] receptor inhibitor). Programmed cell death was analyzed by fluorescence-activated cell sorting (FACS) or Hoechst/propidium iodide staining. Release of ATP was measured using a luciferase-based assay. Mice were made diabetic with streptozotocin. Expression of P2X₇ was assessed by FACS, quantitative PCR, or immunohistochemistry.

RESULTS. Ligation of CD40 in primary RECs did not induce PCD. In contrast, in the presence of primary CD40⁺ Müller cells, CD40 stimulation caused PCD of RECs that was not impaired by L-Nil, α -lipoic acid, or anti-Fas ligand antibody. We found CD40 did not trigger TNF- α or IL-1 β secretion. Primary Müller cells released extracellular ATP in response to CD40 ligation. Inhibition of P2X₇ (A-438079) impaired PCD of RECs; CD40 upregulated P2X₇ in RECs, making them susceptible to ATP/P2X₇-mediated PCD. Diabetic mice upregulated P2X₇ in the retina and RECs in a CD40-dependent manner.

CONCLUSIONS. Cluster of differentiation 40 induces PCD of RECs through a dual mechanism: ATP release by Müller cells and P2X₇ upregulation in RECs. These findings are likely of in vivo relevance since CD40 upregulates P2X₇ in RECs in diabetic mice and CD40 is known to be required for retinal capillary degeneration.

Keywords: retinal endothelial cells, Müller cells, purinergic

Degenerate capillaries are a hallmark of early diabetic retinopathy.¹ They are the result of loss of retinal endothelial cells (RECs) and pericytes with the ensuing transformation into tubes of basement membrane that lack blood flow. The resulting ischemia likely promotes the transition to the proliferative form of diabetic retinopathy characterized by retinal neovascularization.

The development of capillary degeneration is a slow process.^{1,2} It takes 8 months of diabetes for degenerate capillaries to become apparent in streptozotocin-treated rats.² Many studies revealed that RECs undergo programmed cell death (PCD) in the diabetic retina.^{1–6} The fact that only a small fraction of RECs have detectable evidence of PCD at a given time is in keeping with a low rate of cell loss that will eventually result in capillary dropout.

The pathogenesis of retinal capillary degeneration in diabetes has not been fully elucidated, and various mechanisms may contribute to this phenomenon. The increased oxidative and nitrosative stress in the diabetic retina have been linked to PCD of RECs and capillary degeneration.^{7–11} Tumor necrosis factor (TNF)- α and IL-1 increase activation of caspases in RECs, and the genetic or pharmacologic inhibition of IL-1 or TNF- α signaling diminishes capillary degeneration in diabetic mice.^{3,12–14}

Cluster of differentiation 40 (CD40) is a member of the TNF receptor superfamily that has low-level basal expression in retinal Müller cells, RECs, microglia/macrophage, and ganglion neurons.¹⁵ However, CD40 expression in Müller cells, RECs, and microglia/macrophages is upregulated in the diabetic retina.¹⁵ Moreover, CD40 plays a central role in the develop-

ment of early diabetic retinopathy since diabetic CD40^{-/-} mice are protected from capillary degeneration.¹⁵

Delineation of the role of CD40 in retinal capillary degeneration requires examining whether CD40 stimulation causes death of RECs. However, ligation of CD40 in endothelial cells does not typically induce cell death,^{16–18} likely because CD40 activates PI3K/Akt-mediated prosurvival signals.¹⁷ This raised the possibility that CD40 might promote death of RECs by acting through other retinal cells. We examined Müller cells since they are considered key in the pathogenesis of diabetic retinopathy and these cells encircle RECs. We report that while ligation of CD40 in primary human RECs did not increase PCD, CD40 stimulation promoted PCD of RECs when RECs were incubated with human Müller cells. CD40 triggered ATP release by primary Müller cells, upregulated P2X₇ expression in RECs, and caused P2X₇-dependent PCD of REC.

MATERIALS AND METHODS

Cells

Primary human RECs and Müller cells were obtained as described.¹⁵ Retinal endothelial cells were cultured in gelatin-coated flasks containing DMEM plus 10% FBS (HyClone; GE Healthcare Life Sciences, Logan, UT, USA), endothelial cell growth supplement from bovine pituitary (15 µg/mL; Sigma-Aldrich Corp., St Louis, MO, USA) and insulin/transferrin/selenium (Sigma-Aldrich Corp.). Cell identity was confirmed by incorporation of acetylated low-density lipoprotein (>90%). Primary human Müller cells were cultured in DMEM/F12 containing 20% FBS. Cultures were >95% pure for Müller cells (vimentin⁺, CRALBP⁺, and GFAP⁻ by immunofluorescence). Human retinal cells were used between passages 3 to 6. The human Müller cell line MIO-M1 was a gift from Gloria Limb (University College London, London, UK). In coculture experiments, 8 × 10⁴ RECs and 4 × 10⁴ RMCs were cultured per well of 6-well plates.

In Vitro Stimulation

Cells were treated with multimeric human CD154 (CD40 ligand; gift from Richard Kornbluth, Multimeric Biotherapeutics, Inc., La Jolla, CA, USA).¹⁵ As controls we used omission of CD154 or incubation with a nonfunctional CD154 mutant (T147N). We added L-Nil (500 µM; Enzo Life Sciences, Farmingdale, NY, USA); α-lipoic acid (1 mM; Sigma-Aldrich Corp.); A-438079 (10 µM; Tocris Bioscience, Bristol, UK); neutralizing anti-Fas ligand mAb (15 µg/mL; GeneTex, Irvine, CA, USA); or isotype control mAb to cells 1 hour prior to stimulation with CD154. Endothelial cells were also incubated with Bz-ATP (100 µM; Sigma-Aldrich Corp.) during the last 24 hours of incubation with or without CD154, with staurosporine (100 nM; Sigma-Aldrich Corp.) for 6 hours or IFN-γ (100 U/mL; Peprotech, Rocky Hill, NJ, USA) for 72 hours.

Retroviral Vectors and Transductions

Murine stem cell virus-based bicistronic retroviral vector MIEG3 that encodes enhanced green fluorescent protein (EGFP) with or without human CD40 was previously described.¹⁹ Cells were incubated overnight with retrovirus plus polybrene (8 µg/mL, Sigma-Aldrich Corp.). Cells were used at least 72 hours after infection.

Flow Cytometry

Transduced Müller cells were sorted for EGFP expression using a commercial sorter (FACSARIA; BD Biosciences, San Jose, CA,

USA). Müller cells were >95% EGFP⁺ when used in experiments. Retinal cells were incubated with an apoptosis detection kit and commercial stain (Annexin V-PE and 7-AAD; BD Biosciences). Unless otherwise stated, staining was performed between 72 to 100 hours after in vitro coculture of RECs and Müller cells. Cells were analyzed on a flow cytometer (LSR II; BD Biosciences). Commercial software (FlowJo; Tree Star, Inc., Ashland, OR, USA) was used for analysis. Cells were incubated with anti-human ICAM-1 (eBiosciences, San Diego, CA, USA); anti-human Fas ligand (BD Biosciences); or isotype-matched control mAbs, or with anti-P2X₇ Ab (Alomone Labs, Jerusalem, Israel) plus secondary Ab (eBiosciences). Corrected mean fluorescence intensity was calculated by subtracting mean fluorescence for isotype control or secondary Ab alone from values obtained for specific Abs.

Hoechst 33342 and Propidium Iodide Staining

Retinal endothelial cells were incubated with the blue-fluorescence dye Hoechst 33342 (5 µg/mL) and the red-fluorescence dye propidium iodide (1 µg/mL; both from Molecular Probes; Eugene, OR, USA). Nuclear morphology was examined using a fluorescence microscope (Zeiss Axiovert; Carl Zeiss, Inc., Oberkochen, Germany). Percentages of cells with intense blue fluorescence (nuclear condensation) and cells that stained with propidium iodide (dead cells) were determined.

Measurement of Extracellular ATP

Müller cells were prepared for ATP release assays.²⁰ The ecto-ATPase inhibitor β, γ-methylene-ATP (300 µM; Sigma-Aldrich Corp.) was added 15 minutes prior to stimulation with CD154. Extracellular ATP was measured in supernatants using an ATP bioluminescence assay kit (Sigma-Aldrich Corp.). Luminescence was quantified using a luminometer (TD 20/20; Turner Designs, San Jose, CA, USA).²⁰ Concentrations of ATP were calculated using an ATP standard curve.

Cytokine ELISA

Supernatants were used to measure concentrations of human IL-1β and human TNF-α (eBioscience).

Measurement of Nitrite and Superoxide Generation

Nitrite concentrations were calculated using a Griess reaction (Promega Corp., Madison, WI, USA). Superoxide production was examined using the lucigenin assay.⁸

Induction of Diabetes in Mice

Male C57BL/6J, CD40^{-/-} (N° 002928) and P2X₇^{-/-} (N° 005576) mice backcrossed to C57BL/6J (Jackson Laboratory, Bar Harbor, ME, USA) were rendered diabetic by administration of streptozotocin (STZ). Fasted mice of 20 to 25 g body weight received five daily intraperitoneal injections of STZ (55 mg/kg; MP Biomedicals, Solon, OH, USA). Development of diabetes (blood glucose >250 mg/mL) was assessed beginning at 1 week. Blood glucose, glycated hemoglobin, and body weights were similar in diabetic B6, CD40^{-/-}, and P2X₇^{-/-} mice. Studies adhered to the institutional guidelines for humane treatment of animals, “Principles of laboratory animal care” (NIH) and to the ARVO Statement for the Use of Animals in Ophthalmic and Vision Research.




FIGURE 1. Ligation of CD40 in RECs does not cause PCD. (A, B) Primary human RECs were incubated with or without CD154 followed by staining with Annexin V or 7-AAD. (A) Representative dot plot obtained after 24 hours of incubation with or without CD154. (B) We incubated RECs with CD154 as indicated and percentages of cells that stained with Annexin V or 7-AAD were assessed by flow cytometry. Retinal endothelial cells were also incubated with staurosporine for 6 hours. (C, D) We transduced RECs with an empty retroviral vector (MIEG3) or with a retroviral vector that encodes CD40 (MIEG3-CD40). Cells were incubated with or without CD154. Expression of ICAM-1 (C) and percentages of cells that stained with Annexin V or 7-AAD (D) were assessed by flow cytometry. Results are representative of three independent experiments. ** $P < 0.01$; *** $P < 0.001$.

Real-Time Quantitative PCR

Gene expression was assessed using cDNA, SYBR GREEN PCR Master Mix (Applied Biosystems, Foster City, CA, USA), primers for P2X₇ receptor²¹ or 18S rRNA²² and a commercial PCR system (7300 Real Time PCR System; Applied Biosystems). Each cDNA sample was run in triplicate. Samples were normalized according to the content of 18S rRNA.

Immunohistochemistry

Paraffin-embedded sections were incubated with Tomato Lectin DyLight 488 (Vector Laboratories, Burlingame, CA, USA) and either anti-P2X₇ Ab (Acris, Herford, Germany) or ApopTag Red, In situ Apoptosis Detection kit (EMD Millipore, Billerica, MA, USA). Staining specificity of was confirmed by omitting primary Ab. Retinas were analyzed using an automated microscope (Leica DMI 6000B; Leica Microsystems, Inc., Buffalo Grove, IL, USA) equipped for epifluorescence microscopy.

Statistical Analysis

All results were expressed as the mean \pm SEM. Data were analyzed by 2-tailed Student's *t*-test and ANOVA. Differences were considered significant at $P \leq 0.05$.

RESULTS

Ligation of CD40 in Human REC Does Not Result in PCD

Primary RECs incubated with CD154 (CD40 ligand) did not exhibit an increase in the percentages of cells that were

Annexin V⁺ 7-AAD⁻ (early apoptosis) or Annexin V⁺ 7-AAD⁺ (late apoptosis/necrosis; Figs. 1A, 1B). In contrast, staurosporine increased the percentages of Annexin V⁺ 7-AAD⁻ cells (Fig. 1B). Consistent with studies in primary nonhematopoietic cells, human RECs express low levels of CD40 under basal conditions.¹⁵ Thus, to confirm that ligation of CD40 expressed on RECs did not induce PCD, RECs were transduced with an MIEG3-based retroviral vector that encodes human CD40 to increase the percentage of cells that express CD40. This approach allows examining the functional effects of CD40 stimulation in various primary cells including REC.^{23,24} CD40 was functional since, as reported,²³ CD154 upregulated ICAM-1 (Fig. 1C). We found CD154 did not increase in the percentages of Annexin V⁺ 7-AAD⁻ cells or Annexin V⁺ 7-AAD⁺ cells (Fig. 1D). Culturing RECs in normal glucose (5 mmol/L) or high glucose (30 mmol/L) conditions did not affect the results (data not shown). Taken together, CD40 signaling in RECs did not increase PCD.

CD40 Stimulation Results in PCD of Human RECs When Incubated With Müller Cells

We examined whether CD40 induces death of RECs through an indirect mechanism. Given that Müller cells closely associate with RECs, we examined whether CD40 stimulation affected REC survival when these cells were incubated with Müller cells. Müller cells upregulate CD40 in diabetic mice.¹⁵ However, similar to RECs, Müller cells cultured in vitro have minimal levels of CD40 expression under basal conditions.¹⁵ Thus, to test the effect of CD40 ligation in Müller cells, CD40 expression in Müller cells was induced with a retroviral vector.^{23,24} Human Müller cells were transduced with MIEG3 or MIEG3-CD40 retroviral vectors. Expression of CD40 induced by retrovirus was similar to that induced by culturing cells in




FIGURE 2. Stimulation of CD40 causes PCD of RECs in the presence of CD40⁺ Müller cells. (A) Green fluorescence of untransduced RECs, Müller cells transduced with retroviral vector that encodes CD40 plus EGFP (MIEG3-CD40). Green fluorescence of Müller cells transduced with empty retroviral vector was similar to that of Müller cells transduced with CD40-encoding retroviral vector (not shown). (B, C) Primary Müller cells transduced with retroviral vector that encodes CD40 were incubated with RECs in the presence or absence of CD154. Percentages of cells that stained with Annexin V or 7-AAD were assessed by flow cytometry on gated EGFP⁻ cells (REC). Dot plots show representative data obtained at 96 hours of culture. (D, E) Primary Müller cells transduced with MIEG3 or with MIEG3-CD40 (D) or MIO M1 Müller cells transduced with MIEG3-CD40 (E) were incubated with RECs in the presence or absence of CD154 for 96 hours. The percentage of RECs (EGFP⁻) that stained with Annexin V or 7-AAD was assessed by flow cytometry. (F) Primary Müller cells transduced with empty retroviral vector or with retroviral vector that encodes CD40 were incubated with RECs in the presence or absence of CD154. The number of RECs (EGFP⁻) was determined at 120 hours. Results are representative of 3 to 4 independent experiments. **P* < 0.05; ***P* < 0.01.

conditions relevant to diabetes (Portillo et al., Manuscript in preparation). These vectors encode EGFP. Müller cells sorted for EGFP (>95% EGFP⁺) were incubated with primary RECs. These two populations were distinguished by EGFP expression (Fig. 2A). Expression of Annexin V and 7-AAD were examined in gated EGFP⁻ (endothelial cells). CD154 increased the percentages of Annexin V⁺ 7-AAD⁻ (Fig. 2B) that became significant after 72 hours of in vitro incubation (Fig. 2C). This effect was driven by CD40 signaling in Müller cells since no significant change in Annexin V⁺ 7-AAD⁻ cells was detected in cultures that contained Müller cells that were largely CD40⁻ (Fig. 2D). Similar results were obtained with a human Müller cells line (MIO-M1) transduced with the CD40-encoding vector (Fig. 2E). Moreover, the number of endothelial cells decreased when they were incubated with CD40⁺ Müller cells plus CD154 (Fig. 2F). Thus, CD40 stimulation increases PCD of RECs in the presence of retinal Müller cells.

PCD of Human RECs Does Not Appear to be Mediated by NOS2, Oxidative Stress, Fas Ligand, TNF- α , or IL-1 β

Nitrosative stress has been reported to cause PCD.^{7,11} Ligation of CD40 increases nitric oxide production by mouse Müller cells.¹⁵ The coculture of human Müller cells and RECs were incubated with L-Nil, a potent and specific inhibitor of nitric oxide (NO) production, followed by addition of CD154. We found L-NIL did not impair the induction of Annexin V⁺ 7-AAD⁻ RECs in CD40-activated cell cultures (Fig. 3A). Moreover, although CD40 ligation upregulates nitric oxide synthase (NOS2) in human Müller cells,¹⁵ these cells did not exhibit

detectable secretion of nitric oxide when incubated with CD154 for up to 72 hours (<0.39 μ M; not shown). Oxidative stress has been implicated in the development of capillary degeneration in diabetic retinopathy.⁸⁻¹⁰ In vivo administration of the antioxidant α -lipoic acid diminishes PCD of RECs in diabetic rats.⁹ However, incubation with α -lipoic acid failed to impair the increase in Annexin V⁺ 7-AAD⁻ endothelial cells in cultures stimulated with CD154 (Fig. 3B). Regardless of duration of incubation, Müller cells did not increase superoxide production in response to CD154 (control: 116.6 \pm 15.6; CD154: 103.3 \pm 7.9 relative units/mg protein; *P* = 0.4). TNF- α and IL-1 β stimulate cell death signaling in REC.¹² Nonetheless, CD154 did not induce TNF- α or IL-1 β production by Müller cells or RECs (<3.9 pg/mL; not shown). Next, we examined whether CD154 induced PCD of RECs through Fas ligand-Fas interaction. Addition of a blocking anti-Fas ligand antibody failed to affect the induction of Annexin V⁺ and 7-AAD⁻ RECs (Fig. 3C). Moreover, Müller cells remained Fas Ligand negative regardless of duration of incubation with CD154 (not shown). Thus, NOS2, oxidative stress, TNF- α , IL-1 β and Fas ligand do not appear to mediate CD40-mediated PCD of RECs.

Inhibition of the ATP Receptor P2X₇ Prevents PCD of RECs

Extracellular ATP can induce apoptosis or necrosis through binding to the purinergic receptor P2X₇.²⁵⁻²⁸ We examined whether CD40 ligation in primary Müller cells induces release of extracellular ATP. As shown in Figure 4A, Müller cells that expressed CD40 secreted extracellular ATP in response to CD154. Next, the selective P2X₇ receptor inhibitor A-




FIGURE 3. Programmed cell death of RECs is not prevented by an NOS2 inhibitor, an antioxidant, or an anti-Fas ligand mAb. Primary Müller cells transduced with a retroviral vector that encodes CD40 were incubated with RECs. The nitric oxide synthase inhibitor L-NIL (A), antioxidant α -lipoic acid (B) and a blocking anti-Fas ligand mAb (C) were added prior to stimulation with CD154. Percentage of RECs (EGFP⁻) that stained with Annexin V or 7-AAD was assessed by flow cytometry. Results are representative of three independent experiments.

438079^{29,30} was added to the coculture of RECs and Müller cells stimulated with CD154. P2X₇ receptor inhibition prevented the increase in Annexin V⁺ 7-AAD⁻ RECs in cultures stimulated with CD154 (Fig. 4B). Thus, CD40 ligation induces ATP release by primary Müller cells and triggers PCD of bystander RECs that appears dependent on P2X₇.

CD40 Ligation Upregulates P2X₇ Receptor Expression in RECs and Increases Their Susceptibility to PCD Triggered by ATP

The susceptibility to P2X₇ receptor-mediated PCD varies among cells. Addition of CD154 to cocultures of Müller cells and RECs did not enhance PCD of RECs until after 72 hours of incubation. Given that RECs express CD40 under resting conditions,¹⁵ we examined whether the susceptibility of RECs to ATP-mediated PCD is enhanced over time by incubation with CD154. Retinal endothelial cells were cultured with or without CD154, and Bz-ATP (a P2X purinergic agonist) was added during the last 24 hours of culture. Incubation with CD154 caused a significant increase in the percentages of Annexin V⁺ 7-AAD⁻ RECs after addition of Bz-ATP, an effect that was detected after 72 hours of incubation with CD154 (Fig. 5A). Retinal endothelial cells were also stained with Hoechst 33342 and propidium iodide. Increased PCD was confirmed by detecting higher percentages of RECs that stained brightly with Hoechst 33342 (nuclear condensation) and were propidium

iodide negative when these cells were incubated with CD154 for 72 hours and exposed to Bz-ATP (Fig. 5B). An increase in P2X₇ receptor levels promotes the effects of this receptor.³¹ Thus, we examined the effect of CD154 on P2X₇ receptor expression. We used IFN- γ as a positive control. Human RECs expressed low levels of P2X₇ receptor under basal conditions (Fig. 5C). Incubation with CD154 for ≥ 72 hours increased P2X₇ receptor expression in RECs (Fig. 5C). Taken together, CD40 ligation upregulates P2X₇ receptor in RECs and increases the susceptibility of these cells to PCD induced by a P2X₇ receptor agonist.

CD40 Promotes P2X₇ Receptor Upregulation in the Retina and Retinal Endothelial Cells of Diabetic Mice

A small percentage of RECs undergo PCD in the diabetic retina, a phenomenon that leads to development of capillary degeneration.¹⁻⁶ CD40 is required for capillary degeneration in diabetic mice.¹⁵ Indeed, tomato lectin⁺ (endothelial) cells that were TUNEL⁺ were noted in the retinas of diabetic B6 mice (1.15 ± 0.21 TUNEL⁺ tomato lectin⁺ cells/retina section in diabetic mice versus no TUNEL⁺ tomato lectin⁺ cells in nondiabetic controls; $n = 7$; $P < 0.01$), whereas no TUNEL⁺ tomato lectin⁺ cells were detected in diabetic or nondiabetic CD40^{-/-} or P2X₇^{-/-} mice ($n = 6$). To further assess the relevance of P2X₇ receptor, we examined whether expression




FIGURE 4. Programmed cell death of RECs is impaired by inhibition of the P2X₇ receptor. (A) Primary human Müller cells transduced with empty or CD40-encoding MIEG3 retroviral vectors were incubated with CD154. Release of ATP was assessed by a luciferase-based assay. (B) Primary Müller cells transduced with a retroviral vector that encodes CD40 were incubated with RECs. We added the P2X₇ receptor inhibitor A 438079 prior to stimulation with CD154. The percentage of RECs (EGFP⁻) that stained with Annexin V or 7-AAD was assessed by flow cytometry. Results are representative of three independent experiments. ** $P < 0.01$.




FIGURE 5. Cluster of differentiation 154 upregulates P2X₇ receptor in RECs and increases their susceptibility to ATP-driven PCD. **(A)** We incubated RECs with or without CD154 as indicated, followed with incubation with Bz-ATP (100 μM) for 24 hours. We stained RECs with Annexin V plus 7-AAD. **(B)** We cultured RECs with or without CD154 for 72 hours followed by incubation with or without Bz-ATP for 24 hours and stained with Hoechst 33342 plus propidium iodide. *Arrowheads* show RECs that stained intensely with Hoechst 33342 (nuclear condensation). *Bar*: 20 μm. **(C)** We stimulated RECs with or without CD154 or IFN-γ. Expression of P2X₇ receptor in RECs was assessed by flow cytometry. Histograms show P2X₇ receptor expression after incubation in complete medium alone (control), CD154 (96 hours) or IFN-γ (72 hours) of in vitro culture. *Dotted line*: secondary Ab alone. *Thin line*: control. *Thick line*: CD154 or IFN-γ. Results are representative of three independent experiments. **P* < 0.05; ***P* < 0.01; ****P* < 0.001.

of P2X₇ receptor in the retina of diabetic mice is increased in a CD40-dependent manner. Diabetic B6 mice exhibited a marked upregulation of P2X₇ receptor mRNA levels (Fig. 6A). This response was largely dependent on CD40 since no significant P2X₇ receptor upregulation was detected in diabetic CD40^{-/-} mice. Moreover, diabetic B6 mice exhibited increased P2X₇ receptor expression in RECs (Figs. 6B, 6C). Thus, CD40 promotes P2X₇ receptor upregulation in the retina and RECs of diabetic mice.

DISCUSSION

Herein we identified a mechanism by which CD40 can promote PCD of primary endothelial cells. We report that, whereas direct CD40 ligation in RECs did not result in PCD, CD40 stimulation enhanced PCD of RECs when these cells were incubated with Müller cells that express CD40. This effect did not appear to be mediated by NOS2, oxidative stress, TNF-α, IL-1β, or Fas ligand. Ligation of CD40 in primary Müller cells increased secretion of extracellular ATP, while CD40




FIGURE 6. Cluster of differentiation 40 promotes upregulation of P2X₇ receptor in the retina and RECs from diabetic mice. **(A)** At 8 and 18 weeks of diabetes, retinas from diabetic B6, CD40^{-/-} as well as from nondiabetic control animals were collected and used for mRNA extraction. We assessed mRNA levels by real-time quantitative PCR using 18S rRNA as internal control. One nondiabetic B6 mouse was given an arbitrary value of 1 and data are expressed as a fold-increase compared with this animal. Data shown represent mean \pm SEM (6 to 10 animals per group). ** $P < 0.01$; *** $P < 0.001$. **(B)** Retinal sections from B6 and CD40^{-/-} mice at 8 weeks of diabetes and from nondiabetic controls were incubated with anti-P2X₇ receptor Ab plus tomato lectin to label endothelial cells. Original magnification $\times 630$; 6 to 7 mice/group.

ligation in RECs upregulated P2X₇ receptor expression and made them susceptible to ATP-induced PCD. Inhibition of P2X₇ receptor during the coculture of RECs and Müller cells inhibited PCD of RECs. These results may be of in vivo relevance since P2X₇ is upregulated in a CD40-dependent manner in the retina of diabetic mice, RECs in these animals had increased P2X₇ receptor expression and CD40 is known to be required for retinal capillary degeneration.

Extracellular ATP binds two subfamilies of purinergic receptors: the inotropic, ligand-gated P2X receptors and the metabotropic, G protein-coupled P2Y receptors.³² An important distinction between P2X₇ receptor and other P2X receptors is that sustained activation of P2X₇ receptor results in formation of transmembrane pores that are permeable to hydrophilic molecules of up to 900 Da.³³⁻³⁵ Receptor P2X₇ mediates apoptotic or necrotic cell death depending on the cell type, duration of incubation with ATP and ATP concentration.²⁵⁻²⁸ Several lines of evidence support that CD40 stimulation in Müller cells induced PCD of RECs through P2X₇ receptor. First, primary Müller cells secreted extracellular ATP in response to CD40 ligation. Second, CD40 ligation upregulated P2X₇ receptor expression in RECs. Third, the P2X₇ receptor ligand Bz-ATP induced PCD of CD40-stimulated RECs. Fourth, a selective P2X₇ inhibitor prevented PCD of RECs. Addition of extracellular ATP at concentrations of 10 to 100 μ M

have been reported to be required for induction of PCD.^{36,37} The concentrations of extracellular ATP detected in bulk cell-free supernatants from CD40-stimulated Müller cells were lower than these concentrations. However, compared with single administration of extracellular ATP, cell death after repeated exposure to ATP requires lower concentrations of the nucleotide.³⁵ In addition, luciferase-based assays of ATP concentrations in extracellular medium underestimate the concentrations of ATP in the intercellular space.³⁸ Assays based on targeting luciferase to the surface of intact cells revealed higher concentrations of ATP at the cell surface level (100-200 μ M).^{38,39}

Susceptibility to ATP-mediated PCD varies among cell types. Our studies suggest that CD40 plays a dual role in promoting PCD of RECs: it enhances release of extracellular ATP by primary Müller cells and increases the susceptibility of RECs to P2X₇-driven PCD that would overcome the prosurvival signals activated by CD40 ligation. Increased susceptibility to PCD may be mediated by CD40-driven upregulation of the P2X₇ receptor in RECs since increased P2X₇ receptor levels promote the effects of this receptor.³¹ However, it is also possible that CD40 ligation may increase susceptibility to P2X₇ receptor-mediated PCD through modulation of ATP-gated channel expression, ectoATPase activity, and/or coupling to downstream cell signaling pathways that promote cell death. Of

relevance, diabetes or incubation with high concentrations of glucose appears to increase the susceptibility to ATP-driven PCD in retinal microvessels, fibroblasts, and T cells.⁴⁰⁻⁴² Thus, CD40 may contribute to increased susceptibility to ATP-mediated PCD that appears to occur in diabetes.

Müller cells become dysfunctional in diabetic retinopathy⁴³⁻⁴⁵ and are likely to affect other retinal cells since Müller cells span the whole width of the retina and provide structural support to retinal blood vessels and neurons. This study, together with the fact that Müller cells surround retinal capillaries, suggests that CD40-driven ATP release may play an important role in the pathogenesis of PCD of RECs and capillary degeneration in diabetic retinopathy. Indeed, expression of CD40 restricted to Müller cells was sufficient to cause capillary degeneration in diabetic transgenic mice.⁴⁶ Long-term studies in diabetic P2X₇^{-/-} mice may help elucidate the in vivo role of this receptor in capillary degeneration. We recently observed that P2X₇ receptor also promotes expression of proinflammatory cytokines in the diabetic retina.⁴⁶ Ligation of CD40 in human and mouse Müller cell lines caused secretion of ATP that in turn induced P2X₇ receptor-dependent IL-1 β and TNF- α secretion by bystander myeloid cells.⁴⁶ Moreover, diabetic transgenic mice that expressed CD40 restricted to Müller cells exhibited retinal upregulation of IL-1 β and TNF- α was dependent on P2X₇ receptor.⁴⁶ These results together with the present study indicate that the CD40-ATP-P2X₇ receptor pathway plays an important role in induction of inflammatory responses and death of RECs, events considered pathogenic in the development of diabetic retinopathy.

Our studies centered on Müller cells since CD40 expression restricted to these cells is sufficient to cause capillary degeneration in diabetes. However, it is possible that CD40 signaling in other cell types (including REC) may induce ATP release setting in motion PCD of RECs. In addition, while our studies indicate that CD40-induced ATP release followed by P2X₇ receptor signaling in RECs mediates PCD, it is possible that CD40 may also promote death of RECs and capillary degeneration in vivo through additional mechanisms. CD40 enhances retinal leukostasis in diabetic mice, upregulates NOS2, increases retinal protein nitration and upregulates TNF- α and IL-1 β in the retinas of diabetic mice.^{15,46} All these phenomena have been linked to PCD of RECs and retinal capillary degeneration.^{3,7,11-14,47}

In summary, these studies uncovered a mechanism by which CD40 enhances PCD of RECs and suggest that CD40 signaling in Müller cells may be an important contributor to vascular injury in diabetic retinopathy. Our findings may be relevant to other CD40-driven diseases such as atherosclerosis and inflammatory bowel disease. Increased PCD accompanies these disorders and P2X₇ receptor has been suggested to be pathogenic in these diseases.^{48,49} Strong preclinical data indicate that blockade of the CD40-CD154 pathway can be effective therapy for various inflammatory and neurodegenerative disorders. Our studies suggest that novel approaches to inhibit CD40 signaling may be effective in the management of diabetic retinopathy.

Acknowledgments

The authors thank Richard Kornbluth and Gloria Limb for providing reagents and cell lines.

Supported by NIH Grant EY019250 (CSS) and NIH Grant P30 EY11373.

Disclosure: **J.-A.C. Portillo**, None; **Y. Lopez Corcino**, None; **G.R. Dubyak**, None; **T.S. Kern**, None; **S. Matsuyama**, None; **C.S. Subauste**, None

References

- Mizutani M, Kern TS, Lorenzi M. Accelerated death of retinal microvascular cells in human and experimental diabetic retinopathy. *J Clin Invest*. 1996;97:2883-2890.
- Engerman RL, Kern TS. Retinopathy in animal models of diabetes. *Diabetes Metab Rev*. 1995;11:109-120.
- Behl Y, Krothapalli P, Desta T, DiPiazza A, Roy S, Graves DT. Diabetes-enhanced tumor necrosis factor- α production promotes apoptosis and the loss of retinal microvascular cells in type 1 and type 2 models of diabetic retinopathy. *Am J Pathol*. 2008;172:1411-1418.
- Joussen AM, Poulaki V, Mitsiades N, et al. Suppression of Fas-FasL-induced endothelial cell apoptosis prevents diabetic blood-retinal barrier breakdown in a model of streptozotocin-induced diabetes. *FASEB J*. 2002;17:76-78.
- Kern TS, Tang J, Mizutani M, et al. Response of capillary cell death to aminoguanidine predicts the development of retinopathy: comparison of diabetes and galactosemia. *Invest Ophthalmol Vis Sci*. 2000;41:3972-3978.
- Kowluru RA, Odenbach S. Role of interleukin-1 β in the development of retinopathy in rats: effect of antioxidants. *Invest Ophthalmol Vis Sci*. 2004;45:4161-4166.
- Du Y, Smith MA, Miller CM, Kern TS. Diabetes-induced nitrate stress in the retina, and correction by aminoguanidine. *J Neurochem*. 2002;80:771-779.
- Du Y, Miller CM, Kern TS. Hyperglycemia increases mitochondrial superoxide in retina and retinal cells. *Free Radical Biol Med*. 2003;35:1491-1499.
- Kowluru RA, Odenbach S. Effect of long-term administration of alpha-lipoic acid on retinal capillary cell death and the development of retinopathy in diabetic rats. *Diabetes*. 2004;53:3233-3238.
- El-Remessy AB, Al-Shabraway M, Khalifa Y, Tsai NT, Caldwell RB, Liou GI. Neuroprotective and blood-retinal barrier-preserving effects of cannabidiol in experimental diabetes. *Am J Pathol*. 2006;168:235-244.
- Zheng L, Du Y, Miller C, et al. Critical role of inducible nitric oxide synthase in degeneration of retinal capillaries in mice with streptozotocin-induced diabetes. *Diabetologia*. 2007;50:1987-1996.
- Busik JV, Mohr S, Grant MB. Hyperglycemia-induced reactive oxygen species toxicity to endothelial cells is dependent on paracrine mediators. *Diabetes*. 2008;57:1952-1965.
- Joussen AM, Doehmen S, Le ML, et al. TNF- α mediated apoptosis plays an important role in the development of early diabetic retinopathy and long-term histopathological alterations. *Mol Vision*. 2009;15:1418-1428.
- Vincent JA, Mohr S. Inhibition of caspase-1/interleukin-1 β signaling prevents degeneration of retinal capillaries in diabetes and galactosemia. *Diabetes*. 2007;56:224-230.
- Portillo J-AC, Greene JA, Okenka G, et al. CD40 promotes the development of early diabetic retinopathy in mice. *Diabetologia*. 2014;57:2222-2231.
- Dechanet J, Grosset C, Taupin JL, et al. CD40 ligand stimulates proinflammatory cytokine production by human endothelial cells. *J Immunol*. 1997;159:5640-5647.
- Deregibus MC, Buttiglieri S, Russo S, Bussolati B, Camussi G. CD40-dependent activation of phosphatidylinositol 3-kinase/Akt pathway mediates endothelial cell survival and in vitro angiogenesis. *J Biol Chem*. 2003;278:18008-18014.
- Ahmed-Choudhury J, Russell CL, Randhawa S, Young LS, Adams DH, Afford SC. Differential induction of nuclear factor-kappaB and activator protein-1 activity after CD40 ligation is associated with primary human hepatocyte apoptosis or intrahepatic endothelial cell proliferation. *Mol Biol Cell*. 2003;14:1334-1345.

19. Andrade RM, Wessendarp M, Portillo J-AC, et al. TRAF6 signaling downstream of CD40 primes macrophages to acquire anti-microbial activity in response to TNF- α . *J Immunol*. 2005;175:6014–6021.
20. Blum AE, Joseph SM, Przybylski RJ, Dubyak GR. Rho-family GTPases modulate Ca(2+)-dependent ATP release from astrocytes. *Am J Physiol Cell Physiol*. 2008;295:C231–C241.
21. Milano PM, Douillet CD, Riesenman PJ, et al. Intestinal ischemia-reperfusion injury alters purinergic receptor expression in clinically relevant extraintestinal organs. *J Surg Res*. 2008;145:272–278.
22. Portillo J-AC, Van Grol J, Zheng L, et al. CD40 mediates retinal inflammation and neurovascular degeneration. *J Immunol*. 2008;181:8719–8726.
23. Portillo J-A, Schwartz I, Zarini S, et al. Pro-inflammatory responses induced by CD40 in retinal endothelial and Müller cells are inhibited by blocking CD40-Traf23 or CD40-Traf6 signaling. *Invest Ophthalmol Vis Sci*. 2014;55:8590–8597.
24. Portillo JA, Greene JA, Schwartz I, Subauste MC, Subauste CS. Blockade of CD40-TRAF2,3 or CD40-TRAF6 is sufficient to impair pro-inflammatory responses in human aortic endothelial cells and human aortic smooth muscle cells. *Immunology*. 2015;144:21–33.
25. Di Virgilio F, Chiozzi P, Falzoni S, et al. Cytolytic P2X purinoceptors. *Cell Death Differ*. 1998;5:191–199.
26. von Albertini M, Palmetshofer A, Kaczmarek E, et al. Extracellular ATP and ADP activate transcription factor NF- κ B and induce endothelial cell apoptosis. *Biochem Biophys Res Comm*. 1998;248:822–829.
27. Jun DJ, Kim J, Jung SY, et al. Extracellular ATP mediates necrotic cell swelling in SN4741 dopaminergic neurons through P2X7 receptors. *J Biol Chem*. 2007;282:37350–37358.
28. Humphreys BD, Rice J, Kertesz SB, Dubyak GR. Stress-activated protein kinase/JNK activation and apoptotic induction by the macrophage P2X7 nucleotide receptor. *J Biol Chem*. 2000;275:26792–26798.
29. Nelson DW, Gregg RJ, Kort ME, et al. Structure-activity relationship studies on a series of novel substituted 1-benzyl-5-phenyltetrazole P2X7 antagonists. *J Med Chem*. 2006;49:3659–3666.
30. Donnelly-Roberts DL, Jarvis ME. Discovery of P2X7 receptor-selective antagonists offers new insights into P2X7 receptor function and indicates a role in chronic pain states. *Br J Pharmacol*. 2007;151:571–579.
31. Monif M, Reid CA, Powell KL, Smart ML, Williams DA. The P2X7 receptor drives microglial activation and proliferation: a trophic role for P2X7R pore. *J Neurosci*. 2009;29:3781–3791.
32. Eltzschig HK, Sitkovsky MV, Robson SC. Purinergic signaling during inflammation. *New Engl J Med*. 2012;367:2322–2333.
33. Valera S, Hussy N, Evans RJ, et al. A new class of ligand-gated ion channel defined by P2x receptor for extracellular ATP. *Nature*. 1994;371:516–519.
34. Falzoni S, Munerati M, Ferrari D, Spisani S, Moretti S, Di Virgilio F. The purinergic P2Z receptor of human macrophage cells. Characterization and possible physiological role. *J Clin Invest*. 1995;95:1207–1216.
35. Surprenant A, Rassendren F, Kawashima E, North RA, Buell G. The cytolytic P2Z receptor for extracellular ATP identified as a P2X receptor (P2X7). *Science*. 1996;272:735–738.
36. Zanello P, Bronte V, Rosato A, Pizzo P, Di Virgilio F. Responses of mouse lymphocytes to extracellular ATP. II. Extracellular ATP causes cell type-dependent lysis and DNA fragmentation. *J Immunol*. 1990;145:1545–1550.
37. Correale P, Tagliaferri P, Guarrasi R, et al. Extracellular adenosine 5' triphosphate involvement in the death of LAK-engaged human tumor cells via P2X-receptor activation. *Immunol Letters*. 1997;55:69–78.
38. Beigi R, Kobatake E, Aizawa M, Dubyak GR. Detection of local ATP release from activated platelets using cell surface-attached firefly luciferase. *Am J Physiol*. 1999;276:C267–C278.
39. Pellegatti P, Falzoni S, Pinton P, Rizzuto R, Di Virgilio F. A novel recombinant plasma membrane-targeted luciferase reveals a new pathway for ATP secretion. *Mol Biol Cell*. 2005;16:3659–3665.
40. Sugiyama T, Kobayashi M, Kawamura H, Li Q, Puro DG. Enhancement of P2X7-induced pore formation and apoptosis: An early effect of diabetes on the retinal microvasculature. *Invest Ophthalmol Vis Sci*. 2004;45:1026–1032.
41. Elliott JI, Higgins CF. Enhanced P2X7 activity in human fibroblasts from diabetic patients: a possible pathogenetic mechanism for vascular damage in diabetes. *Diabetes*. 2004;53:2012–2017.
42. Solini A, Chiozzi P, Falzoni S, Morelli A, Fellin R, Di Virgilio F. High glucose modulates P2X7 receptor-mediated function in human primary fibroblasts. *Diabetologia*. 2000;43:1248–1256.
43. Lieth E, Barber AJ, Xu B, et al. Glial reactivity and impaired glutamate metabolism in short-term experimental diabetic retinopathy. *Diabetes*. 1998;47:815–820.
44. Li Q, Puro DG. Diabetes-induced dysfunction of the glutamate transporter in retinal Müller cells. *Invest Ophthalmol Vis Sci*. 2002;43:3109–3116.
45. Gerhardinger C, Costa MB, Coulombe MC, Toth I, Hoehn T, Grosu P. Expression of acute-phase response proteins in retinal Müller cells in diabetes. *Invest Ophthalmol Vis Sci*. 2005;46:349–357.
46. Portillo J-AC, Lopez Corcino Y, Miao Y, et al. CD40 in retinal Müller cells induces P2X₇-dependent cytokine expression in macrophages/microglia in diabetic mice and development of early experimental diabetic retinopathy in mice. *Diabetes*. 2016. In press.
47. Joussen AM, Poulaki V, Le ML, et al. A central role for inflammation in the pathogenesis of diabetic retinopathy. *FASEB J*. 2004;18:1450–1452.
48. Piscopiello M, Sessa M, Anzalone N, et al. P2X7 receptor is expressed in human vessels and might play a role in atherosclerosis. *Int J Cardiol*. 2013;168:2863–2866.
49. Neves AR, Castelo-Branco MT, Figliuolo VR, et al. Overexpression of ATP-activated P2X7 receptors in the intestinal mucosa is implicated in the pathogenesis of Crohn's disease. *Inflamm Bowel Dis*. 2014;20:444–457.