

Lawrence Berkeley National Laboratory

Recent Work

Title

E+ POLARIZATION IN $n+n \rightarrow K^+ E^+$ FROM 1.1 TO 2.4 GeV/c

Permalink

<https://escholarship.org/uc/item/65p8d5kx>

Authors

Davies, Donald W.

Abolins, Maris A.

Dahl, Orin I.

et al.

Publication Date

1973

Σ^+ POLARIZATION IN $\pi^+n \rightarrow K^0 \Sigma^+$ FROM 1.1 TO 2.4 GeV/c

Donald W. Davies, Maris A. Abolins,
Orin I. Dahl, Jerome S. Danburg, Paul L. Hoch,
Janos Kirz, Donald H. Miller, and Robert K. Rader

January 17, 1973

Prepared for the U.S. Atomic Energy
Commission under Contract W-7405-ENG-48

For Reference

Not to be taken from this room



DISCLAIMER

This document was prepared as an account of work sponsored by the United States Government. While this document is believed to contain correct information, neither the United States Government nor any agency thereof, nor the Regents of the University of California, nor any of their employees, makes any warranty, express or implied, or assumes any legal responsibility for the accuracy, completeness, or usefulness of any information, apparatus, product, or process disclosed, or represents that its use would not infringe privately owned rights. Reference herein to any specific commercial product, process, or service by its trade name, trademark, manufacturer, or otherwise, does not necessarily constitute or imply its endorsement, recommendation, or favoring by the United States Government or any agency thereof, or the Regents of the University of California. The views and opinions of authors expressed herein do not necessarily state or reflect those of the United States Government or any agency thereof or the Regents of the University of California.

0 0 0 0 3 9 0 0 0 3 2

[To be submitted to Phys. Rev. D, Comments and Addenda]

Σ^+ Polarization in $\pi^+n \rightarrow K^0\Sigma^+$ from 1.1 to 2.4 GeV/c*

Donald W. Davies,[†] Maris A. Abolins,[‡] Orin I. Dahl, Jerome S. Danburg,[§]
Paul L. Hoch,^{||} Janos Kirz,^{**} Donald H. Miller,^{††} and Robert K. Rader^{§§}

Lawrence Berkeley Laboratory
Berkeley, California 94720

January 17, 1973

ABSTRACT

We observe an energy-dependent polarization of the Σ^+ produced in the reaction $\pi^+n \rightarrow K^0\Sigma^+$ at incident beam momenta between 1.1 and 2.4 GeV/c. This data is a significant source of information on the Σ^- polarization in the charge-symmetric reaction $\pi^-p \rightarrow K^+\Sigma^-$.

Previous authors have discussed two reasons for interest in the Σ^- polarization in $\pi^-p \rightarrow K^+\Sigma^-$. First, knowledge of the polarization might be useful for a study of the β decay of the Σ^- .¹ Also, the reactions



are related by charge independence; the two isospin amplitudes can be extracted from the cross sections and polarizations.²⁻⁵

The available data on reaction (2)^{6,7} and reaction (3)⁸⁻¹⁰ includes information on the polarization of the Σ through observation of the Σ decay asymmetry. However, in studies of reaction (1) with unpolarized protons,^{3,9,11} polarization information is unavailable because of the very small asymmetry parameter in the decay $\Sigma^- \rightarrow n\pi^-$ ($\alpha = -0.07 \pm 0.01$).¹² Some limited use has been made of polarized targets;^{1,2} the Σ^- polarization has been measured at 1.74 GeV.²


If charge symmetry (which is weaker than charge independence) holds, the reaction



has properties identical to reaction (1). Then a measurement of the Σ^+ polarization in reaction (4) is equivalent to a measurement of the Σ^- polarization in reaction (1).

The data presented here comes from a 250,000-picture experiment done at the LBL Bevatron in the fall of 1966.¹³ The deuterium-filled 72-inch bubble chamber was exposed to π^+ beams of eight different momenta from 1.1 to 2.4 GeV/c. The film was subsequently scanned twice

for events with a charged-mode decay of a neutral particle (ν) and measured on the Group A Franckenstein measuring machines. The measured events were then processed through the standard geometrical reconstruction and kinematic fitting program SIOUX. Events indicated by the scanner as having a decaying positive particle, as well as a ν , were fitted to all final states possible for that topology, including



Events which had acceptable fits to two or more reactions were examined on the scanning table; the ambiguity was resolved, if possible, by comparing the predicted track ionization for the various hypotheses with the observed bubble density. The end result was 164 events with a "best" fit to reaction (5), only 3 of which remained ambiguous. (See Ref. 13 for the definition of the "best" fit.)

To select those events in which the proton is a "spectator" to the reaction, we require the final-state proton to have a momentum compatible with the internal momentum of the deuteron. All commonly used deuterium wave functions show little probability that the nucleons would have more than 300 MeV/c momentum; therefore events with proton laboratory-frame momentum greater than 300 MeV/c have been omitted from the rest of the analysis.¹³ The remaining 135 events were divided into 6 intervals in the center-of-mass energy, $M(K^0\Sigma^+)$. See Fig. 1 for the c.m. energy distribution.

Fig. 2 shows the production cosine distribution for the 6 c.m. energy intervals. Here 167 events from the $\Sigma^+ \rightarrow n\pi^+$ decay mode have been included to increase the statistics of the production cosine distribution,

although no polarization information can be extracted from these events because of the small decay asymmetry parameter. The distributions of Fig. 2 are consistent with, but not as precise as, the production cosine distributions for the charge-symmetric reaction $\pi^- p \rightarrow K^+ \Sigma^-$.⁹

Any Σ^+ polarization must lie along the normal to the production plane. Fig. 3 shows the cosine distribution of the decay protons in the Σ^+ rest frame plotted against production cosine. Here

$$\cos \theta_{\text{prod}} = \hat{\pi} \cdot \hat{K},$$

$$\hat{n} \propto \hat{\pi} \times \hat{K}, \text{ and}$$

$$\cos \theta_{\text{dk}} = \hat{p} \cdot \hat{n},$$

where $\hat{\pi}$ and \hat{K} are the directions of the beam π^+ and the K^0 in the c.m., \hat{n} is the production normal, and \hat{p} is the direction of the proton in the rest frame of the Σ^+ .

The decay distribution of a spin-1/2 particle must be of the form

$$\frac{dN}{d(\cos \theta_{\text{dk}})} \propto 1 + \alpha P \cos \theta_{\text{dk}}$$

where α is the asymmetry parameter for the decay and P is the polarization. A maximum-likelihood fit was done to the function

$$f(A, \cos \theta_{\text{dk}}) = 1 + A \cos \theta_{\text{dk}}.$$

The value so obtained for A is then our best estimate of the product of the asymmetry parameter and the polarization. The scanning biases are expected to be even in $\cos \theta_{\text{dk}}$ and therefore not to affect the maximum-likelihood estimate.¹⁴

Taking each of the 6 energy intervals separately, and integrating over production angle, we get the polarization given in Table 1 and Fig. 4.

(We have used $\alpha = -1.0^{12}$) There is clear indication of a large net polarization in the lower energy intervals. Noting a difference in the definitions of the normal, we see that our lowest-energy result has the same sign as that of Edgington et al. at 1.74 GeV.²

Because of poor statistics, we cannot divide each of the six energy intervals into bins in production cosine; instead, we have combined the three lower energy intervals and the three upper ones, and calculated the polarization for three bins in production cosine. The results are given in Table 2 and Fig. 5.

REFERENCES AND FOOTNOTES

1. D. Weldon *et al.*, Phys. Rev. 167, 1199 (1968).
2. J. A. Edgington *et al.*, Phys. Rev. 177, 2103 (1969)
3. J. C. Doyle *et al.*, Phys. Rev. 165, 1483 (1968).
4. J. J. Sakurai, Phys. Rev. 107, 908 (1957).
5. T. O. Binford *et al.*, Phys. Rev. 183, 1148 (1969).
6. N. L. Carayannopoulos *et al.*, Phys. Rev. 138, B433 (1965).
7. G. E. Kalmus *et al.*, Phys. Rev. D2, 1824 (1970).
8. Y. S. Kim *et al.*, Phys. Rev. 143, 1028 (1966).
9. O. I. Dahl *et al.*, Phys. Rev. 163, 1430 (1967).
10. T. O. Binford *et al.*, Phys. Rev. 183, 1134 (1969).
11. M. L. Good and R. R. Kofler, Phys. Rev. 183, 1142 (1969).
12. Particle Data Group, Phys. Letters 39B, 1 (1972).
13. This paper supplements a report on strange-particle production in π^+d interactions: Donald W. Davies *et al.*, Phys. Rev. D2, 506 (1970). The data analysis is discussed there in more detail. Information on other aspects of this experiment is given in Jerome S. Danburg *et al.*, Phys. Rev. D2, 2564 (1970) and in Robert K. Rader *et al.*, Phys. Rev. D6, 3059 (1972).
14. Roger O. Bangerter (Ph. D. thesis), LBL Report UCRL-19244 (unpublished), Section IVA.

- * Work supported by the U.S. Atomic Energy Commission
- † Present address: Planetary Atmospheres Section, Jet Propulsion Laboratory, Pasadena, Calif.
- ‡ Present address: Department of Physics, Michigan State University, East Lansing, Mich. 48823
- § Present address: Geophysics Department, Shell Development Company, Houston, Texas
- || Present address: 1735 Highland Place, Berkeley, Calif. 94709
- ** Present address: Dept. of Physics, State University of New York, Stony Brook, NY 11790 (On leave, 1972-73, to Laboratory of Molecular Biophysics, Oxford University, England)
- †† Present address: Dept. of Physics, Northwestern University, Evanston, Ill. 60201
- §§ Present address: Computer-Based Education Research Laboratory, University of Illinois, Urbana, Ill. 61801

TABLE I. Polarization Integrated over Production Angle

C.m. Energy (GeV)	Polarization
1.7 - 1.8	0.90 +0.60 -0.95
1.8 - 1.9	0.90 +0.21 -0.32
1.9 - 2.0	0.40 +0.23 -0.27
2.0 - 2.1	-0.03 +0.26 -0.23
2.1 - 2.2	-0.64 +0.47 -0.38
2.2 - 2.4	0.35 +0.38 -0.50

TABLE II. Polarization as a Function of Production Angle

C.m. Energy (GeV)	Cos θ_{prod}	Polarization
1.7 - 2.0	-1.00 to -0.33	0.53 +0.23 -0.27
"	-0.33 to 0.33	0.72 +0.27 -0.33
"	0.33 to 1.00	0.27 +0.59 -0.68
2.0 - 2.4	-1.00 to -0.33	0.44 +0.22 -0.25
"	-0.33 to 0.33	-0.52 +0.30 -0.25
"	0.33 to 1.00	-0.58 +0.47 -0.34

FIGURE CAPTIONS

- Fig. 1 Distribution of c.m. energy (excluding the spectator proton) for our sample of the reaction $\pi^+d \rightarrow (p)K^0\Sigma^+$, $\Sigma^+ \rightarrow p\pi^0$ (135 events).
- Fig. 2 Production distribution of the Σ^+ for c.m. energies (a) 1.7-1.8, (b) 1.8-1.9, (c) 1.9-2.0, (d) 2.0-2.1, (e) 2.1-2.2, (f) 2.2-2.4 GeV. $\cos \theta_{\text{prod}} = \hat{\pi} \cdot \hat{K}$ (302 events).
- Fig. 3 Decay cosine of $\Sigma^+ \rightarrow p\pi^0$ vs production cosine ($\hat{\pi} \cdot \hat{K}$) for the six c.m. energy intervals of Fig. 2.
- Fig. 4 Polarization integrated over production angle as a function of c.m. energy.
- Fig. 5 Polarization as a function of production angle for c.m. energy (a) 1.7-2.0 GeV and (b) 2.0-2.4 GeV.




Fig. 1





Fig. 2



$\text{Cos}(\theta_{\text{prod}})$

Fig. 3




Fig. 5

LEGAL NOTICE

This report was prepared as an account of work sponsored by the United States Government. Neither the United States nor the United States Atomic Energy Commission, nor any of their employees, nor any of their contractors, subcontractors, or their employees, makes any warranty, express or implied, or assumes any legal liability or responsibility for the accuracy, completeness or usefulness of any information, apparatus, product or process disclosed, or represents that its use would not infringe privately owned rights.

TECHNICAL INFORMATION DIVISION
LAWRENCE BERKELEY LABORATORY
UNIVERSITY OF CALIFORNIA
BERKELEY, CALIFORNIA 94720