

UC Berkeley

UC Berkeley Previously Published Works

Title

TLR2 and endosomal TLR-mediated secretion of IL-10 and immune suppression in response to phagosome-confined *Listeria monocytogenes*

Permalink

<https://escholarship.org/uc/item/65z3j5dc>

Journal

PLOS Pathogens, 16(7)

ISSN

1553-7366

Authors

Nguyen, Brittney N  
Chávez-Arroyo, Alfredo  
Cheng, Mandy I  
et al.

Publication Date

2020

DOI

10.1371/journal.ppat.1008622

Peer reviewed

RESEARCH ARTICLE

# TLR2 and endosomal TLR-mediated secretion of IL-10 and immune suppression in response to phagosome-confined *Listeria monocytogenes*

Brittney N. Nguyen<sup>1</sup>, Alfredo Chávez-Arroyo<sup>1</sup>, Mandy I. Cheng<sup>2</sup>, Maria Krasilnikov<sup>2</sup>, Alexander Louie<sup>3</sup>, Daniel A. Portnoy<sup>2,3\*</sup>

**1** Graduate Group in Microbiology, University of California, Berkeley, Berkeley, California, United States of America, **2** Department of Molecular and Cell Biology, University of California, Berkeley, Berkeley, California, United States of America, **3** Department of Plant and Microbial Biology, University of California, Berkeley, Berkeley, California, United States of America


☯ These authors contributed equally to this work.

✉ Current address: Department of Microbiology Immunology and Molecular Genetics, University of California, Los Angeles, Los Angeles, California United States of America

✉ Current address: Department of Medicine, Cardiovascular Research Institute, University of California, San Francisco, San Francisco, California, United States of America

✉ Current address: Novome Biotechnologies, Inc., South San Francisco, California, United States of America

\* [portnoy@berkeley.edu](mailto:portnoy@berkeley.edu)



**OPEN ACCESS**

**Citation:** Nguyen BN, Chávez-Arroyo A, Cheng MI, Krasilnikov M, Louie A, Portnoy DA (2020) TLR2 and endosomal TLR-mediated secretion of IL-10 and immune suppression in response to phagosome-confined *Listeria monocytogenes*. PLoS Pathog 16(7): e1008622. <https://doi.org/10.1371/journal.ppat.1008622>

**Editor:** Igor Eric Brodsky, University of Pennsylvania, UNITED STATES

**Received:** January 8, 2020

**Accepted:** May 12, 2020

**Published:** July 7, 2020

**Copyright:** © 2020 Nguyen et al. This is an open access article distributed under the terms of the [Creative Commons Attribution License](https://creativecommons.org/licenses/by/4.0/), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

**Data Availability Statement:** All relevant data are within the manuscript and its Supporting Information files.

**Funding:** D.A.P. was supported by National Institutes of Health (<https://www.nih.gov/>) grants 1P01 AI063302, and 1R01 AI027655. B.N.N was supported by National Science Foundation (<https://www.nsf.gov/>), Graduate Research Fellowship DGE 1106400. The funders had no role in study design,

## Abstract

*Listeria monocytogenes* is a facultative intracellular bacterial pathogen that escapes from phagosomes and induces a robust adaptive immune response in mice, while mutants unable to escape phagosomes fail to induce a robust adaptive immune response and suppress the immunity to wildtype bacteria when co-administered. The capacity to suppress immunity can be reversed by blocking IL-10. In this study, we sought to understand the host receptors that lead to secretion of IL-10 in response to phagosome-confined *L. monocytogenes* ( $\Delta hly$ ), with the ultimate goal of generating strains that fail to induce IL-10. We conducted a transposon screen to identify  $\Delta hly$  *L. monocytogenes* mutants that induced significantly more or less IL-10 secretion in bone marrow-derived macrophages (BMMs). A transposon insertion in *Igt*, which encodes phosphatidylglycerol-prolipoprotein diacylglycerol transferase and is essential for the formation of lipoproteins, induced significantly reduced IL-10 secretion. Mutants with transposon insertions in *pgdA* and *oatA*, which encode peptidoglycan N-acetylglucosamine deacetylase and O-acetyltransferase, are sensitive to lysozyme and induced enhanced IL-10 secretion. A  $\Delta hly\Delta pgdA\Delta oatA$  strain was killed in BMMs and induced enhanced IL-10 secretion that was dependent on Unc93b1, a trafficking molecule required for signaling of nucleic acid-sensing TLRs. These data revealed that nucleic acids released by bacteriolysis triggered endosomal TLR-mediated IL-10 secretion. Secretion of IL-10 in response to infection with the parental strain was mostly TLR2-dependent, while IL-10 secretion in response to lysozyme-sensitive strains was dependent on TLR2 and Unc93b1. In mice, the IL-10 response to vacuole-confined *L. monocytogenes* was also dependent on TLR2 and Unc93b1. Co-administration of  $\Delta hly$  and  $\Delta actA$  resulted in suppressed immunity in WT mice,

data collection and analysis, decision to publish, or preparation of the manuscript.

**Competing interests:** The authors have declared that no competing interests exist.

but not in mice with mutations in *Unc93b1*. These data revealed that secretion of IL-10 in response to *L. monocytogenes* infection *in vitro* is mostly TLR2-dependent and immune suppression by phagosome-confined bacteria *in vivo* is mostly dependent on endosomal TLRs.

### Author summary

*Listeria monocytogenes* is a Gram-positive bacterial pathogen that has shown promise as a vaccine-delivery vector because of its ability to stimulate a robust T-cell response. The efficacy of a vaccine is in part tied to how well it avoids inducing the immunosuppressive cytokine IL-10. In this work, we investigated the bacterial and host factors that contribute to secretion of IL-10 and immunosuppression following infection with a strain of *L. monocytogenes* that cannot escape from host cell vacuoles. We identified TLR2 and *Unc93b1*-dependent nucleic-acid-sensing Toll-like receptors as the primary host mediators of IL-10 secretion and immunosuppression. *Unc93b1*-dependent TLRs likely recognize nucleic acids released upon bacterial lysis in phagosomes. Strategies that reduce lytic death of bacteria and subsequent recognition of nucleic acids by endosomal TLRs could be used to improve bacteria-based vaccines.

### Introduction

*Listeria monocytogenes* is a Gram-positive facultative intracellular pathogen that has been widely used as a model to study host immune responses. Infection of mice with *L. monocytogenes* induces the generation of adaptive immune responses that are protective against subsequent infection and are largely mediated by CD8<sup>+</sup> T cells [1,2]. Many innate immune factors contribute to control of primary *L. monocytogenes* infection. The factors that contribute to development of functional CD8<sup>+</sup> effector and memory T cells are less clear [3–6]. It has been posited that intracytosolic growth of *L. monocytogenes* is a prerequisite for the induction of T cell-mediated immunity because  $\Delta hly$  *L. monocytogenes*, which does not produce the virulence factor listeriolysin O and cannot escape phagocytic vacuoles, fails to induce robust protective immunity [7,8]. However, in 2009 Bahjat *et al.* provided evidence that  $\Delta hly$  *L. monocytogenes* fails to induce robust protective immunity because it induces secretion of IL-10 early during infection. When IL-10 signaling was inhibited by administration of anti-IL-10 receptor blocking antibody, the protective capacity of  $\Delta hly$  *L. monocytogenes* was enhanced [9]. Thus, strains that escape phagocytic vacuoles and grow in the cytosol induce generation of protective immunity in part because they avoid inducing IL-10-mediated suppression.

IL-10 is an anti-inflammatory cytokine that acts on many cell types to downregulate inflammation and recruitment of immune cells, thereby limiting immunopathology during the resolution of an immune response [10]. Infection of wildtype mice with *L. monocytogenes* results in high levels of serum IL-10 3–4 days post infection [11,12]. Mice lacking IL-10 clear *L. monocytogenes* faster than wildtype mice, indicating that IL-10 partially suppresses the primary innate immune response to *L. monocytogenes* [13,14]. Importantly, though IL-10 is expressed following primary WT *L. monocytogenes* infection, WT infection still results in the generation of a robust protective immune response.

IL-10 is secreted by many cell types, including macrophages, dendritic cells (DCs), neutrophils, NK cells, T-regulatory cells and B cells [10]. The IL-10 present in serum 3 to 4 days after WT *L. monocytogenes* infection of mice is mostly derived from NK cells [11]. However,  $\Delta hly$  *L.*

*monocytogenes* induces IL-10 secretion that is detectable four hours after infection, and likely comes from macrophages or dendritic cells, which are the first cells infected by *L. monocytogenes* in the spleen [9,15,16]. In macrophages, IL-10 expression can be triggered by activation of the pattern recognition receptors TLR2, TLR3, TLR4, and TLR9 [10,17]. Macrophages secrete high levels of IL-10 in response to the TLR9 agonist CpG, while myeloid DCs secrete much less [17]. Myeloid DCs can secrete IL-10 upon activation of TLR2, TLR4, TLR9, and also the C-type lectins DC-SIGN and Dectin-1 [10,18–20]. Thus, there are diverse pathways leading to IL-10 secretion. The timing of IL-10 secretion and cell types that respond impacts the generation of adaptive immune responses [21].

The bacterial components and signaling pathways that lead to induction of IL-10 following infection with  $\Delta hly$  *L. monocytogenes* have not been experimentally addressed. In this study, we conducted a genetic screen to identify the components of vacuole-confined  $\Delta hly$  *L. monocytogenes* that induce IL-10 secretion from murine bone marrow-derived macrophages. We investigated the host signaling pathways that lead to recognition of  $\Delta hly$  *L. monocytogenes* and secretion of IL-10 in macrophages and mice, and how these host signaling pathways affect vaccination and immune suppression.


## Results

### Genetic screen to identify *L. monocytogenes* mutants that induce enhanced or diminished levels of IL-10

The goal of this study was to identify *L. monocytogenes* determinants that contribute to induction of IL-10 secretion from BMMs. We screened a library of  $\Delta hly$  transposon mutants for their ability to induce IL-10 secretion from BMMs. The transposon library was generated in a flagellin-negative  $\Delta hly$  background ( $\Delta hly\Delta fla$ ) to eliminate the identification of false low-IL-10 mutants resulting from mutations in flagellar components that reduce infection efficiency. Previous work demonstrated that infection of mice with  $\Delta hly$  is immunosuppressive and IL-10 levels are increased in the serum four hours post-infection [9], while infection with WT leads to IL-10 that peaks 3–4 days post-infection [12,13]. To determine whether significant IL-10 is secreted by *L. monocytogenes*-infected BMMs four hours post-infection, BMMs were infected with  $\Delta hly$  and  $\Delta hly\Delta fla$  *L. monocytogenes*, and IL-10 was measured from cell supernatants by ELISA. BMMs infected with both  $\Delta hly$  and  $\Delta hly\Delta fla$  *L. monocytogenes* secreted significantly more IL-10 than uninfected BMMs four hours post-infection (Fig 1A). As *L. monocytogenes* infection leads to significant expression of IL-10 in both BMMs and mice four hours post-infection, we analyzed cell supernatants for IL-10 levels four hours post-infection in our screen.

BMMs were infected in duplicate with 6000 gridded  $\Delta hly\Delta fla$  transposon mutants in 96-well plates (Fig 1B). 200 mutants that induced enhanced or diminished IL-10 levels compared to  $\Delta hly\Delta fla$  were selected for secondary screening. For secondary screening, optical density of bacterial cultures was adjusted to eliminate differences in IL-10 secretion due to growth differences in broth. 59 of the 200 mutants induced significantly enhanced or diminished IL-10 levels compared to  $\Delta hly\Delta fla$  in a 24-well format (S1 Table). The transposons in 58 mutants were phage-transduced from the  $\Delta hly\Delta fla$  background into a  $\Delta hly$  background. In the  $\Delta hly$  background, only two mutants induced less IL-10 secretion and four mutants induced increased IL-10 secretion (Table 1).

Notably, a mutant with a transposon insertion in phosphatidylglycerol-prolipoprotein diacylglycerol transferase (*lgt*), induced nearly no IL-10 secretion, though it entered cells and survived in cells similar to  $\Delta hly$  (Fig 1C). *Lgt* catalyzes the transfer of a lipid moiety from phosphatidylglycerol onto a cysteine residue of prolipoproteins [22–24]. The resulting



**Fig 1. Vacuole-confined *L. monocytogenes* induce IL-10 secretion from BMMs.** (A) BMMs were infected with *L. monocytogenes* lacking LLO ( $\Delta hly$ ) or lacking LLO and flagellin ( $\Delta hly\Delta fla$ ) for four hours. Cell supernatant was collected and IL-10 was measured by ELISA. Data shown are representative of two independent experiments. Mean and SD are shown.  $n = 3$ . Data analyzed using Holm-Sidak's multiple comparisons test. (B) Schematic of screen. 6000  $\Delta hly\Delta fla$  transposon mutants were analyzed for enhanced or diminished IL-10 expression in duplicate in a 96-well plate format. 200 mutants were selected for secondary screening, and were analyzed for IL-10 in triplicate in a 24-well plate format. The mutations from 58 mutants that induced significantly enhanced or diminished IL-10 were transduced into a  $\Delta hly$  background and analyzed for IL-10 secretion. Six mutations that resulted in enhanced or diminished IL-10 secretion were identified. (C and D) Survival of transposon mutants was quantified in BMMs. Mean and SEM are shown. Data are pooled from three coverslips per experiment for three independent experiments.

<https://doi.org/10.1371/journal.ppat.1008622.g001>

**Table 1. IL-10 secretion four hours post-infection of BMMs with deletion mutants and transposon mutants identified in genetic screen.**

Diminished IL-10	Gene Annotation	IL-10 (% $\Delta hly$ )	Strain Number
$\Delta lgt$	<i>lgt</i>	30	DP-L7003
<i>Lmo2482::Tn</i>	<i>lgt</i>	27	DP-L7111
<i>Lmo0709::Tn</i>	hypothetical protein	41	DP-L7080
Enhanced IL-10	Gene Annotation	IL-10 (% $\Delta hly$ )	Strain Number
$\Delta pgdA\Delta oatA$	<i>pgdA/oatA</i>	263	DP-L7004
<i>Lmo0415::Tn</i>	<i>pgdA</i>	177	DP-L7074
<i>Lmo2529::Tn</i>	ATP synthase F0F1 subunit beta	158	DP-L7112
<i>Lmo2634::Tn</i>	<i>ecfT</i>	140	DP-L7116
<i>Lmo1291::Tn</i>	<i>oatA</i>	139	DP-L7089

<https://doi.org/10.1371/journal.ppat.1008622.t001>

lipoproteins are well-known TLR2 agonists [25]. Of the two mutants that induced reduced IL-10 secretion, we focused on understanding the contribution of *lgt* to the induction of IL-10 secretion from BMMs because the other mutant, which had a transposon insertion in *lmo0709*, had reduced infection capability (Fig 1C).


To confirm the role of *lgt* in the reduced IL-10 phenotype, an in-frame deletion of *lgt* was generated in a  $\Delta hly$  background. Deletion of *lgt* did not affect infection efficiency or survival in BMMs (Fig 2B). An *lgt*-deletion mutant induced significantly reduced IL-10 secretion from WT BMMs compared to  $\Delta hly$ . IL-10 levels were restored when *lgt* was complemented in  $\Delta hly$  *L. monocytogenes* under control of the constitutively active pHyper promoter (Fig 2A). These data indicated that  $\Delta hly$  *L. monocytogenes* primarily induces IL-10 secretion in a lipoprotein- and TLR2-dependent manner.

Of the mutants that induced increased IL-10 secretion, mutations in two of the genes, *pgdA* and *oatA*, have previously been shown to render bacteria more susceptible to lysozyme-mediated killing [26]. We hypothesized that these mutants had increased bacterial lysis within host phagosomes, leading to greater activation of endosomal nucleic acid-sensing TLRs. Though the mutations did not contribute to a noticeable survival defect individually (Fig 1D), a strain lacking both *pgdA* and *oatA* ( $\Delta hly\Delta pgdA\Delta oatA$ ) induced significantly more IL-10 than  $\Delta hly$  and had a significant survival defect in BMMs (Fig 2C and 2D), indicative of bacterial death in vacuoles. We hypothesized that lysis of bacteria in the phagocytic vacuole could result in release of bacterial nucleic acids and activation of nucleic acid-sensing TLRs.

### IL-10 secretion in BMMs

While infection with  $\Delta hly$  induced significant IL-10 secretion from WT BMMs,  $\Delta hly$  induced almost no detectable IL-10 secretion in TLR2<sup>-/-</sup> BMMs (Fig 3A), suggesting that bacterial lipoproteins are the major activators of IL-10 secretion in response to infection with  $\Delta hly$ . In contrast,  $\Delta hly\Delta pgdA\Delta oatA$  induced significant levels of IL-10 secretion in TLR2<sup>-/-</sup> BMMs (Fig 3B), suggesting that bacteriolysis can stimulate a second pathway of IL-10 induction. To determine whether this second pathway was dependent on endosomal nucleic acid-sensing TLRs, we infected BMMs with a mutation in the endosomal TLR trafficking protein Unc93b1, which is essential for endosomal TLR signaling [27]. In cells from mice that have a mutation in Unc93b1, nucleic acid-sensing TLRs do not traffic properly from the endoplasmic reticulum to endolysosomes. As a result, signaling through nucleic acid-sensing TLRs is abrogated, but other TLRs retain normal signaling [28,29].  $\Delta hly$  induced similar amounts of IL-10 in WT and Unc93b1<sup>3d/3d</sup> BMMs (Fig 3A), whereas  $\Delta hly\Delta pgdA\Delta oatA$  induced significantly less IL-10 secretion in Unc93b1<sup>3d/3d</sup> BMMs (Fig 3B).






**Fig 2. Induction of IL-10 is mediated by lipoproteins and bacteriolysis.** (A) IL-10 secretion from BMMs four hours-post infection. Mean and SD are shown. Data are representative of two independent experiments. Data analyzed using Holm-Sidak's multiple comparison's test. (B) Survival of *lgt* deletion mutant in BMMs. Data is representative of two independent experiments. Mean and SEM are shown. (C) IL-10 secretion from BMMs four hours-post infection. Mean and SD are shown. Data are representative of three independent experiments. Data analyzed using Holm-Sidak's multiple comparison's test. (D) Survival of  $\Delta hly\Delta pgdA\Delta oatA$  mutant in BMMs. Data are pooled from three independent experiments. Mean and SEM are shown.

<https://doi.org/10.1371/journal.ppat.1008622.g002>

To confirm that IL-10 secretion in response to vacuole-confined *L. monocytogenes* resulted from signaling through TLR2 and endosomal TLRs, we infected BMMs with mutations in both pathways ( $TLR2^{-/-}TLR5^{-/-}Unc93b1^{3d/3d}$ ). These macrophages secreted IL-10 in response to LPS, which is a TLR4 ligand, but not in response to the *L. monocytogenes* mutants that were tested (Fig 3C). These results confirmed that IL-10 secretion from BMMs results from signaling through TLR2 and endosomal TLRs.



**Fig 3. IL-10 secretion from BMMs in response to *L. monocytogenes* infection requires TLR2 and endosomal TLR signaling.** (A) WT, TLR2<sup>-/-</sup>, Unc93b1<sup>3d/3d</sup>, and TLR2<sup>-/-</sup>TLR5<sup>-/-</sup>Unc93b1<sup>3d/3d</sup> BMMs were infected with  $\Delta hly$  *L. monocytogenes*. Cell supernatant was harvested four hours post infection and IL-10 was measured by ELISA. Data shown are representative of two independent experiments. Means were compared to WT using Holm-Sidak's Multiple Comparisons test. (B) WT, TLR2<sup>-/-</sup>, Unc93b1<sup>3d/3d</sup>, and TLR2<sup>-/-</sup>TLR5<sup>-/-</sup>Unc93b1<sup>3d/3d</sup> BMMs were infected with  $\Delta hly\Delta pgdA\Delta oatA$  *L. monocytogenes*. Cell supernatant was harvested four hours post infection and IL-10 was measured by ELISA. Data shown are representative of two independent experiments. Means were compared to WT using Holm-Sidak's Multiple Comparisons test. (C) TLR2<sup>-/-</sup>TLR5<sup>-/-</sup>Unc93b1<sup>3d/3d</sup> BMMs were infected with the indicated strains of *L. monocytogenes*, or treated with lipopolysaccharide (LPS), a TLR4 agonist; CpG ODN 1668 (CpG), a TLR9 agonist; or Pam2CSK4 (PAM), a TLR2 agonist. Data shown are pooled from two independent experiments. Data analyzed using Holm-Sidak's Multiple Comparisons test.

<https://doi.org/10.1371/journal.ppat.1008622.g003>



## Cytokine secretion in mice


To determine whether secretion of IL-10 in mice was dependent on TLR2 and/or endosomal TLRs, mice were infected intravenously and serum IL-10 was quantified four hours post-infection (Fig 4A–4D). Infection of WT mice with  $\Delta hly$  resulted in significant levels of IL-10, but unlike in BMMs  $\Delta hly$  also induced significant amounts of IL-10 in TLR2<sup>-/-</sup> mice (Fig 4B), suggesting that bacterial lipoproteins were not the dominant IL-10-inducing molecules in mice. In contrast, IL-10 secretion following  $\Delta hly$  *L. monocytogenes* infection was reduced in Unc93b1<sup>3d/3d</sup> mice and almost undetectable in TLR2<sup>-/-</sup>TLR5<sup>-/-</sup>Unc93b1<sup>3d/3d</sup> mice (Fig 4C and 4D). Together, these results indicate that endosomal TLRs were the primary mediators of IL-10 secretion in response to  $\Delta hly$  infection in mice, but that TLR2 may also contribute to induction of IL-10.

One function of IL-10 signaling is to limit proinflammatory cytokine secretion. To investigate the relationship between TLRs and proinflammatory cytokine signaling, serum levels of MCP-1 were quantified 24 hours post-infection (Fig 4E–4H). In WT mice,  $\Delta hly$  did not induce secretion of MCP-1, while  $\Delta actA$ , a strain of *L. monocytogenes* that escapes the phagocytic vacuole and replicates in the host cell cytosol, induced significant amounts of MCP-1 (Fig 4E). Likewise,  $\Delta actA$ , but not  $\Delta hly$ , induced significant MCP-1 secretion in TLR2<sup>-/-</sup> mice (Fig 4F). Because early IL-10 expression is thought to limit later expression of proinflammatory cytokines, we hypothesized that  $\Delta hly$  would induce proinflammatory cytokines in Unc93b1<sup>3d/3d</sup> and TLR2<sup>-/-</sup>TLR5<sup>-/-</sup>Unc93b1<sup>3d/3d</sup> mice. However, MCP-1 was not induced in response to  $\Delta hly$  background strains in Unc93b1<sup>3d/3d</sup> and TLR2<sup>-/-</sup>TLR5<sup>-/-</sup>Unc93b1<sup>3d/3d</sup> mice (Fig 4G and 4H). These results indicated that endosomal TLR signaling may be required for  $\Delta hly$  to induce a proinflammatory cytokine response.

Vaccination of mice with  $\Delta actA$  has previously been shown to result in high levels of proinflammatory cytokines, whereas vaccination of mice with  $\Delta hly$  induced high levels of IL-10 and low levels of proinflammatory cytokines in WT mice. Strikingly, co-administration of  $\Delta actA$  and  $\Delta hly$  resulted in high levels of IL-10 and low levels of proinflammatory cytokines in WT mice [9], suggesting that  $\Delta hly$  suppresses the proinflammatory responses normally induced by  $\Delta actA$ . We analyzed the levels of MCP-1 in serum as a measure of the proinflammatory cytokine response. It was previously reported that  $\Delta actA$  induced significant amounts of MCP-1 in WT mice, but that co-administration of  $\Delta actA$  with 1000-fold more  $\Delta hly$  CFU resulted in reduced expression of MCP-1 [9]. We observed that  $\Delta actA$  induced significant amounts of MCP-1 in WT mice, and that co-administration of  $\Delta actA$  and  $\Delta hly$  resulted in low levels of MCP-1 secretion, consistent with previous findings (Fig 4E). MCP-1 levels following co-administration of  $\Delta actA$  and  $\Delta hly$  were partially restored to  $\Delta actA$  alone levels in TLR2<sup>-/-</sup> and Unc93b1<sup>3d/3d</sup> mice, suggesting that the TLR2 and endosomal TLR signaling pathways contribute to suppression of proinflammatory cytokines (Fig 4F and 4G). In TLR2<sup>-/-</sup>TLR5<sup>-/-</sup>Unc93b1<sup>3d/3d</sup> mice (Fig 4H), co-administration of  $\Delta actA$  and  $\Delta hly$  resulted in levels of MCP-1 secretion similar to those induced by  $\Delta actA$  alone. These results indicate that suppression of proinflammatory cytokines by  $\Delta hly$  is mediated by TLR2 and Unc93b1-dependent TLRs.

## Suppression of adaptive immunity

To investigate the relationship between IL-10 signaling pathways and suppression of adaptive immunity, co-vaccination experiments were performed in which  $\Delta actA$  and 1000-fold more  $\Delta hly$  CFU were injected simultaneously as described [9] (Fig 5). In WT mice, vaccination with  $\Delta actA$  led to the induction of robust adaptive immunity that nearly cleared a subsequent lethal dose of WT *L. monocytogenes*. However, the immunity induced by  $\Delta actA$  was suppressed two-




**Fig 4.** *L. monocytogenes* induces IL-10 and MCP-1 secretion in mice. Mice were infected with  $10^8$  CFU of  $\Delta hly$ ,  $10^5$  CFU of  $\Delta actA$ , or a combination of  $10^8$  CFU of  $\Delta hly$  and  $10^5$  CFU of  $\Delta actA$ . WT C57BL/6J mice (A and E), TLR2<sup>-/-</sup> (B and F), Unc93b1<sup>3d/3d</sup> (C and G), and TLR2<sup>-/-</sup>TLR5<sup>-/-</sup>Unc93b1<sup>3d/3d</sup> (D and H) were infected. Serum measurements of IL-10 four hours post-infection (A-D) and MCP-1 twenty-

four hours post-infection (E-H). Data are pooled from two to four independent experiments. Bar represents the median. Data analyzed using Holm-Sidak's Multiple Comparisons test.

<https://doi.org/10.1371/journal.ppat.1008622.g004>

logs by co-administration of  $\Delta hly$  (Fig 5A). This suppression was previously shown to depend on IL-10 [8]. We hypothesized that  $\Delta hly$  would suppress immunity in WT and  $TLR2^{-/-}$  mice,



**Fig 5.** Immune suppression is primarily mediated by endosomal TLR signaling. Mice were infected with  $10^8$  CFU of  $\Delta hly$  background strains or  $10^5$  CFU of  $\Delta actA$ . WT C57BL/6] mice (A),  $TLR2^{-/-}$  (B),  $Unc93b1^{3d/3d}$  (C), and  $TLR2^{-/-}TLR5^{-/-}Unc93b1^{3d/3d}$  (D) were infected. Eight weeks post-vaccination, mice were challenged with  $5 \times 10^5$  WT *L. monocytogenes*. CFU from the spleen were enumerated three days post-challenge. Data are pooled from two to four independent experiments. Bar represents the median. Data analyzed using an unpaired t test.

<https://doi.org/10.1371/journal.ppat.1008622.g005>

in which we observed significant secretion of IL-10 and low levels of MCP-1, but not in Unc93b1<sup>3d/3d</sup> or TLR2<sup>-/-</sup>TLR5<sup>-/-</sup>Unc93b1<sup>3d/3d</sup> mice, in which we observed no IL-10 and high levels of MCP-1. That was indeed the case, as we observed 2-logs of immune suppression in WT and TLR2<sup>-/-</sup> mice (Fig 5A and 5B). However, in Unc93b1<sup>3d/3d</sup> mice co-administration of *ΔactA* with *Δhly* did not significantly reduce protective immunity (Fig 5C). In TLR2<sup>-/-</sup>TLR5<sup>-/-</sup>Unc93b1<sup>3d/3d</sup> mice, we observed a 4-fold reduction in protective immunity (Fig 5D).

## Discussion

The results of this study demonstrate that the secretion of IL-10 by BMMs is largely mediated by TLR2-detection of bacterial lipoproteins. However, analysis of mutants that induced enhanced IL-10 revealed a second pathway that was dependent on triggering of Unc93b1-dependent nucleic-acid sensing TLRs in phagosomes by bacteriolysis. In mice, TLR2 and nucleic acid-sensing TLRs contributed to IL-10 secretion.

The results of our analysis of BMMs indicates that TLR2 and Unc93b1-dependent TLRs represent two independent pathways that lead to IL-10 expression. However, our data in mice indicates that TLR2-signaling does not contribute to immune suppression to the same extent as endosomal TLRs. TLR2-deficient mice expressed significant IL-10 following infection with *Δhly* (Fig 4B). In contrast, infection of Unc93b1<sup>3d/3d</sup> mice resulted in significant but reduced IL-10 levels compared to WT mice (Fig 4C). Only TLR2<sup>-/-</sup>TLR5<sup>-/-</sup>Unc93b1<sup>3d/3d</sup> mice failed to express significant IL-10 after *Δhly* infection (Fig 4D). The reduction in IL-10 correlated with a reduction in immunosuppression, indicating that although TLR2 and endosomal TLRs mediate IL-10 induction, endosomal TLRs may be the major mediators of immune suppression.

The differences in TLR contributions to IL-10 expression in BMMs compared to mice likely reflects differences in their capacity to kill and degrade *L. monocytogenes*, but may also reflect differences in TLR expression levels. It is possible that *L. monocytogenes*-infected cells in mice express less TLR2 than BMMs. For example, mouse peritoneal macrophages express very low levels of surface TLR2 compared to BMMs [30]. Reduced TLR2 expression on *L. monocytogenes*-infected cells in mice would explain why we observed a smaller contribution of TLR2 to IL-10 expression in mice. Additionally, increased bacterial killing in mice likely explains the greater contribution of endosomal TLRs to IL-10 expression in mice compared to BMMs. In mice, CD169+ macrophages, are thought to be the first cells to capture *L. monocytogenes* in the spleen and restrict their multiplication [15,31]. In BMMs, which are differentiated and cultured *ex vivo*, *Δhly* *L. monocytogenes* cannot grow but are not efficiently killed (Fig 2B). In BMMs, in the absence of bacterial killing bacterial lipoproteins are the most abundant TLR stimulus. However, bacterial mutants that had increased lysis in BMMs induced Unc93b1-dependent IL-10 secretion. In contrast to BMMs, peritoneal macrophages taken directly from mice kill vacuole-confined bacteria, suggesting that *L. monocytogenes* may undergo bacteriolysis *in vivo* more than in BMMs [32,33]. In mice, we surmise that increased bacterial killing leads to release of nucleic acids and lipoproteins that are sensed by endosomal TLRs and TLR2, respectively. Therefore, secretion of IL-10 in mice is likely dependent on lipoprotein-dependent TLR2 activation and Unc93b1-dependent nucleic-acid sensing TLRs because of increased bacteriolysis in mice.

It was shown previously that administration of anti-IL-10 receptor blocking antibody restored the protective potential of *Δhly*. However, in this study we observed that, in mice with a defect in endosomal TLR signaling, vaccination with *Δhly* *L. monocytogenes* did not induce IL-10 secretion but still did not lead to the generation of a protective immune response. A proinflammatory response was also lacking in Unc93b1 mutant mice following vaccination

with  $\Delta hly$ , which may explain why a protective immune response was not generated. The differences in immunity following vaccination of WT mice with  $\Delta hly$  and anti-IL10R blocking antibody and vaccination of Unc93b1 mutant mice with  $\Delta hly$  may reflect the fact that anti-IL10R blocking antibody only blocks IL-10 signaling, while mice with defects in endosomal TLR signaling may not be able to generate a proinflammatory response to  $\Delta hly$  and are thus inhibited in generating protective immunity.

Although the results of our study suggest that recognition of bacterial lipoproteins by TLR2 is not the only source of IL-10 *in vivo*, TLR2 remains an attractive pattern recognition receptor to consider for the development of bacterial vaccine delivery systems. *L. monocytogenes* expresses over 30 diacylated lipoproteins that are recognized by TLR2/TLR6 dimers [34–37]. Deletion of *lgt* in WT *L. monocytogenes* results in delayed secretion of proinflammatory cytokines, but only a modest virulence defect in mice [38]. Thus, removing lipoproteins is a promising strategy for improving *L. monocytogenes*-based vaccines by changing the cytokine response without significantly affecting bacterial fitness. One potential method of modulating the response to lipoproteins that has not been explored in *L. monocytogenes* is engineering *L. monocytogenes* lipoproteins with different acylation states. Triacylated lipoproteins are recognized by TLR2/TLR1 heterodimers and elicit a different inflammatory response than diacylated lipoproteins [39]. More recently, three additional classes of lipoprotein modification with differing abilities to stimulate TLR2 have been described in Gram-positive bacteria [34]. It would be interesting to investigate the contributions of differently acylated lipoproteins to the development of adaptive immunity. Modulating the expression of individual lipoproteins could also impact the immune response. It is not clear whether all *L. monocytogenes* lipoproteins contribute to TLR2 activation, or whether *L. monocytogenes* has a subset of lipoproteins that specifically activate TLR2. Though the lipid portion of lipoproteins mediates binding of lipoproteins to TLR2, the sequence of the attached peptides affects the ability of lipoproteins to differentially stimulate cytokine secretion [40]. Therefore, changing the expression of specific lipoproteins could be a strategy for fine-tuning the proinflammatory immune response. In addition, TLR2 is traditionally considered a cell-surface-localized TLR, but there is evidence that lipoproteins can activate TLR2 signaling in the endosomal compartment and that signaling from the endosomal compartment can have different effects than from the plasma membrane [41–45]. How specific lipoproteins and acylation state contribute to activation of TLR2, and how TLR2 activation on the cell surface versus in the lysosome contributes to the cytokine response is relevant for the development of bacterial vaccine vectors and warrants further study.

The results of this study indicated that, in mice, TLR2 and endosomal TLRs sense vacuole-confined *L. monocytogenes*, resulting in the secretion of IL-10. In our genetic screen, we identified mutants that had increased IL-10 secretion in WT, TLR2<sup>-/-</sup>, and Unc93b1<sup>3d/3d</sup> macrophages (S2 Table), including transposon insertions in *pgdA* and *oatA*, genes for which mutations have previously been shown to confer increased lysozyme sensitivity [46]. Enhanced IL-10 expression in response to *L. monocytogenes* mutants was dependent on both TLR2 and endosomal TLRs, as there were no mutants that induced noteworthy IL-10 secretion in TLR2<sup>-/-</sup>TLR5<sup>-/-</sup>Unc93b1<sup>3d/3d</sup> BMMs (S2 Table), suggesting that bacterial lysis releases nucleic acids and lipoproteins that are both sensed by the cell. Though we did not observe a decrease in colony forming units of  $\Delta hly$  or the *pgdA* or *oatA* transposon mutants in macrophages (Fig 1D),  $\Delta hly\Delta pgdA\Delta oatA$  died rapidly (Fig 2D). Therefore, the individual mutants for which we observed increased IL-10 likely have minor increases in lysis that were not easily detected by CFU. However, in combination, the *pgdA* and *oatA* mutations synergized to yield a mutant that was extremely susceptible to lysis in cells and potently induced IL-10. Importantly, these



data suggest that low-level lysis can be detected by TLRs and lead to significant changes in cytokine secretion.

The majority of mutants identified in our screen induced increased IL-10 secretion. Strikingly, while 50 mutants that induced increased IL-10 secretion were identified in our initial screen in  $\Delta hly\Delta fla$  *L. monocytogenes* (S1 Table), only 4 of those mutants induced significantly increased IL-10 secretion in a  $\Delta hly$  background (S2 Table). It is possible that  $\Delta hly$  may be slightly more sensitive to lysis compared to  $\Delta hly\Delta fla$ . Flagellar secretion systems require the activity of hydrolases or lytic transglycosylases to degrade peptidoglycan to allow insertion of flagellar components [47,48]. Thus it is reasonable to suspect that the cell wall of  $\Delta hly$  is more fragile than that of  $\Delta hly\Delta fla$  and is more susceptible to lysis inside vacuoles due to the activity of a flagella-associated peptidoglycan-degrading enzyme. An increase in the basal levels of lysis would increase the basal levels of IL-10 and potentially mask the effects of other mutations that slightly increased lysis. The results of this study suggest that any mutation that promotes bacterial lysis within a phagosome will lead to IL-10 induction.

In this study, we did not investigate whether a particular endosomal nucleic-acid sensing TLR is responsible for induction of IL-10. It is likely that all nucleic-acid sensing TLRs can be activated following bacterial lysis. Bacterial lysis should lead to the release of bacterial contents including: mRNA that can activate TLR7/8, unmethylated CpG chromosomal DNA that can activate TLR9, and ribosomal RNA that can activate TLR13 [49]. Although TLR3 could also recognize double-stranded RNA released upon bacterial lysis, it likely does not contribute to IL-10 secretion because it was previously demonstrated that IL-10 secretion in response to vacuole-confined *L. monocytogenes* is dependent on the signaling adapter MyD88, and TLR3 uses the signaling adapter TRIF [9,49]. It has been previously suggested that bacterial mRNA represents a signature of bacterial viability [50]. Perhaps, then, simultaneous recognition of multiple types of nucleic acid by multiple TLRs is indicative of nonviable bacteria.

The observation that the immune system responds more robustly to bacteria that are alive compared to bacteria that are dead led to the idea that the immune system has ways of monitoring bacterial viability [50–52]. Many signals have been proposed to be “PAMPs per vita” or “vita-PAMPs”—signatures of microbial viability. DNA and RNA have been proposed as possible “PAMP postmortem” (PAMP-PM). In line with the idea of recognition of postmortem PAMPs, the secretion of IL-10 following infection of mice with  $\Delta hly$  could represent a strategy to prevent an unnecessary immune response to bacteria that are already dead, and thus do not pose a threat. To that end, nucleic acid-sensing TLRs are better suited for assessing bacterial viability than TLR2, because living bacteria do not normally release nucleic acids, especially chromosomal DNA, into the surrounding environment. Signaling through nucleic-acid sensing TLRs thus more accurately indicates that a bacterium is dead than sensing of lipoproteins, which can be detected whether a pathogen is alive or dead.

Induction of IL-10 secretion is an important factor to consider in the development of bacterial vaccine vectors. Both live and dead bacterial vaccine vectors have the potential to induce IL-10 secretion. The kinetics of IL-10 secretion may play an important role in determining whether a vaccine will be effective or not. In our study, we identified sensing of nucleic acids as the primary signal for IL-10 induction. For the development of future vaccine strains, strategies to minimize IL-10 induction and immune suppression should be considered. For example, constructs that modify the cell wall could be employed to reduce bacteriolysis in phagosomes. Also, as suggested above, simply deleting flagellin may decrease bacteriolysis. Mutations could also be made to alter lipoproteins and enhance TLR2 activation and proinflammatory cytokine expression. In combination, modifications that reduce IL-10 secretion and modulate proinflammatory cytokines downstream of TLR2 may yield a vaccine strain that has increased potency.



## Materials and methods

### Strain construction

In-frame deletion of genes was performed using allelic exchange as previously described [53].  $\Delta hly\Delta fla$  was generated by deleting *hly* in a  $\Delta flaA$  strain (DP-L5986).  $\Delta hly\Delta lgt$  was generated by deleting *lgt* in a  $\Delta hly$  strain [54].  $\Delta hly\Delta fla\Delta lgt$  was generated by deleting *lgt* in a  $\Delta hly\Delta fla$  strain.  $\Delta hly\Delta pgdA\Delta oatA$  was generated by deleting *hly* in a  $\Delta pgdA\Delta oatA$  strain (DP-L5220).  $\Delta hly\Delta fla\Delta lgt$  was generated by deleting *lgt* in a  $\Delta hly\Delta fla$  strain. The *lgt* complemented strains were generated by integrating a pPL2 vector encoding *lgt* under control of the pHyper promoter (pPL2t-*pHyper-lgt*) into the *L. monocytogenes* genome and selecting for tetracycline-resistant transconjugates [55]. Strains used in this study are listed in S3 Table.

### Transposon library generation

A transposon library was generated in  $\Delta hly\Delta fla$  as previously described [56]. Transposon mutations were transduced into  $\Delta hly$  using U153 phage as previously described [57].

### Bone marrow-derived macrophage culture

BMM growth media was prepared using high glucose DMEM (Thermo Fisher Scientific) with 20% Fetal Bovine Serum (Seradigm), 1% L-glutamine (Corning), 1% Sodium pyruvate (Corning), 14mM 2-Mercaptoethanol (Gibco), and 10% 3T3 cell supernatant (from M-CSF-producing 3T3 cells). Macrophages were prepared from the femurs of C57BL/6J mice. Femurs were isolated, sterilized with 70% ethanol, and crushed with a mortar and pestle in BMM growth media. Cells were strained through a 70 $\mu$ M filter and distributed into ten 150-mm non-TC dishes in 30mL BMM culture medium. An additional 30mL BMM culture medium was added at day 3. After cells were incubated for a total of seven days at 37°C with 5% CO<sub>2</sub>, cells were harvested and frozen at -80°C in BMM culture medium with 10% Fetal Bovine Serum and 10% DMSO (Sigma) added.

### Intracellular growth of *L. monocytogenes* in BMMs

$3 \times 10^6$  BMMs were plated in 60 mm non-TC-treated Petri dishes with 14 12mm glass coverslips in each dish. Dishes were infected with  $5 \times 10^5$  CFU (MOI = 0.17) and intracellular growth curves were performed as described previously [58].

### TLR agonists

For experiments using TLR agonists, cells were incubated with the agonists for the entire duration of the experiment. The sequence of CpG ODN 1668 (Integrated DNA Technologies) is: T\*C\*C\*A\*T\*G\*A\*C\*G\*T\*T\*C\*C\*T\*G\*A\*T\*G\*C\*T, with asterisks indicating phosphorothioate modifications. CpG ODN 1668 was used at a final concentration of 10 $\mu$ M. Pam2CSK4 (InvivoGen, Cat. tlrl-pm2s-1) was used at a final concentration of 100ng/mL. LPS (InvivoGen, Cat. tlrl-eklps) was used at a final concentration of 50ng/mL.

### Quantification of IL-10 from BMMs– 96-well format

$2.6 \times 10^5$  BMMs in 200  $\mu$ L BMM growth media were seeded into wells of a 96-well plate. Bacteria were grown in 1 mL Brain-Heart Infusion Broth containing 200 $\mu$ g/mL streptomycin in 96-well deep well plates at 3°C. Prior to infection, bacteria were pelleted by centrifugation and resuspended in 1 mL PBS. Wells were infected with 8  $\mu$ L of resuspended bacteria. Plates were infected in duplicate. 30 minutes post-infection, cells were washed with warm PBS, and BMM

growth media with 50 µg/mL gentamicin was added. Supernatants were collected and frozen at -80°C until used for analysis. For quantification of IL-10, Mouse IL-10 DuoSet ELISA (R&D Systems) was performed according to manufacturer's instructions.

### Quantification of IL-10 from BMMs– 24-well format

$6 \times 10^5$ – $7 \times 10^5$  BMMs in 500 µL BMM growth media were seeded into wells of a 24-well plate. Bacteria were grown overnight at 30°C at a slant without shaking in 3 mL Brain-Heart Infusion Broth containing 200µg/mL streptomycin. Cultures were then pelleted by centrifugation, and resuspended in phosphate-buffered solution (PBS) to an optical density of 2.0. Wells were infected with 20µL of bacteria, approximately  $8 \times 10^7$  CFU (MOI = 120). Three wells were infected for each bacterial strain. 30 minutes post-infection, cells were washed with warm PBS, and BMM growth media with 50 µg/mL gentamicin was added. Supernatants were collected and frozen at -80°C until used for analysis. For quantification of IL-10, Mouse IL-10 DuoSet ELISA (R&D Systems) was performed according to manufacturer's instructions. Data was analyzed using GraphPad Prism.

### Animal use ethics statement

All animal work was done in strict accordance with university regulations. Protocols were reviewed and approved by the Animal Care and Use Committee at the University of California, Berkeley AUP-2016-05-8811.

### Mice

C57BL/6J mice were purchased from Jackson Laboratories. TLR2<sup>-/-</sup>, Unc93b1<sup>3d/3d</sup>, and TLR2<sup>-/-</sup>TLR5<sup>-/-</sup>Unc93b1<sup>3d/3d</sup> were provided by Greg Barton (UC Berkeley) and were bred in our facility. To investigate the combined effects of TLR2 and endosomal TLR mutations, TLR2<sup>-/-</sup>TLR5<sup>-/-</sup>Unc93b1<sup>3d/3d</sup> were used because TLR2<sup>-/-</sup>Unc93b1<sup>3d/3d</sup> were not available.

### Quantification of cytokines from serum

Eight-to-twelve week old female mice were injected via the tailvein with  $10^8$  CFU of  $\Delta hly$  background strains or  $10^5$  CFU  $\Delta actA$ . Bacteria were grown overnight at 30°C at a slant without shaking in Brain-Heart Infusion Broth containing 200µg/mL streptomycin. Bacteria were then backdiluted 1:20 and grown at 37°C shaking for about two hours until they reached an optical density of 0.5. Cultures were pelleted and resuspended in PBS to the appropriate concentration, such that mice were infected with 200µL. Four and twenty-four hours post infection, blood was collected from the submandibular vein into Microtainer tubes with serum separator additive (BD). Blood was left to rest for 30 minutes before tubes were centrifuged and serum was collected. Collected serum was stored at -20°C until analysis. Analysis of cytokines from serum was performed using Mouse Inflammation Cytokine Bead Arrays (BD, Cat. 552364). Data was analyzed using FlowJo.

### Vaccination and immune suppression

Eight-to-twelve week old C57BL/6 female mice were vaccinated intravenously via the tail vein with  $10^8$  CFU of  $\Delta hly$  background strains or  $10^5$  CFU  $\Delta actA$ . Bacteria were grown overnight at 30°C at a slant without shaking in Brain-Heart Infusion Broth containing 200µg/mL streptomycin. Bacteria were then backdiluted 1:20 and grown at 37°C shaking for about two hours until they reached an optical density of 0.5. Cultures were pelleted and resuspended in PBS to the appropriate concentration, such that mice were vaccinated with 200 µL. Eight weeks post-

vaccination, mice were challenged with  $5 \times 10^4$  CFU WT *L. monocytogenes*. Three days post-challenge, mice were euthanized with CO<sub>2</sub> and cervical dislocation and CFU in the spleens and livers were enumerated.

### Statistical analysis

Data were analyzed using GraphPad Prism 8. \* indicates  $P < 0.05$ ; \*\* indicates  $P < 0.01$ , \*\*\* indicates  $P < 0.001$ , \*\*\*\* indicates  $P < 0.0001$ ; ns indicates no statistical significance.

### Supporting information

**S1 Table. Secretion of IL-10 in response to infection with transposon mutants in  $\Delta hly\Delta fla$  background compared to  $\Delta hly\Delta fla$ .** BMMs were infected with *L. monocytogenes* at an MOI of 120 in a 24-well plate format. Infections were performed in triplicate for each strain. IL-10 secretion from BMMs was measured from the supernatants by ELISA. The mean amount of IL-10 secreted in response to infection with each  $\Delta hly\Delta fla$ -background transposon mutant is reported as a percentage of the mean IL-10 induced by infection with  $\Delta hly\Delta fla$ . (DOCX)

**S2 Table. Secretion of IL-10 in response to infection with transposon mutants in  $\Delta hly$  background compared to  $\Delta hly$ .** WT, TLR2<sup>-/-</sup>, TLR2<sup>-/-</sup>TLR5<sup>-/-</sup>Unc93b1<sup>3d/3d</sup>, and Unc93b1<sup>3d/3d</sup> BMMs were infected with *L. monocytogenes* at an MOI of 120 in a 24-well plate format. Infections were performed in triplicate for each strain. IL-10 secretion from BMMs was measured from the supernatants by ELISA. The mean amount of IL-10 secreted in response to infection with each  $\Delta hly$ -background transposon mutant is reported as a percentage of the mean IL-10 induced by infection with  $\Delta hly$ . <sup>α</sup> BMM background. <sup>μ</sup> IL-10 secretion  $< 80$ pg/mL. <sup>β</sup>  $\Delta hly$  transposon mutant IL-10 values were compared to  $\Delta hly$  IL-10 values using Dunnett's multiple comparisons test, and asterisks indicate level of significance. (DOCX)

**S3 Table. *L. monocytogenes* and *E. coli* strains used in this study.** (DOCX)

### Author Contributions

**Conceptualization:** Brittney N. Nguyen, Alfredo Chávez-Arroyo.

**Data curation:** Brittney N. Nguyen, Alfredo Chávez-Arroyo.

**Formal analysis:** Brittney N. Nguyen, Alfredo Chávez-Arroyo, Mandy I. Cheng, Maria Krasilnikov, Alexander Louie.

**Funding acquisition:** Brittney N. Nguyen, Daniel A. Portnoy.

**Investigation:** Brittney N. Nguyen, Alfredo Chávez-Arroyo, Mandy I. Cheng, Maria Krasilnikov, Alexander Louie.

**Methodology:** Brittney N. Nguyen, Alfredo Chávez-Arroyo, Daniel A. Portnoy.

**Project administration:** Brittney N. Nguyen, Alfredo Chávez-Arroyo, Daniel A. Portnoy.

**Resources:** Brittney N. Nguyen, Alfredo Chávez-Arroyo, Daniel A. Portnoy.

**Supervision:** Brittney N. Nguyen, Alfredo Chávez-Arroyo, Daniel A. Portnoy.

**Validation:** Brittney N. Nguyen, Alfredo Chávez-Arroyo.

**Visualization:** Brittney N. Nguyen, Alexander Louie.

**Writing – original draft:** Brittney N. Nguyen.

**Writing – review & editing:** Brittney N. Nguyen, Alexander Louie, Daniel A. Portnoy.

## References

1. Czuprynski CJ, Brown JF. Effects of purified anti-Lyt-2 mAb treatment on murine listeriosis: comparative roles of Lyt-2+ and L3T4+ cells in resistance to primary and secondary infection, delayed-type hypersensitivity and adoptive transfer of resistance. *Immunology* [Internet]. 1990 Sep [cited 2019 Jul 4]; 71(1):107–12. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/2120126> PMID: 2120126
2. MCGREGOR DD, KOSTER FT, MACKANESS GB. Biological Sciences: The Short Lived Small Lymphocyte as a Mediator of Cellular Immunity. *Nature* [Internet]. 1970 Nov [cited 2019 Jul 4]; 228(5274):855–6. Available from: <http://www.nature.com/articles/228855a0> <https://doi.org/10.1038/228855a0> PMID: 5477011
3. Witte CE, Archer KA, Rae CS, Sauer J-D, Woodward JJ, Portnoy DA. Innate Immune Pathways Triggered by *Listeria monocytogenes* and Their Role in the Induction of Cell-Mediated Immunity. *Adv Immunol* [Internet]. 2012 Jan 1 [cited 2019 Jul 3]; 113:135–56. Available from: <https://www.sciencedirect.com/science/article/pii/B9780123945907000026?via%3Dihub> <https://doi.org/10.1016/B978-0-12-394590-7.00002-6> PMID: 22244582
4. Williams MA, Schmidt RL, Lenz LL. Early events regulating immunity and pathogenesis during *Listeria monocytogenes* infection. *Trends Immunol* [Internet]. 2012 [cited 2019 Jul 4]; 33:488–95. Available from: <https://doi.org/10.1016/j.it.2012.04.007> PMID: 22677184
5. Pamer EG. Immune responses to *Listeria monocytogenes*. *Nat Rev Immunol* [Internet]. 2004; 4(10):812–23. Available from: <https://doi.org/10.1038/nri1461> PMID: 15459672
6. Zenewicz LA, Shen H. Innate and adaptive immune responses to *Listeria monocytogenes*: a short overview. *Microbes Infect* [Internet]. 2007 Aug 1 [cited 2019 Jul 4]; 9(10):1208–15. Available from: <https://www.sciencedirect.com/science/article/pii/S1286457907001840?via%3Dihub> <https://doi.org/10.1016/j.micinf.2007.05.008> PMID: 17719259
7. Berche P, Gaillard JL, Sansonetti PJ. Intracellular growth of *Listeria monocytogenes* as a prerequisite for in vivo induction of T cell-mediated immunity. *J Immunol* [Internet]. 1987 Apr 1 [cited 2019 Jul 4]; 138(7):2266–71. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/3104455> PMID: 3104455
8. Bahjat KS, Liu W, Lemmens EE, Schoenberger SP, Portnoy D a., Dubensky TW, et al. Cytosolic entry controls CD8+T-cell potency during bacterial infection. *Infect Immun*. 2006; 74(11):6387–97. <https://doi.org/10.1128/IAI.01088-06> PMID: 16954391
9. Bahjat KS, Meyer-Morse N, Lemmens EE, Shugart J a., Dubensky TW, Brockstedt DG, et al. Suppression of cell-mediated immunity following recognition of phagosome-confined bacteria. *PLoS Pathog*. 2009; 5(9).
10. Saraiva M, O'Garra A. The regulation of IL-10 production by immune cells. *Nat Rev Immunol* [Internet]. 2010 Mar 15 [cited 2019 Jul 4]; 10(3):170–81. Available from: <http://www.nature.com/articles/nri2711> <https://doi.org/10.1038/nri2711> PMID: 20154735
11. Clark SE, Schmidt RL, McDermott DS, Lenz LL. A Baff3/Nlrp3/IL-18 Axis Promotes Natural Killer Cell IL-10 Production during *Listeria monocytogenes* Infection. *Cell Rep* [Internet]. 2018 May 29 [cited 2019 Jun 26]; 23(9):2582–94. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/29847790> <https://doi.org/10.1016/j.celrep.2018.04.106> PMID: 29847790
12. Clark SE, Filak HC, Guthrie BS, Schmidt RL, Jamieson A, Merkel P, et al. Bacterial Manipulation of NK Cell Regulatory Activity Increases Susceptibility to *Listeria monocytogenes* Infection. Coers J, editor. *PLOS Pathog* [Internet]. 2016 Jun 13 [cited 2019 Aug 19]; 12(6):e1005708. Available from: <https://dx.plos.org/10.1371/journal.ppat.1005708> PMID: 27295349
13. Foulds KE, Rotte MJ, Seder RA. IL-10 Is Required for Optimal CD8 T Cell Memory following *Listeria monocytogenes* Infection. *J Immunol* [Internet]. 2006 [cited 2016 Aug 24]; 177(4):2565–74. Available from: <http://www.jimmunol.org/content/177/4/2565> <https://doi.org/10.4049/jimmunol.177.4.2565> PMID: 16888018
14. Dai WJ, Köhler G, Brombacher F. Both innate and acquired immunity to *Listeria monocytogenes* infection are increased in IL-10-deficient mice. *J Immunol* [Internet]. 1997 Mar 1 [cited 2019 Jun 26]; 158(5):2259–67. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/9036973> PMID: 9036973
15. Perez OA, Yeung ST, Vera-Licona P, Romagnoli PA, Samji T, Ural BB, et al. CD169+ macrophages orchestrate innate immune responses by regulating bacterial localization in the spleen. *Sci Immunol* [Internet]. 2017 Oct 6 [cited 2019 Aug 19]; 2(16):eaah5520. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/28986418> <https://doi.org/10.1126/sciimmunol.aah5520> PMID: 28986418

16. Waite JC, Leiner I, Lauer P, Rae CS, Barbet G, Zheng H, et al. Dynamic imaging of the effector immune response to listeria infection In Vivo. *PLoS Pathog*. 2011 Mar; 7(3).
17. Boonstra A, Rajsbaum R, Holman M, Marques R, Asselin-Paturel C, Pereira JP, et al. Macrophages and myeloid dendritic cells, but not plasmacytoid dendritic cells, produce IL-10 in response to MyD88- and TRIF-dependent TLR signals, and TLR-independent signals. *J Immunol* [Internet]. 2006 Dec 1 [cited 2019 Aug 14]; 177(11):7551–8. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/17114424> <https://doi.org/10.4049/jimmunol.177.11.7551> PMID: 17114424
18. Jang S, Uematsu S, Akira S, Salgame P. IL-6 and IL-10 Induction from Dendritic Cells in Response to *Mycobacterium tuberculosis* Is Predominantly Dependent on TLR2-Mediated Recognition. *J Immunol* [Internet]. 2004 Sep 1 [cited 2019 Jul 4]; 173(5):3392–7. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/15322203> <https://doi.org/10.4049/jimmunol.173.5.3392> PMID: 15322203
19. Geijtenbeek TBH, Van Vliet SJ, Koppel EA, Sanchez-Hernandez M, Vandenbroucke-Grauls CMJE, Appelmek B, et al. Mycobacteria target DC-SIGN to suppress dendritic cell function. *J Exp Med* [Internet]. 2003 Jan 6 [cited 2019 Aug 19]; 197(1):7–17. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/12515809> <https://doi.org/10.1084/jem.20021229> PMID: 12515809
20. Rogers NC, Slack EC, Edwards AD, Nolte MA, Schulz O, Schweighoffer E, et al. Syk-dependent cytokine induction by dectin-1 reveals a novel pattern recognition pathway for C type lectins. *Immunity*. 2005;
21. Couper KN, Blount DG, Riley EM. IL-10: The Master Regulator of Immunity to Infection. *J Immunol*. 2008 May 1; 180(9):5771–7. <https://doi.org/10.4049/jimmunol.180.9.5771> PMID: 18424693
22. Kovacs-Simon A, Titball RW, Michell SL. Lipoproteins of bacterial pathogens. *Infect Immun* [Internet]. 2011 Feb 1 [cited 2017 Sep 11]; 79(2):548–61. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/20974828> <https://doi.org/10.1128/IAI.00682-10> PMID: 20974828
23. Nakayama H, Kurokawa K, Lee BL, Lee BL. Lipoproteins in bacteria: structures and biosynthetic pathways. [cited 2018 Feb 23]; Available from: <http://onlinelibrary.wiley.com/store/10.1111/febs.12041/asset/febs12041.pdf;jsessionid=AB3F9040F3FF4B2E67ED5B99EFC975B4.f04t03?v=1&t=je0hobsk&s=c36e1ee9b7fe3e982143ecb41224d48ac0449292>
24. Sankaran K, Wu HC. Lipid modification of bacterial prolipoprotein. Transfer of diacylglycerol moiety from phosphatidylglycerol. *J Biol Chem* [Internet]. 1994 Aug 5 [cited 2019 Sep 4]; 269(31):19701–6. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/8051048> PMID: 8051048
25. Schenk M, Belisle JT, Modlin RL. TLR2 Looks at Lipoproteins. Vol. 31, *Immunity*. 2009. p. 847–9. <https://doi.org/10.1016/j.immuni.2009.11.008> PMID: 20064441
26. Rae CS, Geissler A, Adamson PC, Portnoy DA. Mutations of the *Listeria monocytogenes* peptidoglycan N-Deacetylase and O-acetylase result in enhanced lysozyme sensitivity, bacteriolysis, and hyperinduction of innate immune pathways. *Infect Immun*. 2011 Sep; 79(9):3596–606. <https://doi.org/10.1128/IAI.00077-11> PMID: 21768286
27. Tabeta K, Hoebe K, Janssen EM, Du X, Georgel P, Crozat K, et al. The Unc93b1 mutation 3d disrupts exogenous antigen presentation and signaling via Toll-like receptors 3, 7 and 9. *Nat Immunol*. 2006 Feb; 7(2):156–64. <https://doi.org/10.1038/ni1297> PMID: 16415873
28. Lee BL, Moon JE, Shu JH, Yuan L, Newman ZR, Schekman R, et al. UNC93B1 mediates differential trafficking of endosomal TLRs. *Elife*. 2013 Feb 19;2013(2).
29. Kim YM, Brinkmann MM, Paquet ME, Ploegh HL. UNC93B1 delivers nucleotide-sensing toll-like receptors to endolysosomes. *Nature*. 2008 Mar 13; 452(7184):234–8. <https://doi.org/10.1038/nature06726> PMID: 18305481
30. Zajd CM, Ziembra AM, Miralles GM, Nguyen T, Feustel PJ, Dunn SM, et al. Bone Marrow-Derived and Elicited Peritoneal Macrophages Are Not Created Equal: The Questions Asked Dictate the Cell Type Used. *Front Immunol* [Internet]. 2020 Feb 21 [cited 2020 Apr 30]; 11:269. Available from: <https://www.frontiersin.org/article/10.3389/fimmu.2020.00269/full> PMID: 32153579
31. Alvarez B, Martínez P, Yuste M, Poderoso T, Alonso F, Domínguez J, et al. Phenotypic and functional heterogeneity of CD169+ and CD163+ macrophages from porcine lymph nodes and spleen. *Dev Comp Immunol*. 2014 May 1; 44(1):44–9. <https://doi.org/10.1016/j.dci.2013.11.010> PMID: 24291017
32. Portnoy DA, Schreiber RD, Connelly P, Tilneys LG.  $\gamma$  Interferon limits access of listeria monocytogenes to the macrophage cytoplasm. *J Exp Med*. 1989 Dec 1; 170(6):2141–6. <https://doi.org/10.1084/jem.170.6.2141> PMID: 2511268
33. Herskovits AA, Auerbuch V, Portnoy DA. Bacterial Ligands Generated in a Phagosome Are Targets of the Cytosolic Innate Immune System. *PLoS Pathog* [Internet]. 2007 Mar [cited 2020 Apr 8]; 3(3):e51. Available from: <https://dx.plos.org/10.1371/journal.ppat.0030051> PMID: 17397264
34. Kurokawa K, Ryu KH, Ichikawa R, Masuda A, Kim MS, Lee H, et al. Novel bacterial lipoprotein structures conserved in low-GC content gram-positive bacteria are recognized by toll-like receptor 2. *J Biol Chem*. 2012 Apr 13; 287(16):13170–81. <https://doi.org/10.1074/jbc.M111.292235> PMID: 22303020



35. Mao G, Zhao Y, Kang X, Li Z, Zhang Y, Wang X, et al. Crystal structure of *E. coli* lipoprotein diacylglyceryl transferase. *Nat Commun* [Internet]. 2016 Jan 5 [cited 2017 Oct 10]; 7:10198. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/26729647> <https://doi.org/10.1038/ncomms10198> PMID: 26729647
36. Baumgärtner M, Kärst U, Gerstel B, Loessner M, Wehland J, Jänsch L. Inactivation of Lgt allows systematic characterization of lipoproteins from *Listeria monocytogenes*. *J Bacteriol* [Internet]. 2007 Jan 15 [cited 2017 Sep 21]; 189(2):313–24. Available from: <http://jb.asm.org/cgi/doi/10.1128/JB.00976-06> PMID: 17041050
37. Bierre H, Cossart P. *Listeria monocytogenes* Surface Proteins: from Genome Predictions to Function. *Microbiol Mol Biol Rev*. 2007 Jun 1; 71(2):377–97. <https://doi.org/10.1128/MMBR.00039-06> PMID: 17554049
38. Machata S, Tchatalbachev S, Mohamed W, Jänsch L, Hain T, Chakraborty T. Lipoproteins of *Listeria monocytogenes* Are Critical for Virulence and TLR2-Mediated Immune Activation. *J Immunol* [Internet]. 2008 [cited 2017 Sep 11]; 181(3). Available from: <http://www.jimmunol.org/content/181/3/2028.short>
39. Nguyen M-T, Uebele J, Kumari N, Nakayama H, Peter L, Ticha O, et al. Lipid moieties on lipoproteins of commensal and non-commensal staphylococci induce differential immune responses. *Nat Commun* [Internet]. 2017 Dec 21 [cited 2018 Apr 18]; 8(1):2246. Available from: <http://www.nature.com/articles/s41467-017-02234-4> <https://doi.org/10.1038/s41467-017-02234-4> PMID: 29269769
40. Sawahata R, Shime H, Yamazaki S, Inoue N, Akazawa T, Fujimoto Y, et al. Failure of mycoplasma lipoprotein MALP-2 to induce NK cell activation through dendritic cell TLR2. *Microbes Infect*. 2011 Apr; 13(4):350–8. <https://doi.org/10.1016/j.micinf.2010.12.003> PMID: 21172450
41. Motoi Y, Shibata T, Takahashi K, Kanno A, Murakami Y, Li X, et al. Lipopeptides are signaled by Toll-like receptor 1, 2 and 6 in endolysosomes. *Int Immunol* [Internet]. 2014 Oct 1 [cited 2019 Nov 19]; 26(10):563–73. Available from: <https://academic.oup.com/intimm/article-lookup/doi/10.1093/intimm/dxu054> PMID: 24860120
42. Marre ML, Petnicki-Ocwieja T, DeFrancesco AS, Darcy CT, Hu LT. Human integrin  $\alpha 3 \beta 1$  regulates TLR2 recognition of lipopeptides from endosomal compartments. *PLoS One*. 2010; 5(9):1–14.
43. Nilsen NJ, Vladimer GI, Stenvik J, Orning MPA, Zeid-Kilani M V., Bugge M, et al. A role for the adaptor proteins TRAM and TRIF in toll-like receptor 2 signaling. *J Biol Chem*. 2015 Feb 6; 290(6):3209–22. <https://doi.org/10.1074/jbc.M114.593426> PMID: 25505250
44. Brandt KJ, Fickentscher C, Kruithof EKO, de Moerloose P. TLR2 Ligands Induce NF- $\kappa$ B Activation from Endosomal Compartments of Human Monocytes. Fritz JH, editor. *PLoS One* [Internet]. 2013 Dec 12 [cited 2019 Nov 19]; 8(12):e80743. Available from: <https://doi.org/10.1371/journal.pone.0080743> PMID: 24349012
45. Underhill D, Ozinsky A, Hajjar A, Stevens A, Wilson C, Bassetti M, et al. The Toll-like receptor 2 is recruited to macrophage phagosomes and discriminates between pathogens. *Nature*. 1999; 402(S6763):39–43.
46. Burke TP, Loukitcheva A, Zemansky J, Wheeler R, Boneca IG, Portnoy DA. *Listeria monocytogenes* is resistant to lysozyme through the regulation, not the acquisition, of cell wall-modifying enzymes. *J Bacteriol*. 2014; 196(21):3756–67. <https://doi.org/10.1128/JB.02053-14> PMID: 25157076
47. Herlihey FA, Osorio-Valeriano M, Dreyfus G, Clarke AJ. Modulation of the lytic activity of the dedicated autolysin for flagellum formation SlfF by flagellar rod proteins FlgB and FlgF. *J Bacteriol*. 2016; 198(13):1847–56. <https://doi.org/10.1128/JB.00203-16> PMID: 27114466
48. Scheurwater E, Reid CW, Clarke AJ. Lytic transglycosylases: Bacterial space-making autolysins. Vol. 40, *International Journal of Biochemistry and Cell Biology*. 2008. p. 586–91. <https://doi.org/10.1016/j.biocel.2007.03.018> PMID: 17468031
49. O'Neill LAJ, Golenbock D, Bowie AG. The history of Toll-like receptors-redefining innate immunity. Vol. 13, *Nature Reviews Immunology*. 2013. p. 453–60. <https://doi.org/10.1038/nri3446> PMID: 23681101
50. Mourao-Sa D, Roy S, Blander JM. Vita-PAMPs: Signatures of microbial viability. In: *Advances in Experimental Medicine and Biology*. Springer New York LLC; 2013. p. 1–8.
51. Blander JM, Barbet G. Exploiting vita-PAMPs in vaccines. *Curr Opin Pharmacol* [Internet]. 2018 Aug 1 [cited 2019 Feb 4]; 41:128–36. Available from: <https://www.sciencedirect.com/science/article/pii/S1471489218300304?via%3Dihub> <https://doi.org/10.1016/j.coph.2018.05.012> PMID: 29890457
52. Vance RE, Isberg RR, Portnoy DA. Patterns of Pathogenesis: Discrimination of Pathogenic and Non-pathogenic Microbes by the Innate Immune System. Vol. 6, *Cell Host and Microbe*. 2009. p. 10–21. <https://doi.org/10.1016/j.chom.2009.06.007> PMID: 19616762
53. Camilli A, Tilney LG, Portnoy DA. Dual roles of plcA in *Listeria monocytogenes* pathogenesis. *Mol Microbiol* [Internet]. 1993 Apr [cited 2019 Nov 24]; 8(1):143–57. Available from: <http://doi.wiley.com/10.1111/j.1365-2958.1993.tb01211.x> PMID: 8388529



54. Jones S, Portnoy DA. Characterization of *Listeria monocytogenes* pathogenesis in a strain expressing perfringolysin O in place of listeriolysin O. *Infect Immun* [Internet]. 1994 Dec 1 [cited 2018 Aug 25]; 62(12):5608–13. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/7960143> PMID: 7960143
55. Lauer P, Chow MYN, Loessner MJ, Portnoy DA, Calendar R. Construction, characterization, and use of two *Listeria monocytogenes* site-specific phage integration vectors. *J Bacteriol*. 2002; 184(15):4177–86. <https://doi.org/10.1128/jb.184.15.4177-4186.2002> PMID: 12107135
56. Zemansky J, Kline BC, Woodward JJ, Leber JH, Marquis H, Portnoy DA. Development of a mariner-based transposon and identification of *Listeria monocytogenes* determinants, including the peptidyl-prolyl isomerase PrsA2, that contribute to its hemolytic phenotype. *J Bacteriol*. 2009 Jun; 191(12):3950–64. <https://doi.org/10.1128/JB.00016-09> PMID: 19376879
57. Hodgson DA. Generalized transduction of serotype 1/2 and serotype 4b strains of *Listeria monocytogenes*. *Mol Microbiol*. 2000; 35(2):312–23. <https://doi.org/10.1046/j.1365-2958.2000.01643.x> PMID: 10652092
58. Portnoy DA, Jacks PS, Hinrichs DJ. Role of hemolysin for the intracellular growth of *Listeria monocytogenes*. *J Exp Med* [Internet]. 1988 Apr 1 [cited 2018 Aug 25]; 167(4):1459–71. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/2833557> <https://doi.org/10.1084/jem.167.4.1459> PMID: 2833557