

UC Santa Cruz

UC Santa Cruz Previously Published Works

Title

Secondary Metabolites from the Marine Sponge Genus Phyllospongia

Permalink

<https://escholarship.org/uc/item/6600n2ws>

Journal

Marine Drugs, 15(1)

ISSN

1660-3397

Authors

Zhang, Huawei
Dong, Menglian
Wang, Hong
[et al.](#)

Publication Date

2017

DOI

10.3390/md15010012

Peer reviewed

Review

Secondary Metabolites from the Marine Sponge Genus *Phyllospongia*

Huawei Zhang ^{1,*}, Menglian Dong ¹, Hong Wang ¹ and Phillip Crews ²

¹ School of Pharmaceutical Sciences, Zhejiang University of Technology, Hangzhou 310014, China; baixl2012@163.com (M.D.); yingzhoudengyuan@sina.com (H.W.)

² Department of Chemistry & Biochemistry, University of California-Santa Cruz, Santa Cruz, CA 95064, USA; pcrews@ucsc.edu

* Correspondence: hwzhang@zjut.edu.cn; Tel.: +86-571-8832-0903

Academic Editor: Vassilios Roussis

Received: 15 November 2016; Accepted: 29 December 2016; Published: 6 January 2017

Abstract: *Phyllospongia*, one of the most common marine sponges in tropical and subtropical oceans, has been shown to be a prolific producer of natural products with a broad spectrum of biological activities. This review for the first time provides a comprehensive overview of secondary metabolites produced by *Phyllospongia* spp. over the 37 years from 1980 to 2016.

Keywords: marine sponge; *Phyllospongia* sp.; secondary metabolites; bioactivity

1. Introduction

Marine sponges, as very primitive animals, are widely distributed in the oceans from tropic to polar regions. Growing evidence indicates that these animals are the most prolific source of natural products as pharmaceutical leads [1–3]. Marine sponges possess a large variety of secondary metabolites with diverse chemical structures, such as terpenoids [4], macrolides [5], and sterols [6]. Therefore, it has greatly attracted the attention of natural product chemists and pharmaceutical experts around the world to carry out chemical research for the new drug discovery.

Phyllospongia (Porifera, Demospongiae, Dictyoceratida, Thorectidae) is one of the most common marine sponges in tropical and subtropical areas, including the Indian Ocean, the Great Barrier Reef, Papua New Guinea, the South China Sea, the South Pacific, and the Red Sea. Chemical investigation of *Phyllospongia* spp. has been extensively carried out and has given rise to a great array of bioactive secondary metabolites. In order to better understand and rationally exploit the marine sponge genus *Phyllospongia*, relevant research references reported between 1980 and 2016 are summarized in this review for the first time.

2. Natural Products from *Phyllospongia* spp.


By 2016, a total of 132 various secondary metabolites (1–132) had been isolated and characterized from the marine sponge genus *Phyllospongia*, including *P. dendyi*, *P.* (syn. *Carteriospongia*) *foliascens*, *P. lamellosa*, *P. madagascarensis*, *P. papyracea*, *Carteriospongia* (syn. *Phyllospongia*) *flabellifera*, and other unidentified *Phyllospongia* spp. (Table 1). In terms of their chemical structures, most *Phyllospongia* sponge-derived natural products are sesterterpenoids, especially scalaranes [7], which are classified as C₂₅ (scalarane), C₂₆ (homoscalarane), and C₂₇ (bishomoscalarane) [8]. Bioassay results indicated that some chemicals have pronounced biological activities and can be used as lead drugs. These secondary metabolites are summarized below according to biological origin.

Table 1. *Phyllospongia* sponge-derived natural products.

| Organism | Locality | Compound | References |
|---|--|--|-------------------------|
| <i>Phyllospongia dendyi</i> | the Great Barrier Reef Andaman and Nicobar Islands | furodendin (1), sesterterpenes (2, 3) scalaranes (4–6), sesterterpene (7) | [9] [10] |
| | Palau Islands | bromophenols (8–12) and diphenyl ethers (13–22) | [11,12] |
| <i>P. (syn. Carteriospongia) foliascens</i> | the Okinawa | foliaspongins (23), dehydrofoliaspongins (24), phyllifoliaspongins (25), dihydrofurospingins-2 (26), furospingins-1 (27) | [13,14] |
| | the South China Sea | phyllaketals (28), phyllofenones A (29), phyllofenones B (30), phyllofolactones A–B (31–32), phyllohemiketals A (33) and B (34), acetoxy phyllofolactone A (35), phyllactones A–G (36–42), phyllofolactones C–D (43–44), phyllofolactone L (45), phyllofenones D–E (46–47), phyllofolactone F–G (48–49), carteriofenones A–K (50–60), analogue (61), phyllofolactone M (62), sterol (63) | [7,15–25] |
| | Papua New Guinea the Indonesia Sea the South Pacific | 20,24-dimethylscalaranes (64–70) scalaranes (71–75) | [26,27] [28] |
| | the Great Barrier Reef unknown | 12- <i>epi</i> -phyllifolactone B (76) scalaranes (77–79) furanoterpene (80) | [29] [30,31] [32] |
| <i>P. lamellosa</i> | the Indo-West Pacific Ocean | phyllolactones A–E (81–85) | [33] |
| | the Egyptian Red Sea | phyllifoliaspongins A–E (86–90), sesterterpenes (76, 91, 92) | [8] |
| <i>P. madagascarensis</i> | Northern Madagascar | sesterterpenes (93–95) | [34] |
| <i>P. papyracea</i> | Hainan Island | scalaranes (96–98) | [35] |
| | Papua New Guinea Sangihe Island | bishomoscalarane sesterterpenes (99–104) phyllactone H (105) | [36] [37] |
| <i>C. (syn. P.) flabellifera</i> | the South Pacific Ocean | flabelliferins A (106) and B (107) | [38] |
| | the Great Barrier Reef | sesterterpenoid (108) | [39] |
| Unclassified <i>P. spp.</i> | the Indonesia Sea | scalaranes (109–114) | [40] |
| | Northern Madagascar | homoscalarane sesterterpenes (115–119) | [41] |
| | the South China Sea | fatty acids (120, 123), steroids (121, 122, 124–127), phylloamide A (128) | [42,43] |
| | Philippines | carteriosulfonic acids A–C (129–131) | [44] |
| | Fiji | sesterterpene (132) | [45] |

2.1. *Phyllospongia dendyi*

One new C₂₂ furanoterpene, furodendin (1), was characterized from *P. dendyi* collected near Cairns on the Great Barrier Reef together with two known C₂₆ tetracyclic sesterterpenes (2, 3) in 1980 [9]. One specimen of *P. dendyi* collected from the coasts of Andaman and Nicobar Islands in the Indian Ocean was found to produce three novel scalaranes (4–6) and a known sesterterpene (7) [10]. Fifteen brominated compounds, bromophenols (8–12) [11] and diphenyl ethers (13–22) [12], were detected in *P. dendyi* grown at Palau Islands (Chart 1). To the best of our knowledge, *P. dendyi* is the only sponge that metabolizes brominated compounds among the *Phyllospongia* genus. Bioassay results suggested that these bromides have a broad spectrum of biological activities, including antimacrolgal activity [11], inhibitory effects on the assembly of microtubule proteins, and the meiotic maturation of starfish oocytes [12].

**Chart 1.** *Cont.*




Chart 1. Chemical structures of secondary metabolites (1–22) from *P. dendyi*.

2.2. *Phyllospongia* (syn. *Carteriospongia*) *foliascens*

Marine sponge *P. foliascens* is the most productive maker of secondary metabolites among all known *Phyllospongia* spp. Chemical investigation indicates that most of these compounds belong to scalarane sesterterpenoid. *P. foliascens* grows in many marine areas, such as Okinawa, the South China Sea, Papua New Guinea, Indonesia, the South Pacific near Vanuatu, and the Great Barrier Reef. Interestingly, the same species of marine sponge collected from different areas possesses various scalarane sesterterpenoids.

After Kikuchi et al. firstly reported the isolation of one novel anti-inflammatory scalarane bishomosesterterpene foliaspongins (**23**) from the Okinawan *P. foliascens* in 1981 [13], two new scalarane-type bishomosesterterpenes, dehydrofoliaspongins (**24**) and phyllofoliaspongins (**25**), and two new furanoterpenes, dihydrofurospongins-2 (**26**) and furospongins-1 (**27**), were also characterized from Okinawan *P. foliascens* (Chart 2). Compound **25** exhibited a broad spectrum of pharmacological effects, such as cytotoxic, anti-thrombocyte, and vasodilative activities [14].




Chart 2. Chemical structures of secondary metabolites (23–27) from *P. foliascens*.

Natural products from the marine sponge *P. foliascens* collected from the South China Sea are abundant. Since 1989, up to 36 compounds have been reported. In 1991, Fu and his coworkers isolated and identified 7 compounds: phylloketal (**28**) [15], phyllofenone A (**29**) [16], phyllofenone B (**30**), phyllofolactones A–B (**31–32**) [17], and phyllohemiketal A–B (**33–34**) [18]. Among these secondary metabolites, compound **29** showed weak antifungal activity against *Candida pseudotropicalis*, while compound **30** displayed cytotoxic activity against the P388 murine leukemia cell line with an IC₅₀ value of 5 µg/mL. Acetoxy phyllofolactone A (**35**) [19] and phyllactones A–E (**36–40**) [20] were found and characterized in 1992. Compounds **36** and **37** had moderate in vitro cytotoxicity against KB cells with the same IC₅₀ value of 20 µg/mL. Interestingly, phyllactones F–G (**41–42**) [21] and phyllofolactones C–D (**43–44**) [22] were also detected in the same specimen. Phyllofolactone L (**45**) and phyllofenone D–E (**46–47**) belong to ascalarane sesterterpenoids [23]. Their chemical structures were elucidated on

the basis of spectroscopic analysis. A biological assay showed that only compound **47** had moderate cytotoxic activity against the leukemia cell line P388, with an IC_{50} value of 6.5 $\mu\text{g}/\text{mL}$. Two new sesquiterpenes, phyllofolactone F (**48**) and phyllofolactone G (**49**), together with phyllofenone D (**46**) and phyllofenone E (**47**), were isolated and identified by chromatographic methods and modern analytical methods [7]. Carteriofenones A–C (**50–52**) were 20,24-bishomo-25-norscalaranes and carteriofenones D–K (**53–60**) and one analogue (**61**) belong to 20,24-bishomoscalaranes [24]. Moreover, phyllofolactone M (**62**) and a new sterol, (24*E*)-5 α ,6 α -epoxystigmasta-7,24(28)-dien-3 β -ol (**63**), together with a known sesterterpene, phyllofolactone B (**32**), were detected in the same sample collected from the South China Sea [25] (Chart 3).




Chart 3. Chemical structures of secondary metabolites (**28–63**) from *P. foliascens*.

Seven 20,24-dimethylsclalarane derivatives (**64–70**) were characterized from *P.* (syn. *Carteriospongia*) *foliascens* gathered from Papua New Guinea. Compounds **64** and **70** were C-14 anomers [26,27]. Chemical investigation of *P.* (syn. *C.*) *foliascens* collected from the Indonesian sea afforded five sclalarane sesterterpenoids (**71–75**), which possessed Ras Converting Endoprotease (RCE) inhibitory effect except **72** [28]. Specimens of *P.* (syn. *C.*) *foliascens* from the South Pacific could metabolite 12-*epi*-phyllolactone B (**76**) [29], while the sample collected from the Great Barrier Reef produced sclalarane sesterterpenoids (**77–79**) [30,31]. Additionally, one furanoterpene (**80**) was also isolated from the marine sponge *P.* (syn. *C.*) *foliascens* [32] (Chart 4).




Chart 4. Chemical structures of secondary metabolites (**64–80**) from *P.* (syn. *Carteriospongia*) *foliascens*.

2.3. *Phyllospongia lamellosa*

Until now, there have only been two reports on the chemical study of the marine sponge *P. lamellosa* in the database Web of Sciences [33]. Five 20,24-bishomosclalarane sesterterpenes, phyllolactones A–E (**81–85**), were characterized from *P. lamellosa* collected in the Indo-West Pacific Ocean. Bioassay results indicated that these secondary metabolites possessed an inhibitory effect on human immunodeficiency virus type 1 (HIV-1) envelope-mediated fusion in vitro with IC₅₀ values of about 2 μM [34]. From the same marine sponge derived from the Egyptian Red Sea, five new sclalarane sesterterpenes, phyllospongins A–E (**86–90**), were detected together with four known derivatives (**76**, **91**, **92** and **99**) [8] (Chart 5). Compounds **87–91** had potent cytotoxic activity against HCT-116 as the positive control doxorubicin, while **90** showed cytotoxic activity against MCF-7.




Chart 5. Chemical structures of secondary metabolites (81–92) from *P. lamellosa*.

2.4. *Phyllospongia madagascarensis*

To the best of our knowledge, only three natural products (93–95) (Chart 6) have been found in the marine sponge *P. madagascarensis*, which was grown near the northwest coast of Madagascar [35]. Interestingly, the chemical structure of 94 possesses seven-membered oxacycle.




Chart 6. Chemical structures of secondary metabolites (93–95) from *P. madagascarensis*.

2.5. *Phyllospongia papyracea*

Ten small molecules (96–105) (Chart 7) were isolated from *P. papyracea* collected on Hainan Island in the South China Sea, Papua New Guinea, and Sangihe Island in the Indonesian Sea [36–38]. Cytotoxic tests suggested that compound 96 had an in vitro cytotoxic effect on the leukemia cancer cell line P388 with an IC₅₀ value of 5 μ g/mL, and 99–104 were inactive against the β -catenin and transcription factor 4 (Tcf4) complex. Phyllactone H (105) was found in the marine sponge derived from Sangihe Island and possessed in vitro moderate cytotoxicities against cell lines A549, MCF-7, and HeLa with IC₅₀ values of no more than 25 μ M.




Chart 7. Cont.




Chart 7. Chemical structures of secondary metabolites (96–105) from *P. papyracea*.

2.6. *Carteriospongia* (syn. *Phyllospongia*) *flabellifera*

The marine sponge *C.* (syn. *P.*) *flabellifera* is usually distributed in a wide corridor of the Indo-Pacific Ocean. A chemical study of the marine sponge collected in the South Pacific Ocean led to the isolation of two new small molecules, flabelliferins A (**106**) and B (**107**) [39]. Compound **106** had a rare 25-homocheilanthane carbon skeleton, and **107** exhibited inhibitory effect on the human colon tumor cell lines KM12 and COLO205. A new sesterterpenoid derivative (**108**) was also characterized from *C. flabellifera* collected around the Great Barrier Reef [40] (Chart 8).




Chart 8. Chemical structures of secondary metabolites (106–108) from *C.* (syn. *Phyllospongia*) *flabellifera*.

2.7. Other *Phyllospongia* spp.

Up to 24 secondary metabolites (**109–132**) (Chart 9) were isolated and identified from other unclassified *Phyllospongia* spp. Compounds **109–114** from an Indonesian marine sponge displayed 30%–95% inhibition of the growth of KB cells at 10 $\mu\text{g}/\text{mL}$ [41]. Compounds **115–119** were produced by the *Phyllospongia* sp. collected from Northern Madagascar. Compounds **115**, **116**, and **119** possessed strong in vitro cytotoxic activities against human ovarian cancer cell line A2780 with IC_{50} values of 0.26, 0.28, and 0.65 μM , respectively, while **117** and **118** had moderate activities with IC_{50} values of 4.5 and 8.7 μM , respectively. Compound **116** exhibited a strong inhibitory effect on the human lung non-small cell line H522-T1 with an IC_{50} value of 0.61 μM [42]. Compounds **120–127** [43] and phylloamide A (**128**) [44] were also isolated from the South China Sea sponge. Carteriosulfonic acids A–C (**129–131**) [45] and **132** [46] were respectively isolated from two specimens collected at Philippines and Fiji. Bioassay results suggested that **128–130** had an inhibitory effect on the growth of glycogen synthase kinase-3 β (GSK-3 β), with IC_{50} values of 12.5, 6.8, and 6.8 μM , respectively.




Chart 9. Cont.




Chart 9. Chemical structures of secondary metabolites (109–132) from other *Phyllospongia* spp.

3. Conclusions

Natural products from the marine sponge genus *Phyllospongia* have been well studied over the last 37 years. Their frameworks are diverse, including terpenoid, macrolide, sterol, and ceramide. Moreover, the most common structure is sesterterpenoid, which has diverse biological activities. With the increasing development of oceanographic technology leading to the isolation of new *Phyllospongia* species from marine environments, more secondary metabolites with novel chemical structure(s) and/or potent bioactivities will be found in the near future.

Acknowledgments: This project was financially supported by the Zhejiang Natural Science Foundation of China (LY16H300007).

Author Contributions: H.Z. conceived and wrote the review; M.D. searched and collected the references; H.W. made an effective analysis of the document; P.C. offered important advice to improve the review.

Conflicts of Interest: The authors declare no conflict of interest.

References

1. Keyzers, R.A.; Davies-Coleman, M.T. Anti-inflammatory metabolites from marine sponges. *Chem. Soc. Rev.* **2005**, *34*, 355–365. [[CrossRef](#)] [[PubMed](#)]
2. Xu, W.H.; Ding, Y.; Jacob, M.R.; Agarwal, A.K.; Clark, A.M.; Ferreira, D.; Liang, Z.; Li, X. Puupehanol, a sesquiterpene-dihydroquinone derivative from the marine sponge *Hyrtios* sp. *Bioorg. Med. Chem. Lett.* **2009**, *19*, 6140–6143. [[CrossRef](#)] [[PubMed](#)]

3. Boonlarppradab, C.; Faulkner, D.J. Eurysterols A and B, cytotoxic and antifungal steroidal sulfates from a marine sponge of the genus *Euryspongia*. *J. Nat. Prod.* **2007**, *70*, 846–848. [[CrossRef](#)] [[PubMed](#)]
4. Gupta, P.; Sharma, U.; Schulz, T.C.; McLean, A.B.; Robins, A.J.; West, L.M. Bicyclic C₂₁ terpenoids from the marine sponge *Clathria compressa*. *J. Nat. Prod.* **2012**, *75*, 1223–1227. [[CrossRef](#)] [[PubMed](#)]
5. Camille, C.; Bugni, T.S.; Feng, X.; Kay, H.M.; Orendt, A.M.; Irel, C.M. Tedanolide C: A potent new 18-membered-ring cytotoxic macrolide isolated from the Papua New Guinea marine sponge *Ircinia* sp. *J. Org. Chem.* **2006**, *71*, 2510–2513.
6. Abdelmohsen, U.R.; Cheng, C.; Reimer, A.; Kozjak-Pavlovic, V.; Ibrahim, A.K.; Rudel, T.; Hentschel, U.; Edrada-Ebel, R.; Ahmed, S.A. Antichlamydial sterol from the Red Sea sponge *Callyspongia* aff. *implexa*. *Planta Med.* **2015**, *81*, 382–387. [[CrossRef](#)] [[PubMed](#)]
7. Wan, Y.Q.; Su, J.Y.; Zeng, L.M.; Liu, Q. The secondary metabolites of marine sponge *Phallospongia foliascens*. *Acta Sci. Nat. Univ. Sunyatseni* **1998**, *37*, 81–84.
8. Hassan, M.H.A.; Rateb, M.E.; Hetta, M.; Abdelaziz, T.A.; Sleim, M.A.; Jaspars, M.; Mohammed, R. Scalarane sesterterpenes from the Egyptian Red Sea sponge *Phyllospongia lamellosa*. *Tetrahedron* **2015**, *71*, 577–583. [[CrossRef](#)]
9. Kazlauskas, R.; Murphy, P.T.; Wells, R.J. Furodendin, a C₂₂ degraded terpene from the sponge *Phyllospongia dendyi*. *Experientia* **1980**, *36*, 814–815. [[CrossRef](#)]
10. Rao, C.B.; Kalidindi, R.S.; Trimurtulu, G.; Rao, D.V. Metabolites of Porifera, Part III. New 24-Methylscalaranes from *Phyllospongia dendyi* of the Indian Ocean. *J. Nat. Prod.* **1991**, *54*, 364–371. [[CrossRef](#)]
11. Hattori, T.; Konno, A.; Adachi, K.; Shizuri, Y. Four new bioactive bromophenols from the Palauan sponge *Phyllospongia dendyi*. *Fish. Sci.* **2001**, *67*, 899–903. [[CrossRef](#)]
12. Liu, H.W.; Namikoshi, M.; Meguro, S.; Nagai, N.; Kobayashi, H.; Yao, X. Isolation and characterization of polybrominated diphenyl ethers as inhibitors of microtubule assembly from the marine sponge *Phyllospongia dendyi* collected at Palau. *J. Nat. Prod.* **2004**, *67*, 472–474. [[CrossRef](#)] [[PubMed](#)]
13. Kikuchi, H.; Tsukitani, Y.; Shimizu, I.; Kobayashi, M.; Kitagawa, I. Foliasspongin, an antiinflammatory bishomosesterterpene from the marine sponge *Phyllospongia foliascens* (Pallas). *Chem. Pharm. Bull.* **1981**, *29*, 1492–1494. [[CrossRef](#)]
14. Kitagawa, I.; Kobayashi, M.; Lee, N.K.; Oyama, Y.; Kyogoku, Y. Marine Natural Products XX. Bioactive scalarane-type bishomosesterterpenes from the Okinawan marine sponge *Phyllospongia foliascens*. *Chem. Pharm. Bull.* **1989**, *37*, 2078–2082. [[CrossRef](#)]
15. Fu, X.; Zeng, L.M.; Su, J.Y.; Francis, J.S. A new sesterterpene from the sponge *Phyllospongia foliascens*. *Chin. Chem. Lett.* **1991**, *2*, 543–544.
16. Zeng, L.M.; Fu, X.; Su, J.Y. Phyllofenone A, a new scalarane sesterterpene from the sponge *Phyllospongia foliascens* (Pallas). *Chem. Res. Chin. Univ.* **1991**, *7*, 100–106.
17. Zeng, L.M.; Fu, X.; Su, J.Y. Novel bishomoscalarane sesterterpenes from the sponge *Phyllospongia foliascens*. *Chem. Res. Chin. Univ.* **1991**, *54*, 421–427. [[CrossRef](#)]
18. Fu, X.; Zeng, L.M.; Su, J.Y.; Chen, S.; Snyder, J.K. Bishomoscalarane sesterterpenes from the sponge *Phyllospongia foliascens*. *Chem. Res. Chin. Univ.* **1991**, *12*, 1486–1487.
19. Fu, X.; Zeng, L.M.; Su, J.Y. Acetoxy phyllofolactone A—A novel bishomoscalarane sesterterpene from the South China Sea sponge *Phyllospongia foliascens*. *Chem. J. Chin. Univ.* **1992**, *13*, 628–629.
20. Fu, X.; Zeng, L.M.; Su, J.Y.; Païs, M.; Potier, P. Scalarane-type bishomosesterterpenes from the sponge *Phyllospongia foliascens*. *J. Nat. Prod.* **1992**, *55*, 1607–1613. [[CrossRef](#)] [[PubMed](#)]
21. Fu, X.; Zeng, L.M.; Su, J.Y.; Païs, M.; Potier, P. Two new sesterterpenes from a South China Sea Sponge. *J. Nat. Prod.* **1993**, *56*, 1985–1988. [[CrossRef](#)]
22. Fu, X.; Zeng, L.M.; Su, J.Y.; Païs, M.; Potier, P. Phyllofolactones C and D, two new minor homoscalarane sesterterpenes from the Chinese sponge *Phyllospongia foliascens*. *J. Nat. Prod.* **1999**, *62*, 644–646. [[CrossRef](#)] [[PubMed](#)]
23. Zhang, H.J.; Tang, H.F.; Yi, Y.H.; Lin, H.W. Scalarane sesterterpenes from the Chinese sponge *Phyllospongia foliascens*. *Helv. Chim. Acta* **2009**, *92*, 762–767. [[CrossRef](#)]
24. Cao, F.; Wu, Z.H.; Shao, C.L.; Pang, S.; Liang, X.Y.; de Voogd, N.J.; Wang, C.Y. Cytotoxic scalarane sesterterpenoids from the South China Sea sponge *Carteriospongia foliascens*. *Org. Biomol. Chem.* **2015**, *13*, 4016–4024. [[CrossRef](#)] [[PubMed](#)]

25. Zhang, H.J.; Yi, Y.H.; Yang, F.; Chen, W.S.; Lin, H.W. Sesterterpenes and a new sterol from the marine sponge *Phyllospongia foliascens*. *Molecules* **2010**, *15*, 834–841. [CrossRef] [PubMed]
26. Braekman, J.C.; Daloze, D.; Kaisin, M.; Moussiaux, B. Ichthyotoxic sesterterpenoids from the new guinean sponge *Carteriospongia foliascens*. *Tetrahedron* **1985**, *41*, 4603–4614. [CrossRef]
27. Declercq, J.P.; Meerssche, M.V.; Braekman, J.C.; Daloze, D. A new 20,24-dimethylsclerane derivative isolated from the sponge *Carteriospongia foliascens*, C₃₀H₄₂O₆. *Acta Cryst.* **1985**, *41*, 1222–1224. [CrossRef]
28. Williams, D.E.; Hollander, I.; Feldberg, L.; Frommer, E.; Mallon, R.; Tahir, A.; van Soest, R.; Andersen, R.J. Sclerane-based sesterterpenoid RCE-protease inhibitors isolated from the Indonesian marine sponge *Carteriospongia foliascens*. *J. Nat. Prod.* **2009**, *72*, 1106–1109. [CrossRef] [PubMed]
29. Reddy, M.V.R.; Venkateswarlu, Y.; Rao, J.V. A new bishomoscclerane sesterterpene 12-*epi*-phyllolactone-B from *Phyllospongia foliascens*. *Indian J. Chem. B* **1993**, *32*, 1196–1197.
30. Barron, P.F.; Quinn, R.J.; Tucker, D.J. Structural elucidation of a novel sclerane derivative by using high-field (14.1T) NMR. Spectroscopy. *Aust. J. Chem.* **1991**, *44*, 995–999. [CrossRef]
31. Quinn, R.J.; Tucker, D.J. Two novel bisalkylated norscleranes from the sponge *Carteriospongia foliascens*. *Aust. J. Chem.* **1989**, *42*, 751–755. [CrossRef]
32. Masaru, K.; Ramadas, M.; Osamu, M.; Nittala, S. Marine terpenes and terpenoids XVI: Revised structure of the marine furanoterpene (+)-furospingonin-1. *J. Chem. Res.-Synop.* **1992**, *11*, 366–367.
33. Available online: https://apps.webofknowledge.com/Search.do?product=WOS&SID=V2B6tzoudOkWopSGj2w&search_mode=GeneralSearch&prID=b0222ef8-1c6f-4f79-b3bf-ed19663b27c6 (accessed on 3 January 2017).
34. Chang, L.C.; Otero-Quintero, S.; Nicholas, G.M.; Bewley, C.A. Phyllolactones A–E: New bishomoscclerane sesterterpenes from the marine sponge *Phyllospongia lamellosa*. *Tetrahedron* **2001**, *57*, 5731–5738. [CrossRef]
35. Ponomarenko, L.P.; Kalinovsky, A.I.; Stonik, V.A. New sclerane-based sesterterpenes from the sponge *Phyllospongia madagascarensis*. *J. Nat. Prod.* **2004**, *67*, 1507–1510. [CrossRef] [PubMed]
36. Lan, W.J.; Li, H.J. New sesterterpenoids from the marine sponge *Phyllospongia papyracea*. *Helv. Chim. Acta* **2007**, *90*, 1218–1222. [CrossRef]
37. Li, H.J.; Amagata, T.; Tenney, K.; Crews, P. Additional sclerane sesterterpenes from the sponge *Phyllospongia papyracea*. *J. Nat. Prod.* **2007**, *70*, 802–807. [CrossRef] [PubMed]
38. Zhang, H.W.; Crews, P.; Tenney, K.; Valeriote, F.A. Cytotoxic phyllactone analogs from the marine sponge *Phyllospongia papyracea*. *Med. Chem.* **2016**, *12*. [CrossRef]
39. Diyabalanage, T.; Ratnayake, R.; Bokesch, H.R.; Ransom, T.T.; Henrich, C.J.; Beutler, J.A.; McMahon, J.B.; Gustafson, K.R. Flabelliferins A and B, sesterterpenoids from the South Pacific sponge *Carteriospongia flabellifera*. *J. Nat. Prod.* **2012**, *75*, 1490–1494. [CrossRef] [PubMed]
40. Francis, J.S.; James, C.C. Sesterterpenes from a Pacific sponge, *Carteriospongia flabellifera*. *J. Nat. Prod.* **1988**, *51*, 745–748.
41. Michael, C.R.; Junichi, T.; de Voogd, N.; Higa, T. New sclerane class sesterterpenes from an Indonesian sponge, *Phyllospongia* sp. *J. Nat. Prod.* **2001**, *62*, 1838–1842.
42. Harinantenaina, L.; Brodie, P.J.; Maharavo, J.; Bakary, G.; TenDyke, K.; Shen, Y.; Kingston, D.G.I. Antiproliferative homoscclerane sesterterpenes from two Madagascan sponges. *Bioorg. Med. Chem.* **2013**, *21*, 2912–2917. [CrossRef] [PubMed]
43. Wan, Y.Q.; Zeng, L.M.; Su, J.Y. Studies on the sponge *Phyllospongia* sp. (II). *Acta Sci. Nat. Univ. Sunyatseni* **1997**, *36*, 116–118.
44. Wan, Y.Q.; Su, J.Y.; Zeng, L.M. Ceramide from the marine sponge *Phyllospongia* sp. *Acta Sci. Nat. Univ. Sunyatseni* **1997**, *36*, 113–115.
45. McCulloch, M.W.B.; Bugni, T.S.; Concepcion, G.P.; Coombs, G.S.; Harper, M.K.; Kaur, S.; Mangalindan, G.C.; Mutizwa, M.M.; Veltri, V.A.; Virshup, D.M.; et al. Carteriosulfonic acids A–C, GSK-3β inhibitors from a *Carteriospongia* sp. *J. Nat. Prod.* **2009**, *72*, 1651–1656. [CrossRef] [PubMed]
46. Croft, K.D.; Ghisalberti, E.L.; Skelton, B.W.; White, A.H. Structural study of a new dialkylated sclerane from a *Carteriospongia* sp. *Chem. Soc. Perkin Trans.* **1983**, *41*, 155–159. [CrossRef]

