

ISOLATION AND CHARACTERIZATION OF A TRITERPENOID
ALCOHOL FROM THE GREEN RIVER SHALE


W. Henderson and G. Steel

September 1971

Prepared for the U.S. Atomic Energy Commission
under Contract W-7405-ENG-48

TWO-WEEK LOAN COPY

*This is a Library Circulating Copy
which may be borrowed for two weeks.
For a personal retention copy, call
Tech. Info. Division, Ext. 5545*



RECEIVED
LAWRENCE
BERKELEY LABORATORY
LIBRARY AND
DOCUMENTS SECTION

DISCLAIMER

This document was prepared as an account of work sponsored by the United States Government. While this document is believed to contain correct information, neither the United States Government nor any agency thereof, nor the Regents of the University of California, nor any of their employees, makes any warranty, express or implied, or assumes any legal responsibility for the accuracy, completeness, or usefulness of any information, apparatus, product, or process disclosed, or represents that its use would not infringe privately owned rights. Reference herein to any specific commercial product, process, or service by its trade name, trademark, manufacturer, or otherwise, does not necessarily constitute or imply its endorsement, recommendation, or favoring by the United States Government or any agency thereof, or the Regents of the University of California. The views and opinions of authors expressed herein do not necessarily state or reflect those of the United States Government or any agency thereof or the Regents of the University of California.

LBL-1380

1972-428

Reproduced with permission of
Chemical Communications,
The Chemical Society, London
November 1972

Isolation and Characterization of a Triterpenoid Alcohol from the Green River Shale

By W. HENDERSON* and G. STEEL

(Department of Chemistry, University of California, Berkeley, California 94720)

Reprinted from

Chemical Communications 1971

The Chemical Society, Burlington House, London W1V 0BN

Isolation and Characterization of a Triterpenoid Alcohol from the Green River Shale

By W. HENDERSON* and G. STEEL




(Department of Chemistry, University of California, Berkeley, California 94720)

Summary A triterpenoid alcohol (tetrahymanol) has been isolated from the Green River Shale (60 million years).

SEVERAL publications have shown the presence of fully reduced steroids and triterpenoids (steranes and triterpanes) in the Green River Shale (Eocene, 60×10^6 years old).¹⁻³ The identities of many of these compounds have been established by capillary g.l.c., capillary g.c.-m.s.,³ and by X-ray crystallography.² Although absolute stereochemistries in most cases have not been established, the following structural identifications have been made: cholestane, ergostane, stigmastane, gammacerane, and a hopane-type structure. The presence of these compounds in the Green River Shale has been taken as substantial proof that the original source of the organic content of the sediment was biological and probably dominantly algal.⁴ It has been suggested that these molecules were derived from naturally occurring phytosterols, and higher plant and micro-organism triterpenoids which, through the processes of deposition, consolidation and maturation of the sediment, were reduced to the cycloalkanes described above. No evidence has been produced which would confirm this postulate by identifying the precursors of the cycloalkanes, *i.e.*, sterols and triterpenoid alcohols, from this sediment. However, it has been reported that the Messel Oil Shale (Eocene; with a milder geological history than the Green River Formation) does contain triterpenoid alcohols although the authors found no evidence for triterpenoid alcohols in the Green River Shale.^{5,6} We now report the isolation and identification of a triterpenoid alcohol from the Green River Shale.

The sediment (350 g) from the Mahogany Ledge was broken into small pieces, pulverized in a disc mill (Angstrom Inc., Chicago, Illinois) and the powder extracted with a mixture of benzene-methanol (3:1 v/v) with ultrasonication. After centrifugation the supernatant

solution was separated, evaporated to dryness, and the total extract chromatographed on a silica gel column eluting successively with n-heptane, mixtures of 5-40% diethyl ether in n-heptane, diethyl ether, and finally ethyl acetate. The fractions collected were monitored by analytical t.l.c. using a solvent mixture of n-heptane, diethyl ether, and methanol (18:6:1 v/v). By comparison of the extract fractions with the mobilities of reference compounds some of the fractions obtained using 40% diethyl ether in n-heptane showed constituents with the correct mobilities to be sterol or triterpenoid in character. The hydroxylated compounds were selectively removed from extraneous materials by preparative t.l.c. (n-heptane:benzene 1:1 v/v).



as mobile phase) in the form of trimethylsilyl ethers. After purification, fractions as trimethylsilyl ethers were examined by g.l.c. on two different liquid phases (*viz.* JXR and OV-17) and finally by g.c.-m.s.⁷

The mass spectra obtained showed that one of the main derivatives had a molecular ion (M) at m/e 500 and showed other ions at m/e 410 ($M - 90$), 395 ($M - 105$), and 73 characteristic of the trimethylsilyl ether of an alcohol. The two most abundant ions in the spectrum were found at m/e 191 and m/e 189, respectively, which were deduced to result from a fragmentation of a symmetrical nucleus such as is found in the pentacyclic triterpenoid alcohol tetrahymanol (I; shown here as the trimethylsilyl ether).

TABLE

G.l.c. retention data for extracts and reference compounds as trimethylsilyl ethers

Parent compound	I(JXR) ^a	I(OV-17) ^b
Alcohol from Green River Shale ..	3520	3725
Alcohol from <i>Tetrahymena pyriformis</i>	3520	3725
Authentic tetrahymanol	3520	3730

^a 10 ft \times $\frac{1}{8}$ in stainless-steel column packed with 3% JXR on 60-80 mesh Gas Chrom Q at 265°.

^b 6 ft \times $\frac{1}{8}$ in stainless-steel column packed with 3% OV-17 on 60-80 mesh Gas Chrom Q at 270°.

In order to confirm this tentative identification a sample of the protozoan, *Tetrahymena pyriformis* (from which tetrahymanol had previously been isolated and identified⁶) was obtained and cultured in this laboratory. Extraction

and liquid-solid chromatography followed by t.l.c. of the trimethylsilyl ether fraction yielded a mixture of components dominated by tetrahymanol trimethylsilyl ether. Comparison of the g.l.c. retention indices of this component with those of the alcohol isolated from the Green River Shale showed them to be identical. Further confirmation was achieved by comparison of the retention data (Table) and mass spectrum of an authenticated sample of tetrahymanol trimethylsilyl ether with those of the compounds obtained from *Tetrahymena pyriformis* and the Green River Shale.

The triterpane gammacerane has already been identified in the hydrocarbon fraction from Green River Shale,^{2,3} and this new evidence showing the presence of the corresponding saturated alcohol indicates that hydroxylated triterpenoids can exist intact over a significant period of geological time and provides information on the mechanisms of diagenesis and maturation of organic natural products in a sedimentary environment. Qualitative evidence indicates the presence of other, as yet unidentified, possible triterpenoid alcohols in the Green River Shale.

We acknowledge a gift of pure tetrahymanol from Dr. F. B. Mallory, Dept. of Chemistry, Bryn Mawr College, Penn. This work was supported by a grant from NASA.

(Received, September 2nd, 1971; Com. 1538.)

¹ G. Eglinton, P. M. Scott, T. Belsky, A. L. Burlingame, and M. Calvin, *Science*, 1964, **145**, 263.

² I. R. Hills, E. V. Whitehead, D. E. Anders, J. J. Cummings, and W. E. Robinson, *Chem. Comm.*, 1966, 752.

³ W. Henderson, V. Wollrab, and G. Eglinton, *Adv. Org. Geochem.*, 1968, **31**, 181.

⁴ W. H. Bradley, *Geol. Soc. Amer. Bull.*, 1966, **77**, 1333.

⁵ P. Albrecht and G. Ourisson, *Science*, 1969, **163**, 1192.

⁶ G. Mattern, P. Albrecht, and G. Ourisson, *Chem. Comm.*, 1970, 1570.

⁷ W. Henderson, W. E. Reed, G. Steel, and M. Calvin, *Nature*, 1971, **231**, 308.

⁸ F. B. Mallory, J. T. Gordon, and R. L. Conner, *J. Amer. Chem. Soc.*, 1963, **85**, 1362.

LEGAL NOTICE

This report was prepared as an account of work sponsored by the United States Government. Neither the United States nor the United States Atomic Energy Commission, nor any of their employees, nor any of their contractors, subcontractors, or their employees, makes any warranty, express or implied, or assumes any legal liability or responsibility for the accuracy, completeness or usefulness of any information, apparatus, product or process disclosed, or represents that its use would not infringe privately owned rights.

TECHNICAL INFORMATION DIVISION
LAWRENCE BERKELEY LABORATORY
UNIVERSITY OF CALIFORNIA
BERKELEY, CALIFORNIA 94720