

UC Irvine

Clinical Practice and Cases in Emergency Medicine

Title

Drug-Induced Immune Thrombocytopenia Caused by Sulfamethoxazole-Trimethoprim: A Case Report

Permalink

<https://escholarship.org/uc/item/6959f45r>

Journal

Clinical Practice and Cases in Emergency Medicine, 0(0)

Authors

Powell, Alec

Weindruch, Louisa

Thornton, Jonathan

Publication Date

2026-10-31

DOI

10.5811/cpcem.65094

Copyright Information

This work is made available under the terms of a Creative Commons Attribution License, available at <https://creativecommons.org/licenses/by/4.0/>

Peer reviewed

Drug-Induced Immune Thrombocytopenia Caused by Sulfamethoxazole-Trimethoprim: A Case Report

Alec Powell, DO
Louisa Weindruch, DO
Jonathan Thornton, MD

Baylor University Medical Center, Department of Emergency Medicine, Dallas, Texas

Section Editor: Melanie Heniff, MD, JD

Submission history: Submitted May 6, 2026; Revision received August 10, 2026; Accepted August 10, 2026

Electronically published October 30, 2026

Full text available through open access at http://escholarship.org/uc/uciem_cpccm

DOI: 10.5811/cpccm.65094

Introduction: Drug-induced immune thrombocytopenia (DITP) is a rare process in which specific drugs induce antibodies against platelet antigens, resulting in the destruction, consumption, or impaired production of platelets.^{1,3} Sulfamethoxazole-trimethoprim can result in DITP.²

Case Report: This is a case of a 64-year-old man who developed severe thrombocytopenia after sulfamethoxazole-trimethoprim use. He presented to the emergency department with oral hemorrhagic lesions, mucositis, and petechiae. Laboratory evaluation revealed a low platelet count. Drug-induced immune thrombocytopenia was suspected, and the sulfamethoxazole-trimethoprim was discontinued. The patient's platelet count responded rapidly after discontinuation of the offending drug and administration of dexamethasone.

Conclusion: This case highlights the importance of early recognition of DITP and the avoidance of re-exposure to the causative agent. Prompt drug withdrawal and adjunctive corticosteroid therapy can facilitate complete clinical recovery. [Clin Pract Cases Emerg Med. 2026;X(X):XXX-XXX.]

Keywords: *drug-induced immune thrombocytopenia; thrombocytopenia; sulfamethoxazole-trimethoprim; mucosal bleeding; case report.*

INTRODUCTION

Drug-induced immune thrombocytopenia (DITP) is caused by drug-specific antibodies that develop against platelet antigens, most commonly the glycoprotein IIb/IIIa complex (GPIIb/IIIa) and the glycoprotein Ib/IX complex (GPIb/IX),¹ which can result in suppression of hematopoiesis, megakaryocytopoiesis inhibition, platelet destruction, and platelet consumption.^{1,3} This potentially life-threatening hematologic condition can be caused by drugs such as including quinine, carbamazepine, vancomycin, eptifibatide, sulfamethoxazole, and trimethoprim.² We present a case of sulfamethoxazole-trimethoprim-induced thrombocytopenia presenting with oral mucosal bleeding, mucositis, petechiae, and a critically low platelet count in a patient who had a similar presentation with a prior exposure. This case highlights a rare complication of a commonly prescribed medication and

demonstrates the importance of prompt recognition in the emergency department (ED) setting.

CASE REPORT

A 64-year-old man with type 2 diabetes mellitus, hypertension, hyperlipidemia, and chronic urinary retention presented to the ED with several hemorrhagic oral lesions and a rash on his bilateral lower extremities. Four days prior to presentation, his chronic indwelling Foley catheter had been removed at an outside hospital. The patient, believing he had developed a urinary tract infection, started taking leftover sulfamethoxazole-trimethoprim belonging to his wife. Within three days, he developed hemorrhagic lesions on his tongue and a petechial rash involving his lower extremities. He reported dark stools two days prior to presentation but denied hematemesis, melena, and hematuria at presentation. He recalled a prior reaction to sulfa drugs that resulted in an

episode of “low blood counts.”

The patient was well-appearing, in no acute distress, afebrile, and hemodynamically stable. On physical exam, he had hemorrhagic blisters on the tongue and lips (Image 1), mucositis, and petechiae involving the bilateral lower extremities (Image 2). No scleral icterus, lymphadenopathy, or hepatosplenomegaly was observed.

Laboratory evaluation was most notable for significant thrombocytopenia, with a platelet count of less than 1×10^3 platelets/ μL (reference range, $150\text{--}400 \times 10^3/\mu\text{L}$). Hemoglobin and hematocrit were within normal limits.

Dermatology consult found the presentation not to be consistent with Stevens-Johnson syndrome or toxic epidermal necrolysis. The hematology team suspected drug-induced immune thrombocytopenia and recommended dexamethasone 40 mg intravenously (IV) in the ED, followed by daily doses while inpatient. Platelet transfusion and IV immunoglobulin were deferred due to absence of severe symptoms and active bleeding.

During his hospital stay, the patient's platelet count and clinical findings improved rapidly after discontinuation of sulfamethoxazole-trimethoprim (Table). His hospital course was complicated by mild steroid-induced hyperglycemia, which was managed conservatively. The patient's platelet count was adequately responding; therefore, hematology agreed without the decision to withhold the last dose of dexamethasone given the hyperglycemia. The patient was discharged on hospital day five after complete platelet recovery with instructions for repeat labs outpatient one week after discharge.




Image 1. Hemorrhagic blisters seen on patient's buccal mucosa and lateral tongue (arrows).

CPC-EM Capsule

What do we already know about this clinical entity?

Drug-induced immune thrombocytopenia (DITP) is a rare process by which exposure to certain drugs results in production of antibodies against platelet antigens.

What makes this presentation of disease reportable?

We present a case of sulfamethoxazole-trimethoprim-induced thrombocytopenia presenting with oral mucosal bleeding and petechiae.

What is the major learning point?

Several medications can cause DITP and lead to significant thrombocytopenia. Early recognition, treatment, and avoidance of re-exposure is crucial.

How might this improve emergency medicine practice?

This case highlights a rare complication of a commonly prescribed medication and demonstrates the importance of prompt recognition in the emergency care setting.

DISCUSSION

Sulfamethoxazole-trimethoprim-induced immune thrombocytopenia is a well-known cause of DITP; use of other medications such as quinine, quinidine, vancomycin, penicillin, rifampin, ibuprofen, mirtazapine, oxaliplatin, suramin, carbamazepine, and ceftriaxone can cause this condition.⁴ A study by Kaufman et al reported 38 cases of DITP per million users of sulfamethoxazole-trimethoprim.⁵ It is most commonly caused by the production of immunoglobulin G antibodies that bind to platelet antigens, specifically GPIIb/IIIa and GPIb/IX.¹

Thrombocytopenia is defined as a platelet count of less than $150 \times 10^3/\mu\text{L}$ ²; however, in DITP, the platelet nadir is frequently less than $20 \times 10^3/\mu\text{L}$.¹ Onset typically occurs 5 to 10 days following drug exposure, resulting in the clinical manifestations of thrombocytopenia.¹ Classic clinical signs include purpuric lesions on the skin and mucous membranes, hematuria, and gastrointestinal hemorrhage.³ Diagnostic confirmation is aided by platelet recovery after discontinuation and recurrence upon accidental re-exposure. In this patient, acute platelet reduction manifested shortly after initiating sulfamethoxazole-trimethoprim, consistent with a documented




Image 2. Extensive petechial rash on the patient's lower extremities.

historical episode following exposure to the same agent.

When the specific causative agent cannot be definitively identified, discontinuation of all nonessential medications is warranted.¹ As platelet count improves, the symptoms of thrombocytopenia will begin to subside, usually within 1 to 2 days (4 to 5 half-lives) after stopping the medication.³ Steroids can be used as an adjunct to increase the rate of platelet recovery. Intravenous immunoglobulin (IVIG) and platelet transfusions are typically reserved for severe bleeding, such as intracranial or intrapulmonary hemorrhage, or for persistent thrombocytopenia.^{1,3} Platelet transfusion may not show any benefit if the drug antibodies are still in the patient's system.¹

Table. Platelet count and hospital course.

Hospital Day	Platelet Count ($\times 10^3/\mu\text{L}$)	Clinical Status
1	Less than 1	Oral mucosal hemorrhagic lesions, petechiae
2	11	Oral lesions improving
3	43	Petechiae resolving
4	85	Continued improvement
5	163	Complete recovery

μL , microliters.

CONCLUSION

This case demonstrates the clinical presentation of drug-induced immune thrombocytopenia and the recovery process, focusing on the rate of recovery and treatments that may accelerate it. It is important for clinicians to recognize drug-induced immune thrombocytopenia and consider new medications as possible causes. Immediate cessation of the offending agent is the primary therapeutic intervention for DITP. Steroids, IVIG, and platelets may serve as beneficial adjuncts depending on the clinical picture.

Patient consent has been obtained and filed for the publication of this case report.

Address for Correspondence: Louisa Weindruch, DO, Baylor University Medical Center, Department of Emergency Medicine, 3500 Gaston Ave, Dallas, TX 75246. Email: Louisa.Weindruch@bswhealth.org.

Conflicts of Interest: By the *CPC-EM* article submission agreement, all authors are required to disclose all affiliations, funding sources and financial or management relationships that could be perceived as potential sources of bias. The authors disclosed none.

Copyright: © 2026 Powell et al. This is an open access article distributed in accordance with the terms of the Creative Commons Attribution ([CC BY 4.0](http://creativecommons.org/licenses/by/4.0/)) License. See: <http://creativecommons.org/licenses/by/4.0/>

REFERENCES

1. Arnold DM, Kukaswadia S, Nazi I, et al. A systematic evaluation of laboratory testing for drug-induced immune thrombocytopenia. *J Thromb Haemost.* 2013;11(1):169-176.
2. Aster RH, Curtis BR, McFarland JG, et al. Drug-induced immune thrombocytopenia: pathogenesis, diagnosis, and management. *J Thromb Haemost.* 2009;7(6):911-918.
3. Cimperman C, Whitesel E, Ray A, et al. Drug-induced immune thrombocytopenia secondary to trimethoprim-sulfamethoxazole. *Case Rep Infect Dis.* 2026;5522047.
4. Kaufman DW, Kelly JP, Johannes CB, et al. Acute thrombocytopenic purpura in relation to the use of drugs. *Blood.* 1993;82(9):2714-2718.
5. Marini I, Uzun G, Jamal K, et al. Treatment of drug-induced immune thrombocytopenias. *Haematologica.* 2022;107(6):1264-1277.