UC Davis

UC Davis Previously Published Works

Title

FishShapes v1: Functionally relevant measurements of teleost shape and size on three dimensions

Permalink

https://escholarship.org/uc/item/69c9z03v

Journal

Ecology, 103(12)

ISSN

0012-9658

Authors

Price, Samantha A Friedman, Sarah T Corn, Katherine A et al.

Publication Date

2022-12-01

DOI

10.1002/ecy.3829

Peer reviewed

DATA PAPER



FishShapes v1: Functionally relevant measurements of teleost shape and size on three dimensions

Samantha A. Price¹ | Sarah T. Friedman² | Katherine A. Corn² | Olivier Larouche¹ | Kasey Brockelsby² | Anna J. Lee² | Maya Nagaraj² | Nick G. Bertrand² | Mailee Danao² | Megan C. Coyne² | John R. Estrada² | Rachel Friedman³ | Evan Hoeft² | Mikayla Iwan¹ | Dominique Gross¹ | Jo Hsuan Kao² | Brian Landry² | Monica J. Linares² | Carley McGlinn¹ | Jennifer A. Nguyen² | Allison G. Proffitt² | Sierra Rodriguez¹ | Maxwell R. Rupp² | Erin Y. Shen² | Victoria Susman² | Angelly J. Tovar² | Laura L. J. Vary² | Katerina L. Zapfe¹ | Peter C. Wainwright² |

Correspondence

Samantha A. Price Email: sprice6@clemson.edu

Present address

Olivier Larouche, Department of Biology and Biochemistry, University of Houston, Houston, Texas, USA.

Kasey Brockelsby, Department of Evolution, Ecology, and Behavior, School of Integrative Biology, University of Illinois, Urbana-Champaign, Urbana, Illinois, USA.

Maya Nagaraj, Division of Evolutionary Biology, Ludwig-Maximilians-Universität, Munich, Planegg-Martinsried, Germany.

Nick G. Bertrand, Bureau of Reclamation, Sacramento, California, USA.

Megan C. Coyne, VCA Animal Referral and Emergency Center of Arizona, Mesa, Arizona, USA.

Laura L. J. Vary, College of Earth, Ocean, and Atmospheric Sciences, Oregon State University, Corvallis, Oregon, USA.

Abstract

Teleost fishes account for 96% of all fish species and exhibit a spectacular variety of body forms. Teleost lineages range from deep bodied to elongate (e.g., eels, needlefish), laterally compressed (e.g., ribbonfish) to globular (e.g., pufferfish), and include uniquely shaped lineages such as seahorses, flatfishes, and ocean sunfishes. Adaptive body shape convergence within fishes has long been hypothesized but the nature of the relationships between fish form and ecological and environmental variables remain largely unknown at the macroevolutionary scale. To facilitate the investigation of the interacting factors influencing teleost body shape evolution we measured eight functionally relevant linear traits on adult-sized specimens along with specimen mass. Linear measurements of standard length, maximum body depth, maximum fish width, lower jaw length, mouth width, head depth, minimum caudal peduncle depth, and minimum caudal peduncle width were taken in millimeters with calipers, or tape measures for oversized specimens. We measured these traits on a total of 16,523 specimens (1-3 specimens per species) at the Smithsonian National Museum of Natural History and took approximately 7000 person hours of data collection to complete. The data went through a three-step error-checking process to clean and validate the data and then species averages were calculated. We present the complete specimen data set,

This is an open access article under the terms of the Creative Commons Attribution License, which permits use, distribution and reproduction in any medium, provided the original work is properly cited.

¹Department of Biological Sciences, Clemson University, Clemson, South Carolina, USA

²Department of Evolution & Ecology, University of California Davis, Davis, California, USA

³Independent Researcher

^{© 2022} The Authors. *Ecology* published by Wiley Periodicals LLC on behalf of The Ecological Society of America.

2 of 2 PRICE ET AL.

Katerina L. Zapfe, Department of Genomics and Bioinformatics, University of North Carolina at Charlotte, Charlotte, North Carolina, USA.

Funding information

National Science Foundation, Grant/Award Numbers: DEB 1556953 & 1830127

Handling Editor: William K. Michener

which encompasses approximately one-fifth of extant teleost species diversity, spanning ~90% of teleost families and ~96% of orders. The species and family names are compatible with the taxonomy used by FishBase and the order information with the phylogenetically informed taxonomy of Betancur-R and colleagues published in 2014. This dataset is licensed under Creative Commons CC0 1.0 Universal (CC0 1.0) but please cite this paper when using the data or a subset of it.

KEYWORDS

body depth, body width, caudal peduncle depth, caudal peduncle width, ecomorphology, fishes, head depth, lower jaw length, mouth width, standard length, Teleostei

CONFLICT OF INTEREST

The authors declare no conflict of interest.

DATA AVAILABILITY STATEMENT

The complete data set is available as Supporting Information and is also available from Dryad at https://doi.org/10.5061/dryad.vt4b8gtvf.

ORCID

Samantha A. Price https://orcid.org/0000-0002-1389-8521

Sarah T. Friedman https://orcid.org/0000-0003-0192-5008

Katherine A. Corn https://orcid.org/0000-0002-9437-1683

Olivier Larouche https://orcid.org/0000-0003-0335-0682

Peter C. Wainwright https://orcid.org/0000-0003-0498-4759

SUPPORTING INFORMATION

Additional supporting information can be found online in the Supporting Information section at the end of this article.

How to cite this article: Price, Samantha A., Sarah T. Friedman, Katherine A. Corn, Olivier Larouche, Kasey Brockelsby, Anna J. Lee, Maya Nagaraj, et al. 2022. "FishShapes v1: Functionally Relevant Measurements of Teleost Shape and Size on Three Dimensions." *Ecology* 103(12): e3829. https://doi.org/10.1002/ecy.3829