

UC San Diego

UC San Diego Previously Published Works

Title

Humanized UGT1 Mice, Regulation of UGT1A1, and the Role of the Intestinal Tract in Neonatal Hyperbilirubinemia and Breast Milk-Induced Jaundice

Permalink

<https://escholarship.org/uc/item/69d1n6cn>

Journal

Drug Metabolism and Disposition, 46(11)

ISSN

0090-9556

Authors

Chen, Shujuan

Tukey, Robert H

Publication Date

2018-11-01

DOI

10.1124/dmd.118.083212

Peer reviewed

Special Section – New Models in Drug Metabolism and Transport—Minireview

Humanized *UGT1* Mice, Regulation of *UGT1A1*, and the Role of the Intestinal Tract in Neonatal Hyperbilirubinemia and Breast Milk-Induced Jaundice

Shujuan Chen and Robert H. Tukey

Laboratory of Environmental Toxicology (R.H.T.) and Department of Pharmacology (S.C., R.H.T.), University of California, San Diego, La Jolla, California

Received June 20, 2018; accepted August 3, 2018

ABSTRACT

Neonatal hyperbilirubinemia and the onset of bilirubin encephalopathy and kernicterus result in part from delayed expression of UDP-glucuronosyltransferase 1A1 (*UGT1A1*) and the ability to metabolize bilirubin. It is generally believed that acute neonatal forms of hyperbilirubinemia develop due to an inability of hepatic *UGT1A1* to metabolize efficiently bilirubin for clearance through the hepatobiliary tract. Newly developed mouse models designed to study bilirubin metabolism have led to new insight into the role of the intestinal tract in controlling neonatal hyperbilirubinemia. Humanization of mice with the *UGT1* locus (*hUGT1* mice) and the *UGT1A1* gene provide a unique tool to study the onset of hyperbilirubinemia

since the human *UGT1A1* gene is developmentally regulated during the neonatal period in *hUGT1* mice. A new mechanism outlying developmental expression of intestinal *UGT1A1* is presented and its implications in the control of neonatal hyperbilirubinemia discussed. New findings linking breast milk protection against necrotizing enterocolitis and intestinal control of *UGT1A1* may help explain the contribution of breast milk toward the development of neonatal hyperbilirubinemia. Our findings outline a new model that includes an active intestinal ROS/ $\text{I}\kappa\text{B}$ kinase/nuclear receptor corepressor 1 loop that can be applied to an understanding of breast milk-induced jaundice.

Introduction

UDP-glucuronosyltransferase (*UGT*) 1A1, the only transferase capable of conjugating bilirubin (Bosma et al., 1994), is developmentally delayed in newborn children (Burchell et al., 1989). Thus, approximately 80% of newborns have some form of hyperbilirubinemia (Watchko, 2009; Maisels, 2015). Most cases have a benign outcome, except in situations when the rapid onset of severe neonatal hyperbilirubinemia (SNH) is not monitored nor prevented. Shortly after birth, an increase in red blood cell turnover occurs where heme is released from hemoglobin and further degraded by heme oxygenase to carbon monoxide and biliverdin, which is further reduced by biliverdin

reductase to bilirubin (Tenhunen, 1976; Dennery et al., 2001; Ayer et al., 2016) (Fig. 1). Once in the circulation, unconjugated bilirubin, which is hydrophobic, is readily absorbed into tissues such as the liver. Hyperbilirubinemia develops when a rise in total serum bilirubin (TSB) levels exceeds the capacity of hepatic *UGT1A1* to drive bilirubin conjugation, an event that leads to the excretion of the glucuronide by multidrug resistance protein 2 (Kullak-Ublick et al., 2000) into the biliary channels for deposit in the gastrointestinal tract. Thus bilirubin glucuronidation by *UGT1A1* is the rate limiting step in bilirubin excretion.

Major risk factors that lead to the onset of SNH include accelerated hemolysis (Wong and Stevenson, 2015) brought on by Rhesus disease and ABO incompatibility, glucose-6-phosphate dehydrogenase deficiency, infections, in addition to premature birth and breast milk (Bhutani et al., 2013; Maisels et al., 2014; Olusanya et al., 2015; Wong and Stevenson, 2015; Cunningham et al., 2016). As bilirubin accumulates early after birth and exceeds the capacity to bind to serum proteins, unconjugated bilirubin accumulates in brain tissue.

This work is supported in part by National Institutes of Health National Institute of Environmental Health Sciences [Grants ES010337 and ES024818], National Institute of General Medical Sciences [Grants GM086713 and GM126074], National Institute of Allergy and Infectious Diseases [Grant AI135677].

<https://doi.org/10.1124/dmd.118.083212>.

ABBREVIATIONS: ABE, acute bilirubin encephalopathy; AhR, aryl hydrocarbon receptor; ARE, antioxidant response elements; BM, breast milk; BMJ, breast milk jaundice; CAR, constitutive active receptor; CBE, chronic bilirubin encephalopathy; DEX, dexamethasone; GSH, glutathione; HMO, human milk oligosaccharides; IEC, intestinal epithelial cell; IKK, $\text{I}\kappa\text{B}$ kinase; ITC, isothiocyanates; KSD, kernicterus spectrum disorder; LXR α/β , liver X receptors alpha and beta; NAC, *N*-acetylcysteine; NCoR1, nuclear receptor corepressor 1; NEC, necrotizing enterocolitis; NF, nuclear factor; Nrf2, nuclear E2-factor related factor 2; PAMP, pathogen-associated molecular pattern; Pb, phenobarbital; PEITC, phenethyl isothiocyanate; PPAR α , peroxisome proliferator-activated receptor alpha; PXR, pregnane X receptor; ROS, reactive oxygen species; SI, small intestine; SMRT, silencing mediator of retinoid and thyroid hormone receptor; SNH, severe neonatal hyperbilirubinemia; TLR, Toll-like receptor; TSB, total serum bilirubin; *UGT*, UDP-glucuronosyltransferase; XNR, xenobiotic nuclear receptors.




Fig. 1. Bilirubin production and metabolism. Bilirubin is a byproduct of heme degradation when phagocytic cells, primarily macrophages, remove aged red blood cells (RBC) from the circulation. The production of biliverdin through heme oxygenase is the first major step, after which bilirubin is produced through biliverdin reductase. Bilirubin, on the other hand, is oxidized to biliverdin to form the biliverdin reductase cycle that functions as an important antioxidant in protecting lipid oxidation. Depending on the part of the body in which the breakdown occurs, such as Kupffer cells of the liver or cells in the spleen and bone marrow, the produced bilirubin eventually is released into the circulation, where bilirubin is taken up by plasma albumin protein and transported throughout the body. Once bilirubin reaches the liver, bilirubin enters the hepatocyte either passively or actively through OATP transporters. In the hepatocyte, bilirubin is metabolized through UGT1A1-catalyzed glucuronidation to form either mono- or di-glucuronides. The conjugates are secreted into bile via the multidrug resistance protein 2 (MRP2), and they in turn can be hydrolyzed by β -glucuronidase in the intestinal tract and then undergo enterohepatic circulation. Conventionally, liver is considered as the organ for bilirubin detoxification. It is worth noting that bilirubin is metabolized solely by UGT1A1, which becomes the rate limiting step. Therefore, insufficient UGT1A1 enzyme activity may lead to the rapid build-up of free bilirubin, a condition called hyperbilirubinemia or jaundice.

Extreme levels of TSB can lead to early or acute bilirubin encephalopathy (ABE) (Watchko, 2016), presented early as lethargy and poor feeding, but can progress to hypo- and hypertonia, high-pitched crying, muscle spasms, opisthotonus, seizures, and even death (Bhutani and Johnson-Hammerman, 2015). The more chronic form, which proceeds ABE, is termed kernicterus (Kaplan et al., 2011). Affected individuals are characterized by choreoathetotic cerebral palsy, dystonic/athetoid movement disorders, hearing loss, ocular motor defects, hypotonia, and ataxia, and it has been linked to cerebellar involvement (Dennerly et al., 2001; Tabarki et al., 2002; Kaplan et al., 2011; Bhutani and Johnson-Hammerman, 2015). Kernicterus was originally classified as the pathologic finding of yellow staining (icterus) of deep nuclei in the brain. Improved imaging technology has allowed for visualization of bilirubin damage to specific regions, e.g., the globus pallidus and subthalamic nucleus. Thus, there are two major periods after birth where SNH can lead to toxicity. ABE is designated for the early signs of toxicity, and chronic bilirubin encephalopathy (CBE) covers the long-term consequences of an acute but resolved process. Because the clinical signs of CBE are broad, it has been recommended to abandon the use of CBE for the term kernicterus spectrum disorder (KSD) (Le Pichon et al., 2017).

SNH, which leads to ABE and KSD, has been observed worldwide (Kaplan et al., 2011), with the highest incidences being recorded in sub-Saharan Africa and South Asia (Bhutani et al., 2013; Olusanya et al., 2016). A report from the Nigerian Society of Neonatal Medicine suggested that extreme hyperbilirubinemia accounts for at least 5% of all neonatal mortalities in Nigeria (Olusanya et al., 2016). Recent evidence shows that SNH, estimated to impact over 1 million children every year, is associated with substantial mortality and permanent morbidities (Bhutani et al., 2013; Olusanya et al., 2018). SNH accounted for over 1300 newborn deaths per 100,000 globally, ranking as the seventh most dangerous contributor toward infant deaths (Olusanya et al., 2018). If global deaths linked to SNH in children under 5 years of age are estimated, the mortality rate in Indonesia is calculated to exceed 40% of all deaths. While conventional therapies entail aggressive blue light therapy or blood transfusions once extreme hyperbilirubinemia has been diagnosed (Dennerly et al., 2001), the inability to conjugate effectively bilirubin resulting from developmental delay in expression of UGT1A1 can ultimately give rise to ABE and other KSDs (Johnson et al., 1999; Yueh et al., 2017). While genetic predisposition and

environmental influences play key roles in driving TSB levels to potentially toxic levels, an understanding of the developmentally regulated molecular and cellular mechanisms that underlie the regulation of UGT1A1 expression have not been elucidated. With the development of new animal models designed to examine expression of the human *UGT1A* genes, progress has been made in characterizing the regulatory events that underlie the expression of the *UGT1A1* gene during development. Our understanding of the events that control expression of the human *UGT1A1* gene has provided new insight toward the mechanisms leading to hyperbilirubinemia and disease.

Results

Humanized *UGT1* Mice and Neonatal Hyperbilirubinemia.

Cellular and molecular events that underlie the regulation of the UGTs, especially those encoded by the human *UGT1* gene family, have been studied in humanized transgenic mice that express the human *UGT1* locus (*TgUGT1* mice) in a *Ugt1*-null background (*Ugt1*^{-/-} mice) (Fujiwara et al., 2010). In humans, there are nine *UGT1A* genes, including *UGT1A1*, that are encoded by the *UGT1* locus (Clarke et al., 1997; Mackenzie et al., 1997; Gong et al., 2001). To examine the importance of the human *UGT1* locus and the *UGT1A1* gene in control of serum bilirubin, an inactivating mutation was engineered into exon 4 of the murine *Ugt1a1* gene, generating *Ugt1*^{-/-} mice (Nguyen et al., 2008). The UGT1A proteins are encoded by five exons, with coding sequence for exons 2–5 being identical. Located on chromosome 1, the organization of the murine *Ugt1* locus is encoded by >150 kDa of DNA, with exons 2–5 located at the 3'-end and flanked consecutively by an array of nine cassette exons that encode the N-terminal portion of each UGT1A protein. Thus the *UGT1* locus in humans and the *Ugt1* locus in mice encodes nine functional *UGT1A* proteins, each composed of a unique N-terminal region encoded by one of the 5-flanking exons while sharing an identical C-terminal region encoded by exons 2–5. When heterozygous *Ugt1*^{+/-} mice were bred, offspring with a *Ugt1*^{-/-} genotype were born with a striking orange skin color that resulted from severely elevated UCB levels that developed into lethal SNH between 4 and 7 days after birth. To recover the lethality resulting from nonfunctional murine UGT1A1 expression, *TgUGT1* mice that carried

and expressed the entire *UGT1* locus (Chen et al., 2005) were crossed with *Ugt1*^{+/-} mice to generate *TgUGT1/Ugt1*^{+/-} mice. These founders were then backcrossed to generate *TgUGT1/Ugt1*^{-/-} mice or humanized *UGT1* (*hUGT1*) mice. Expression of the human *UGT1A1* gene rescued the lethality attributed to SNH in *Ugt1*^{-/-} mice (Fig. 2).

Developmental and tissue specific regulation of the *UGT1A* genes in *hUGT1* mice (Chen et al., 2005; Senekeo-Effenberger et al., 2007; Fujiwara et al., 2010; Yueh et al., 2017) has been shown to reflect similar patterns of expression as characterized in human tissues (Strassburg et al., 1997a,b, 1998a,b; Tukey and Strassburg, 2000), allowing us to leverage the use of *hUGT1* mice in experiments designed to characterize the mechanisms leading to tissue specific, inducible and developmental expression of the human *UGT1A* genes. An important discovery was the finding that the human *UGT1A1* gene, which is developmentally delayed in newborn children, is also developmentally delayed or repressed in *hUGT1* neonatal mice (Fujiwara et al., 2012). Since *UGT1A1* is the sole enzyme responsible for the metabolism of serum bilirubin, this delay in hepatic *UGT1A1* gene expression leads to SNH in neonatal *hUGT1* mice, with lethality affecting 10% of the newborns, as displayed by dramatic motor impairment with seizure-like movements. The brains of *hUGT1* mice that develop seizures concentrate bilirubin in the tissues, which is evident by an intense yellow or icteric appearance (Fujiwara et al., 2010). The icteric pattern is not regionally localized as found in the globus pallidus and subthalamic nucleus in human brain tissue, although we have classified this phenotype in neonatal *hUGT1* mice as kernicterus. The peak development of TSB levels in neonatal *hUGT1* mice occurs at days 10–14 after birth and sharply declines through the next 5 days. Throughout the neonatal period, which we

classify as the first 3 weeks after birth, liver *UGT1A1* expression is repressed. The clearance of TSB in the later stages of the neonatal period corresponds with developmental induction of intestinal *UGT1A1* expression (Fujiwara et al., 2010) (Fig. 2).

During this developmental phase, dietary or environmentally driven activation of the *UGT1A1* gene and induction of *UGT1A1* in *hUGT1* mice results in metabolism of bilirubin and a sharp decrease in TSB levels. Mapping nuclear receptor binding sites associated with the *UGT1A1* gene has indicated that transcriptional activation can be achieved by activated subfamily 3 steroid receptors in addition to subfamily 1 xenobiotic nuclear receptors (XNRs) (Fig. 3). Sugatani et al. (2001) first characterized the phenobarbital (Pb) responsive enhancer module in the human *UGT1A1* gene that binds to the constitutive active receptor (CAR). When *hUGT1* neonatal mice are treated with Pb, liver *UGT1A1* is induced and TSB levels are dramatically reduced (Chen et al., 2012; Shibuya et al., 2013; Yoda et al., 2017; Yueh et al., 2017). However, in *hUGT1/Car*^{-/-} mice, Pb administration does not induce *UGT1A1* and has no impact on neonatal TSB levels, confirming genetically that CAR regulates the *UGT1A1* gene. In *hUGT1* mice, the *UGT1A1* gene can be transcriptionally induced by other XNRs, including the pregnane X receptor (PXR) (Chen et al., 2012), the peroxisome proliferator-activated receptor alpha (PPAR α) (Senekeo-Effenberger et al., 2007), the liver X receptors alpha and beta (LXR α/β ; unpublished observations), and the environmental sensors, the aryl hydrocarbon receptor (AhR) (Yueh et al., 2003; Chen et al., 2005; Bonzo et al., 2007), and Nrf2 (Yueh and Tukey, 2007; Yoda et al., 2017). The oral or intraperitoneal administration of ligands capable of activating these receptors in neonatal *hUGT1* mice leads to the




Fig. 2. Humanized *UGT1* mice and neonatal hyperbilirubinemia. *Ugt1*^{-/-} mice were generated by interrupting common exon 4 with a neomycin gene, impairing the function of the entire *Ugt1* locus. *Ugt1*^{-/-} newborn mice are easily identified by their distinct orange skin color, which results from the accumulation of unconjugated bilirubin. The loss of the *Ugt1* locus and *UGT1A1* from the germ line is lethal, with all the *Ugt1*^{-/-} mice dying within 2 weeks after birth. When we crossed mice carrying the entire human *UGT1* gene locus as a transgene with mice carrying the *Ugt1*-null allele, humanized *UGT1* mice (*hUGT1*) were generated. The presence of the human *UGT1A1* gene rescues the lethality associated with the absence of murine *UGT1A1*. All newborn *hUGT1* mice develop severe neonatal hyperbilirubinemia (SNH). Total serum bilirubin (TSB) levels in *hUGT1* newborns increase after birth, reach peak levels at approximately 2 weeks after birth with the levels returning to a normal range after weaning. The majority of the *hUGT1* mice can mature to adulthood. Approximately 10% of newborn *hUGT1* mice develop seizures and die. These mice accumulate high concentrations bilirubin in brain tissue. The development of SNH results from delayed liver *UGT1A1* expression. *UGT1A1* expression in the small intestine is associated with saving the lethality of *hUGT1* mice and lowering of TSB levels.




Fig. 3. Nuclear receptor (NR)-mediated activation, repression, and derepression of human *UGT1A1* gene and the impact on hyperbilirubinemia in *hUGT1* mice. Binding elements of various NRs, including PPAR α , AhR, Nrf2, CAR, PXR, and LXR α , have been identified in the promoter region upstream of the transcription start site of the *UGT1A1* gene. Upon ligand binding, these NRs can recruit coactivators and initiate downstream *UGT1A1* gene expression. When *hUGT1* neonates were exposed to different NR agonists, lowered serum bilirubin levels were observed (except for PPAR γ). In the absence of ligand, some NRs associate with NR corepressors (such as NCoR1 and SMRT) and recruit chromatin-modifying enzyme HDACs to form a repressive complex and limit transcriptional activation. Disrupting of the repressive complex may lead to the derepression of gene expression. Humanized *UGT1* mice deficient in PXR (*hUGT1/Pxr $^{-/-}$* and *hUGT1/Pxr $^{-/-}$ /Car $^{-/-}$*), but not CAR (*hUGT1/Car $^{-/-}$*), are associated with a reduction of total serum bilirubin. When selectively silenced, the corepressors SMRT, NCoR1, or HDACs in the primary hepatocytes isolated from *hUGT1* mice, *UGT1A1* gene activation was observed.

induction of liver or intestinal *UGT1A1* followed by reductions in TSB levels. The reduction in TSB levels early in the neonatal window assures that the mice avoid bilirubin-induced neurologic dysfunction (Fig. 3).

PXR is a Steroid Target, Repressor Protein, and Regulator of *UGT1A1*. The importance of PXR as a regulator of endocrine control, drug metabolism, and hyperbilirubinemia started to emerge when we predicted that PXR controlled UGT-directed drug metabolism during gestation. For example, circulating TSB levels are lower in women during pregnancy, indicating that bilirubin metabolism is accelerated through induced *UGT1A1* (Bacq et al., 1996). Labetalol, an antihypertensive drug that is metabolized primarily by *UGT1A1*, shows increased clearance in the second and third trimesters of pregnancy compared with the postpartum period (Jeong et al., 2008). Pregnancy results in dramatic surges in hormones during gestational development, with increases in steroids such as estrogens, progestins, and the glucocorticoids. Since PXR had been predicted to serve as a low affinity endogenous target for circulating steroid pools that would eventually result in physiologic changes such as drug metabolism (Blumberg et al., 1998), we examined the impact of PXR on control of the *UGT1A* genes during pregnancy in *hUGT1*1* and *hUGT1*28* mice (Fujiwara et al., 2010). Adult *hUGT1*28* mice are hyperbilirubinemic, with TSB levels that average 1 mg/dl compared with undetectable levels in *hUGT1*1* mice. During pregnancy and late gestation, the TSB levels in *hUGT1*28* mice averaged 0.4 mg/dl, over 50% lower than in nonpregnant mice. During late gestation, the *UGT1A* proteins are induced in liver. Activation of PXR was also documented by CHIP analysis, demonstrating increased binding of PXR associated with the *UGT1A1* gene during late pregnancy (Chen et al., 2012). To examine the role of PXR as a modulator of drug metabolism, we crossed *hUGT1*1* mice with *Pxr $^{-/-}$* mice and

examined expression of the *UGT1* locus in pregnant *hUGT1/Pxr $^{-/-}$* mice. In comparison with *hUGT1* mice, *UGT1A1*, *-IA3*, *-IA4*, *-IA6*, and *-IA9* gene expression in *hUGT1/Pxr $^{-/-}$* mice are greatly reduced during the later stages of gestation. When the glucocorticoid dexamethasone (DEX) was administered to *hUGT1* mice, each of the five *UGT1A* genes expressed in the liver was induced. Although activation of the glucocorticoid receptor by DEX was shown previously to activate *UGT1A1* reporter gene constructs, DEX had no effect on induction of *UGT1A1* in *hUGT1/Pxr $^{-/-}$* mice, indicating that the induction of the *UGT1A* genes by glucocorticoids is facilitated solely by activation of PXR in vivo.

Transcriptional Silencing of the *UGT1A1* Gene. In breeding experiments to create *hUGT1/Pxr $^{-/-}$* mice, analysis of TSB levels in neonatal *hUGT1* and *hUGT1/Pxr $^{-/-}$* mice demonstrated that PXR plays a key role in controlling TSB levels during development. In neonatal *hUGT1/Pxr $^{-/-}$* mice, liver *UGT1A1* is induced when compared with its expression in *hUGT1* mice (Chen et al., 2012). The increased levels of liver *UGT1A1* in *hUGT1/Pxr $^{-/-}$* mice lead to reduced TSB levels. This finding suggests that in the absence of endogenous ligand, the physiologic role of PXR leads to repression of the *UGT1A1* gene during early development. More importantly, blocking activation of the *UGT1A1* gene by PXR is controlled in liver through neonatal developmental processes, since its deletion in adult *hUGT1/Pxr $^{-/-}$* mice has no impact on *UGT1A1* gene expression, except for pregnant mice. Thus PXR is involved in transcriptional silencing of the *UGT1A1* gene (Fig. 3).

In combination with nuclear receptors, transcriptional silencing, or repression is largely achieved with two prototypical nuclear repressor corepressors, the silencing mediator of retinoid and thyroid hormone

receptor (SMRT), and the nuclear receptor corepressor 1 (NCoR1) (Francis et al., 2003; Mottis et al., 2013), suggesting that NCoR1 and/or SMRT participate in control of *UGT1A1* expression during development. Mechanistically, corepressor proteins interact with XNRs, such as RXR, LXR, PPAR, and PXR, to control collectively gene expression patterns by recruiting chromatin-modifying enzymes that limit nucleosomal DNA accessibility and transcriptional activation. In the absence of either the nuclear receptor or the corepressor, additional coactivators cooperate with a different set of chromatin-modifying enzymes to promote transcriptional activation. In experiments targeted to delete NCoR1/SMRT in various tissues or cell lines, deletion of the corepressor spontaneously leads to nonligand activation of the XNRs, facilitating transcriptional activation of XNR target genes (Li et al., 2013). Often transcriptional activation of downstream target genes by spontaneously activated XNRs provides a genomic signature that can be used to identify the exact XNR involved in the response. To determine if NCoR1 and SMRT are engaged in repression of the *UGT1A1* gene in liver, knockdown of NCoR1 and SMRT in primary neonatal *hUGT1* hepatocytes by siRNA leads to derepression of *UGT1A1* (Chen et al., 2017). Similar effects were seen after HDAC1 or HDAC3 ablation, indicating that NCoR1 and SMRT are implicated along with HDAC1/HDAC3 in repressing expression of *UGT1A1* gene expression. While knockdown of genes in tissue culture is an effective tool to identify potential regulatory events, targeted tissue specific knockout of genes in vivo provides additional evidence of the biochemical and physiologic impact of their role in normal homeostasis. With evidence that PXR along with NCoR1 and SMRT is tied to neonatal repression of liver *UGT1A1* expression, experiments were initiated using conditional deletion of NCoR1 to examine the molecular mechanisms linking neonatal development with liver and intestinal *UGT1A1* gene expression (Fig. 3).

An important role for NCoR1 in regulating *UGT1A1* gene expression in neonates has emerged (Chen et al., 2017). We generated floxed *Ncor1*

mice under a *hUGT1* background (*TgUGT1/Ugt1^{-/-}/Ncor1^{F/F}*), named as *F/F^{UN}* mice, and then further knocked out NCoR1 in liver (Δ *HEP^{UN}* mice) and intestinal tissue (Δ *IEC^{UN}* mice) of *hUGT1* mice. When TSB levels were examined in newborn mice, wild-type *F/F^{UN}* and Δ *HEP^{UN}* mice exhibited similar patterns of neonatal hyperbilirubinemia as observed in *hUGT1* mice. Deletion of NCoR1 in liver tissue had no impact on hepatic *UGT1A1* gene expression. Strikingly, following intestinal-specific deletion of NCoR1, the development of neonatal hyperbilirubinemia was completely diminished in Δ *IEC^{UN}* mice. The impact of deleting intestinal NCoR1 led to the derepression of intestinal *UGT1A1* expression with significant induction of protein occurring in both the longitudinal axis [i.e., all sections of the small intestine (SI)] as well as the crypt-villi axis. Mechanistically, derepression or induction of *UGT1A1* following NCoR1 knockout was observed only during neonatal developmental, since derepression of *UGT1A1* did not occur in adult Δ *IEC^{UN}* mice. This interesting observation indicates that NCoR1 plays a key role as a corepressor protein in developmentally important processes linked to intestinal homeostasis. These processes may be directly, or indirectly, linked to derepression of the *UGT1A1* gene (Fig. 4).

When we conducted transcriptional profiling of intestinal tissue by implementing RNA-seq analysis of *F/F^{UN}* and Δ *IEC^{UN}* mice, gene ontology analysis identified the metabolic pathway as the most significantly altered pathway in Δ *IEC^{UN}* mice, followed by oxidative phosphorylation and drug metabolism. Deleting NCoR1 promoted expression of PPAR α and LXR α target genes, indicating that NCoR1 is docked on these genes during neonatal development. NCoR1 deletion had an extensive impact on fatty acid and energy metabolism, in addition to lipid and cholesterol metabolism. Messenger RNA levels of TCA cycle genes and oxidative phosphorylation genes were coordinately increased. These accelerated metabolic pathways indicated that NCoR1 was repressing key pathways essential for epithelial cell maturation and synthesis. In Δ *IEC^{UN}* neonatal mice, the average length of the small




Fig. 4. NCoR1 represses human *UGT1* gene expression during development. *NCoR1^{F/F}* mice were crossed into *hUGT1* to generate *F/F^{UN}* mice, which in turn were crossed with mice carrying Cre recombinase and knocked out NCoR1 specifically in liver (Δ *HEP^{UN}*) and intestinal tissue (Δ *IEC^{UN}*). Serum bilirubin levels were examined in neonatal mice, Δ *HEP^{UN}* mice exhibited a similar pattern of neonatal hyperbilirubinemia as *hUGT1* (*F/F^{UN}*) mice, but this pattern was completely diminished with NCoR1 knockout in the intestinal tract of Δ *IEC^{UN}* mice. The change of bilirubin accumulation was also evidenced in fat tissues. Induction of intestinal *UGT1A1* in Δ *IEC^{UN}* mice was observed and was most prominent throughout the developmental stage but did not occur after weaning or in adulthood. Data are expressed as mean \pm SEM, ** $p < 0.01$, *** $p < 0.001$, **** $p < 0.0001$, Student's t test. Two-way ANOVA analysis for Δ *IEC^{UN}* vs *F/F^{UN}* ($p < 0.0001$).

intestines was ~8% longer and 15% heavier than in F/F^{UN} littermates, indicating increased intestinal proliferation. Histology and electron microscopy findings indicated greatly matured brush border and microvilli in ΔIEC^{UN} . When we examined cellular proliferation signals, immunostaining of Ki-67, a cellular marker for proliferation, was increased in both ΔIEC^{UN} duodenum and jejunum. In support of this finding, bromodeoxyuridine staining, used to detect proliferating cells, showed a progressive increase in positive epithelial cells. These findings demonstrated that NCoR1 deletion was the catalyst to accelerate cell proliferation and promote epithelial cell proliferation. In support of cellular proliferation, upregulation of intestinal maturation Sucrase isomaltase, a brush border glucosidase that only exists in differentiated intestinal tissue and is a marker for enterocyte maturation, was significantly induced in ΔIEC^{UN} mice, along with other maturation markers that included the duodenal-specific *Akp3* gene and the jejunal-specific *Krt20* gene. These studies confirmed that neonatal NCoR1 expression serves to repress gene expression, including UGT1A1, while mechanistically controlling the developmental changes underlying intestinal maturation from the neonatal stage to the adult tissue.

NCoR1 Regulation and Expression of Intestinal UGT1A1. Intestinal NCoR1 may be the centerpiece in controlling bilirubin metabolism in intestinal epithelial cells (IECs) during the neonatal window. The functional ability of NCoR1 to promote its repressive properties with nuclear receptors as well as its potential to activate gene transcription have been linked to phosphorylation events (Mottis et al., 2013; Jo et al., 2015), controlled by a number of different kinases (Fernández-Majada et al., 2007; Huang et al., 2009; Choi et al., 2013). For example, the loss of nuclear NCoR1 in malignant melanoma was associated with IKK-dependent phosphorylation of specific NCoR1 serine residues (Gallardo et al., 2015). When NCoR1 is phosphorylated, it dissociates from chromatin and migrates to the cytoplasm, a move that stimulates transcriptional derepression events. Findings from our studies have also established close ties between IKK activation, NCoR1 function, and reductions in TSB levels in *hUGT1* mice (Chen et al., 2017). Intestinal IKK β is critical in controlling basal UGT1A1-dependent metabolism of bilirubin. When IKK β is deleted in IECs in *hUGT1* mice (*hUGT1/Ikk β ^{ΔIEC}* mice), basal expression of intestinal UGT1A1 in neonatal mice drops, leading to elevations in TSB levels that are lethal (Liu et al., 2016). When NCoR1 is dephosphorylated, as potentially exists when IKK β is deleted, we hypothesize that the NCoR1/NR complex tightens its binding to chromatin to improve transcriptional repression, as observed by greater reduction in IEC expression of UGT1A1. This hypothesis was tested by overexpressing active IKK β in IECs, an event that led to derepression of intestinal maturation genes, like the aforementioned responses we documented when NCoR1 was deleted (Chen et al., 2017). The ability of constitutively activated IKK β to mimic the actions of NCoR1 deletion was also demonstrated when *hUGT1* neonatal mice were orally treated with cadmium (Cd^{2+}), acting both as a potent inducer of oxidative stress and an activator of the IKK β (NF- κ B) pathway (Liu et al., 2016). Cadmium treatment resulted in intestinal induction of UGT1A1, a sharp reduction in TSB levels, and induction of intestinal maturation genes, events that are observed when NCoR1 is deleted. Cd^{2+} -dependent induction of intestinal UGT1A1 could be driven by activation of IKK, through which it can directly, or through alternative mechanisms, contribute to the production of reactive oxygen species (ROS). Interestingly, both processes could play a role in targeting the phosphorylation of NCoR1, an event that would result in activation of the intestinal *UGT1A1* gene.

Oxidative Stress and Regulation of UGT1A1. Knowing that constitutive expression of intestinal IKK β is critical in maintaining adequate expression levels of the *UGT1A1* gene, most likely through

activating and modifying NCoR1, we examined the impact of intestinal oxidative stress toward induction of UGT1A1. The rationale for this experiment stemmed from findings that the IKK β /NF- κ B pathway responds to redox changes, such as alterations in ROS. To carry out this experiment and eliminate the role of IKK β , *hUGT1* and *hUGT1/Ikk β ^{ΔIEC}* mice were treated orally with either Cd^{2+} or arsenic (As^{3+}), both known to induce significant oxidative stress (Santra et al., 2000; Ercal et al., 2001; Li et al., 2002; Liu et al., 2003; López et al., 2006). Untreated *hUGT1/Ikk β ^{ΔIEC}* mice displayed reduced intestinal UGT1A1 expression and elevated TSB levels, with most of the mice dying from SNH (Liu et al., 2016). When neonatal *hUGT1* mice were orally treated with either Cd^{2+} or As^{3+} , intestinal UGT1A1 was induced and TSB levels were lowered (Liu et al., 2016; Yoda et al., 2017). Likewise, when *hUGT1/Ikk β ^{ΔIEC}* mice were treated with Cd^{2+} or As^{3+} , oxidative stress was induced in the intestines, commensurate with induction of UGT1A1 and a lowering of the TSB levels. While untreated neonatal *hUGT1/Ikk β ^{ΔIEC}* mice accumulated toxic levels of TSB that were lethal, oral treatment with Cd^{2+} or As^{3+} led to induction of intestinal UGT1A1 and lowering of TSB levels (Fujiwara et al., 2012; Liu et al., 2016). Arsenic, an inducer of oxidative stress, leads to the rapid activation of p38 mitogen-activated protein kinase (Yoda et al., 2017) and extracellular signal-regulated kinase, both sensors of oxidative stress. Interestingly, oral As^{3+} treatment confirmed that IEC maturation marker genes were also induced. In addition, expression of *Glb1*, *Nox4*, and *Lrp2* genes, which are downregulated at the suckling to weaning transition, was reduced, mimicking the same pattern observed in ΔIEC^{UN} mice. The activation of MAPK and extracellular signal-regulated kinase by As^{3+} may be stimulating kinase activity in the IECs, promoting derepression of gene expression in an NCoR1-dependent fashion.

A preemptive assumption of oxidative stress in tissues and cells is the existence or evolution of a potent defense system to counter the potential toxicities associated with the generation of ROS. The formation of oxidative stress, induced by natural diet or through environmental exposure, sets into motion a unique signaling cascade that activates gene families that regulate cytoprotective proteins to provide an active defense against cellular damage. The central mediator of this defense is the nuclear E2-factor related factor 2 (Nrf2), which exists as an Nrf2-Keap1 complex that is sensitive to ROS (Wasserman and Fahl, 1997). Activation of Nrf2 sets into motion transcriptional activation of target genes following binding to unique enhancer sequences identified as antioxidant response elements (AREs) (Wasserman and Fahl, 1997; Nguyen et al., 2000). The ARE sequences are clustered with the AhR xenobiotic response element flanking the *UGT1A1* gene and bind activated Nrf2 (Yueh and Tukey, 2007). The antioxidant response can target the *UGT1A1* gene, in addition to other genes responsive to ROS, such as NAD(P)H:quinine oxidoreductase 1 (McWalter et al., 2004), glutathione-S-transferase, and heme oxygenase (Alam et al., 1999). Thus induction of these genes following oxidative stress provides a molecular footprint that can be leveraged to identify substances that activate the Nrf2-Keap1 complex in providing antioxidant protection.

CAR, Oxidative Stress, and UGT1A1. Some of the more potent inducers of oxidative stress are the isothiocyanates (ITCs). Regarded as an important component of our diet, cruciferous vegetables (i.e., broccoli, Brussels sprouts, cauliflower, watercress) have been linked to a lower incidence of cancer development (Murillo and Mehta, 2001) because they produce antioxidative metabolites that induce cytoprotective proteins, such as the UGTs, believed to protect tissue from dietary or environmental toxicity such as cancer. These vegetables are rich in glucosinolates, which are hydrolyzed by plant myrosinases following chewing or by microflora in the gastrointestinal tract to form ITCs (Shapiro et al., 2001a,b, 2006), the active anticarcinogenic metabolites.

Isothiocyanates activate the Nrf2-Keap1 complex, initiating a series of events leading to induction of ARE containing target genes (Nguyen et al., 2003). We rationalized that ITC treatment would induce intestinal and potentially liver UGT1A1 in *hUGT1* mice and serve as an efficient modulator of neonatal hyperbilirubinemia by inducing UGT1A1 and lowering TSB levels. When neonatal *hUGT1* mice were treated orally with phenethyl isothiocyanate (PEITC), TSB levels were reduced to normal after 24 hours (Yoda et al., 2017). Induction of UGT1A1 was dramatic in both liver and small intestines, but the impact on the classic Nrf2-target genes *Nqo1* and *Gsta1/2* was not significant. We showed previously that oral As^{3+} , which generates a potent antioxidant response, has a dramatic impact on *Gsta1* and *Gsta2* induction while also upregulating several of the cytochrome P450 2 and 4 subfamilies (Liu et al., 2016), indicating that the antioxidant response potentially initiated by PEITC treatment could be influencing the activation of additional signaling pathways. A screen of potential target genes induced by XNRs resulted in prominent induction of *Cyp2b10*/CYP2B10 in both liver and SI, indicating that CAR may be involved in the PEITC induction of UGT1A1 (Fig. 5).

To examine the component of CAR linking induction of UGT1A1 and CYP2B10 by PEITC, *hUGT1/Car^{-/-}* were bred to generate CAR knockout *hUGT1/Car^{-/-}* and heterozygous *hUGT1/Car^{+/-}* littermates. Both *hUGT1/Car^{-/-}* and *hUGT1/Car^{+/-}* neonatal mice develop hyperbilirubinemia at the same rate. When treated with an oral dose of PEITC, UGT1A1 and CYP2B10 were induced in liver and SI of *hUGT1/Car^{+/-}* mice, with a total reduction in TSB levels. The induction of hepatic UGT1A1 was significant and most likely drove the metabolism of bilirubin, leading to the reduction in TSB levels. In liver tissue from PEITC-treated *hUGT1/Car^{-/-}* mice, there was no expression of CYP2B10 or UGT1A1, indicating that CAR was essential for induction by PEITC. In SI, the induction of CYP2B10 was also absent, yet TSB levels were significantly reduced. In the absence of CAR, there was induction of UGT1A1 in the SI, which we expect was responsible for the reduction in TSB levels. While the induction by PEITC in SI could be attributed to activation of Nrf2, there was no significant transcriptional activation of other ARE target genes. The possibility exists that PEITC induction of intestinal UGT1A1 in neonatal mice is being driven by a mechanism that is independent from CAR and Nrf2 activation (Fig. 5).

Confirmation that CAR underlies CYP2B10 and UGT1A1 induction by PEITC was validated in adult *hUGT1/Car^{+/-}* and *hUGT1/Car^{-/-}* mice. In liver and SI, oral PEITC treatment induced CYP2B10 in a

strictly CAR-dependent fashion. Significant induction of CYP2B10 by PEITC in these tissues was generated only in *hUGT1/Car^{+/-}* mice, with no response in *hUGT1/Car^{-/-}* mice in either tissue, leaving us to speculate that induction of CYP2B10 by PEITC involved activation of CAR. A different pattern of induction was observed with UGT1A1 expression. In liver, UGT1A1 expression was greatly reduced in *hUGT1/Car^{-/-}* mice, clearly linking CAR activation with UGT1A1 expression. However, in SI, PEITC treatment effectively induced UGT1A1 in both *hUGT1/Car^{+/-}* and *hUGT1/Car^{-/-}* mice, indicating that alternative CAR processes, such as activation of Nrf2, are responsible for the induction. In addition, Nrf2 target genes were induced in both liver and SI, confirming that the actions of PEITC are also initiating an ARE response. These findings indicate that two important signaling events, the activation of CAR and Nrf2, are linked to induction of UGT1A1 (Fig. 5).

Independently, the activation of CAR by ligands or the activation of Nrf2 by oxidative stress can facilitate the induction of UGT1A1. However, with PEITC treatment, evidence suggests that CAR and Nrf2 are linked in the induction of liver UGT1A1. To separate an antioxidative response initiated by PEITC from activation of CAR and induction of UGT1A1, we first administered adult *hUGT1* mice *N*-acetylcysteine (NAC) for 3 weeks in their drinking water. Cysteine is the rate-limiting substrate for the synthesis of glutathione (GSH), and thus supplementation of NAC promotes the resynthesis of GSH and the neutralization of free radicals (Rizzardini et al., 1994). GSH has antioxidant properties in addition to facilitating the activity of enzymatic antioxidant systems, including GSH peroxidase and GSH reductase. After 2 weeks of NAC exposure, the mice were treated with PEITC, and antioxidant responses were measured. In the absence of NAC exposure, PEITC treatment leads to robust induction of Nrf2 target genes *Nqo1*, *Gsta1*, and *Gsta2*, along with *UGT1A1* and *Cyp2b10*. In mice exposed to NAC and treated with PEITC, *Cyp2b10* gene expression was completely suppressed, along with *Nqo1*, *Gsta1*, and *Gsta2*. While PEITC induces oxidative stress that targets Nrf2 target genes, induction of UGT1A1 and CYP2B10 is dependent upon CAR. These findings indicate that induction of liver UGT1A1 by PEITC is dependent upon both Nrf2 and CAR, while the generation of oxidative stress is a prerequisite toward CAR activation and induction of UGT1A1 and CYP2B10 (Fig. 5).

Breast Milk-Induced Hyperbilirubinemia. Breast milk (BM) plays a significant role in neonatal hyperbilirubinemia (Newman and Gross, 1963; Fujiwara et al., 2015), termed breast milk jaundice (BMJ) and is




Fig. 5. PEITC regulates *UGT1A1* gene expression through ROS-mediated nuclear receptor signaling. Consumption of cruciferous vegetables leads to the generation of reactive oxygen species (ROS). ROS activates Nrf2 nuclear translocation, thus facilitating Nrf2-antioxidant response element (ARE) signaling and the transcriptional activation of the susceptible target genes. In addition, ROS are linked to the activation of CAR in both liver and small intestines, which results in the induction of *Cyp2b10* and *UGT1A1*. The addition of *N*-acetylcysteine (NAC) blocks ROS production, nullifying both CAR and Nrf2 activation.

characterized by prolonged increases in unconjugated bilirubin. Infants with BMJ have elevated TSB levels above 10 mg/dl that can reach levels above 20 mg/dl, approaching the range that can induce bilirubin-induced neurologic toxicities. The prolonged increases in TSB levels during the neonatal window in *hUGT1* mice indicate that breast milk may be participating in the development of hyperbilirubinemia. When neonatal *hUGT1* mice were fed human breast milk, the mice continued to develop jaundice (Fujiwara et al., 2012). Clinical studies have demonstrated that newborns with jaundice given a formula diet have lower levels of TSB than breast-fed newborns, indicating that breast milk facilitates the repression of UGT1A1 (Schneider, 1986; Huang et al., 2004). To examine if we could replicate this effect in *hUGT1* mice, neonatal mice were given a diet of baby formula for 72 hours (Fujiwara et al., 2012). After formula treatment, TSB levels were reduced over 95% compared with breast-fed mice. Formula induced UGT1A1 in intestinal tissue, which correlated with the reductions in TSB levels. Of significance, formula also induced the CAR target gene *Cyp2b10* and PXR target gene *Cyp3a11*, indicating that formula targeted activation of CAR and PXR. When formula was given to neonatal *hUGT1/Car^{-/-}* or *hUGT1/Pxr^{-/-}* mice, intestinal UGT1A1, *Cyp2b10*, and *Cyp3a11* were induced in both strains, indicating formula was not activating either CAR or PXR. However, formula feeding led to activation of $\text{I}\kappa\text{B}/\text{NF-}\kappa\text{B}$ target genes, linking activation of $\text{NF-}\kappa\text{B}$ with induction of intestinal UGT1A1. Given the finding that formula leads to activation of $\text{NF-}\kappa\text{B}$ and induction of intestinal UGT1A1 in neonatal *hUGT1* mice, it can be speculated that the repressive effects of BM are tied with inhibition of the $\text{IKK}/\text{NF-}\kappa\text{B}$ pathway.

While $\text{I}\kappa\text{B}$ kinase (IKK) is an upstream component of $\text{NF-}\kappa\text{B}$ signaling, it is linked to propagating the cellular response to inflammation (Karin, 2009) and plays an important role in neonatal intestinal UGT1A1 expression (Fujiwara et al., 2012). Human breast milk, especially colostrum, is rich in human milk oligosaccharides (HMOs) (Bode, 2015; Lin et al., 2017) that bind to Toll-like receptors (TLRs) and prevent activation of the receptors by pathogen-associated molecular patterns (PAMPs), which are expressed by microflora (He et al., 2016a; Newburg et al., 2016). When TLRs are activated, they initiate intracellular signaling events that result in regulation of IKK and $\text{NF-}\kappa\text{B}$. In comparison with term babies and adults, preterm children express higher concentrations of TLRs, and if activated, could induce an $\text{IKK}/\text{NF-}\kappa\text{B}$ inflammatory response that results in sepsis or necrotizing enterocolitis (NEC) (He et al., 2016a,b; Niño et al., 2016). Components of breast milk block intestinal TLR activation by PAMPs, which in turn blocks downstream IKK phosphorylation. Since we have demonstrated that targeted deletion of intestinal $\text{IKK}\beta$ in *hUGT1* mice represses newborn intestinal UGT1A1, we can speculate that IKK activity is directly linked to intestinal UGT1A1 expression. This concept is further supported by findings that when intestinal TLRs are activated in neonatal *hUGT1* mice by agents, such as LPS, or IKK is directly activated by Cd^{2+} , intestinal UGT1A1 is induced followed by a reduction in TSB levels. Thus the delayed expression of intestinal UGT1A1 can be traced to breast milk inhibition of TLR activation, which controls downstream activation of IKK. This concept can be highly relevant toward understanding the underlying mechanism associated with BMJ, since NCoR1 activity controls intestinal maturation and is directly linked to induction of intestinal UGT1A1. This also may explain why a diet of formula, which lacks TLR binding glycans, allows downstream activation of the $\text{IKK}/\text{NCoR1}$ loop, induction of intestinal UGT1A1, and a reduction in TSB levels. Equation would naturally dilute the concentrations of the TLR binding glycans allowing activation of the TLRs by PAMPs, followed by derepression of NCoR1 and induction of intestinal UGT1A1.

Discussion

The human *UGT1A1* gene has evolved as a key target for naturally produced humoral agents as well as dietary and environmental compounds through expression in both liver and intestinal tissues to control the levels of accumulating TSB in newborns. While current dogma has produced an image that UGT1A1 plays a principle role in drug metabolism, expression of the human *UGT1A1* gene as a transgene in mice has allowed the development of experiments to document its role as a key regulator of TSB levels. The importance of the *UGT1A1* gene cannot be underestimated since silencing mutations in the gene are lethal, leading to the accumulation of toxic levels of TSB that result in encephalitis followed by neurologic damage that is eventually lethal. While most of mutations in genes associated with drug metabolism could be classified as “disease enhancing mutations” (Evans and Relling, 1999, 2004; Relling and Evans, 2015), meaning that they are not detrimental to the host but under stress conditions or in the presence of saturating substrates can result in toxicity, the mutations in the *UGT1A1* gene can be classified as “disease causing mutations,” indicating that the *UGT1A1* gene is essential for life. Since the hundreds of mutations in the other 19 *UGT* genes can be classified as disease enhancing mutations (i.e., not important for life), UGT1A1 can be thought of as the only relevant UGT needed for survival. Studies in *hUGT1* mice have concluded that regulation of the *UGT1A1* gene can be controlled by virtually all the key XNRs, which include PXR (Xie et al., 2003; Chen et al., 2005, 2012), CAR (Sugatani et al., 2005; Cai et al., 2010), PPAR α (Senekeo-Effenberger et al., 2007), LXR α/β (unpublished observations), as well as the environmental sensors AhR (Yueh et al., 2005; Bonzo et al., 2007) and Nrf2 (Yueh and Tukey, 2007). In adding to this diversity, changes in the redox state of the tissue that result in oxidative stress and activation of sensors, such as ROS-sensitive MAPKs, are directly linked to induction of the *UGT1A1* gene (Fujiwara et al., 2012; Liu et al., 2016; Yoda et al., 2017). There are few other genes that have been linked to drug metabolism that command this range of regulation by humoral and environmental stimuli. An understanding of these events may lead to a greater appreciation of the contribution BM plays in the etiology of BM-induced jaundice.

An important finding that will shed light on the mechanisms leading to neonatal BMJ is the importance of UGT1A1 expression in the intestines and its contribution to the metabolism and clearance of accumulating serum bilirubin. The expression of UGT1A1 is abundant through the intestinal tract (Tukey and Strassburg, 2000, 2001; Tukey et al., 2002), and that conserved expression is observed in *hUGT1* mice (Chen et al., 2005). Given that the intestines become introduced immediately after birth with breast milk, this provides a conduit for the colonization of microflora that produces TLR ligands such as PAMPs. The generation of PAMPs and activation of TLRs can have a negative impact by initiating an $\text{IKK}/\text{NF-}\kappa\text{B}$ -directed inflammatory insult, which can lead to serious health issues such as NEC (Egan et al., 2016; Good et al., 2016; He et al., 2016a; Niño et al., 2016). To combat this insult, human breast milk is rich in complex oligosaccharides that antagonize TLR activation, block PAMP binding, and prevent the acceleration of an inflammatory insult. In addition, breast milk contains antioxidants that protect the tissue from oxidative damage, an important secondary defense mechanism that helps prevent the induction of an inflammatory episode. In neonatal *hUGT1* mice, a diet of human breast milk maintains a state of severe hyperbilirubinemia, like what is seen with nursing neonatal *hUGT1* mice (Fujiwara et al., 2012). In *hUGT1* mice, hepatic UGT1A1 expression is developmentally delayed, a major factor that leads to elevated TSB levels. In addition, intestinal UGT1A1 expression is also repressed. Our findings indicate that developmental repression of hepatic and intestinal UGT1A1 expression is not random




Fig. 6. NCoR1 as a potential target for both neonatal hyperbilirubinemia and necrotizing enterocolitis (NEC). Intestinal UGT1A1 is essential in bilirubin detoxification and is directly linked to neonatal hyperbilirubinemia. We demonstrated that NCoR1 represses intestinal *UGT1A1* gene expression in a developmental-dependent manner that is limited within the breastfeeding stage, suggesting that NCoR1 may function as an important underlying mechanism for BMJ. It is known that human milk oligosaccharides (HMOs) inhibit Toll-like receptors (TLRs), a family of membrane receptors upstream of NCoR1. When activated, TLRs activate IKKs and c-Jun signaling cascades, leading to turnover of NCoR1 from the target promoters and subsequent gene transcription. It hints at the possibility that inhibition of TLR function by HMOs leads to the heightened NCoR1 repression on the *UGT1A1* gene, resulting in an increase in TSB levels. In addition, we now know that induction of intestinal UGT1A1 during the neonatal period by any means (i.e., oxidative stress, ligand-activated nuclear receptors, formula) will lower TSB values, adding support for a mechanism that implicates BM in facilitating reduced expression of UGT1A1. The establishment of the microflora starts immediately following birth. Pathogen-associated molecular patterns (PAMPs) can activate TLRs and induce an inflammatory response and gene transcription through NCoR1 derepression. This response may lead to important health concerns including necrotizing enterocolitis (NEC), a devastating disease that affects mostly premature infants. Countering this, premature infants on breast milk have demonstrated less NEC in comparison with formula feeding.

but is a controlled event, with repression of the intestinal *UGT1A1* gene being influenced by breast milk. In addition, we now know that induction of intestinal UGT1A1 during the neonatal period by any means (i.e., oxidative stress, ligand activated nuclear receptors, formula) will lower TSB values, adding support for a mechanism that implicates breast milk in facilitating reduced expression of UGT1A1 (Fig. 6).

Combined, our findings along with other key observations can help develop a working mechanism implicating BM with neonatal jaundice. Proof of this model relies on two key findings: First, HMOs, which are abundant in breast milk, antagonize intestinal TLRs, preventing downstream intestinal inflammation. This is critical because it is now known that activation of IKK during the neonatal window leads to derepression of the *UGT1A1* gene and reductions in TSB levels. This mechanism can explain the actions of formula toward the induction of UGT1A1 and reduction of TSB levels. Formula does not contain HMOs; thus, a diet of formula would lead to a dramatic reduction in intestinal HMO concentrations, allowing TLR activation by microflora-generated PAMPs that would ultimately lead to IKK activation and derepression of the *UGT1A1* gene. Such an action would lead to reduction in TSB levels. This mechanism may also explain the observations that formula leads to NEC, which is observed to occur predominantly in preterm infants who

have been enterally fed (Niño et al., 2016). It is known that preterm infants are born with a far greater concentration of TLRs in the intestines (Nanthakumar et al., 2011; Glaser and Speer, 2013), a fact that would increase the sensitivity of developing NEC if the TLRs are activated. Second, the antioxidant potential of breast milk protects the intestines from oxidative stress-induced episodes. Events that may lead to the production of ROS would be detected by intestinal sensors, resulting in activation of MAPK pathways that culminate in derepression of the *UGT1A1* gene. In both events, breast milk plays an important role in maintaining repression of the intestinal *UGT1A1* gene (Fig. 6).

Development of transgenic and *hUGT1* mice coupled with genetic modification of transcriptional and other regulatory proteins in these mice has provided new insight into the mechanisms that control the tissue-specific and inducible expression of the genes linked to the *UGT1* locus. Since the human *UGT1A1* gene is regulated in *hUGT1* mice in a developmental- and tissue-specific fashion like its expression in humans, we have taken advantage of its developmental delay to uncover unique insight into the mechanisms leading to neonatal hyperbilirubinemia. With an appreciation that the intestines can be an easy target for the delivery agents that would induce the *UGT1A1* gene, future efforts using these mice could be exploited to develop effective therapeutics to treat newborns that are predisposed toward the development of SNH.

Authorship Contributions

Participated in research design: Chen, Tukey.

Performed data analysis: Chen, Tukey.

Wrote or contributed to the writing of the manuscript: Chen, Tukey.

References

- Alam J, Stewart D, Touchard C, Boinapally S, Choi AM, and Cook JL (1999) Nrf2, a Cap'n'Collar transcription factor, regulates induction of the heme oxygenase-1 gene. *J Biol Chem* **274**: 26071–26078.
- Ayer A, Zarjou A, Agarwal A, and Stocker R (2016) Heme oxygenases in cardiovascular health and disease. *Physiol Rev* **96**:1449–1508.
- Bacq Y, Zarka O, Bréchet JF, Mariotte N, Vol S, Tichet J, and Weill J (1996) Liver function tests in normal pregnancy: a prospective study of 103 pregnant women and 103 matched controls. *Hepatology* **23**:1030–1034.
- Bhutani VK and Johnson-Hamerman L (2015) The clinical syndrome of bilirubin-induced neurologic dysfunction. *Semin Fetal Neonatal Med* **20**:6–13.
- Bhutani VK, Zupursky A, Blencowe H, Khanna R, Sgro M, Ebbesen F, Bell J, Mori R, Slusher TM, Fahmy N, et al. (2013) Neonatal hyperbilirubinemia and Rhesus disease of the newborn: incidence and impairment estimates for 2010 at regional and global levels. *Pediatr Res* **74** (Suppl 1):86–100.
- Blumberg B, Sabbagh W, Jr, Juguilon H, Bolado J, Jr, van Meter CM, Ong ES, and Evans RM (1998) SXR, a novel steroid and xenobiotic-sensing nuclear receptor. *Genes Dev* **12**:3195–3205.
- Bode L (2015) The functional biology of human milk oligosaccharides. *Early Hum Dev* **91**: 619–622.
- Bonzo JA, Bélanger A, and Tukey RH (2007) The role of chrysin and the arh receptor in induction of the human UGT1A1 gene in vitro and in transgenic UGT1 mice. *Hepatology* **45**:349–360.
- Bosma PJ, Seppen J, Goldhoorn B, Bakker C, Oude Elferink RP, Chowdhury JR, Chowdhury NR, and Jansen PL (1994) Bilirubin UDP-glucuronosyltransferase 1 is the only relevant bilirubin glucuronidating isoform in man. *J Biol Chem* **269**:17960–17964.
- Burchell B, Coughtrie M, Jackson M, Harding D, Fournel-Gigleux S, Leakey J, and Hume R (1989) Development of human liver UDP-glucuronosyltransferases. *Dev Pharmacol Ther* **13**: 70–77.
- Cai H, Nguyen N, Peterkin V, Yang YS, Hotz K, La Placa DB, Chen S, Tukey RH, and Stevens JC (2010) A humanized *UGT1* mouse model expressing the *UGT1A1**28 allele for assessing drug clearance by UGT1A1-dependent glucuronidation. *Drug Metab Dispos* **38**:879–886.
- Chen S, Beaton D, Nguyen N, Seneke-Effenberger K, Brace-Sinnokrak E, Argikar U, Rummel RP, Trotter J, Barbier O, Ritter JK, et al. (2005) Tissue-specific, inducible, and hormonal control of the human UDP-glucuronosyltransferase-1 (*UGT1*) locus. *J Biol Chem* **280**:37547–37557.
- Chen S, Lu W, Yueh MF, Rettenmeier E, Liu M, Paszek M, Auwerx J, Yu RT, Evans RM, Wang K, et al. (2017) Intestinal NCoR1, a regulator of epithelial cell maturation, controls neonatal hyperbilirubinemia [published correction appears in *Proc Natl Acad Sci USA* (2017) 114:E4115]. *Proc Natl Acad Sci USA* **114**:E1432–E1440.
- Chen S, Yueh MF, Evans RM, and Tukey RH (2012) Pregnane-x-receptor controls hepatic glucuronidation during pregnancy and neonatal development in humanized UGT1 mice. *Hepatology* **56**:658–667.
- Choi HK, Yoo JY, Jeong MH, Park SY, Shin DM, Jang SW, Yoon HG, and Choi KC (2013) Protein kinase A phosphorylates NCoR to enhance its nuclear translocation and repressive function in human prostate cancer cells. *J Cell Physiol* **228**:1159–1165.
- Clarke DJ, Cassidy AJ, See CG, Povey S, and Burchell B (1997) Cloning of the human UGT1 gene complex in yeast artificial chromosomes: novel aspects of gene structure and subchromosomal mapping to 2q37. *Biochem Soc Trans* **25**:S562.
- Cunningham AD, Hwang S, and Mochly-Rosen D (2016) Glucose-6-phosphate dehydrogenase deficiency and the need for a novel treatment to prevent kernicterus. *Clin Perinatol* **43**: 341–354.

- Dennery PA, Seidman DS, and Stevenson DK (2001) Neonatal hyperbilirubinemia. *N Engl J Med* **344**:581–590.
- Egan CE, Sodhi CP, Good M, Lin J, Jia H, Yamaguchi Y, Lu P, Ma C, Branca MF, Weyand S, et al. (2016) Toll-like receptor 4-mediated lymphocyte influx induces neonatal necrotizing enterocolitis. *J Clin Invest* **126**:495–508.
- Ercal N, Gurer-Orhan H, and Aykin-Burns N (2001) Toxic metals and oxidative stress part I: mechanisms involved in metal-induced oxidative damage. *Curr Top Med Chem* **1**:529–539.
- Evans WE and Relling MV (1999) Pharmacogenomics: translating functional genomics into rational therapeutics. *Science* **286**:487–491.
- Evans WE and Relling MV (2004) Moving towards individualized medicine with pharmacogenomics. *Nature* **429**:464–468.
- Fernández-Majada V, Aguilera C, Villanueva A, Vilardeff F, Robert-Moreno A, Aytés A, Real FX, Capella G, Mayo MW, Espinosa L, et al. (2007) Nuclear IKK activity leads to dysregulated notch-dependent gene expression in colorectal cancer. *Proc Natl Acad Sci USA* **104**:276–281.
- Francis GA, Fayard E, Picard F, and Auwerx J (2003) Nuclear receptors and the control of metabolism. *Annu Rev Physiol* **65**:261–311.
- Fujiwara R, Chen S, Karin M, and Tukey RH (2012) Reduced expression of UGT1A1 in intestines of humanized UGT1 mice via inactivation of NF- κ B leads to hyperbilirubinemia. *Gastroenterology* **142**:109–118.e2.
- Fujiwara R, Maruo Y, Chen S, and Tukey RH (2015) Role of extrahepatic UDP-glucuronosyltransferase 1A1: advances in understanding breast milk-induced neonatal hyperbilirubinemia. *Toxicol Appl Pharmacol* **289**:124–132.
- Fujiwara R, Nguyen N, Chen S, and Tukey RH (2010) Developmental hyperbilirubinemia and CNS toxicity in mice humanized with the UDP glucuronosyltransferase 1 (UGT1) locus. *Proc Natl Acad Sci USA* **107**:5024–5029.
- Gallardo F, Padrón A, García-Carbonell R, Rius C, González-Perez A, Arumi-Uria M, Iglesias M, Nonell L, Bellosillo B, Segura S, et al. (2015) Cytoplasmic accumulation of NCoR in malignant melanoma: consequences of altered gene repression and prognostic significance. *Oncotarget* **6**:9284–9294.
- Glaser K and Speer CP (2013) Toll-like receptor signaling in neonatal sepsis and inflammation: a matter of orchestration and conditioning. *Expert Rev Clin Immunol* **9**:1239–1252.
- Gong QH, Cho JW, Huang T, Potter C, Gholami N, Basu NK, Kubota S, Carvalho S, Pennington MW, Owens IS, et al. (2001) Thirteen UDPglucuronosyltransferase genes are encoded at the human UGT1 gene complex locus. *Pharmacogenetics* **11**:357–368.
- Good M, Sodhi CP, Yamaguchi Y, Jia H, Lu P, Fulton WB, Martin LY, Prindle T, Nino DF, Zhou Q, et al. (2016) The human milk oligosaccharide 2'-fucosyllactose attenuates the severity of experimental necrotizing enterocolitis by enhancing mesenteric perfusion in the neonatal intestine. *Br J Nutr* **116**:1175–1187.
- He Y, Lawlor NT, and Newburg DS (2016a) Human milk components modulate toll-like receptor-mediated inflammation. *Adv Nutr* **7**:102–111.
- He Y, Liu S, Kling DE, Leone S, Lawlor NT, Huang Y, Feinberg SB, Hill DR, and Newburg DS (2016b) The human milk oligosaccharide 2'-fucosyllactose modulates CD14 expression in human enterocytes, thereby attenuating LPS-induced inflammation. *Gut* **65**:33–46.
- Huang MJ, Kua KE, Teng HC, Tang KS, Weng HW, and Huang CS (2004) Risk factors for severe hyperbilirubinemia in neonates. *Pediatr Res* **56**:682–689.
- Huang W, Ghisletti S, Perissi V, Rosenfeld MG, and Glass CK (2009) Transcriptional integration of TLR2 and TLR4 signaling at the NCoR derepression checkpoint. *Mol Cell* **35**:48–57.
- Jeong H, Choi S, Song JW, Chen H, and Fischer JH (2008) Regulation of UDP-glucuronosyltransferase (UGT) 1A1 by progesterone and its impact on labetalol elimination. *Xenobiotica* **38**:62–75.
- Jo YS, Ryu D, Maida A, Wang X, Evans RM, Schoonjans K, and Auwerx J (2015) Phosphorylation of the nuclear receptor corepressor 1 by protein kinase B switches its corepressor targets in the liver in mice. *Hepatology* **62**:1606–1618.
- Johnson LH, Brown AK, and Bhutani VK (1999) BIND—a clinical score for bilirubin induced neurologic dysfunction in newborns. *Pediatrics* **199**:746–747.
- Kaplan M, Bromiker R, and Hammerman C (2011) Severe neonatal hyperbilirubinemia and kernicterus: are these still problems in the third millennium? *Neonatology* **100**:354–362.
- Karin M (2009) NF- κ B as a critical link between inflammation and cancer. *Cold Spring Harb Perspect Biol* **1**:a000141.
- Kullak-Ublick GA, Beuers U, and Paumgartner G (2000) Hepatobiliary transport. *J Hepatol* **32** (1 Suppl):3–18.
- Le Pichon JB, Riordan SM, Watchko J, and Shapiro SM (2017) The neurological sequelae of neonatal hyperbilirubinemia: definitions, diagnosis and treatment of the kernicterus Spectrum disorders (KSDs). *Curr Pediatr Rev* **13**:199–209.
- Li M, Cai JF, and Chiu JF (2002) Arsenic induces oxidative stress and activates stress gene expressions in cultured lung epithelial cells. *J Cell Biochem* **87**:29–38.
- Li P, Spann NJ, Kaikkonen MU, Lu M, Oh DY, Fox JN, Bandyopadhyay G, Talukdar S, Xu J, Lagakos WS, et al. (2013) NCoR repression of LXR α restricts macrophage biosynthesis of insulin-sensitizing omega 3 fatty acids. *Cell* **155**:200–214.
- Lin AE, Autran CA, Szyzka A, Escajadillo T, Huang M, Godula K, Prudden AR, Boons GJ, Lewis AL, Doran KS, et al. (2017) Human milk oligosaccharides inhibit growth of group B *Streptococcus*. *J Biol Chem* **292**:11243–11249.
- Liu L, Trimarchi JR, Navarro P, Blasco MA, and Keefe DL (2003) Oxidative stress contributes to arsenic-induced telomere attrition, chromosome instability, and apoptosis. *J Biol Chem* **278**:31998–32004.
- Liu M, Chen S, Yueh MF, Fujiwara R, Konopnicki C, Hao H, and Tukey RH (2016) Cadmium and arsenic override NF- κ B developmental regulation of the intestinal UGT1A1 gene and control of hyperbilirubinemia. *Biochem Pharmacol* **110**:111:37–46.
- López E, Arce C, Oset-Gasque MJ, Cañadas S, and González MP (2006) Cadmium induces reactive oxygen species generation and lipid peroxidation in cortical neurons in culture. *Free Radic Biol Med* **40**:940–951.
- Mackenzie PI, Owens IS, Burchell B, Bock KW, Bairoch A, Bélanger A, Fournel-Gigleux S, Green M, Hum DW, Iyanagi T, et al. (1997) The UDP glycosyltransferase gene superfamily: recommended nomenclature update based on evolutionary divergence. *Pharmacogenetics* **7**:255–269.
- Maisels MJ (2015) Managing the jaundiced newborn: a persistent challenge. *CMAJ* **187**:335–343.
- Maisels MJ, Clune S, Coleman K, Gendelman B, Kendall A, McManus S, and Smyth M (2014) The natural history of jaundice in predominantly breastfed infants. *Pediatrics* **134**:e340–e345.
- McWalter GK, Higgins LG, McLellan LI, Henderson CJ, Song L, Thornalley PJ, Itoh K, Yamamoto M, and Hayes JD (2004) Transcription factor Nrf2 is essential for induction of NAD(P)H:quinone oxidoreductase 1, glutathione S-transferases, and glutamate cysteine ligase by broccoli seeds and isothiocyanates. *J Nutr* **134** (12 Suppl):3499S–3506S.
- Mottis A, Mouchiroud L, and Auwerx J (2013) Emerging roles of the corepressors NCoR1 and SMRT in homeostasis. *Genes Dev* **27**:819–835.
- Murillo G and Mehta RG (2001) Cruciferous vegetables and cancer prevention. *Nutr Cancer* **41**:17–28.
- Nanthakumar N, Meng D, Goldstein AM, Zhu W, Lu L, Uauy R, Llanos A, Claud EC, and Walker WA (2011) The mechanism of excessive intestinal inflammation in necrotizing enterocolitis: an immature innate immune response. *PLoS One* **6**:e17776.
- Newburg DS, Ko JS, Leone S, and Nanthakumar NN (2016) Human milk oligosaccharides and synthetic galactosyloligosaccharides contain 3', 4-, and 6'-galactosyllactose and attenuate inflammation in human T84, NCM-460, and H4 cells and intestinal tissue ex vivo. *J Nutr* **146**:358–367.
- Newman AJ and Gross S (1963) Hyperbilirubinemia in breast-fed infants. *Pediatrics* **32**:995–1001.
- Nguyen N, Bonzo JA, Chen S, Chouinard S, Kelner MJ, Hardiman G, Bélanger A, and Tukey RH (2008) Disruption of the *ugt1* locus in mice resembles human Crigler-Najjar type I disease. *J Biol Chem* **283**:7901–7911.
- Nguyen T, Huang HC, and Pickett CB (2000) Transcriptional regulation of the antioxidant response element. Activation by Nrf2 and repression by MafK. *J Biol Chem* **275**:15466–15473.
- Nguyen T, Sherratt PJ, and Pickett CB (2003) Regulatory mechanisms controlling gene expression mediated by the antioxidant response element. *Annu Rev Pharmacol Toxicol* **43**:233–260.
- Niño DF, Sodhi CP, and Hackam DJ (2016) Necrotizing enterocolitis: new insights into pathogenesis and mechanisms. *Nat Rev Gastroenterol Hepatol* **13**:590–600.
- Olusanya BO, Osibanjo FB, Mabogunje CA, Slusher TM, and Olowe SA (2016) The burden and management of neonatal jaundice in Nigeria: a scoping review of the literature. *Niger J Clin Pract* **19**:1–17.
- Olusanya BO, Osibanjo FB, and Slusher TM (2015) Risk factors for severe neonatal hyperbilirubinemia in low and middle-income countries: a systematic review and meta-analysis. *PLoS One* **10**:e0117229.
- Olusanya BO, Teeple S, and Kassebaum NJ (2018) The contribution of neonatal jaundice to global child mortality: findings from the GBD 2016 Study. *Pediatrics* **141**:e20171471.
- Relling MV and Evans WE (2015) Pharmacogenomics in the clinic. *Nature* **526**:343–350.
- Rizzardini M, Carelli M, Cabello Porras MR, and Cantoni L (1994) Mechanisms of endotoxin-induced haem oxygenase mRNA accumulation in mouse liver: synergism by glutathione depletion and protection by N-acetylcysteine. *Biochem J* **304**:477–483.
- Santra A, Maiti A, Chowdhury A, and Mazumder DN (2000) Oxidative stress in liver of mice exposed to arsenic-contaminated water. *Indian J Gastroenterol* **19**:112–115.
- Schneider AP, II (1986) Breast milk jaundice in the newborn. A real entity. *JAMA* **255**:3270–3274.
- Seneko-Effenberger K, Chen S, Brace-Sinnokrak E, Bonzo JA, Yueh MF, Argikar U, Kaeding J, Trotter J, Remmel RP, Ritter JK, et al. (2007) Expression of the human UGT1 locus in transgenic mice by 4-chloro-6-(2,3-xylylidino)-2-pyrimidinylthioacetic acid (WY-14643) and implications on drug metabolism through peroxisome proliferator-activated receptor alpha activation. *Drug Metab Dispos* **35**:419–427.
- Shapiro GI, Supko JG, Patterson A, Lynch C, Lucca J, Zaccarola PF, Muzikansky A, Wright JJ, Lynch TJ, Jr, and Rollins BJ (2001a) A phase II trial of the cyclin-dependent kinase inhibitor flavopiridol in patients with previously untreated stage IV non-small cell lung cancer. *Clin Cancer Res* **7**:1590–1599.
- Shapiro TA, Fahey JW, Dinkova-Kostova AT, Holtzclaw WD, Stephenson KK, Wade KL, Ye L, and Talalay P (2006) Safety, tolerance, and metabolism of broccoli sprout glucosinolates and isothiocyanates: a clinical phase I study. *Nutr Cancer* **55**:53–62.
- Shapiro TA, Fahey JW, Wade KL, Stephenson KK, and Talalay P (2001b) Chemoprotective glucosinolates and isothiocyanates of broccoli sprouts: metabolism and excretion in humans. *Cancer Epidemiol Biomarkers Prev* **10**:501–508.
- Shibuya A, Itoh T, Tukey RH, and Fujiwara R (2013) Impact of fatty acids on human UDP-glucuronosyltransferase 1A1 activity and its expression in neonatal hyperbilirubinemia. *Sci Rep* **3**:2903.
- Strassburg CP, Manns MP, and Tukey RH (1997a) Differential down-regulation of the *UDP-glucuronosyltransferase 1A* locus is an early event in human liver and biliary cancer. *Cancer Res* **57**:2979–2985.
- Strassburg CP, Manns MP, and Tukey RH (1998a) Expression of the *UDP-glucuronosyltransferase 1A* locus in human colon. Identification and characterization of the novel extrahepatic UGT1A8. *J Biol Chem* **273**:8719–8726.
- Strassburg CP, Nguyen N, Manns MP, and Tukey RH (1998b) Polymorphic expression of the *UDP-glucuronosyltransferase UGT1A* gene locus in human gastric epithelium. *Mol Pharmacol* **54**:647–654.
- Strassburg CP, Oldhafer K, Manns MP, and Tukey RH (1997b) Differential expression of the *UGT1A* locus in human liver, biliary, and gastric tissue: identification of UGT1A7 and UGT1A10 transcripts in extrahepatic tissue. *Mol Pharmacol* **52**:212–220.
- Sugatani J, Kojima H, Ueda A, Kakizaki S, Yoshinari K, Gong QH, Owens IS, Negishi M, and Sueyoshi T (2001) The phenobarbital response enhancer module in the human bilirubin UDP-glucuronosyltransferase UGT1A1 gene and regulation by the nuclear receptor CAR. *Hepatology* **33**:1232–1238.
- Sugatani J, Sueyoshi T, Negishi M, and Miwa M (2005) Regulation of the human UGT1A1 gene by nuclear receptors constitutive active/androstane receptor, pregnane X receptor, and glucocorticoid receptor. *Methods Enzymol* **400**:92–104.
- Tabarki B, Khalifa M, Yacoub M, Tlili K, and Essoussi AS (2002) Cerebellar symptoms heralding bilirubin encephalopathy in Crigler-Najjar syndrome. *Pediatr Neurol* **27**:234–236.
- Tenhunen R (1976) The enzymatic conversion of heme to bilirubin in vivo. *Ann Clin Res* **8** (Suppl 17):2–9.
- Tukey RH and Strassburg CP (2000) Human UDP-glucuronosyltransferases: metabolism, expression, and disease. *Annu Rev Pharmacol Toxicol* **40**:581–616.
- Tukey RH and Strassburg CP (2001) Genetic multiplicity of the human UDP-glucuronosyltransferases and regulation in the gastrointestinal tract. *Mol Pharmacol* **59**:405–414.
- Tukey RH, Strassburg CP, and Mackenzie PI (2002) Pharmacogenomics of human UDP-glucuronosyltransferases and irinotecan toxicity. *Mol Pharmacol* **62**:446–450.
- Wasserman WW and Fahl WE (1997) Functional antioxidant responsive elements. *Proc Natl Acad Sci USA* **94**:5361–5366.
- Watchko JF (2009) Identification of neonates at risk for hazardous hyperbilirubinemia: emerging clinical insights. *Pediatr Clin North Am* **56**:671–687.

- Watchko JF (2016) Bilirubin-induced neurotoxicity in the preterm neonate. *Clin Perinatol* **43**: 297–311.
- Wong RJ and Stevenson DK (2015) Neonatal hemolysis and risk of bilirubin-induced neurologic dysfunction. *Semin Fetal Neonatal Med* **20**:26–30.
- Xie W, Yeuh MF, Radominska-Pandya A, Saini SPS, Negishi Y, Bottroff BS, Cabrera GY, Tukey RH, and Evans RM (2003) Control of steroid, heme, and carcinogen metabolism by nuclear pregnane X receptor and constitutive androstane receptor. *Proc Natl Acad Sci USA* **100**: 4150–4155.
- Yoda E, Paszek M, Konopnicki C, Fujiwara R, Chen S, and Tukey RH (2017) Isothiocyanates induce UGT1A1 in humanized UGT1 mice in a CAR dependent fashion that is highly dependent upon oxidative stress. *Sci Rep* **7**:46489.
- Yeuh MF, Bonzo JA, and Tukey RH (2005) The role of Ah receptor in induction of human UDP-glucuronosyltransferase 1A1. *Methods Enzymol* **400**:75–91.
- Yeuh MF, Chen S, Nguyen N, and Tukey RH (2017) Developmental, genetic, dietary, and xenobiotic influences on neonatal hyperbilirubinemia. *Mol Pharmacol* **91**:545–553.
- Yeuh MF, Huang YH, Hiller A, Chen S, Nguyen N, and Tukey RH (2003) Involvement of the xenobiotic response element (XRE) in Ah receptor-mediated induction of human UDP-glucuronosyltransferase 1A1. *J Biol Chem* **278**:15001–15006.
- Yeuh MF and Tukey RH (2007) Nrf2-Keap1 signaling pathway regulates human *UGT1A1* expression in vitro and in transgenic UGT1 mice. *J Biol Chem* **282**:8749–8758.

Address correspondence to: Dr. Robert H. Tukey, UC San Diego, Department of Pharmacology, Leichtag Biomedical Research Building, 9500 Gilman Drive, La Jolla, CA 92093-0722. E-mail: rtukey@ucsd.edu; or Dr. Shujuan Chen, UC San Diego, Department of Pharmacology, Leichtag Biomedical Research Building, 9500 Gilman Drive, La Jolla, CA 92093-0722. E-mail: s18chen@ucsd.edu
