

# UC Davis

## UC Davis Previously Published Works

### Title

In-vitro and in-vivo imaging of MMP activity in cartilage and joint injury

### Permalink

<https://escholarship.org/uc/item/69n5h4sr>

### Journal

Biochemical and Biophysical Research Communications, 460(3)

### ISSN

0006-291X

### Authors

Fukui, Tomoaki  
Tenborg, Elizabeth  
Yik, Jasper HN  
[et al.](#)

### Publication Date

2015-05-01

### DOI

10.1016/j.bbrc.2015.03.100

Peer reviewed



# HHS Public Access

Author manuscript

*Biochem Biophys Res Commun.* Author manuscript; available in PMC 2016 May 08.

Published in final edited form as:

*Biochem Biophys Res Commun.* 2015 May 8; 460(3): 741–746. doi:10.1016/j.bbrc.2015.03.100.

## ***In-vitro* and *in-vivo* imaging of MMP activity in cartilage and joint injury**

**Tomoaki Fukui, Elizabeth Tenborg, Jasper H. N. Yik, and Dominik R. Haudenschild\***

Lawrence J. Ellison Musculoskeletal Research Center, Department of Orthopaedic Surgery, University of California Davis Medical Center, 4635 Second Avenue Suite 2000, Sacramento CA 95817, USA

### **Abstract**

Non-destructive detection of cartilage-degrading activities represents an advance in osteoarthritis (OA) research, with implications in studies of OA pathogenesis, progression, and intervention strategies. Matrix metalloproteinases (MMPs) are principal cartilage degrading enzymes that contribute to OA pathogenesis. MMPsense750 is an *in-vivo* fluorimetric imaging probe with the potential to continuously and non-invasively trace real-time MMP activities, but its use in OA-related research has not been reported. Our objective is to detect and characterize the early degradation activities shortly after cartilage or joint injury with MMPsense750. We determined the appropriate concentration, assay time, and linear range using various concentrations of recombinant MMPs as standards. We then quantified MMP activity from cartilage explants subjected to either mechanical injury or inflammatory cytokine treatment *in-vitro*. Finally, we performed *in vivo* MMP imaging of a mouse model of post-traumatic OA. Our *in-vitro* results showed that the optimal assay time was highly dependent on the MMP enzyme. In cartilage explant culture media, mechanical impact or cytokine treatment increased MMP activity. Injured knees of mice showed significantly higher fluorescent signal than uninjured knees. We conclude that MMPsense750 detects human MMP activities and can be used for *in-vitro* study with cartilage, as well as *in-vivo* studies of knee injury, and can offering real-time insight into the degradative processes that occurring within the joint before structural changes become evident radiographically.

### **Keywords**

cartilage; *in-vivo* imaging; MMP activity; MMPsense750; osteoarthritis

---

© 2015 Published by Elsevier Inc.

\*Corresponding author: Dominik R. Haudenschild Lawrence J. Ellison Musculoskeletal Research Center, Department of Orthopaedic Surgery, University of California Davis Medical Center, 4635 Second Avenue Suite 2000, Sacramento CA 95817, USA Tel: +1-916-734-5015 Fax: + 1-916-734-5750. DRHaudenschild@ucdavis.edu.

**Publisher's Disclaimer:** This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final citable form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

The authors declare that they have no competing interests.

## Introduction

Osteoarthritis (OA) is a degenerative disease of the whole joint organ characterized by cartilage degradation,[1] and the number of OA patients continues to increase, estimated at nearly 27 million in the United States.[2] and there is no effective treatment to prevent OA or restore joints after the onset of OA. At this time, the gold standard for clinical OA diagnosis and evaluation are morphologic assessments, such as radiography[3,4,5], computed tomography (CT)[5,6], and magnetic resonance imaging (MRI)[3,5,6]. These imaging technologies primarily reveal the morphological changes that become evident at the later stages of OA, but do not offer insight into the process of cartilage degradation. As the field of OA research moves toward OA prevention it is becoming important to also measure the biological processes responsible for joint degradation, processes that precede the morphological and structural changes.

It is generally accepted that enzymatic activities contribute to cartilage degradation and loss in OA, and that elevated enzymatic activity precedes morphological joint space narrowing. [7,8] The ability to non-destructively image and quantify enzymatic activity would be an important tool to assess OA initiation and progression, and the efficacy of intervention strategies. While primary OA is considered idiopathic, in post-traumatic OA (PTOA) the time point of OA initiation (trauma) can be easily identified, and this is therefore an appropriate model to study the enzymatic activities during the early phases of OA.

Matrix metalloproteinases (MMPs) are a family of zinc-dependent degradative proteinases with roles in the enzymatic cartilage degradation and OA progression. MMP-mediated degradation of type II collagen fibrils is considered one of the first irreversible steps in OA pathogenesis (reviewed in [9]), and the presence of MMPs correlates with OA symptoms, including joint effusion and pain[10,11]. Although serum level of MMP-3 is used as a biomarker for rheumatoid arthritis (RA), there is no clinically established MMP biomarker for OA[12]. Most studies investigating MMP activity in OA rely on assays such as ELISA[13,14] or Western Blotting[14,15], and RT-PCR[14,16] to estimate protein levels and mRNA expression, respectively. However, these assays are not suitable for *in-vivo* use, and can only measure the amount of MMP protein but do not directly assay MMP activity. Direct measurements of MMP activity include zymography[14,16,17] and more recently fluorimetric MMP assays[14,16], but again these assays are generally not suited for *in-vivo* imaging. The development of a method to visualize MMP activities *in-vivo* hence may offer new insight into OA initiation and treatment efficacy.

An *in-vivo* fluorimetric probe was recently developed that allows non-destructive imaging of activity from a broad spectrum of MMPs (MMPSense™750 FAST PerkinElmer, Inc., Boston, MA). This near-infrared fluorescent probe is a peptide substrate that enables the detection of MMP activities by exhibiting fluorescent signal when cleaved by MMPs[13,15,18]. *In-vivo* imaging with this reagent has the potential to continuously measure real-time MMP activities noninvasively. This probe has been successfully used to detect tumor progression[13,18] or ischemia reperfusion in brain[15] with *in-vivo* mouse model, but to our best knowledge there is no previous report investigating MMPSense750 for the assessment of MMP activity in cartilage or joint injury. Moreover, although this

probe is expected to be utilized for human patients in clinical setting in future, no paper has studied the kinetics of this substrate using human MMPs. The objective of the present study is to investigate the potential of MMPsense750 for detection of human MMPs and to use in an assay with cartilage explants, as well as an *in-vivo* mouse model of knee injury leading to PTOA.

## Methods

### Assessment of optimal MMPsense750 concentration

We first wanted to determine the appropriate MMPsense750 concentration for *in-vitro* studies using purified recombinant human MMPs. Human MMP-3, -9, and -13 were chosen based on their established importance in OA[11,19,20]. To achieve comparable results between the different enzymes, the amount of active enzyme in each assay was normalized using the specific activity (Units of enzyme activity per weight) provided by the manufacturer (Supplementary information 1). MMPsense750 was added to media containing the active proteases, the reactions were incubated at 37°C, and the resulting fluorescent signal was measured at different time points as described in detail below.

Recombinant human MMP-3 (Enzo Life Sciences, Farmingdale, USA) was reconstituted to various concentrations in assay buffer consisting of 50mM sodium acetate, 10mM CaCl<sub>2</sub>, 150mM NaCl and 0.05% Brij-35 at pH 6.0.[9] Recombinant human MMP-9 and MMP-13 (Enzo Life Sciences) were reconstituted in the assay buffer consisting of 50mM HEPES, 10mM CaCl<sub>2</sub> and 0.05% Brij-35 at pH 7.5.[21]

MMPsense750 (24 nmol per vial) was reconstituted in 1200µl sterile phosphate-buffered saline (Invitrogen) as recommended by the manufacturer, then added into the MMP solutions at 0.2, 0.7 and 2.0µM final concentration. Imaging was performed on an IVIS-Spectrum imaging system at multiple time points for up to 72 hours after adding MMPsense750.

### Cartilage explants

Cartilage explants were harvested from the weight-bearing area of the femoral articular surfaces of bovine stifle knee joints purchased from a local slaughterhouse (Petaluma, CA). A 6mm dermal biopsy punch was used to isolate cartilage cylinders, which were then cut to 2mm height from the articular surface using a custom jig. Explants were cultured for 3 days in DMEM with 10% FBS and 1% penicillin-streptomycin (all from Invitrogen, Carlsbad, CA) at 37°C and 5% CO<sub>2</sub>. Six joints were used, and 1 or 2 explants from each joint was randomly assigned to one of three treatment groups; IL-1β, mechanical injury, or control. There was no significant difference among the cartilage weights of each group. The IL-1β group was treated with 10ng/ml IL-1β (R&D Systems, Minneapolis, MN). The explants in the mechanical injury group were mechanically compressed with an Instron 8511.20 digital servo-hydraulic mechanical testing device using displacement control. A compressive preload of ~0.5N was applied, and then the explant was loaded to 30% strain at a strain rate of 100%/s, held at 30% strain for 100ms, then unloaded. Following compression, all loaded explants were transferred to fresh culture medium and returned to an incubator at 37°C and

5% CO<sub>2</sub> until the termination of the experiment. In the control group, the explants were given a preload of ~0.5N and then returned to the culture media. The culture media were replenished at day 3. The media were collected at 3 and 6 days after IL-1 $\beta$  stimulation or mechanical injuries. MMPsense750 was added to a final concentration of 0.7 $\mu$ M, and the fluorescence measured at 60 minutes and 24 hours after adding MMPsense.

### Animal model of joint injury

Eight adult male BALB/cByJ mice (9-week-old at time of injury) were obtained from Jackson Laboratory (Bar Harbor, Maine). All animals were maintained and used in accordance with National Institutes of Health guidelines on the care and use of laboratory animals. This study was approved by our Institutional Animal Care and Use Committee (IACUC). The right knees of the mice were injured with a single mechanical compression as previously described in our PTOA model [22]. Briefly, the tibial compression system consists of two custom-built loading platens; the bottom platen that holds the knee flexed, and the top platen that holds the heel. The platens were aligned vertically and positioned within an electromagnetic materials testing machine (Bose ElectroForce 3200) (Eden Prairie, MN). Mice were anesthetized using isoflurane inhalation, then the right leg of each mouse was subjected to a single dynamic axial compression (1mm/s loading rate) to a target load of 12N. This causes a transient anterior subluxation of the tibia, which injures the anterior cruciate ligament and leads to PTOA within 8 weeks. The contra-lateral uninjured knees served controls, and were used to normalize the data within each animal.

All mice received an injection of 2nmol of MMPsense750 via the orbital sinus at 24-hours post-injury, and IVIS imaging was performed 24 hours after the injection (48-hours post-injury). Mice were euthanized immediately after the imaging and both knees were dissected for isolation of total RNA and analysis of mRNA expression.

### Quantitative real-time RT-PCR

Total RNA was extracted from injured and uninjured knees using the miRNeasy Mini Kit (Qiagen Valencia, CA) and reverse transcribed by the QuantiTect Reverse Transcription Kit (Qiagen). 2 $\mu$ l of cDNA was used for quantitative RT-PCR (in a final volume of 10 $\mu$ l) performed in triplicate in a 7900HT RT-PCR system with gene-specific probes according to the manufacturer's conditions. Results were normalized to the 18s rRNA and calculated as fold-change in mRNA expression relative to the untreated control, using the 2<sup>-CT</sup> method. The probes used are shown in Supplementary information 2.

### IVIS imaging

An IVIS Spectrum imaging system (Perkin Elmer) was utilized to monitor fluorescent signal of MMPsense750. For imaging of media cultured with cartilage explants, the samples were placed in black plates. For *in-vivo* imaging of mice, the hairs from the lower trunk and both legs were removed, and then imaging was performed under general anesthesia by isoflurane inhalation. The excitation and emission wavelengths were set to 745 and 800nm, respectively. The fluorescence intensities were analyzed by Living Image software 4.2(Perkin Elmer). Grid type and circle type of regions of interest (ROI) were set for plates and mice, respectively. Average radiant efficiency [p/s/cm<sup>2</sup>/sr] / [ $\mu$ W/cm<sup>2</sup>] in the ROI was

measured as an index of intensity of fluorescent signal. For the experiment with MMP enzymes, normalized average radiant efficiency was calculated by subtracting the value of average radiant efficiency with 0(m)U of MMPs from that with each concentration of MMPs and used for the assessment. For the purpose of clarity, we use the term “fluorescence intensity” to indicate normalized average radiant efficiency in the ROI.

### Statistical analysis

The results were statistically analyzed using a software package (GraphPad Prism; MDF Software, Inc.). Values of all measurements were expressed as the mean with error bars representing the 95% confidence interval. Correlation between MMP concentration and MMPSense signal was evaluated with Pearson correlation coefficient. Differences of MMPSense signal with MMP enzymes and *in-vivo* study were analyzed by paired t tests. Comparison of MMPSense signal between different concentrations of MMPs and of MMPSense signal with cartilage explants were analyzed by unpaired t test. Comparison of mRNA expression was analyzed by Wilcoxon signed rank test.

## Results

### Determination of the optimal MMPSense750 concentration for *in-vitro* experiments

To examine how well the normalized fluorescent signal correlated to the MMP activity at each time point, Pearson's correlation coefficients (R) were calculated for each combination of MMPSense and time (Figure 1). For all MMPs, the best correlation between MMP activity and fluorescent signal occurred at the higher concentrations of MMPSense probe, 0.7 or 2.0 $\mu$ M. Perhaps more surprisingly the best time to measure fluorescence intensity was highly dependent on the types of MMP enzyme. For MMP-13, high correlations were observed as early as 15 minutes, while MMP-3 started to become significant after 60 minutes, and MMP-9 not until after 24 hours. At the later time points the MMP activity and fluorescent signal were highly correlated for all three MMP enzymes.

To estimate the detection limit of the assay at each concentration of MMPSense750 over time, we statistically analyzed the differences between fluorescent intensity for each concentration increase of MMP (Supplementary Figure 1). The intermediate concentrations of MMP generated significantly different fluorescence signal in all conditions, but there were variations in the lower and upper detection limits. With respect to the upper detection limit, we examined the fluorescent signal over time for each enzyme and the higher MMPSense probe concentrations (Figure 2). While higher concentrations of MMPSense yielded greater absolute fluorescence signals, the statistical analysis showed no benefit of the 2 $\mu$ M compared to 0.7 $\mu$ M MMPSense probe.

Taken together, these results indicate that 0.7 $\mu$ M of MMPSense750 at a 24 hour time point would yield the best assay to measure the activities of the three MMPs over the greatest range of concentrations

### MMP activity in IL-1 $\beta$ -treated cartilage explants

Three days of IL-1 $\beta$  treatment caused a significant increase in fluorescence intensity of culture media, indicating elevated MMP activity in cartilage explants (Figure 3A). Specifically, the fluorescence signal of the IL-1 $\beta$  group at day 3 was significantly greater than that of the control group both at 60 minutes and 24 hours after adding MMPSense750. At day 6, fluorescence intensity in the IL-1 $\beta$ -treated group was significantly greater than that of the control group at 24 hours (but not 60 minutes) after adding MMPSense750.

### MMP activity in mechanically injured cartilage explants

As observed with the IL-1 $\beta$  treatment, the fluorescence of the mechanical injury group at day 3 was significantly greater than that of the control group at both 60 minutes and 24 hours after adding MMPSense750. At day 6, the trends are similar to the IL-1 $\beta$  treatment, with greater fluorescence at the 24 hour, although this did not reach statistical significance (Figure 3B).

### MMP activity *in-vivo* after knee injury

The non-surgical joint injury caused a substantial increase in the fluorescence intensity in the injured right knee relative to the uninjured left knee of the same animal, indicating that injury increased the local MMP activity (Figure 4A). The real-time RT-PCR results showed elevated mRNA expression of MMP-3 in the injured knee at this time point (48 hours after injury), while the expression of MMP-9 and -13 were not statistically different in the injured and contralateral limb at this time point (Figure 4B).

## Discussion

Imaging technologies used in OA primarily measure structural morphology rather than the biological processes that contribute to joint degradation. The results in the present study demonstrate that the MMPSense750 is useful with human MMPs, providing insight into the parameters to consider when interpreting the data, and show a good response of the assay in *in-vitro* studies of cartilage explants and in a mouse model of joint injury. Although there are a few previous reports using MMPSense680, precedent product of MMPSense, to investigate MMP activity in OA or RA with human cartilage or mouse model,[23,24,25] this article is the first studying OA-related assay with cartilage and joint injury using MMPSense750. This is an important contribution because MMPSense680 and MMPSense750 have different substrate specificities and different *in-vivo* kinetics, and MMPSense750 has the advantage that it enables a shorter time between injection and imaging (6 hours versus 24 hours for MMPSense680). Although human MMPs, bovine cartilage and mice were used in the current study, MMPs share a high degree of orthology among most vertebrates [26,27] and we do not expect significant species-related differences in the substrate-enzyme interactions.

With all MMPs tested, the fluorescent signal increased as the MMP activity increased. Interestingly, the reaction kinetics of the enzymes were different for the three MMPs. Namely, recombinant human MMP-13 caused a rapid increase in fluorescence within 15 minutes even at low enzyme concentrations, and longer incubations past 24 hours decreased

the assay linearity. In contrast, human MMP-9 required at least 24 hours to show a dose-dependent increase in substrate activation, and longer incubations out to 3 days improved the assay linearity. Human MMP-3 was intermediate, showing linear response after 1 hour out to 2 days, but decreasing at 3 days. The experiments were all performed within the reported tissue half-life of 72 hours for the MMPsense750 reagent. MMPsense produces fluorescent signal upon MMP-mediated hydrolysis, and the discrepancy of the detection time between MMPs may suggest the kinetics of cleavage differs between MMPs. We examined MMP-3, -9 and -13 in the current study, but according to the manufacturer MMPsense750 can also detect the activities of MMP-2, -7, and -12. In summary, it is important to recognize that a fluorescent signal indicates activities from multiple MMPs with differing sensitivities and reaction kinetics.

The result from cartilage explants assay provides a novel non-destructive method to quantify the MMP activity. At day 6, significantly higher fluorescence was detected in the IL-1 $\beta$  group when compared with the control group, while there was no significant difference between the mechanical injury group and the control group at both 60 minutes and 24 hours after adding MMPsense. A possible reason for this difference could be the continuous presence of IL-1 $\beta$  during the 6 days, compared to a single mechanical injury at day 0.

In the *in-vivo* mouse model, injured knees showed significantly higher signals of MMPsense750 than the contralateral uninjured knees, indicating that MMP activity was elevated 48 hours after injury. This provides a novel real-time non-destructive imaging method to quantify knee injury response and the progression of cartilage degradation and OA based on MMP activity. Interestingly, when we examined MMP mRNA expression at 48 hours after injury, we found that only MMP-3 was still elevated. In a separate study, we found that mRNA up-regulation of MMP expression after injury peaked at 4 hours after injury and returned to baseline after 24 hours using the same animal model (data not shown). MMPs are secreted as inactive proenzymes that are later activated in the extracellular matrix, which may explain the apparent discrepancy between the mRNA expression and the protease activity and highlight the importance of quantifying the enzymatic activity.

A limitation of the imaging technology is that we did not have sufficient resolution to determine the exact tissue source of MMP activity. It is likely that MMPs are active in multiple tissues, including cartilage, bone, and synovium. In our explant experiments we were able to detect activity of MMP secreted by cartilage, but not within the cartilage itself. Based on these observations we speculate that the *in-vivo* source of fluorescent signal might be joint tissues other than cartilage, although this does not preclude the MMP activity in the cartilage. In future experiments we would like to localize the source of MMP activity more precisely.

In conclusion, we established experimental parameters to use the MMPsense750 imaging reagent to quantify MMP activity *in-vitro* in cartilage explants, and *in-vivo* in a mouse joint injury model. The advantages of MMPsense750 over other techniques to evaluate MMP activity include its non-destructive nature, enabling repeated measurements on the same samples. This provides an imaging opportunity to monitor the destructive enzymatic



processes that contribute to OA progression, and complements traditional imaging technologies that quantify the resulting structural changes.

## Supplementary Material

Refer to Web version on PubMed Central for supplementary material.

## Acknowledgements

This work was funded by NIH/NIAMS grant AR063348 to Dominik R. Haudenschild.

Imaging work was performed at the Center for Molecular and Genomic Imaging (CMGI), University of California, Davis. We would like to acknowledge Douglas Rowland and Jennifer Fung for help with the in-vivo and in-vitro imaging on the IVIS-Spectrum instrument.


## References

1. Loeser RF, Goldring SR, Scanzello CR, Goldring MB. Osteoarthritis: a disease of the joint as an organ. *Arthritis Rheum.* 2012; 64:1697–1707. [PubMed: 22392533]
2. Losina E, Weinstein AM, Reichmann WM, Burbine SA, Solomon DH, Daigle ME, Rome BN, Chen SP, Hunter DJ, Suter LG, Jordan JM, Katz JN. Lifetime risk and age at diagnosis of symptomatic knee osteoarthritis in the US. *Arthritis Care Res (Hoboken).* 2013; 65:703–711. [PubMed: 23203864]
3. Lozito TP, Tuan RS. Endothelial cell microparticles act as centers of matrix metalloproteinase-2 (MMP-2) activation and vascular matrix remodeling. *J Cell Physiol.* 2012; 227:534–549. [PubMed: 21437907]
4. Yoshida K, Barr RJ, Galea-Soler S, Aspden RM, Reid DM, Gregory JS. Reproducibility and Diagnostic Accuracy of Kellgren-Lawrence Grading for Osteoarthritis Using Radiographs and Dual-Energy X-ray Absorptiometry Images. *J Clin Densitom.* 2014
5. Guermazi A, Eckstein F, Hellio Le Graverand-Gastineau MP, Conaghan PG, Burstein D, Keen H, Roemer FW. Osteoarthritis: current role of imaging. *Med Clin North Am.* 2009; 93:101–126, xi. [PubMed: 19059024]
6. Williams A, Smith JR, Allaway D, Harris P, Liddell S, Mobasheri A. Carprofen inhibits the release of matrix metalloproteinases 1, 3, and 13 in the secretome of an explant model of articular cartilage stimulated with interleukin 1beta. *Arthritis Res Ther.* 2013; 15:R223. [PubMed: 24373218]
7. Troeberg L, Nagase H. Proteases involved in cartilage matrix degradation in osteoarthritis. *Biochim Biophys Acta.* 2012; 1824:133–145. [PubMed: 21777704]
8. Verma P, Dalal K. ADAMTS-4 and ADAMTS-5: key enzymes in osteoarthritis. *J Cell Biochem.* 2011; 112:3507–3514. [PubMed: 21815191]
9. Di Cesare PE, Haudenschild DR, Samuels J, Abramson SB, Firestein GS, Kelley WN. Pathogenesis of Osteoarthritis. *Kelley's Textbook of Rheumatology.* 2013:1617–1635.
10. Cattano NM, Driban JB, Balasubramanian E, Barbe MF, Amin M, Sitler MR. Biochemical comparison of osteoarthritic knees with and without effusion. *BMC Musculoskelet Disord.* 2011; 12:273. [PubMed: 22122951]
11. Mapp PI, Walsh DA, Bowyer J, Maciewicz RA. Effects of a metalloproteinase inhibitor on osteochondral angiogenesis, chondropathy and pain behavior in a rat model of osteoarthritis. *Osteoarthritis Cartilage.* 2010; 18:593–600. [PubMed: 20067755]
12. Yamanaka H, Matsuda Y, Tanaka M, Sendo W, Nakajima H, Taniguchi A, Kamatani N. Serum matrix metalloproteinase 3 as a predictor of the degree of joint destruction during the six months after measurement, in patients with early rheumatoid arthritis. *Arthritis Rheum.* 2000; 43:852–858. [PubMed: 10765930]
13. Hollis CP, Weiss HL, Evers BM, Gemeinhart RA, Li T. In Vivo Investigation of Hybrid Paclitaxel Nanocrystals with Dual Fluorescent Probes for Cancer Theranostics. *Pharm Res.* 2013


14. Hufeland M, Schunke M, Grodzinsky AJ, Imgenberg J, Kurz B. Response of mature meniscal tissue to a single injurious compression and interleukin-1 in vitro. *Osteoarthritis Cartilage*. 2013; 21:209–216. [PubMed: 23069857]
15. Barber PA, Rushforth D, Agrawal S, Tuor UI. Infrared optical imaging of matrix metalloproteinases (MMPs) up regulation following ischemia reperfusion is ameliorated by hypothermia. *BMC Neurosci*. 2012; 13:76. [PubMed: 22742423]
16. Moon MH, Jeong JK, Lee YJ, Seol JW, Jackson CJ, Park SY. SIRT1, a class III histone deacetylase, regulates TNF-alpha-induced inflammation in human chondrocytes. *Osteoarthritis Cartilage*. 2013; 21:470–480. [PubMed: 23257246]
17. Schure R, Costa KD, Rezaei R, Lee W, Laschinger C, Tenenbaum HC, McCulloch CA. Impact of matrix metalloproteinases on inhibition of mineralization by fetuin. *J Periodontal Res*. 2013; 48:357–366. [PubMed: 23058002]
18. Waschkau B, Faust A, Schafers M, Bremer C. Performance of a new fluorescence-labeled MMP inhibitor to image tumor MMP activity in vivo in comparison to an MMP- activatable probe. *Contrast Media Mol Imaging*. 2013; 8:1–11. [PubMed: 23109387]
19. Vaatainen U, Lohmander LS, Thonar E, Hongisto T, Agren U, Ronkko S, Jaroma H, Kosma VM, Tammi M, Kiviranta I. Markers of cartilage and synovial metabolism in joint fluid and serum of patients with chondromalacia of the patella. *Osteoarthritis Cartilage*. 1998; 6:115–124. [PubMed: 9692066]
20. Misko TP, Radabaugh MR, Highkin M, Abrams M, Friese O, Gallavan R, Bramson C, Hellio Le Graverand MP, Lohmander LS, Roman D. Characterization of nitrotyrosine as a biomarker for arthritis and joint injury. *Osteoarthritis Cartilage*. 2013; 21:151–156. [PubMed: 23025928]
21. Lu YC, Evans CH, Grodzinsky AJ. Effects of short-term glucocorticoid treatment on changes in cartilage matrix degradation and chondrocyte gene expression induced by mechanical injury and inflammatory cytokines. *Arthritis Res Ther*. 2011; 13:R142. [PubMed: 21888631]
22. Christiansen BA, Anderson MJ, Lee CA, Williams JC, Yik JH, Haudenschild DR. Musculoskeletal changes following non-invasive knee injury using a novel mouse model of post-traumatic osteoarthritis. *Osteoarthritis Cartilage*. 2012; 20:773–782. [PubMed: 22531459]
23. Jones EF, Schooler J, Miller DC, Drake CR, Wahnische H, Siddiqui S, Li X, Majumdar S. Characterization of human osteoarthritic cartilage using optical and magnetic resonance imaging. *Mol Imaging Biol*. 2012; 14:32–39. [PubMed: 21384207]
24. Zhou J, Chen Q, Lanske B, Fleming BC, Terek R, Wei X, Zhang G, Wang S, Li K, Wei L. Disrupting the Indian hedgehog signaling pathway in vivo attenuates surgically induced osteoarthritis progression in Col2a1-CreERT2; Ihhf1/fl mice. *Arthritis Res Ther*. 2014; 16:R11. [PubMed: 24428864]
25. Galligan CL, Fish EN. Circulating fibrocytes contribute to the pathogenesis of collagen antibody-induced arthritis. *Arthritis Rheum*. 2012; 64:3583–3593. [PubMed: 22729466]
26. Jackson BC, Nebert DW, Vasiliou V. Update of human and mouse matrix metalloproteinase families. *Hum Genomics*. 2010; 4:194–201. [PubMed: 20368140]
27. Massova I, Kotra LP, Fridman R, Mobashery S. Matrix metalloproteinases: structures, evolution, and diversification. *FASEB J*. 1998; 12:1075–1095. [PubMed: 9737711]

1. MMPsense750 is near-infrared fluorescent probe which can detect MMP activity.
2. MMPsense750 can detect human MMP-3, -9, and -13.
3. The reaction kinetics with MMPsense750 were different for the three MMPs.
4. MMPsense750 can visualize real time MMP activity in mouse injured knees.
5. MMPsense750 is a convenient tool to evaluate real-time MMP activity non-invasively.

**R<sup>2</sup>(R: Pearson Correlation Coefficient)**




**Figure 1. Correlation between MMP Activity and MMPsense750 Signal**  
 R<sup>2</sup> (R: Pearson correlation coefficient) as indices of correlation of the relationship between MMPs concentration and fluorescent signals at each time point were shown in tables and figures. R<sup>2</sup> greater than 0.8 were shown in the table with bold letters and shaded background.



**Figure 2. Substrate Activation over Time**

The normalized average radiant efficiency of the greatest and the second greatest concentration of each MMP with 0.7 or 2.0 μM of MMPSense750. p-values vs. the second greatest concentrations at the same time points. (n=3)





\*:  $p < 0.05$ , \*\*:  $p < 0.01$ , \*\*\*:  $p < 0.001$

### Figure 3. Increased MMP Activity in Models of Cartilage Explant Injury




The average radiant efficiency in culture media at day 3 of the IL- $\beta$  group (A) and the mechanical injury group (B) measured at 60 minutes and 24 hours after adding MMPsense was significantly greater than that of the control group. As for the media collected at day 6, the average radiant efficiency of the IL-1 $\beta$  group measured at 24 hours was significantly greater than the other, while there was no significant difference in the radiant efficiency measured at 60 minutes of the IL-1 $\beta$  group and at 60 minutes and 24 hours of the mechanical injury group compared to the control group. M. Inj.: mechanical injury group, Ctrl: control group. (n=10)

**A. *in-vivo* imaging**



\*: p<0.05  
 \*\*: p<0.01  
 \*\*\*: p< 0.001

**B. Real-time RT-PCR**



**Figure 4. Increased MMP Activity and mRNA in a Mouse Model of Joint Injury**

A. Representative IVIS imaging of knees after MMPsense750 injection are shown. The image is a merged picture of fluorescent signal in color and a grayscale picture of the mouse. Blue circles on both knees are ROI with same shape and size as each other. The graph shows that average radiant efficiency in injured knees was significantly higher than the controls. (n=8)

B. mRNA expression of MMP-3 in injured knees were significantly greater than in uninjured knees. No significant difference in MMP-9 and MMP-13 mRNA expression was detected between the knees at this time point. (n=8)