

# UC San Diego

## SIO Reference

### **Title**

Determination of the on-zero asymptote of an exponential decay function

### **Permalink**

<https://escholarship.org/uc/item/6cb7w69p>

### **Author**

Richardson, W H

### **Publication Date**

1958-07-01

Visibility Laboratory  
University of California  
Scripps Institution of Oceanography  
San Diego 52, California

DETERMINATION OF THE NON-ZERO ASYMPTOTE  
OF AN EXPONENTIAL DECAY FUNCTION

W. H. Richardson


1 July 1958  
Index Number NS 714-100


Bureau of Ships  
Contract NObs-72092

SIO REFERENCE 58-36

Approved:

Approved for Distribution:

  
Seibert Q. Duntley, Director  
Visibility Laboratory

  
Roger Revelle, Director  
Scripps Institution of Oceanography

# Determination of the Non-zero Asymptote of an Exponential Decay Function\*

William Hadley Richardson

Scripps Institution of Oceanography, University of California

La Jolla, California

It frequently becomes necessary in physical investigations to deal with exponential decay functional forms where the asymptote is displaced from zero. The method of least squares as normally applied cannot be used, and other methods are complicated by the presence of the non-zero asymptote. However, if the data is based on equal progressive increments of the independent variable, the asymptote may be readily found by the following method. The method may also be applied to data involving unequal increments provided that the data can be reduced to effectively equal incremental form. Given a set of data points and knowing or assuming that they should fall on a curve of the type:

$$y = a e^{bx} + c$$

Where  $b$  is less than 0

Divide data into three groups, each of which centers about  $x_1, x_2, x_3$  so that  $x_2 - x_1 = x_3 - x_2$  and  $2x_2 = x_1 + x_3$

Then take the geometric mean of each group and consider this mean the

$y_1(x_1), y_2(x_2), y_3(x_3)$  respectively.

$$y_i = a e^{bx_i} \quad \text{or} \quad e^{bx_i} = \frac{y_i - c}{a}$$

---

\*Contribution from the Scripps Institution of Oceanography, New Series No. \_\_\_\_\_.  
This paper represents results of research which has been supported by the  
Bureau of Ships, U. S. Navy.

$$\left(\frac{y_2 - c}{a}\right)^{\frac{x_1}{x_2}} = \frac{y_1 - c}{a} \quad \text{or} \quad a^{\frac{x_2 - x_1}{x_2}} = \frac{y_1 - c}{(y_2 - c)^{\frac{x_1}{x_2}}}$$

$$\left(\frac{y_2 - c}{a}\right)^{\frac{x_3}{x_2}} = \frac{y_3 - c}{a} \quad \text{or} \quad a^{\frac{x_2 - x_3}{x_2}} = \frac{y_3 - c}{(y_2 - c)^{\frac{x_3}{x_2}}}$$

Since  $x_2 - x_1 = x_3 - x_2$  :  $a^{\frac{x_2 - x_1}{2}} = a^{\frac{x_3 - x_2}{x_2}} = \frac{1}{a^{\frac{x_2 - x_3}{x_2}}}$

and  $\frac{(y_2 - c)^{\frac{x_3}{x_2}}}{y_3 - c} = \frac{y_1 - c}{(y_2 - c)^{\frac{x_1}{x_2}}}$  ; or  $(y_2 - c)^{\frac{x_3 + x_1}{x_2}} = (y_1 - c)(y_3 - c)$

since  $x_3 + x_1 = 2x_2$  :  $(y_2 - c)^2 = (y_1 - c)(y_3 - c)$

$$y_2^2 - 2cy_2 + c^2 = y_1 y_3 - cy_1 - cy_2 + c^2$$

$$c(y_1 + y_3 - 2y_2) = y_1 y_3 - y_2^2$$

$$c = \frac{y_1 y_3 - y_2^2}{y_1 + y_3 - 2y_2}$$

EXAMPLE: In this case a series of attenuation constants varying with depth has been determined. It appeared that the constants diminished exponentially with depth. Applying the method to the following data:

j	x	k	
1	0	0.505	$k_1 = (0.505 \times 0.463)^{1/2} = 0.483,544$
2	40	0.463	
3	80	0.455	$k_2 = (0.455 \times 0.444)^{1/2} = 0.449,466$
4	120	0.444	
5	160	0.440	
6	200	0.437	$k_3 = (0.440 \times 0.437)^{1/2} = 0.438,497$

$$k_{\infty} = \frac{k_1 k_3 - k_2^2}{k_1 + k_3 - 2k_2} = \frac{0.483,544 \times 0.438,497 - 0.449,466^2}{0.483,544 + 0.438,497 - 2 \times 0.449,466} = \frac{0.010,013}{0.023,109} = 0.433$$

Notes:

1. This method may be used wherever three or more points are known, satisfying the condition that the resulting  $x_2 - x_1 = x_3 - x_2$
2. Knowing the three meanpoints  $y_1, y_2$  and  $y_3$  and the asymptote, the equation to fit these points may be found by subtracting the asymptote from the values of  $y_i$  and applying any of various methods, including the method of least squares.
3. The principal source of error is the subtractive process in the numerator and particularly in the denominator of the formula. Hence it is advisable to carry as ~~many~~ many figures in the geometric means as is practicable.