

UC Irvine

UC Irvine Previously Published Works

Title

Aluminum and copper in drinking water enhance inflammatory oroxidative events specifically in the brain

Permalink

<https://escholarship.org/uc/item/6cf5g9rg>

Journal

J. Neuroimmunology, 176

Author

Bondy, Stephen Bondy C

Publication Date

2006

Peer reviewed

Aluminum and copper in drinking water enhance inflammatory or oxidative events specifically in the brain

Angelica Becaria^a, Debomoy K. Lahiri^b, Stephen C. Bondy^a, DeMao Chen^b, Ali Hamadeh^a, Huihui Li^a, Russell Taylor^a, Arezoo Campbell^{a,*}

^a Department of Community and Environmental Medicine, Center for Occupational and Environmental Health Sciences, University of California, Irvine, Irvine, CA 92697-1825, USA

^b Department of Psychiatry, Institute of Psychiatric Research, Indiana University School of Medicine, Indianapolis, IN 42602-4887, USA

Received 5 January 2006; received in revised form 30 March 2006; accepted 31 March 2006

Abstract

Inflammatory and oxidative events are up-regulated in the brain of AD patients. It has been reported that in animal models of AD, exposure to aluminum (Al) or copper (Cu) enhanced oxidative events and accumulation of amyloid beta (A β) peptides. The present study was designed to evaluate the effect of a 3-month exposure of mice to copper sulfate (8 μ M), aluminum lactate (10 or 100 μ M), or a combination of the salts. Results suggest that although Al or Cu may independently initiate inflammatory or oxidative events, they may function cooperatively to increase APP levels.

© 2006 Elsevier B.V. All rights reserved.

Keywords: Aluminum; Alzheimer's disease; Amyloid beta; Copper; Neuroinflammation; Oxidative stress

1. Introduction

Cu is an essential metal and a component of many enzymatic reactions. However, this redox active metal can also mediate the formation of reactive oxygen species (ROS) and this can have adverse consequences. Al is a trivalent cation incapable of redox changes and unlike Cu, has no known biological role. Both metals have been associated with neurological impairments.

Al has been shown to play a causal role in dialysis encephalopathy (Alfrey et al., 1976) and epidemiological studies suggest a possible link between exposure to this metal and a higher prevalence of AD (McLachlan et al., 1996; Rondeau et al., 2000; Flaten, 2001). This association is dependent on the duration of Al exposure and only becomes significant if an individual has resided in an area with high Al in drinking water (>100 μ g/L) for several

years. Homeostatic alterations in brain Cu levels have also been implicated in the pathogenesis of several neurological disorders including Alzheimer's, Parkinson's and prion diseases (Torsdottir et al., 1999; Mercer, 2001; Brown and Kozlowski, 2004). The toxicity of excess Cu is best demonstrated by the heritable disorder, Wilson's disease, in which the liver is unable to excrete the metal and thus there is tissue accumulation. In this condition, liver and brain are the most compromised areas (Kitzberger et al., 2005).

The present study investigated the potential synergistic effects of chronic exposure to Cu, Al or both metals in promoting inflammatory and oxidative events in mouse brains. The design was based on the following observations: Al present in the drinking water enhanced inflammatory markers in the CNS (Campbell et al., 2004). Furthermore, in transgenic mouse models of AD, dietary Al increased markers of lipid peroxidation and A β levels (Pratico et al., 2002a,b). In isolated systems, Al can potentiate the oxidative stress produced by transition metals such as iron (Bondy and Kirstein, 1996) or Cu (Bondy et al., 1998;

* Corresponding author. Tel.: +1 949 824 3439; fax: +1 949 824 2070.
E-mail address: aghadimi@uci.edu (A. Campbell).

Becaria et al., 2003) and Cu present in trace amounts (0.12 ppm) in the drinking water has been implicated in the accelerated deposition of A β and promotion of oxidative events in the brain of a rabbit model of AD (Sparks and Schreurs, 2003).

Animals were exposed for 12 weeks to Al lactate (10 or 100 μ M), Cu sulfate (8 μ M), or both metals concomitantly, and oxidative and inflammatory markers were evaluated. To determine if exposure to the metals contribute to potentiation of AD pathology, levels of APP, and its constituent peptides, A β _{1–40} and A β _{1–42}, were also determined. The lower Al exposure is in the range of concentrations reported to be associated with an increased risk of AD. The present EPA maximum recommended standard for Cu in drinking water is 1.3 ppm (Buchanan et al., 1999). However, levels as high as 7.8 ppm has been found in tap water (Spitalny et al., 1984). The copper (8 μ M) exposure used in this study is equivalent to 2 ppm.

While Al caused an increase in CNS proinflammatory cytokines, both metals, independently, enhanced oxidative markers in the brain. Except for an enhancement of APP levels, there was no other synergistic interaction between the two metals.

2. Materials and methods

Unless otherwise noted, all chemicals were purchased from Sigma (St. Louis, MO).

2.1. Animal maintenance and exposure

Two-month-old male B6C3F1 mice (a hybrid between C57BL/6 and C3H) were obtained from Charles River Laboratories (Wilmington, MA). They were housed four per cage and maintained on a 12-h light/dark cycle in a temperature controlled (20 \pm 1 $^{\circ}$ C) room. Food and water were provided ad lib. Mice were divided randomly into 6 groups: (1) control, (2) copper sulfate (8 μ M), (3) aluminum lactate (10 μ M), (4) aluminum lactate (10 μ M) plus copper sulfate (8 μ M), (5) aluminum lactate (100 μ M), and (6) aluminum lactate (100 μ M) plus copper sulfate (8 μ M). The metal salts were freshly prepared and added to double-distilled water twice weekly. The drinking water of the control animals was also double distilled. The animals were exposed for 12 weeks. The weight of the animals or the amount of water consumed was not different between groups.

2.2. Preparation of samples

Animals were anesthetized with pentobarbital (65 mg/kg, i.p.). After blood was collected from the right ventricle, the mice were perfused transcardially with phosphate buffer saline to remove blood from all organs. The brain and spleen were removed and quickly frozen. The blood was

centrifuged at 1000 \times g for 10 min and the serum was collected and stored at -80° C.

Cytoplasmic fractions were prepared using the method of Lahiri and Ge (2000). Tissue from each animal was weighed and homogenized in 2 ml of an ice-cold buffer consisting of (10 mM HEPES (pH 7.9), 10 mM KCl, 0.1 mM EDTA, 0.1 mM EGTA, 1 mM DTT, 0.5 mM PMSF and 0.5% NP-40). The suspension was incubated on ice for 10 min and centrifuged (1500 \times g) at 4 $^{\circ}$ C for 1 min. The supernatant containing the cytoplasmic constituents was collected and 200 μ l of 5 \times protease inhibitor cocktail was added. Samples were aliquoted and stored at -80° C.

2.3. Competitive enzyme immunoassay

Levels of TNF- α or IL-1 α were determined using a competitive enzyme immunoassay kit from Neogen Corp. (Lexington, KY) for the detection of the total protein in the cytoplasmic tissue fractions. Briefly, 100 μ l of the brain or spleen cytoplasmic fraction, along with murine TNF- α or IL-1 α antibody were added to plates precoated with goat anti-rabbit antibody and incubated at room temperature for 3 h. Murine TNF- α or IL-1 α conjugate was added and incubated for 30 min. The plate was washed and incubated for another 30 min with streptavidin–alkaline phosphatase. Then 200 μ l of chromogen was added to the plate and the color generated was determined with a spectrophotometric plate reader set at 490 nm. Levels of TNF- α or IL-1 α were determined in the serum using sandwich enzyme immunoassay kits (Neogen Corp., Lexington, KY) for the detection of free forms of the proteins.

Levels of IL-4 were measured in the brain, serum, or spleen using a kit from Biosource (Camarillo, CA). Samples (50 μ l) were added to plates coated with a monoclonal antibody against mouse IL-4. After addition of a biotin conjugate, plates were incubated for 2 h. The plate was washed and streptavidin–HRP was added to the plate. After 30-min incubation, the plate was washed and chromogen was added. The color generated was measured at 450 nm.

2.4. Immunohistochemistry assay

Animals were anesthetized with pentobarbital (65 mg/kg, i.p.) and then perfused transcardially with saline, followed by 4% paraformaldehyde. The brains were excised and placed in 4% paraformaldehyde. The tissue was prepared as previously described (Campbell et al., 2004). For nNOS staining, the sections were stained free-floating. After hydrogen peroxide treatment, samples were blocked in serum, and then immunostained with a 1:300 dilution of primary rabbit anti-nNOS antibody (Immunostar; #24431), a goat anti-rabbit secondary antibody, and finally an avidin–biotin–HRP complex (Elite Vectastain ABC kit, Vector, Burlingame, CA). Incubation times were 24 h for the primary antibody, 30 min for the secondary antibody, and 1 h for the avidin–biotin–HRP complex. Sections were

treated with hydrogen peroxide with DAB to visualize the antibody binding sites and mounted on gelatinized glass slides. The nNOS images were captured by a Nikon DXM1200F HIQE Color CCD camera attached to an Eclipse TE2000 fluorescent microscope.

2.5. Malondialdehyde (MDA) assay

The levels of MDA were determined spectrophotometrically by following a modified protocol developed by Gerard-Monnier et al. (1998). This procedure measures free MDA as opposed to the commonly used TBARS (thio-barbituric acid-reactive substances) method which measures total MDA in biological samples. The modified protocol is specific for this lipid peroxidation product because 4-hydroxyalkenals do not produce significant color under the conditions of the assay. To further minimize their reaction, an antioxidant was added. 150 μ l of brain cytoplasmic fractions, 10 μ l of the antioxidant probucol and 690 μ l of the chromogenic reagent phenylmethyl indole were added together. Hydrochloric acid was then added to the mixture and the tubes were incubated for 1 h at 45 °C. A clear supernatant was obtained by centrifugation at 10,000 \times g for 10 min. The supernatant was transferred to a cuvette and absorbance was measured at 586 nm.

2.6. Assay for A β Levels

A sensitive ELISA test was used for measuring levels of A β_{1-40} and A β_{1-42} by using a previously described assay (Lahiri et al., 1998). Highly specific antibodies were used to detect A β_{1-40} and A β_{1-42} species. An affinity-purified anti-A β rabbit IgG was used as a capture antibody and an affinity-purified HRP-conjugated anti-A β rabbit IgG Fab fragment was used as a detection antibody. Tetramethylbenzidine (TMB) was used as the chromogen.

2.7. Assay for APP levels

APP was measured by Western blotting technique as previously described (Lahiri et al., 2004). Briefly, 30 μ g of total brain protein was separated on a 12% polyacrylamide gel containing sodium dodecyl sulfate (SDS-PAGE). Proteins from the gel were electrophoretically transferred to a Nytran membrane. Each immunoblot was independently probed with a mouse monoclonal antibody, 22C11, which was raised against the N-terminus of APP, and mouse monoclonal antibody against the beta-actin protein. The 22C11 clone identifies all forms of APP found in cell membranes and the APP-like proteins (APLP1 and APLP2). A biotinylated horse antimouse IgG was used as the secondary antibody. Immunochemical detection was performed and density of each immunoreactive band was measured by using UVP-GDS5000 Gel Documentation system (San Gabriel, CA, USA). Signals from APP bands were normalized with beta-actin bands. Data was analyzed

using Scion Image software (Scion Corporation, Frederick, MD, USA).

2.8. Statistical analysis

The difference among groups was assessed using one-way ANOVA, followed by the Tukey test. Results were considered significant at $P \leq 0.05$.

3. Results

3.1. Cytokine levels

The levels of TNF- α were elevated in the brains of animals exposed to Al (100 μ M) or Al (100 μ M)+Cu (8




Fig. 1. Brain cytokine levels. Levels of TNF- α (A), IL-1 α (B), or IL-4 (C), in brain cytoplasmic fractions after treatment with aluminum lactate and/or copper sulfate in drinking water for 12 weeks [Con=Control; Cu=Cu (8 μ M); Al10=Al (10 μ M); Al10+Cu=Al (10 μ M) plus Cu (8 μ M); Al100=Al (100 μ M); Al100+Cu=Al (100 μ M) plus Cu (8 μ M)]. Levels of TNF- α and IL-1 α were greatest in the Al100 and Al100+Cu exposed animals. The levels of IL-4 were increased in the Al10, Al10+Cu and Al100+Cu groups. After exposure to Cu alone, levels of IL-4 were not significantly altered. Bars represent mean of six individual determinations \pm S.E. ($P \leq 0.05$).




Fig. 2. Serum cytokine levels. Levels of TNF- α (A) or IL-1 α (B) in serum after exposure to aluminum lactate and/or copper sulfate in drinking water for 12 weeks [Con=Control; Cu=Cu (8 μ M); Al10=Al (10 μ M); Al10+Cu=Al (10 μ M) plus Cu (8 μ M); Al100=Al (100 μ M); Al100+Cu=Al (100 μ M) plus Cu (8 μ M)]. Levels of TNF- α were depressed in treatment groups except Al10+Cu. IL-1 α levels were increased after exposure to Al10, Al10+Cu, Al100, or Al100+Cu. The changes in the Al100 and Al100+Cu exposed animals were significantly greater than those of the Al10 and Al10+Cu groups. Bars represent mean of six individual determinations \pm S.E. ($P \leq 0.05$).

μ M) but there was no significant difference between these two treatment groups (Fig. 1A). To determine whether this response was mediated by systemic pro-inflammatory changes, serum and spleen levels were also measured (Figs. 2A and 3A). In the serum, TNF- α levels decreased in all metal-exposed groups compared to the control animals. This effect was not significant in the Al (10 μ M)+Cu (8 μ M) group, which had high variance (Fig. 2A). Following Al treatment, levels of TNF- α in spleen decreased in a dose-dependent manner. Exposure to Al together with Cu further enhanced the decrease in TNF- α in this tissue (Fig. 3A). The IL-1 α response in the brain was similar to the TNF- α response in that the levels of this cytokine were greater after exposure to Al (100 μ M) or Al (100 μ M)+Cu (8 μ M). Again, there was no significant difference between the two treatment groups (Fig. 1B).

In contrast to TNF- α , levels of IL-1 α dose-dependently increased in the serum of Al-treated animals and co-exposure to Cu did not modulate this effect (Fig. 2B). Basal IL-1 α levels were greater in the spleen than in serum or brain. Exposure to Cu, Al (10 μ M) or Al (100 μ M) did not significantly affect the levels of the cytokine in this tissue. However co-exposure with Cu and Al increased the levels of IL-1 α at both concentrations used. Exposure to Al (100 μ M) together with Cu (8 μ M) caused a greater response

compared to treatment to either Cu or Al (100 μ M) alone (Fig. 3B).

Levels of the anti-inflammatory cytokine IL-4 were also determined in the brain, blood and spleen. The levels were undetectable in the serum. Compared to control, brain levels significantly increased after exposure to Al (10 μ M), Al (10 μ M)+Cu (8 μ M) or Al (100 μ M)+Cu (8 μ M). After exposure to Cu or Al (100 μ M), there was no




Fig. 3. Spleen cytokine levels. Levels of TNF- α (A), IL-1 α (B), or IL-4 (C) in the spleen after exposure to aluminum lactate and/or copper sulfate in drinking water for 12 weeks [Con=Control; Cu=Cu (8 μ M); Al10=Al (10 μ M); Al10+Cu=Al (10 μ M) plus Cu (8 μ M); Al100=Al (100 μ M); Al100+Cu=Al (100 μ M) plus Cu (8 μ M)]. Levels of TNF- α were decreased after exposure to Al10+Cu, Al100, or Al100+Cu. After Cu exposure, the levels of TNF- α were not significantly different from the control or Al-treated animals. In the Cu-treated animals, TNF- α levels were significantly higher than those of the Al10+Cu and Al100+Cu groups. After exposure to Al100+Cu, TNF- α levels were lower when compared to the Al100 exposed animals. In the spleen, levels of IL-1 α were increased in the Al10+Cu and Al100+Cu groups. Levels of IL-1 α were further increased in the Al100+Cu group over the elevation found with 100 μ M Al alone. IL-4 levels were decreased in Al10, Al10+Cu, Al100, and Al100+Cu-treated animals. Bars represent mean of six individual determinations \pm S.E. ($P \leq 0.05$).




Fig. 4. Brain lipid peroxidation. MDA levels in brain cytoplasmic fraction after exposure to aluminum lactate and/or copper sulfate in drinking water for 12 weeks [Con=Control; Cu=Cu (8 µM); Al10=Al (10 µM); Al10+Cu=Al (10 µM) plus Cu (8 µM); Al100=Al (100 µM); Al100+Cu=Al (100 µM) plus Cu (8 µM)]. MDA levels were increased after exposure to Cu, Al10+Cu, Al100, or Al100+Cu. The Al10+Cu group had higher levels of MDA than that of the group receiving 10 µM Al alone. In contrast, the Al100+Cu group had a lower MDA content than that of the Al100 group. Bars represent mean of six individual determinations ± S.E. ($P \leq 0.05$).




Fig. 5. nNOS immunoreactivity. (A) nNOS positive neurons in the cortex of mice after treatment with aluminum lactate and/or copper sulfate in drinking water for 12 weeks. Bar=100 µm. (B) Quantification of the nNOS immunoreactive neurons [Con=Control; Cu=Cu (8 µM); Al100=Al (100 µM); Al100+Cu=Al (100 µM) plus Cu (8 µM)]. The number of nNOS immunoreactive neurons was elevated in the Cu, Al100, and Al100+Cu groups. The number of nNOS positive neurons was greater in the Al100 group relative to the Al100+Cu group. Bars represent mean of three individual determinations ± S.E. ($P \leq 0.05$).

significant increase in IL-4 levels in the brain compared to control. Treatment with Cu in combination with Al (100 µM) significantly increased the levels of this cytokine relative to exposure to Al (100 µM) alone (Fig. 1C). In the spleen, levels of IL-4 decreased after exposure to Al at both concentrations. This response was not modulated by the further addition of Cu to the drinking water (Fig. 3C).

3.2. Oxidative markers

The lipid peroxidation end product malondialdehyde (MDA) was significantly elevated in the brain tissue after exposure to Cu (8 µM), or Al (100 µM), or Al (10 µM)+Cu (8 µM), or Al (100 µM)+Cu (8 µM). Al (10 µM) did not significantly increase the levels of MDA but together with Cu caused a response similar to that found after exposure to Cu alone. MDA levels were greatest after exposure to Al (100 µM) and co-exposure with Cu significantly decreased this response (Fig. 4).

The number of nNOS-immunoreactive neurons increased in the frontal cortex of mice exposed to Cu (8 µM), or Al (100 µM), or Al (100 µM)+Cu (8 µM). Treatment with Al and Cu in combination decreased the response relative to exposure to Al (100 µM) alone (Fig. 5).

3.3. APP and Aβ levels

Aluminum (100 µM) or copper (8 µM) exposure did not lead to an increase in the levels of Aβ₁₋₄₀ and Aβ₁₋₄₂ (Fig. 6). In contrast, levels of APP were significantly increased




Fig. 6. Brain Aβ levels. Levels of Aβ₁₋₄₀ (A) or Aβ₁₋₄₂ (B) in the brain of mice exposed to copper, aluminum, or both salts [Con=Control; Cu=Cu (8 µM); Al100=Al (100 µM); Al100+Cu=Al (100 µM) plus Cu (8 µM)]. Bars represent mean of six individual determinations ± S.E.




Fig. 7. CNS concentrations of APP. Relative levels of APP, compared with control, in the brain cytoplasmic fraction of animals exposed to aluminum lactate and/or copper sulfate in drinking water for 12 weeks [Con=Control; Cu=Cu (8 μ M); Al10=Al (10 μ M); Al10+Cu=Al (10 μ M) plus Cu (8 μ M); Al100=Al (100 μ M); Al100+Cu=Al (100 μ M) plus Cu (8 μ M)]. APP levels were elevated in the Al100 and Al100+Cu-treated animals. The levels of the protein were greatest in the Al100+Cu group. Bars represent mean of six individual determinations \pm S.E. ($P \leq 0.05$).

after exposure to Al (100 μ M) and copper co-exposure further enhanced this effect (Fig. 7).

4. Discussion

Inflammatory events are present in neurodegenerative disorders and appear to contribute to progression of the disease (Campbell, 2004). This is illustrated by the finding that significant increase in CSF TNF- α levels precedes development of AD in patients with mild cognitive impairment (Tarkowski et al., 2003). We have previously reported that Al present in drinking water dose-dependently increased TNF- α and IL-1 α levels selectively in mouse brains (Campbell et al., 2004). Furthermore, in primary human neural cells, Al exposure induces expression of inflammatory genes (Lukiw et al., 2005). Since both Al and Cu have been implicated in the pathology of AD (Pratico et al., 2002a,b; Sparks and Schreurs, 2003), we investigated whether co-exposure to both metals in the drinking water further exacerbate the effects of individual metals. Cu did not modulate the Al-induced pro-inflammatory events in the brain and this was paralleled by similar findings using an isolated human glioblastoma cell line (Becaria et al., 2003).

The elevation in TNF- α appears to be unique to brain tissue since the concentrations in both serum and spleen decreased after exposure to the metal salts. This provides evidence that the inflammatory response within the CNS consequent to chronic treatment with low levels of Al differs qualitatively from that of the systemic immune response. Furthermore, the Al-induced neuroinflammatory response cannot be attributed to penetration of plasma TNF- α into the brain. In contrast, the Al-induced enhancement of IL-1 α levels in the brain may be influenced by systemic events since serum concentrations of this cytokine were also increased after exposure to Al. The spleen of the mice contained the greatest concentration of IL-1 α (75–107)

compared to the serum (0.8–1.9) or brain (2.3–4.7) levels (values are expressed as (ng/ml)/mg protein and are in the range of lowest to highest concentrations found). Thus, IL-1 α may have a homeostatic function in the spleen. After exposure to Al and Cu, levels of this cytokine were increased in cytoplasmic fractions from this organ.

Al hydroxide (alum) is the most commonly used adjuvant in vaccines. Intraperitoneal injection of alum caused spleen accumulation of IL-4 producing cells. These cells expressed granulocyte, monocyte/macrophage markers (Jordan et al., 2004). Since activated microglia also express IL-4 (Park et al., 2005), the present study investigated how a different duration and route of exposure to Al would affect the levels of this cytokine in the brain and spleen. Chronic exposure to low levels of Al lactate in drinking water decreased the levels of IL-4 in the spleen. Cu did not have an effect and did not modulate the response to Al. In the brain, IL-4 has been shown to function as an anti-inflammatory cytokine that reduces mediators of brain inflammation such as TNF- α , IL-1 β and iNOS (Ledebor et al., 2000). There was a significant increase in IL-4 levels after exposure to Al (10 μ M), or to Cu plus Al (10 or 100 μ M). The increase in IL-4 may be attenuating TNF- α and IL-1 α levels after exposure to low levels of Al (10 μ M). However after exposure to higher concentrations of Al (100 μ M) plus Cu, the enhanced IL-4 response was not able to mediate the increase in the pro-inflammatory cytokines. The interplay between pro- and anti-inflammatory cytokines released by CNS cells may determine the over-all inflammatory milieu of the brain after an environmental exposure to metals.

Although oxidative and inflammatory events are independent phenomena, they are related. During inflammation, an increase in NO production may increase the oxidative status of the CNS. The neuronal isoform of nitric oxide synthase (nNOS) is highly responsive to changes in oxidative stress (Liu et al., 2002). Perinatal exposure to Al (5 mM) in the drinking water increased the number of nNOS immunoreactive neurons in the frontal cortex of rat pups (Kim, 2003). In the brain tissue, MDA and nNOS levels were elevated after exposure to either Al (100 μ M) or Cu (8 μ M). Al and other trivalent metals which have no redox capacity in biological systems may stimulate lipid peroxidation by interacting with membrane phospholipids, causing changes in the physical properties of the membrane such as rigidification and phase separation (Verstraeten et al., 1997).

Although in vitro, Al interacts with Cu to increase its oxidative potential (Bondy et al., 1998; Becaria et al., 2003; Di and Bi, 2003) this was not the case in the present study. After exposure to Al (100 μ M) plus Cu, MDA and nNOS levels decreased compared to the response obtained with Al (100 μ M) alone and were similar to those of the Cu-exposed group. This may be due to the dual nature of Cu. Although it is a redox-active metal, it is also a cofactor in the antioxidant enzyme Cu/Zn superoxide dismutase.

To determine to what extent the inflammatory and oxidative changes may influence amyloid pathology, the levels of APP, A β_{1-40} , and A β_{1-42} were determined. Al has been shown to aggregate A β in vitro (Exley et al., 1993; Kawahara et al., 1994; Bondy and Truong, 1999) and in vivo (Miu et al., 2003). Dietary exposure to the metal exacerbates oxidative stress, A β deposition, and plaque formation in the brain of transgenic mice over-expressing APP (Pratico et al., 2002a,b). The binding of Cu to A β also promotes amyloid aggregation (Atwood et al., 2000) and trace amounts of Cu in the drinking water led to accelerated deposition of A β and promotion of oxidative events in the brain of a rabbit model of Alzheimer's disease (Sparks and Schreurs, 2003). In the brain of these animals, fibrillar forms of A β increased predominantly (Sparks, 2004). In the present study, the concentrations of A β peptides were not significantly altered after exposure to the metals. However, Al (100 μ M) increased levels of APP and co-exposure with Cu further enhanced this response. This finding is in agreement with a previous study showing that in primary human neural cells, Al or iron exposures up-regulated gene expression for APP and that the combination of the metals was more effective compared to the response to the metals individually (Alexandrov et al., 2005).

The events leading to the common sporadic form of AD are unknown. The pathophysiology of AD is complex and most likely involves multiple distinct and overlapping pathways of neuronal damage. In addition to the presence of the pathological hallmarks of the disease (senile plaques and neurofibrillary tangles), AD brains exhibit other abnormalities: loss of synapses, gliosis, microglial activation, signs of inflammation and oxidative damage (Pratico et al., 2002b). Current evidence suggests that both oxidative stress and neuroinflammation are early events in the pathogenesis of AD. The relationship between inflammatory responses and free radical generation is complex and although they may be separate events, the two are correlated. Thus, in a transgenic animal model of AD, long-term treatment with the anti-inflammatory drug indomethacin, in combination with the anti-oxidant, vitamin E, improves amyloid plaque pathology much more efficiently than the individual compounds (Yao et al., 2004).

In the present study, it was postulated that Al and Cu co-exposure would lead to exacerbation of oxidative and inflammatory factors. While in CNS, Al, at the higher concentration, acted as a pro-oxidant and a pro-inflammatory agent, the administration of the metal with Cu did not synergistically modulate the response obtained with Al alone. However, Cu did potentiate the Al-induced increase in brain APP levels. Thus, the results suggest that in the CNS, the oxidative stress and inflammation caused by Al or Cu may be, to some extent, separate events. However, the metals may work in concert to increase levels of APP. To what extent this increase may contribute to the pathological cascade associated with AD is at present unknown. Since both Cu and Al have been shown to accelerate amyloid

plaques in animal models of AD, the increase in APP may eventually lead to exacerbation of amyloid pathology.

Acknowledgement

This study was supported by grants from the National Institute of Health (ES 7992, AG-16794, AG18379 and AG18884).

References

- Alexandrov, P.N., Zhao, Y., Pogue, A.I., Tarr, M.A., Kruck, T.P., Percy, M.E., Cui, J.G., Lukiw, W.J., 2005. Synergistic effects of iron and aluminum on stress-related gene expression in primary human neural cells. *J. Alzheimer's Dis.* 8, 117–127.
- Alfrey, A.C., leGendre, G.R., Kaehny, W.D., 1976. The dialysis encephalopathy syndrome. Possible aluminum intoxication. *N. Engl. J. Med.* 294, 184–188.
- Atwood, C.S., Scarpa, R.C., Huang, X., Moir, R.D., Jones, W.D., Fairlie, D.P., Tanzi, R.E., Bush, A.I., 2000. Characterization of copper interactions with Alzheimer amyloid beta peptides: identification of an attomolar-affinity copper binding site on amyloid beta1–42. *J. Neurochem.* 75, 1219–1233.
- Becaria, A., Campbell, A., Bondy, S.C., 2003. Aluminum and copper interact in the promotion of oxidative but not inflammatory events: implications for Alzheimer's disease. *J. Alzheimer's Dis.* 5, 31–38.
- Bondy, S.C., Kirstein, S., 1996. The promotion of iron-induced generation of reactive oxygen species in nerve tissue by aluminum. *Mol. Chem. Neuropathol.* 27, 185–194.
- Bondy, S.C., Truong, A., 1999. Potentiation of beta-folding of β -amyloid peptide 25–35 by aluminum salts. *Neurosci. Lett.* 267, 25–28.
- Bondy, S.C., Guo-Ross, S.X., Truong, A.T., 1998. Promotion of transition metal-induced reactive oxygen species formation by beta-amyloid. *Brain Res.* 799, 91–96.
- Brown, D.R., Kozlowski, H., 2004. Biological inorganic and bioinorganic chemistry of neurodegeneration based on prion and Alzheimer diseases. *Dalton Trans.* 13, 1907–1917.
- Buchanan, S.D., Diseker, R.A., Sinks, T., Olson, D.R., Daniel, J., Flodman, T., 1999. Copper in drinking water, Nebraska, 1994. *Int. J. Occup. Environ. Health* 5, 256–261.
- Campbell, A., 2004. Inflammation, neurodegenerative diseases, and environmental exposures. *Ann. N.Y. Acad. Sci.* 1035, 117–132.
- Campbell, A., Becaria, A., Sharman, K., Bondy, S.C., 2004. Chronic exposure to aluminum in drinking water increases inflammatory parameters selectively in the brain. *J. Neurosci. Res.* 75, 565–572.
- Di, J., Bi, S., 2003. Aluminum ions accelerated the oxidative stress of copper-mediated melanin formation. *Spectrochim. Acta, Part A: Mol. Biomol. Spectrosc.* 59, 3075–3083.
- Exley, C., Price, N.C., Kelly, S.M., Birchall, J.D., 1993. An interaction of beta-amyloid with aluminum in vitro. *FEBS Lett.* 324, 293–295.
- Flaten, T.P., 2001. Aluminium as a risk factor in Alzheimer's disease, with emphasis on drinking water. *Brain Res. Bull.* 55, 187–196.
- Gerard-Monnier, D., Erdelmeier, I., Regnard, K., Moze-Henry, N., Yadan, J.C., Chaudiere, J., 1998. Reactions of 1-methyl-2-phenylindole with malondialdehyde and 4-hydroxyalkenals. Analytical applications to a colorimetric assay of lipid peroxidation. *Chem. Res. Toxicol.* 11, 1176–1183.
- Jordan, M.B., Mills, D.M., Kappler, J., Marrack, P., Cambier, J.C., 2004. Promotion of B cell immune responses via an alum-induced myeloid cell population. *Science* 304, 1808–1810.
- Kawahara, M., Muramoto, K., Kobayashi, K., Mori, H., Kuroda, Y., 1994. Aluminum promotes the aggregation of Alzheimer's amyloid beta-protein in vitro. *Biochem. Biophys. Res. Commun.* 198, 531–535.

- Kim, K., 2003. Perinatal exposure to aluminum alters neuronal nitric oxide synthase expression in the frontal cortex of rat offspring. *Brain Res. Bull.* 61, 437–441.
- Kitzberger, R., Madl, C., Ferenci, P., 2005. Wilson disease. *Metab. Brain Dis.* 20, 295–302.
- Lahiri, D.K., Ge, Y., 2000. Electrophoretic mobility shift assay for the detection of specific DNA–protein complex in nuclear extracts from the cultured cells and frozen autopsy human brain tissue. *Brain Res.* 5, 257–265.
- Lahiri, D.K., Farlow, M.R., Sambamurti, K., 1998. The secretion of amyloid beta-peptides is inhibited in the tacrine-treated human neuroblastoma cells. *Brain Res. Mol. Brain Res.* 62, 131–140.
- Lahiri, D.K., Chen, D., Ge, Y.W., Bondy, S.C., Sharman, E.H., 2004. Dietary supplementation with melatonin reduces levels of amyloid beta-peptides in the murine cerebral cortex. *J. Pineal Res.* 36, 224–231.
- Ledeboer, A., Breve, J.J., Poole, S., Tilders, F.J., Van Dam, A.M., 2000. Interleukin-10, interleukin-4, and transforming growth factor-beta differentially regulate lipopolysaccharide-induced production of pro-inflammatory cytokines and nitric oxide in co-cultures of rat astroglial and microglial cells. *Glia* 30, 134–142.
- Liu, P.K., Robertson, C.S., Valadka, A., 2002. The association between neuronal nitric oxide synthase and neuronal sensitivity in the brain after brain injury. *Ann. N.Y. Acad. Sci.* 962, 226–241.
- Lukiw, W.J., Percy, M.E., Kruck, T.P., 2005. Nanomolar aluminum induces pro-inflammatory and pro-apoptotic gene expression in human brain cells in primary culture. *J. Inorg. Biochem.* 99, 1895–1898.
- McLachlan, D.R.C., Bergeron, C., Smith, J.E., Boomer, D., Rifat, S.L., 1996. Risk for neuropathologically confirmed Alzheimer's disease and residual aluminum in municipal drinking water employing weighted residential histories. *Neurology* 335, 1037.
- Mercer, J.F., 2001. The molecular basis of copper-transport diseases. *Trends Mol. Med.* 7, 64–69.
- Miu, A.C., Andreescu, C.E., Vasii, R., Olteanu, A.I., 2003. A behavioral and histological study of the effects of long-term exposure of adult rats to aluminum. *Int. J. Neurosci.* 113, 1197–1211.
- Park, K.W., Lee, D.Y., Joe, E.H., Kim, S.U., Jin, B.K., 2005. Neuroprotective role of microglia expressing interleukin-4. *J. Neurosci. Res.* 81, 397–402.
- Pratico, D., Uryu, K., Sung, S., Tang, S., Trojanowski, J.Q., Lee, V.M., 2002a. Aluminum modulates brain amyloidosis through oxidative stress in APP transgenic mice. *FASEB J.* 16, 1138–1140.
- Pratico, D., Clark, C.M., Liun, F., Rokach, J., Lee, V.Y., Trojanowski, J.Q., 2002b. Increase of brain oxidative stress in mild cognitive impairment: a possible predictor of Alzheimer disease. *Arch. Neurol.* 59, 972–976.
- Rondeau, V., Commenges, D., Jacqmin-Gadda, H., Dartigues, J.F., 2000. Relation between aluminum concentrations in drinking water and Alzheimer's disease: an 8-year follow-up study. *Am. J. Epidemiol.* 152, 59–66.
- Sparks, D.L., 2004. Cholesterol, copper, and accumulation of thioflavine S-reactive Alzheimer's-like amyloid beta in rabbit brain. *J. Mol. Neurosci.* 24, 97–104.
- Sparks, D.L., Schreurs, B.G., 2003. Trace amounts of copper in water induce beta-amyloid plaques and learning deficits in a rabbit model of Alzheimer's disease. *Proc. Natl. Acad. Sci. U. S. A.* 100, 11065–11069.
- Spitalny, K.C., Brondum, J., Vogt, R.L., Sargent, H.E., Kappel, S., 1984. Drinking-water-induced copper intoxication in a Vermont family. *Pediatrics* 74, 1103–1106.
- Tarkowski, E., Andreasen, N., Tarkowski, A., Blennow, K., 2003. Intrathecal inflammation precedes development of Alzheimer's disease. *J. Neurol. Neurosurg. Psychiatry* 74, 1200–1205.
- Torsdottir, G., Kristinsson, J., Sveinbjornsdottir, S., Snaedal, J., Johannesson, T., 1999. Copper, ceruloplasmin, superoxide dismutase and iron parameters in Parkinson's disease. *Pharm. Toxicol.* 85, 239–243.
- Verstraeten, S.V., Nogueira, L.V., Schreier, S., Oteiza, P.I., 1997. Effect of trivalent metal ions on phase separation and membrane lipid packing: role in lipid peroxidation. *Arch. Biochem. Biophys.* 338, 121–127.
- Yao, Y., Chinnici, C., Tang, H., Trojanowski, J.Q., Lee, V., Pratico, D., 2004. Brain inflammation and oxidative stress in a transgenic mouse model of Alzheimer-like brain amyloidosis. *J. Neuroinflamm.* 1, 21.