UC Irvine

UC Irvine Previously Published Works

Title

Highly regioselective monothionation of squaric and semisquaric acid derivatives

Permalink

<https://escholarship.org/uc/item/6g2296fw>

Journal

Synthesis, 1997(1)

ISSN

0039-7881

Authors

Müller, M
Heileman, MJ
Moore, HW
[et al.](#)

Publication Date

1997

DOI

10.1055/s-1997-1496

Copyright Information

This work is made available under the terms of a Creative Commons Attribution License, available at <https://creativecommons.org/licenses/by/4.0/>

Peer reviewed

Highly Regioselective Monothiation of Squaric and Semisquaric Acid Derivatives

Martin Müller,^a Matthew J. Heileman,^b Harold W. Moore,^{*b} Ernst Schaumann,^{*a} Gunadi Adiwidjaja^c

^a Institut für Organische Chemie, Technische Universität Clausthal, Leibnizstraße 6, D-38678 Clausthal-Zellerfeld, Germany

Fax +49(5323)722858; E-mail: ernst.schaumann@tu-clausthal.de

^b Department of Chemistry, University of California-Irvine, Irvine, CA 92717, USA

Fax +1(714)8242210; E-mail hal.moore@uci.edu

^c Mineralogisch-Petrographisches Institut, Universität Hamburg, Grindelallee 48, D-20146 Hamburg, Germany

Received 2 July 1996; revised 13 August 1996

An efficient synthesis of 4-thioxocyclobut-2-enones by the action of 0.5 molar equivalents of Lawesson's reagent on the corresponding diones is reported.

The chemistry of squaric acid has found considerable interest over the last thirty years.^{1,2} A noteworthy application is the conversion of dialkyl squarates into 3,4-disubstituted cyclobut-3-ene-1,2-diones which allows a plethora of further transformations via vinylketene intermediates.^{3,4} Consequently, analogous transformations from the corresponding vinylthioketenes can be envisaged from 4-thioxocyclobut-2-enones. In fact, a few monothiones with 3-dialkylamino or symmetrical 3,4-bis(dialkylamino) substitution have been reported,⁵⁻⁹ but for the former only indirect evidence on the site of thiation was obtained. Phosphorus pentasulfide appeared to be the sulfur transfer reagent of choice, though it may lead to bithiation. In contrast, ethoxycarbonyl isothiocyanate allows selective monothiation, but works only for 3,4-bis(dialkylamino)cyclobut-3-ene-1,2-diones ("squaric acid diamides").^{7,10} 2,4-Bis(4-methoxyphenyl)-1,3,2,4-dithiadiphosphetane 2,4-disulfide (**2**) ("Lawesson's reagent") which usually allows thiation reactions under mild conditions and in well-defined stoichiometry¹¹ had already proven to be advantageous in the thiation of simple enones¹² calling for its application also in squaric acid chemistry.

(a) 0.5 mol. equiv **2**, CH₂Cl₂, r.t., 12 h

Scheme

Dichloromethane solutions of 4-aryl- or 4-butylsemisquaric acid derivatives **1a-e** or squaric acid derivatives **1f,g** were treated with 0.5 molar equivalents of Lawesson's reagent **2** to yield 4-thioxocyclobut-2-enones **3** in good to moderate yields. The reaction is highly regioselective as neither the regioisomeric monothione nor dithionated products could be detected. Flash chromatography had to be used to separate thiones **3** from phosphorus containing impurities, since excessive base washing resulted in decomposition of **3**. An unambiguous

Table 1. Yields, Physical Properties and IR Spectroscopic Data of 4-Thioxocyclobut-2-enones **3** Prepared^a

3	R ¹	R ²	Yield (%)	mp (°C)	IR (KBr or neat) ν [C=O, C=C (cm ⁻¹)]
a	Oi-Pr	Ph	59	144–145	1770, 1605
b	Oi-Pr	Bu	76	oil	1785, 1605
c	Oi-Pr	4-MeOC ₆ H ₄	47	139–140	1765, 1603
d	Oi-Pr	4-Me ₂ NC ₆ H ₄	33	146 (dec)	1760, 1598
e	NEt ₂	Ph	76	95	1770, 1642
f	NBn ₂	OEt	65	80	1760, 1640
g	N(CH ₂) ₂ O	OEt	49	210	1770, 1660

^a Satisfactory microanalyses obtained for **3a, c, e-g**: C $\pm$ 0.27, H $\pm$ 0.11, N $\pm$ 0.08, S $\pm$ 0.12 except for S in **3f**: +0.46 (S not measured for **3a, c**); for **3b-e** exact mass with maximum deviation $\pm$ 0.0007 (**3d**).

proof of the site of thiation in **1** met with difficulties. Charge control of the reaction should favor attack of the reagent on the carbonyl group of the highly delocalized 3-(amino or alkoxy) enone moiety to give **3** with a vinylogous thioamide or thionoester unit, but also thiation of nonactivated thiones by **2** is a well-established reaction.¹³ Formation of **3** would be in line with previous claims on the regioselectivity of the reaction,^{5,6,8} but the failure to thionate dialkyl squarates with **2**¹⁰ shed doubt on the thiation of the 3-(isopropoxy) enone moiety in **1a-e**. Moreover, NMR data gave no clue as to the position of the sulfur; in particular, a 2D-COLOC-NMR spectrum of **3b** revealed coupling of the 1-methylene protons of the butyl substituent with both the carbonyl and the thiocarbonyl carbon. Therefore, an X-ray structural investigation of a single crystal of **3a** was carried out and finally confirmed the suggested structure (Figure). Pronounced electron delocalization within the S4-C4-C3-C2-O21 unit is shown by a relatively short C3-C4 distance as compared to the C1-C2 bond (Figure) and also in comparison to the enone moiety in a 3-(ethoxy)cyclobutenone [C=C 1.36 Å, C-C(=O) 1.452 Å].¹⁴ The four-membered ring is planar (maximum deviation 0.1 pm) and the benzene ring adopts an almost coplanar position [angle between planes 3.7(2)°].

Having established structure **3a**, the same regioselectivity is assumed for derivatives **3b-d** and, similarly, also for **3e-g** assuming the preferred thiation of the even more polarized vinylogous thioamide unit.

NMR spectra were recorded on a Bruker ARX-400 or G.E. Omega 500 spectrometer using CDCl₃ and TMS as internal standard. IR spectra were recorded on a Perkin-Elmer 1600 FT-IR spectrometer.

Figure. X-ray structure of **3a** in the ORTEP presentation. Selected bond distances [Å] and bond angles [°]: S4–C4 1.627(3), O1–C1 1.201(4), O21–C2 1.298(3), O21–C21 1.484(3), C1–C2 1.490(4), C1–C4 1.510(4), C2–C3 1.395(4), C3–C31 1.443(4), C3–C4 1.449(4); C2–C1–C4 86.0(2), C3–C2–C1 92.4(2), C2–C3–C4 92.1(2), C3–C4–C1 89.5(2)²²

MS were recorded on a VG Analytic 7070E instrument. Elemental analyses were performed by Pharmazeutisches Institut, TU Braunschweig. Merck silica gel 60 was used for chromatography. THF was distilled from diphenylketyl sodium just prior to use. Solvents for chromatography were distilled prior to use. Diones **1** could be obtained from dialkyl squarates using literature procedures: 3-isopropoxy-4-phenyl- (**1a**),¹⁵ 3-butyl-4-isopropoxy- (**1b**),¹⁶ 3-diethylamino-4-phenyl- (**1e**),² 3-(dibenzylamino)-4-ethoxy- (**1f**),¹⁷ and 3-ethoxy-4-morpholinocyclobut-3-ene-1,2-dione (**1g**).¹⁸ 3-Isopropoxy-4-(4-methoxyphenyl)cyclobut-3-ene-1,2-dione (**1c**) was prepared analogously to ref 15; yield 73%; mp 99–100°C (Lit. 101–102°C). ¹H NMR: δ = 1.54 (d, 6H, J = 6.2 Hz, *i*-Pr-CH₃), 3.88 (s, 3H, OCH₃), 5.59 (sept, 1H, J = 6.2 Hz, *i*-Pr-CH), 7.00, 8.03 (each d, 2H, J = 8.8 Hz, C₆H₄). 3-(4-Dimethylaminophenyl)-4-isopropoxycyclobut-3-ene-1,2-dione (**1d**) was obtained analogously to ref 7; yield 49%; yellow solid; mp 151–152°C. ¹H NMR: δ = 1.52 (d, 6H, J = 6.2 Hz, *i*-Pr-CH₃), 3.09 (s, 6H, NCH₃), 5.57 (sept, 1H, J = 6.2 Hz, *i*-Pr-CH), 6.71, 7.94 (each d, 2H, J = 8.9 Hz, C₆H₄). ¹³C NMR: δ = 23.0 (*i*-Pr-CH₃), 40.0 (NCH₃), 78.8 (*i*-Pr-CH), 111.4, 115.6, 129.9, 153.0 (C₆H₄), 174.1 (C-3), 190.3, 191.1, 193.8 (C-1, C-2, C-4).

Thionation of Cyclobutenediones **1**; General Procedure:

To a solution of **1** (5 mmol) in CH₂Cl₂ (3 mL) was added Lawesson's reagent **2** (1.011 g, 2.5 mmol). The reaction mixture was stirred overnight at r.t. EtOAc (50 mL) was added and the solution was washed with 10% aq NaOH (2 × 10 mL), sat aq NaHCO₃ (10 mL), and brine (2 × 10 mL). The organic layer was dried (MgSO₄), the solvents were removed in vacuo, and the residue was purified by flash chromatography (silica gel, CH₂Cl₂/hexanes 3:2). For yields and properties see Tables 1–3.

Crystal Structure Analysis of **3a**:

Rotating crystal and precession photographs of a crystal with dimensions 0.3 × 0.2 × 0.1 mm gave approximate lattice constants

Table 2. ¹H NMR Spectroscopic Data [(CDCl₃/TMS) δ , J (Hz)] of 4-Thioxocyclobut-2-enones **3** Prepared

3	δ_{2-XCH}	δ_{2-XCCH}	δ_{Ar-H}	Other Signals
a	5.59 (sept, J = 6.2)	1.60 (d, J = 6.2)	7.49–7.59, 8.42–8.46 (m)	–
b	5.37 (sept, J = 6.2)	1.49 (d, J = 6.2)	–	0.93 (t, J = 7.4), 1.38 (qt, J = 7.4), 1.70 (tt, J = 7.4), 2.72 (t, J = 7.4)
c	5.56 (sept, J = 6.2)	1.59 (d, J = 6.2)	7.01, 8.43 (each d, J = 9.2))	3.89 (s)
d	6.33 (sept, J = 6.2)	1.52 (d, J = 6.2)	6.70, 7.92 (each d, J = 8.8)	3.10 (s)
e	3.49, 3.90 (q, J = 7.3)	1.14, 1.33 (t, J = 7.3)	7.40–7.49 (m), 7.69 (d, J = 8.4)	–
f	4.48, 4.73	–	7.22–7.28, 7.37–7.46 (m)	1.49 (t, J = 7.2), 5.28 (q, J = 7.2)
g	3.67, 3.83, 3.92 (mc)	–	–	1.48 (t, J = 7.1), 5.20 (q, J = 7.1)

Table 3. ¹³C NMR Spectroscopic Data (CDCl₃/TMS, δ) of 4-Thioxocyclobut-2-enones **3** Prepared

3	$\delta_{C=O}$	$\delta_{C=S}$	δ_{C-2}	δ_{C-3}	δ_{2-XC}	δ_{2-XCC}	δ_{Ar-C}	Other Signals
a	196.8	227.6	185.2	176.2	81.4	22.8	128.0, 128.4, 128.7, 132.9	–
b	200.6	232.4	189.2	186.6	80.3	22.6 ^a	–	13.5, 22.7 ^a , 25.2, 27.2, 55.5
c	195.7	227.8	184.9	175.8	81.2	23.0	114.3, 121.6, 130.6, 163.4	55.5
d	175.2	202.9	173.7	163.9	77.6	30.1	107.5, 109.9, 123.7, 144.0	44.8
e	185.2	217.3	180.7	171.3	45.1, 45.4	12.8, 14.4	127.5, 128.5, 129.0, 130.2	–
f	184.3	206.9	179.9	174.7	51.7, 52.1	–	128.4, 128.8, 128.9, 129.0, 129.2, 129.3, 133.79, 133.82	16.3, 70.2
g	183.9	206.6	179.6	172.7	47.6, 47.8	–	–	16.3, 66.5, 66.8, 70.0

^a Signal assignments may be reversed.

and the preliminary space group. Refinement of the lattice constants led to the following cell dimensions: $a = 8.057(1)$, $b = 9.410(1)$, $c = 9.442(1)$ Å; $\alpha = 92.71(1)^\circ$, $\beta = 113.13(1)^\circ$, $\gamma = 111.76(1)^\circ$; $V = 595.61$ Å³; space group PI ; $Z = 2$; $d_{\text{calc}} = 1.295$ g cm⁻³. Intensity measurements were carried out with an Enraf-Nonius single-crystal diffractometer CAD 4-SDP using Cu- K_α radiation monochromated with graphite in the θ - 2θ scan technique at ambient temperature. The intensity data were not corrected for absorption. The 2431 measured intensities were reduced to 2258 symmetry-independent reflexions.¹⁹ The structure was solved by the direct-methods program SIR-92.²⁰ The E maps revealed the positions of all the heavy atoms. After the refinement of these positions the H atoms were found from a Fourier synthesis.²¹ In the final refinement, convergence was achieved at $R = 0.0757$.²²

This work was supported by funds from the Fonds der Chemischen Industrie, Frankfurt, and from the NATO Scientific and Environmental Affairs Division (Collaborative Research Grant No. 920016) as well as by a gift of squaric acid from Hüls AG.

- (1) Schmidt, A.H.; Ried, W. *Synthesis* **1978**, 1.
Knorr, H.; Ried, W. *Synthesis* **1978**, 649.
Schmidt, A.H.; Ried, W. *Synthesis* **1978**, 869.
Schmidt, A.H.; *Synthesis* **1980**, 961.
- (2) Frauenrath, H. In *Houben-Weyl*, 4th ed., Vol. E15/2; Kropf, H.; Schaumann, E.; Eds.; Thieme: Stuttgart, 1993; pp 1391, 1479.
- (3) Moore, H. W.; Yerxa, B. R. *Chemtracts* **1992**, 5, 273; *Adv. Strain Org. Chem.* **1995**, 4, 81.
- (4) Liebeskind, L. S. *Tetrahedron* **1989**, 45, 1989.
- (5) Seitz, G.; Morck, H.; Mann, K.; Schmiedel, R. *Chem. Ztg.* **1974**, 98, 459.
- (6) Seitz, G.; Arndt, G. *Synthesis* **1976**, 445.
- (7) Seitz, G.; Sutrisno, R. *Synthesis* **1978**, 831.
- (8) Seitz, G.; Morck, H.; Schmiedel, R.; Sutrisno, R. *Synthesis* **1979**, 361.
- (9) Seitz, G. *Thioxocarbon Dianions and their Derivatives*, In *Oxo-carbons*; West, R., Ed.; Academic: New York, 1980.
- (10) Müller, M., Ph. D. thesis, Clausthal 1994.
- (11) Cava, M.P.; Levinson, M.I. *Tetrahedron* **1985**, 41, 5061.
- (12) Walter, W.; Proll, T. *Synthesis* **1979**, 941
Dannhardt, G.; Geyer, Y.; Obergrusberger, P.; Ziereis, K. *Chem. Ztg.* **1987**, 111, 237.
Tubéry, F.; Grierson, D.S.; Husson, H.-P. *Tetrahedron Lett.* **1987**, 28, 6461.
- (13) Rauchfuss, T.B.; Zank, G.A. *Tetrahedron Lett.* **1986**, 27, 3445.
- (14) Trifonov, L.S.; Orahovats, A.S.; Prewo, R.; Bieri, J.H.; Heim-gartner, H. *J. Chem. Soc., Chem. Commun.* **1986**, 708.
- (15) Liebeskind, L.S.; Fengl, R.W.; Wirtz, K.R.; Shawe, T.T. *J. Org. Chem.* **1988**, 53, 2482. Liebeskind, L.S., personal communication.
- (16) Edwards, J.P.; Krysan, D.J.; Liebeskind, L.S. *J. Org. Chem.* **1993**, 58, 3942.
- (17) Thorpe, J.E. *J. Chem. Soc. B* **1968**, 435.
- (18) Roth, H.J.; Sporleder, H. *Arch. Pharm. (Weinheim, Ger.)* **1970**, 363, 886.
- (19) MOLEN, Enraf-Nonius, Delft 1995.
- (20) Giacovazzo, C. *Acta Crystallogr. Sect. A* **1983**, 39, 685.
- (21) Sheldrick G. SHELXL-93, Univ. Göttingen 1993.
- (22) Further details of the crystal structure determination are available on request from the Fachinformationszentrum Karlsruhe, Gesellschaft für wissenschaftlich-technische Information mbH, D-76344 Eggenstein-Leopoldshafen, on quoting the depository number CSD-405425, the authors, and the journal citation.