

UC Irvine

UC Irvine Previously Published Works

Title

The complete mitochondrial genome of the horned lizard *Phrynosoma blainvillii* (Squamata: Phrynosomatidae) from California, USA.

Permalink

<https://escholarship.org/uc/item/6q47s32t>

Journal

Mitochondrial DNA Part B, 2(2)

Authors

Ayala, Laura

Boo, Ga

Boo, Sung

et al.

Publication Date

2017-11-25

DOI

10.1080/23802359.2017.1407706

Peer reviewed

The complete mitochondrial genome of the horned lizard *Phrynosoma blainvillii* (Squamata: Phrynosomatidae) from California, USA

Laura Ayala^a, Ga Hun Boo^b, Sung Min Boo^c, Brandon Cluff^d, Gage H. Dayton^e, Leah Edwards^a, Armando Garcia^a, Mariana Gomez^a, Thomas A. Jimenez^a, Jeffery R. Hughey^a, Timothy Huynh^a, Natalia I. Marquez^a, Ana I. Meza^a, Alexis D. Munoz^f, Cristina Perez^a, Dominic Pina^a, Michael Porras^a, Paola E. Razo^a, Alexis Rosales^a, Bailee Rusconi^a, Noel Salvador^a, Alicia Steinhardt^a, Ruben Tinajero^a, Nancy Wheat^a, Frances L. Wong^a and Ann Wright^a; Hartnell College Genomics Group*

^aDivision of Mathematics, Science and Engineering, Hartnell College, Salinas, CA, USA; ^bUniversity Herbarium, University of California, Berkeley, CA, USA; ^cDepartment of Biology, Chungnam National University, Daejeon, Korea; ^dDepartment of Ecology and Evolutionary Biology, University of California, Santa Cruz, CA, USA; ^eUniversity of California Natural Reserves, Santa Cruz, CA, USA; ^fDepartment of Molecular, Cell and Developmental Biology, University of California, Santa Cruz, CA, USA

ABSTRACT

Analysis of *Phrynosoma blainvillii* Gray from Marina, Monterey County, California, using 150 bp paired-end Illumina sequences (Illumina, San Diego, CA) resulted in the assembly of its complete mitogenome. The mitogenome is 16,946 bp in length and contains a putative origin of light strand replication (OL), control region, 22 tRNA, 2 rRNA, and 13 protein-coding genes. Its content and organization are similar to other Squamata. Phylogenetic analysis of *P. blainvillii* resolves it in a clade with *P. sherbrookei* Nieto-Montes de Oca, Arenas-Moreno, Beltrán-Sánchez & Leaché, sister in position to *Uma notata* Baird. Mitochondrial marker analysis of *P. blainvillii* from Marina shows that it belongs to a coastal Santa Lucia Mountain Range haplogroup that is distinct from other populations of *P. blainvillii* in California.

ARTICLE HISTORY

Received 8 November 2017
Accepted 17 November 2017

KEYWORDS

Horned lizard; mitogenome; *Phrynosoma blainvillii*; Phrynosomatidae; special concern; Squamata

The family Phrynosomatidae consists of North American spiny lizards classified into nine genera and 148 species (Leaché et al. 2015a). Three complete mitogenomes have been published for the family: *Uta stansburiana* Baird & Girard (Leaché et al. 2015b), *Sceloporus occidentalis* Baird & Girard (Kumazawa 2004), and *Urosaurus nigricaudus* Cope (Bernardo et al. 2016). One of the 9 genera, *Phrynosoma* Wiegmann, includes 17 species of horned lizards (Nieto-Montes de Oca et al. 2014). Here, we describe the complete mitogenome of *Phrynosoma blainvillii*, a California Department of Fish and Wildlife 'Species of Special Concern', distributed from Northern California to Northern Baja California, Mexico (Leaché et al. 2009).

DNA was extracted from *P. blainvillii* (specimen voucher deposited in the Herpetological Collection – Museum of Vertebrate Zoology at UC Berkeley: MVZ:Herp:283805) collected from the UCSC Fort Ord Natural Reserve, Marina, California using the DNeasy Blood and Tissue Kit (Qiagen, Valencia, CA). The 150 bp paired-end library construction and sequencing was performed by myGenomics, LLC (Alpharetta, GA), yielding 4,978,820 reads. The mitogenome was assembled *de novo* using MEGAHIT (Li et al. 2015) and by mapping the reads against *U. stansburiana* (GenBank NC_027261) with the Medium-Low Sensitivity/Fast setting in Geneious R11 (Biomatters Limited, Auckland, New Zealand).

The genes were annotated using MITOS (Bernt et al. 2013) and adjusted in Sequin (<https://www.ncbi.nlm.nih.gov/Sequin/index.html>). Alignment of the mitogenome to other Squamata was performed with MAFFT (Katoh and Standley 2013). The maximum likelihood analysis was executed using complete mitogenome sequences with T-REX (Boc et al. 2012) and the GTR+gamma model with 1000 fast bootstraps. The tree was visualized with TreeDyn 198.3 at Phylogeny.fr (Dereeper et al. 2008).

The mitogenome of *P. blainvillii* (GenBank MG387969) is 16,946 bp in length and has a base composition of 33.85% A, 26.60% T, 13.07% G, and 26.48% C. It contains 22 tRNA (trnL and trnS are duplicated), two rRNA (rnl, rns), and 13 electron transport and oxidative phosphorylation genes. Eleven of the 13 genes initiate with the ATG start codon, however *cox1* and *nad1* initiate with GTG. Most of the genes terminate with TAA, but *nad2* terminates with TAG, *nad6* with AGG, and *cox1*, *cox2*, and *nad4* with AGA. The *nad6* gene and eight tRNAs encode on the light strand, while the others encode on the heavy strand. The putative OL is located between trnN and trnC, and is 30 bp in length, and the control region is 1540 bp. Phylogenetic analysis of *P. blainvillii* resolves it with *P. sherbrookei* in a fully supported clade sister to the genus *Uma* Baird (Figure 1). This relationship is similar to the findings of previous workers (Reeder and Wiens 1996;

CONTACT Jeffery R. Hughey  jhughey@hartnell.edu  Division of Mathematics, Science and Engineering, Hartnell College, Salinas, CA, USA

*All authors contributed equally to the analysis and writing of this paper.

© 2017 The Author(s). Published by Informa UK Limited, trading as Taylor & Francis Group.

This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0/>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.




Figure 1. Maximum-likelihood phylogram of *P. blainvillii* and representative Phrynosomatidae and Iguanidae mitogenomes. Numbers along branches are bootstrap supports based on 1000 nreps (<75% support not shown). The legend below represents the scale for nucleotide substitutions.

Wilgenbusch and de Queiroz 2000). Comparison of the complete *P. blainvillii* mitogenome to published *P. blainvillii* sequences (Leaché et al. 2009) found nearly identical sequences of *nad1* and *nad2* from a specimen from the Santa Lucia Mountain Range, but divergent sequences compared to a specimen of *P. blainvillii* from the nearby Gabilan Mountain Range. These data show that the Fort Ord, Marina and Santa Lucia populations of *P. blainvillii* represent a distinct coastal mitochondrial haplogroup.

Acknowledgements

Specimen collection was made under Gage H. Dayton's California Department of Fish and Wildlife Scientific Collecting Permit #633.

Disclosure statement

The authors report no conflicts of interest. The authors alone are responsible for the content and writing of this article.

Funding

This work was funded by a 2017–2018 Innovation Grant from Hartnell College to Jeffery R. Hughey.

References

Bernardo PH, Aguilera-Miller EF, Álvarez-Castañeda ST, Cruz FRMDL, Murphy RW. 2016. The complete mitochondrial genome of the black-tailed brush lizard *Urosaurus nigricaudus* (Reptilia, Squamata, Phrynosomatidae). *Mitochondrial DNA A DNA Mapp Seq Anal.* 27:4023–4025.

Bernt M, Donath A, Jühling F, Externbrink F, Florentz C, Fritzsche G, Pütz J, Middendorf M, Stadler PF. 2013. MITOS: improved *de novo* metazoan mitochondrial genome annotation. *Mol Phylogenet Evol.* 69:313–319.

Boc A, Diallo AB, Makarenkov V. 2012. T-REX: a web server for inferring, validating and visualizing phylogenetic trees and networks. *Nucleic Acids Res.* 40:W573–W579.

Dereeper A, Guignon V, Blanc G, Audic S, Buffet S, Chevenet F, Dufayard JF, Guindon S, Lefort V, Lescot M, et al. 2008. Phylogeny. fr: robust phylogenetic analysis for the non-specialist. *Nucleic Acids Res.* 36:W465–W469.

Katoh K, Standley DM. 2013. MAFFT multiple sequence alignment software version 7: improvements in performance and usability. *Mol Biol Evol.* 30:772–780.

Kumazawa Y. 2004. Mitochondrial DNA sequences of five squamates: phylogenetic affiliation of snakes. *DNA Res.* 11:137–144.

Leaché AD, Banbury BL, Felsenstein J, Nieto-Montes de Oca A, Stamatakis A. 2015a. Short tree, long tree, right tree, wrong tree: new acquisition bias corrections for inferring SNP phylogenies. *Syst Biol.* 64:1032–1047.

Leaché AD, Chavez AS, Jones LN, Grummer JA, Gottscho AD, Linkem CW. 2015b. Phylogenomics of phrynosomatid lizards: conflicting signals from sequence capture versus restriction site associated DNA sequencing. *Genome Biol Evol.* 7:706–719.

Leaché AD, Koo MS, Spencer CL, Papenfuss TJ, Fisher RN, McGuire JA. 2009. Quantifying ecological, morphological, and genetic variation to delimit species in the coast horned lizard species complex (*Phrynosoma*). *Proc Natl Acad Sci USA.* 106:12418–12423.

Li D, Liu CM, Luo R, Sadakane K, Lam TW. 2015. MEGAHIT: an ultra-fast single-node solution for large and complex metagenomics assembly via succinct de Bruijn graph. *Bioinformatics.* 31:1674–1676.

Nieto-Montes de Oca A, Arenas-Moreno D, Beltrán-Sánchez E, Leaché AD. 2014. A new species of horned lizard (genus *Phrynosoma*) from Guerrero, Mexico, with an updated multilocus phylogeny. *Herpetologica.* 70:241–257.

Reeder TW, Wiens JJ. 1996. Evolution of the lizard family phrynosomatidae as inferred from diverse types of data. *Herpetol Mon.* 10:43–84.

Wilgenbusch J, de Queiroz K. 2000. Phylogenetic relationships among the phrynosomatid sand lizards inferred from mitochondrial DNA sequences generated by heterogeneous evolutionary processes. *Syst Biol.* 49:592–612.