

UCLA

UCLA Previously Published Works

Title

Enhancing Primary Care Experiences for Homeless Patients with Serious Mental Illness: Results from a National Survey.

Permalink

<https://escholarship.org/uc/item/6gm5c1bx>

Authors

Gabrielian, Sonya

Jones, Audrey

Hoge, April

et al.

Publication Date

2021

DOI


10.1177/2150132721993654


Copyright Information

This work is made available under the terms of a Creative Commons Attribution-NonCommercial License, available at <https://creativecommons.org/licenses/by-nc/4.0/>

Peer reviewed

Enhancing Primary Care Experiences for Homeless Patients with Serious Mental Illness: Results from a National Survey

Journal of Primary Care & Community Health
Volume 12: 1–10
© The Author(s) 2021
Article reuse guidelines:
sagepub.com/journals-permissions
DOI: 10.1177/2150132721993654
journals.sagepub.com/home/jpc


Sonya Gabrielian^{1,2} , Audrey L. Jones^{3,4}, April E. Hoge⁵, Aerin J. deRussy⁵, Young-il Kim^{5,6}, Ann Elizabeth Montgomery^{5,7}, John R. Blosnich^{8,9}, Adam J. Gordon^{3,4}, Lillian Gelberg^{1,2}, Erika L. Austin^{5,7}, David Pollio¹⁰, Sally K. Holmes⁵, Allyson L. Varley⁵, and Stefan G. Kertesz^{5,6}

Abstract

Objectives: Patients experiencing homelessness (PEH) with serious mental illness (SMI) have poor satisfaction with primary care. We assessed if primary care teams tailored for homeless patients (Homeless-Patient Aligned Care Teams (H-PACTs)) provide this population with superior experiences than mainstream primary care and explored whether integrated behavioral health and social services were associated with favorable experiences. **Methods:** We surveyed VA PEH with SMI (n = 1095) to capture the valence of their primary care experiences in 4 domains (Access/Coordination, Patient-Clinician Relationships, Cooperation, and Homeless-Specific Needs). We surveyed clinicians (n = 52) from 29 H-PACTs to elucidate if their clinics had embedded mental health, addiction, social work, and/or housing services. We counted these services in each H-PACT (0-4) and classified H-PACTs as having high (3-4) versus low (0-2) service integration. We controlled for demographics, housing history, and needs in comparing H-PACT versus mainstream experiences; and experiences in high versus low integration H-PACTs. **Results:** Among respondents, 969 (91%) had complete data and 626 (62%) were in H-PACTs. After covariate adjustment, compared to mainstream respondents, H-PACT respondents were more likely ($P < .01$) to report favorable experiences (AORs = 1.7-2.1) and less likely to report unfavorable experiences (AORs = 0.5-0.6) in all 4 domains. Of 29 H-PACTs, 27.6% had high integration. High integration H-PACT respondents were twice as likely as low integration H-PACT respondents to report favorable access/coordination experiences (AOR = 1.7). **Conclusions:** Homeless-tailored clinics with highly-integrated services were associated with better care experiences among PEH with SMI. These observational data suggest that tailored primary care with integrated services may improve care perceptions among complex patients.

Keywords

primary care, homelessness, veterans, patient experience, serious mental illness

Dates received 30 October 2020; revised 10 January 2021; accepted 15 January 2021.

Introduction

Patients with serious mental illness (SMI, including psychotic disorders and bipolar disorders) are 10 to 20 times more likely to experience homelessness than the general population.¹ Unfortunately, patients experiencing homelessness (PEH) with SMI have high rates of chronic disease, increased morbidity/mortality, fragmented service use, poor primary care experiences, and social isolation.²⁻¹⁰ Regardless of housing status, patients with SMI often struggle to navigate medical care^{2,7}; they may report dissatisfaction with access to primary care and the coordination of services received.¹¹ In light of perceived discrimination reported by

PEH in primary care,^{12,13} and the central role of primary care in addressing the mortality gap for persons with SMI,¹⁴ there is a pressing need to identify clinic paradigms that optimize care experiences for PEH with SMI.

There are several reasons why primary care experiences are critical for these patients. First, these experiences are associated with medication adherence.¹⁵⁻¹⁷ Second, negative primary care perceptions often contribute to suboptimal service engagement,¹⁸ leading to increased Emergency Department utilization and hospitalizations. Third, positive primary care experiences are linked to improved outcomes for some chronic conditions.^{2,6,7} As such, healthcare



systems have increasingly embraced patient experience as a key indicator of quality of care.^{2,19,20}

Though some programs could optimize primary care experiences for PEH with SMI, doing so requires identifying and implementing relevant service design features. For example, the Health Care for the Homeless program²¹ has run for >3 decades; some of these programs deliver homeless-tailored services while others do not. To date, some surveys support the value of tailored primary care for PEH.^{6,22-24} However, research to understand which service design features influence patient experience is limited; this resource-intensive work requires collecting patient- and staff-level information from large numbers of clinics that vary in meaningful ways.

The Veterans Administration (VA) provides one opportunity to fill this gap. In 2012, the VA implemented Homeless-Patient Aligned Care Teams (H-PACTs), patient-centered medical homes tailored for PEH.²⁵ These teams operate at >60 VAs; dependent on the site, “tailored” care may include service delivery in non-traditional locations (eg, streets), practices that increase access (eg, walk-in services), tangible services (eg, clothing), and various levels of integration of social services and behavioral health care.²⁵ Ultimately, H-PACTs were developed to engage the VA’s most vulnerable PEH, including those with SMI.

This paper examines primary care experiences reported by PEH with SMI. First, using patient surveys, we assessed if H-PACTs provide this population with superior care experiences than mainstream primary care. Then, adding data from a survey of clinic staff, we examined whether one aspect of service design—having behavioral health and social services embedded within primary care—contributes to superior H-PACT experiences. Ultimately, this study aimed to inform primary care paradigms that enhance care experiences for PEH with SMI.

Methods

These analyses are part of the Primary Care Homeless Services Tailoring study,²⁶ a national survey of PEH who

receive primary care at 26 VAs with both H-PACTs and mainstream primary care (“mainstream”), with the goal of offering an observational comparison of care experiences in these 2 settings. Study procedures were approved by VA’s Central Institutional Review Board.

Setting

The VA is an integrated healthcare system that offers primary care, mental health services, and specialty medical and surgical care. VA primary care is delivered in patient-centered medical homes,^{27,28} with patients assigned to primary care providers embedded in teams with nurse care managers and ancillary staff. All primary care teams (PACTs) aim to deliver patient-centered and comprehensive primary care, including screening and preventive services.²⁹ Some PEHs receive primary care in mainstream PACTs while others receive care in H-PACTs, which are designed to address social determinants of health and facilitate housing.²⁴

Participants

Survey recruitment is detailed elsewhere.²⁶ VA patients were eligible if they: (a) received ≥ 2 primary care visits at a study site; (b) had evidence of homelessness between May 2015 and November 2017³⁰ (ICD-9/ICD-10 diagnoses of homelessness or VA-specific indicators of receipt of homeless services) in VA’s national electronic medical record (EMR) and c) were assigned to a single primary care team. Among eligible patients ($n = 57220$), sampling was stratified by facility and type of primary care (H-PACT vs mainstream); 14340 patients (derived from projected response rates and power calculations) were randomly selected at a 2:1 H-PACT to mainstream ratio. Participants were excluded if they had no available contact information or were deceased prior to the start of the survey. A professional survey organization, Strategic Research Group of Columbus, Ohio, conducted recruitment and survey collection in 4 waves of 4 to 6 weeks/each between March and October

¹VA Greater Los Angeles Health Care System, Los Angeles, CA, USA

²University of California Los Angeles, Los Angeles, CA, USA

³VA Salt Lake City Health Care System, Salt Lake City, UT, USA

⁴University of Utah School of Medicine, Salt Lake City, UT, USA

⁵Birmingham VA Medical Center, Birmingham, AL, USA

⁶University of Alabama at Birmingham School of Medicine, Birmingham, AL, USA

⁷University of Alabama at Birmingham School of Public Health, Birmingham, AL, USA

⁸VA Pittsburgh Healthcare System, Pittsburgh, PA, USA

⁹University of Southern California, Los Angeles, CA, USA

¹⁰University of Alabama at Birmingham College of Letters and Sciences, Birmingham, AL, USA

Corresponding Author:

Sonya Gabrielian, Greater Los Angeles Health Care System, Los Angeles, 11301 Wilshire Boulevard, Building 210A, Los Angeles, CA 90073, USA.

Email: sonya.gabrielian@va.gov

2018, cross-referencing patients' VA contact information with data from a commercial address verification database (MelissaData). Each wave included a pre-notification letter, a survey with \$1 pre-incentive, a reminder postcard, and a second survey for non-responders. The survey organization called non-responders up to 5 times during a 1-month period with the option to complete the survey by telephone. PEH who completed the survey ($n=5766$, 40.2%) received \$10. Next, using VA administrative data, we identified respondents who had at least 1 ICD-9/ICD-10 code for schizophrenia spectrum disorders, bipolar spectrum disorders, or other psychotic disorders (Supplemental Material Online)³¹ in VA's national EMR between May 2015 and November 2017 ($n=1095$, 19% of respondents).

We also conducted telephone-based surveys with the lead nurse and prescriber (primary care physician or physician extender) at participating H-PACTs; 52 of 58 (89.7%) lead nurses and prescribers from 29 H-PACTs were surveyed.

Conceptual Framework

This study was guided by the Behavioral Model for Vulnerable Populations,³² which models health care use among vulnerable populations and is widely used with PEH^{33,34} and persons with SMI.^{35,36} This framework identifies factors that *predispose* individuals to access services (demographics, homelessness chronicity), which interact with *enabling* factors (primary care clinic type, clinic characteristics) and *needs* for services to influence *behaviors* (service use) and *outcomes* (patient experience).³²

Measures

Covariates. To adjust for measurable differences among respondents' who received care in H-PACT versus mainstream primary care, the patient survey assessed factors that *predispose* patients to use services, including demographics (age, race, ethnicity, marital status, educational attainment), housing history (≥ 1 night in the past 6 months spent outside or in a place not meant for sleeping), and chronic homelessness (≥ 4 episodes of homelessness in the past 3 years or ≥ 1 episode of homelessness of ≥ 1 year).³⁷ Patients' *need* for primary care was captured with a self-reported count of 8 medical conditions administered in the survey (diabetes, hypertension, coronary artery disease, myocardial infarction, stroke, asthma, emphysema, arthritis, derived from the Medical Expenditure Panel Survey's satisfaction studies)³⁸ and SMI diagnoses were identified using VA EMR data. The presence of current alcohol or drug problems was assessed via survey with the Two-Item Conjoint Screening test.³⁹

Enabling characteristics. We also examined factors that enable or impede service use. Our key comparison was care

experiences in H-PACT versus mainstream clinics, determined from VA's national EMR.

The clinician survey assessed clinic characteristics that could contribute to differences in patient experience by primary care setting. These analyses focused on the presence or absence of mental health, addiction treatment, social work, and housing services within the H-PACTs. We asked clinicians to identify if each of the 4 types of services were provided: within H-PACT; outside H-PACT but within brief walking distance; or not provided within H-PACT or within brief walking distance. We classified responses as within or outside H-PACT, as services outside H-PACT but within walking distance seemed substantively different than services within the H-PACT. As surveys were conducted with 2 clinicians per H-PACT, we cross-checked responses by facility. If different responses were provided by the nurse and prescriber, we assigned the more conservative response (ie, outside H-PACT).

To capture the contributions of service integration on care experiences, we counted the number of services in each H-PACT (0-4), then classified H-PACTs as having high (3-4) or low (0-2) service integration based on the number of embedded services.

Outcome. Primary care experience was assessed in the patient survey, using the Primary Care Quality-Homeless (PCQ-H) questionnaire,⁴⁰ a 33-item instrument developed and validated for PEH. The PCQ-H is detailed elsewhere⁴⁰; it uses Likert scales (1-4) to capture primary care experience in 4 domains: access/coordination; patient-clinician relationship; perceived cooperation among clinicians; and homeless-specific needs. In each domain, the PCQ-H offers a categorical indicator for favorable experiences based on the top tertile of respondents and a categorical indicator for unfavorable experiences based on the lowest tertile of respondents.

Analyses

The sample consisted of PEH with SMI and complete data on study variables ($n=969$, 91%). Analyses were performed in Stata version 15.1. First, we compared respondents assigned to H-PACT (626, 64.6%) versus mainstream (343, 35.4%). We used chi-square and analysis of variance tests to assess between-group differences in predisposing, enabling, and need factors, then the outcome of favorable and unfavorable primary care experiences.

Next, we used multivariable logistic regressions to test for differences between H-PACT versus mainstream respondents in favorable and unfavorable experiences. These models controlled for study covariates and were weighted for non-response (the inverse probability of response, modeled from VA clinical records data).

To explore whether service integration was a factor in primary care experience, we reran the multiple logistic regression models comparing respondents from 3 groups: high integration H-PACTs (3-4 embedded services); low integration H-PACTs (0-2 embedded services); or mainstream. We used Wald tests to assess pairwise clinic differences (eg, high integration H-PACTs vs low integration H-PACTs) in favorable and unfavorable experiences.

Sensitivity analyses. We assessed if study patterns persisted with alternate definitions of service integration. Focused on the subsample of H-PACT respondents ($n=626$), we first used mixed effect logistic regressions to test for associations between the count of embedded services (0-4) with favorable and unfavorable experiences. The models included fixed effects for number of embedded services and respondent covariates, and a random effect for site. Second, we tested whether specific embedded services were associated with favorable and unfavorable experiences. The models included a fixed effect indicator for the embedded service and respondent covariates, and a random effect for site.

Results

Table 1 describes H-PACT versus mainstream respondents across *predisposing* and *need* factors. H-PACT respondents differed from their mainstream counterparts ($P < .05$) in several ways. Specifically, H-PACT respondents were younger (12.1%/25.1% were ≥ 65 years), less likely to be female (6.9%/15.5%), and less likely to be married (13.1%/19.8%). More H-PACT versus mainstream respondents had a history of chronic homelessness (27.6%/12.5%) and self-reported drug problems (21.7%/16.3%). About one-third of respondents (37.6%) had diagnoses of psychotic disorders; 56.0% had bipolar illness. There were no significant between-group differences in the self-reported number of chronic medical conditions; fewer ($P < .05$) H-PACT respondents self-reported diabetes than their mainstream counterparts (22.4%/28.6%).

Table 2 presents *the outcome* of primary care experiences by PCQ-H domain and clinic type. Compared to mainstream respondents, H-PACT respondents had higher rates of favorable primary care experiences and lower rates of unfavorable experiences in all 4 PCQ-H domains ($P < .05$). Differences in rates of favorable experiences were particularly pronounced in access/coordination (45.3%/28.4%), followed by homeless-specific needs (39.9%/25.1%), the patient-clinician relationship (45.2%/33.8%), and cooperation (38.0%/30.9%). Differences in rates of unfavorable experiences were largest in homeless-specific needs (42.7%/59.3%), followed by the patient-clinician relationships (28.9%/38.9%), access/coordination (27.7%/37.5%), then cooperation (27.9%/37.3%).

Table 3 displays these data adjusted for *predisposing* characteristics (age, gender, race, ethnicity, marital status, educational attainment, and housing history) and *need* (count of self-reported medical problems and presence of current alcohol/drug problems). Compared to the mainstream group, H-PACT respondents were more than twice as likely to report favorable experiences in access/coordination and homeless-specific needs (adjusted odds ratios (AOR)=2.2/2.1); and nearly twice as likely to report favorable patient-clinician relationships and cooperation (AOR=1.9/1.7). Similarly, compared to their mainstream peers, H-PACT respondents were about half as likely (AOR 0.5-0.6) to have unfavorable experiences in each PCQ-H domain.

Considering respondents from high integration H-PACTs, low integration H-PACTs, and mainstream, Table 4 presents pairwise differences in favorable and unfavorable experiences. High integration H-PACT respondents were 1.7 times as likely to have favorable access/coordination ($P < .05$) than low integration H-PACT respondents. No other statistically significant findings were revealed in comparing high versus low integration H-PACT respondents. In all 4 domains, high integration H-PACT respondents were significantly ($P < .05$) more likely than their mainstream peers to report favorable and/or less likely to report unfavorable experiences. For example, high integration H-PACT respondents were 3.5 times as likely to report favorable access/coordination and 2.1 times as likely to report favorable homeless-specific needs than mainstream respondents. Even low integration H-PACT respondents were about 2 times as likely (AOR=1.8-2.1) as mainstream respondents to have favorable experiences in all 4 domains ($P < .05$).

Regarding service integration in H-PACTs, surveys with nurses and prescribers revealed that most clinics had embedded social work ($n=22$, 75.9%) and mental health care ($n=17$, 58.6%). Fewer clinics had integrated housing ($n=8$, 27.6%) and addiction treatment ($n=3$, 10.3%) services. Just over a quarter (27.6%) of participating H-PACTs had high service integration (3-4 embedded services); most (72.4%) had low service integration (0-2 embedded services).

Table 5 depicts our sensitivity analyses. In analyses that considered the count of embedded services, the number of services integrated into H-PACTs was positively associated with favorable experiences in access/coordination (AOR=1.4). When specific embedded services were examined, only housing services were significantly associated ($P < .05$) with respondent experiences within H-PACTs. Specifically, respondents receiving services in H-PACTs with embedded housing services were more than 2 times (AOR=2.4) as likely as respondents in H-PACTs lacking embedded housing services to report favorable experiences with access/coordination. None of the other embedded services (mental health, social work, addiction) were statistically associated with favorable/unfavorable care experiences in any domain.

Table 1. Characteristics of Homeless-Experienced Respondents with Serious Mental Illness Who are Assigned to H-PACT vs VA Mainstream Clinics.

Predisposing and need variables	H-PACT (n = 626, 64.6%)		Mainstream (n = 343, 35.4%)		Total (N = 969, 100.0%)		P-value
	n (%)		n (%)		n (%)		
Predisposing							
Age*							<.001
18-54 years	195 (31.2)		96 (28.0)		291 (30.0)		
55-64 years	355 (56.7)		161 (46.9)		516 (53.2)		
≥65 years	76 (12.1)		86 (25.1)		162 (16.7)		
Female gender*	43 (6.9)		53 (15.5)		149 (9.9)		<.001
Race/ethnicity							.683
Non-Hispanic white	238 (38.0)		127 (37.0)		365 (37.7)		
Non-Hispanic black	209 (33.4)		113 (32.9)		322 (33.2)		
Hispanic, any race	80 (12.8)		39 (11.4)		119 (12.3)		
Other	99 (15.8)		64 (18.7)		163 (16.8)		
Marital status*							.015
Married or partnered	82 (13.1)		68 (19.8)		150 (15.5)		
Previously married	329 (52.6)		175 (51.0)		504 (52.0)		
Single, never married	215 (34.4)		100 (29.2)		315 (32.5)		
>12 years of education	381 (60.9)		220 (64.1)		601 (62.0)		.315
Housing history							
≥1 day unsheltered in past 6 months	103 (16.5)		47 (13.7)		150 (15.5)		.258
History of chronic homelessness*	173 (27.6)		43 (12.5)		216 (22.3)		<.001
Need							
SMI diagnoses†							
Schizophrenia spectrum disorders	238 (38.0)		126 (36.7)		364 (37.6)		.693
Bipolar spectrum disorders	346 (55.3)		197 (57.4)		543 (56.0)		.517
Other psychotic disorders*	214 (34.2)		94 (27.4)		308 (31.8)		.030
Substance use disorders							
Alcohol problem	196 (31.3)		102 (29.7)		298 (30.8)		.612
Drug problem*	136 (21.7)		56 (16.3)		192 (19.8)		.044
Medical diagnoses							
Diabetes*	140 (22.4)		97 (28.6)		237 (24.6)		.034
Hypertension	316 (50.6)		180 (52.9)		496 (51.4)		.480
Coronary artery disease	68 (10.9)		37 (10.8)		105 (10.9)		.963
Myocardial infarction	52 (8.3)		31 (9.1)		83 (8.6)		.688
Stroke	52 (8.3)		29 (8.6)		81 (8.4)		.900
Asthma	96 (15.4)		67 (19.6)		163 (16.9)		.093
Emphysema	58 (9.3)		37 (10.8)		95 (9.8)		.437
Arthritis	292 (46.9)		178 (52.4)		470 (48.8)		.104
	Mean	SD	Mean	SD	Mean	SD	
# of medical diagnoses from list above	1.7	1.5	1.9	1.6	1.8	1.5	.053

* $P < .05$, P -values obtained from chi-square tests of differences between patients in H-PACT and mainstream clinics.

†SMI diagnoses reflect diagnoses in the administrative data associated with visits in the 24 months preceding May 2015 to November 2017; participants may have ≥1 associated SMI diagnosis.

Discussion

In this national survey of PEH with SMI receiving VA primary care, we found that assignment to primary care clinics tailored for homeless patients was associated with more favorable experiences than mainstream primary care. For

PEH with SMI who received homeless-tailored primary care, the presence of specific behavioral health services was not associated with the valence of experiences. Rather, having more embedded services, that is, highly integrated clinics, was associated with favorable perceptions of clinic access/coordination.

Table 2. Primary Care Experiences Among Homeless-Experienced Respondents with Serious Mental Illness Assigned to H-PACT vs Mainstream Clinics.

Primary Care Quality-Homeless (PCQ-H) Questionnaire Domain	HPACT (n = 626, 64.6%)	Mainstream (n = 343, 35.4%)	Total (N = 969, 100.0%)	P-value
	n (%)	n (%)	n (%)	
Favorable experiences*				
Accessibility and coordination [†]	278 (45.3)	94 (28.4)	372 (39.4)	<.001
Patient-clinician relationship [†]	279 (45.2)	114 (33.8)	393 (41.2)	.001
Perceived cooperation among clinician [†]	211 (38.0)	96 (30.9)	307 (35.4)	.004
Homeless-specific needs [†]	236 (39.9)	77 (25.1)	313 (34.8)	<.001
Unfavorable experiences*				
Accessibility and coordination [†]	170 (27.7)	124 (37.5)	294 (31.1)	.002
Patient-clinician relationship [†]	178 (28.9)	131 (38.9)	309 (32.4)	.002
Perceived cooperation among clinicians [†]	155 (27.9)	116 (37.3)	271 (31.3)	.004
Homeless-specific needs [†]	253 (42.7)	182 (59.3)	435 (48.4)	<.001

*Favorable experiences include the top tertile of respondents; unfavorable experiences include the lowest tertile of respondents. The middle tertile is not displayed in this table.

[†]P < .05, P-value obtained from chi-square tests of differences between respondents in H-PACTs and mainstream clinics. The chi-square tests were run separately for favorable and unfavorable experiences in each domain, with analyses limited to respondents with less than 2 missing items in that domain: access/coordination (n = 945); relationship (n = 954); cooperation (n = 867); homeless-specific needs (n = 899).

Table 3. Logistic Regression of Primary Care Experiences among Homeless-Experienced Respondents with Serious Mental Illness in H-PACT vs Mainstream Clinics.

Primary care quality-homeless (PCQ-H) questionnaire domain	HPACT (Adjusted* %)	Mainstream (Adjusted* %)	AOR*	95% CI
Favorable experiences [†]				
Accessibility and coordination	46.2	28.0	2.2	1.6, 3.1
Patient-clinician relationship	46.8	31.7	1.9	1.4, 2.6
Perceived cooperation among clinicians	40.1	28.6	1.7	1.2, 2.4
Homeless-specific needs	40.2	24.5	2.1	1.5, 2.9
Unfavorable experiences [†]				
Accessibility and coordination	26.4	38.4	0.6	0.4, 0.8
Patient-clinician relationship	26.5	42.4	0.5	0.3, 0.6
Perceived cooperation among clinicians	25.6	38.8	0.5	0.4, 0.7
Homeless-specific needs	41.9	59.1	0.5	0.4, 0.7

*Logistic regression models, run separately for each domain, were weighted for non-response (calculated as inverse probability of response) and controlled for predisposing characteristics (age, gender, race, ethnicity, marital status, educational attainment, housing history) and need (count of 8 self-reported medical conditions and presence of current alcohol or drug problems). The adjusted percentages are model-derived predicted probabilities, holding covariates at their mean values. All AORs were statistically significant at P < .05.

[†]Favorable experiences include the top tertile of respondents; unfavorable experiences include the lowest tertile of respondents. The middle tertile is not displayed in this table.

While some prior studies have examined primary care tailoring for PEH, little attention has been given to primary care approaches for PEH with SMI. In this sample, PEH with SMI who received homeless-tailored primary care overwhelmingly endorsed better care experiences than their peers who received mainstream primary care. Tailoring features (eg, small panels, longer appointments) may be particularly important for vulnerable PEH, including those with SMI.

However, to serve this population with 2 vulnerabilities—SMI and homelessness—some may question the relative value of primary care tailored for PEH versus integration of primary care into mental health clinics.^{41,42} To this end,

though SMI diagnoses predispose patients to homeless experiences,^{43–46} only a minority of patients with SMI become homeless. However, PEH with SMI have incredibly high needs and health disparities.^{2–10} Little is known about similarities and differences in pathways to homelessness for patients with and without psychiatric illness. One study suggests that PEH with mental illness have similar pathways to homelessness as their peers without psychiatric problems.⁴⁴ That is, homeless patients as a unified cohort may have more in common, regardless of diagnoses, than homeless versus housed patients with SMI. This idea supports the use of homeless-tailored primary care for PEH with SMI.

Table 4. Logistic Regression of Primary Care Experiences among Homeless-Experienced Respondents with Serious Mental Illness in High Integration H-PACTs vs Low Integration H-PACTs vs Mainstream Clinics.

Primary Care Quality-Homeless (PCQ-H) Questionnaire Domain	High integration H-PACT vs low-integration H-PACT		High integration H-PACT vs mainstream		Low integration H-PACT vs mainstream	
	AOR [†]	95% CI	AOR [†]	95% CI	AOR [†]	95% CI
Favorable experiences [‡]						
Accessibility and coordination	1.7*	1.1, 2.6	3.5*	2.2, 5.6	2.0*	1.5, 2.8
Patient-clinician relationship	0.9	0.6, 1.3	1.7*	1.1, 2.8	2.0*	1.5, 2.7
Perceived cooperation among clinicians	0.8	0.5, 1.2	1.4	0.8, 2.3	1.8*	1.3, 2.5
Homeless-specific needs	1.0	0.7, 1.6	2.1*	1.3, 3.4	2.1*	1.5, 3.0
Unfavorable experiences [‡]						
Accessibility and coordination	0.8	0.5, 1.3	0.5*	0.3, 0.8	0.6*	0.4, 0.8
Patient-clinician relationship	1.1	0.7, 1.8	0.5*	0.3, 0.8	0.5*	0.3, 0.6
Perceived cooperation among clinicians	1.1	0.7, 1.8	0.6*	0.3, 0.9	0.5*	0.4, 0.7
Homeless-specific needs	1.1	0.7, 1.6	0.5*	0.3, 0.8	0.5*	0.3, 0.7

* $P < .05$.[†]Logistic regression models, run separately for each domain, were weighted for non-response (calculated as inverse probability of response) and controlled for predisposing characteristics (age, gender, race, ethnicity, marital status, educational attainment, housing history) and need (count of 8 self-reported medical conditions and presence of current alcohol or drug problems).[‡]Favorable experiences include the top tertile of respondents; unfavorable experiences include the lowest tertile of respondents. The middle tertile is not displayed in this table.**Table 5.** Sensitivity Analyses Testing Associations of Number of Services, then Specific Embedded Services, with Favorable and Unfavorable Primary Care Experiences among Homeless-Experienced Respondents with Serious Mental Illness in H-PACTs.

Service integrated into H-PACT	Accessibility and coordination		Patient-clinician relationship		Perceived cooperation among clinicians		Homeless-specific needs	
	AOR [†]	95% CI	AOR [†]	95% CI	AOR [†]	95% CI	AOR [†]	95% CI
Favorable experiences [§]								
Number of services [†]	1.4*	1.1, 1.7	1.1	0.8, 1.3	1.0	0.8, 1.2	1.1	0.9, 1.3
Specific services [‡]								
Mental health	1.6	0.9, 2.8	1.0	0.6, 1.7	1.0	0.6, 1.6	1.1	0.7, 1.7
Addiction treatment	2.0	0.8, 4.8	0.6	0.2, 1.3	0.7	0.3, 1.7	1.0	0.5, 2.1
Social work	1.8	1.0, 3.3	1.3	0.7, 2.4	1.1	0.6, 1.8	1.0	0.6, 1.7
Housing services	2.4*	1.4, 4.2	1.5	0.8, 2.7	1.1	0.6, 1.9	1.4	0.9, 2.3
Unfavorable experiences [§]								
Number of services [†]	0.8	0.6, 1.0	1.0	0.8, 1.2	1.0	0.8, 1.2	0.9	0.7, 1.1
Specific services [‡]								
Mental health	0.9	0.5, 1.6	1.1	0.7, 1.8	0.9	0.6, 1.5	0.9	0.5, 1.5
Addiction treatment	0.6	0.2, 1.7	0.9	0.4, 2.0	0.9	0.4, 2.0	1.1	0.5, 2.4
Social work	0.6	0.3, 1.2	0.8	0.5, 1.5	0.9	0.6, 1.6	0.7	0.4, 1.2
Housing services	0.5	0.3, 1.0	1.0	0.5, 1.7	1.0	0.6, 1.7	0.8	0.4, 1.4

* $P < .005$.[†]Estimates derived from mixed effect logistic regressions. Each model included fixed effects for number of embedded services (0-4) and patient predisposing characteristics (age, gender, race, ethnicity, marital status, educational attainment, housing history) and need (count of 8 self-reported medical conditions and presence of current alcohol or drug problems), and site random effects.[‡]Estimates derived from mixed effect logistic regressions. Each model included fixed effects indicator for the embedded service and patient predisposing characteristics (age, gender, race, ethnicity, marital status, educational attainment, housing history) and need (count of 8 self-reported medical conditions and presence of current alcohol or drug problems), and site random effects.[§]Favorable experiences include the top tertile of respondents; unfavorable experiences include the lowest tertile of respondents. The middle tertile is not displayed in this table.

We were surprised that the presence of integrated behavioral health care (eg, addiction services) was not associated with more favorable experiences. One

possibility is that the logistic demands of PEH dictate that the number of services available at a single site is more important than the types of services offered; this finding

is buttressed by our models. Also, despite common enthusiasm for colocation or integration of primary and mental health care,⁴⁷ patients with SMI likely have a diversity of preferences. Some patients with SMI value specialty mental health care in ways that requires its identification as distinct from primary care.⁴⁸

Our finding that high-integration H-PACTs were associated with more favorable access/coordination experiences may have service design implications, particularly as it relates to implementing case management in primary care.^{49,50} From its inception, H-PACT was described as a one-stop model of care, where multiple and sometimes competing needs could be addressed in a single setting.⁵¹ Our data suggest that PEH with SMI who receive primary care in “one-stop shops” with highly integrated services⁵²—extending beyond the basics of primary care-mental health integration⁵³ to encompass social services—may have better care experiences. A recent study showed that enhancing case management in VA primary care was related to decreased Emergency Department utilization⁵⁴; our findings echo the benefits of amplifying this role within primary care to manage care coordination for complex patients. At the same time, managers have to determine if the benefits of implementing these services exceed the costs incurred.

This study had limitations. First, the VA has robust medical, psychiatric, and social services for PEH; extrapolating these findings to other settings and populations requires caution. However, these findings could prove useful for the Health Care for the Homeless program²¹ or other settings that serve complex patients. Second, though our data derive from the largest survey of PEH to date, with a response rate (40.2%) that is about double what is reported for this population in VA’s standard patient experience methodology,⁵⁵ our survey respondents reflect a population that is engaged in care, with potentially fewer social vulnerabilities than other PEH with SMI. Third, these data reflect an observational study of PEH with SMI utilizing homeless-tailored versus mainstream primary care, as opposed to a randomized controlled trial that assigned PEH with SMI to homeless-tailored primary care services versus treatment as usual. The survey measures attempted to control a wide range of variables associated with patient-reported experience, but unmeasured confounders are not controlled. Fourth, in identifying relevant service design features, we focused on integration of behavioral health and social services; additional clinic features (eg, staffing, field-based services) are worthy of study. Last, though patient experience is associated with important outcomes for PEH with SMI, a central treatment goal for patients with SMI is to improve community functioning^{56–58}; the relationships between patient experience and functioning are unexplored.

Conclusions

Experiences of homelessness and SMI diagnoses convey synergistic risks for morbidity and mortality. To address health disparities faced by PEH with SMI, we must identify and scale viable primary care models that are well-received by this population. This study suggests that, at least within the VA, PEH with SMI who receive care in homeless-tailored primary care clinics had more favorable care experiences than their peers who received primary care in mainstream settings. More favorable patient experiences do favor continued care engagement, but whether optimal health outcomes result from such engagement remains to be seen. Further research could explore other features of and adaptations to H-PACT that optimize care for PEH with SMI, moving beyond patient experience to explore additional important outcomes, including medication adherence, substance use disorder outcomes, and functioning.

Declaration of Conflicting Interests

The author(s) declared no potential conflicts of interest with respect to the research, authorship, and/or publication of this article.

Funding

The author(s) disclosed receipt of the following financial support for the research, authorship, and/or publication of this article: This work was supported by VA HSR&D IIR 15-095-2 (PI: Kertesz). Dr. Gabrielian is supported in part by VA HSR&D Career Development Award 15-074. Dr. Jones is supported by the National Center for Advancing Translational Sciences of the National Institutes of Health under (award numbers UL1TR002538 and KL2TR002539). The views expressed in this article are those of the authors alone and do not represent the views of the United States Department of Veterans Affairs, the National Institutes of Health, or the United States Government.

ORCID iD

Sonya Gabrielian  <https://orcid.org/0000-0002-1790-4271>

Supplemental Material

Supplemental material for this article is available online.

References

1. Kuno E, Rothbard AB, Averyt J, Culhane D. Homelessness among persons with serious mental illness in an enhanced community-based mental health system. *Psychiatr Serv*. 2000;51:1012-1016.
2. Desai RA, Stefanovics EA, Rosenheck RA. The role of psychiatric diagnosis in satisfaction with primary care: data from the department of veterans affairs. *Med Care*. 2005;43:1208-1216.
3. Lester H, Tritter JQ, England E. Satisfaction with primary care: the perspectives of people with schizophrenia. *Fam Pract*. 2003;20:508-513.

4. Abt Associates and the National Center on Homelessness Among Veterans. *Veteran Homelessness: A Supplemental Report to the 2010 Annual Homeless Assessment Report to Congress*; 2011. Accessed on December 4, 2020. <http://www.va.gov/HOMELESS/docs/2010AHARVeteransReport.pdf>
5. Gallagher TC, Andersen RM, Koegel P, Gelberg L. Determinants of regular source of care among homeless adults in Los Angeles. *Med Care*. 1997;35:814-830.
6. Chrystal JG, Glover DL, Young AS, et al. Experience of primary care among homeless individuals with mental health conditions. *PLoS One*. 2015;10:e0117395.
7. Kilbourne AM, McCarthy JF, Post EP, et al. Access to and Satisfaction with care comparing patients with and without serious mental illness. *Int J Psychiatry Medicine*. 2006;36:383-399.
8. Nielssen O, Jones N, Fong H, Nielssen A, Staples L, Large M. Comparison of homeless clinic attenders with and without psychotic illness. *Aust N Z J Psychiatry*. 2020;54:195-201.
9. Karadzhev D, Yuan Y, Bond L. Coping amidst an assemblage of disadvantage: a qualitative metasynthesis of first-person accounts of managing severe mental illness while homeless. *J Psychiatr Ment Health Nurs*. 2020;27:4-24.
10. Gilmoor A, Vallath S, Regeer B, Bunders J. "If somebody could just understand what I am going through, it would make all the difference": conceptualizations of trauma in homeless populations experiencing severe mental illness. *Transcult Psychiatry*. 2020;57:455-467.
11. Kaufman EA, McDonnell MG, Cristofalo MA, Ries RK. Exploring barriers to primary care for patients with severe mental illness: frontline patient and provider accounts. *Issues Ment Health Nurs*. 2012;33:172-180.
12. Skosireva A, O'Campo P, Zerger S, Chambers C, Gapka S, Stergiopoulos V. Different faces of discrimination: perceived discrimination among homeless adults with mental illness in healthcare settings. *BMC Health Serv Res*. 2014;14:376.
13. Ramsay N, Hossain R, Moore M, Milo M, Brown A. Health care while homeless: barriers, facilitators, and the lived experiences of homeless individuals accessing health care in a Canadian regional municipality. *Qual Health Res*. 2019;29:1839-1849.
14. Perrin J, Reimann B, Capobianco J, Wahrenberger JT, Sheitman BB, Steiner BD. A model of enhanced primary care for patients with severe mental illness. *N C Med J*. 2018;79:240-244.
15. Fortuna RJ, Nagel AK, Rocco TA, Legette-Sobers S, Quigley DD. Patient experience with care and its association with adherence to hypertension medications. *Am J Hypertens*. 2018;31:340-345.
16. Clever SL, Ford DE, Rubenstein LV, et al. Primary care patients' involvement in decision-making is associated with improvement in depression. *Med Care*. 2006;44:398-405.
17. Bauer AM, Parker MM, Schillinger D, et al. Associations between antidepressant adherence and shared decision-making, patient-provider trust, and communication among adults with diabetes: diabetes study of Northern California (DISTANCE). *J Gen Intern Med*. 2014;29:1139-1147.
18. Bauer LK, Baggett TP, Stern TA, O'Connell J, Shtasel D. Caring for homeless persons with serious mental illness in general hospitals. *Psychosomatics*. 2013;54:14-21.
19. Delaney KR, Loucks J, Ray R, et al. Delineating quality indicators of inpatient psychiatric hospitalization. *J Am Psychiatr Nurses Assoc*. 2020;4:107839032097136-11.
20. Jun J, Stern K, Djukic M. Integrative review of the interventions for improving patients' experiences revealed in quality improvement projects. *J Patient Exp*. 2020;7:882-892.
21. Zlotnick C, Zerger S, Wolfe PB. Health care for the homeless: what we have learned in the past 30 years and what's next. *Am J Public Health*. 2013;103(suppl 2):S199-S205.
22. Loeb DF, Binswanger IA, Candrian C, Bayliss EA. Primary care physician insights into a typology of the complex patient in primary care. *Ann Fam Med*. 2015;13:451-455.
23. Kertesz SG, Holt CL, Steward JL, et al. Comparing homeless persons' care experiences in tailored versus nontailored primary care programs. *Am J Public Health*. 2013;103(suppl 2):S331-S339.
24. Jones AL, Hausmann LRM, Kertesz SG, et al. Providing positive primary care experiences for homeless veterans through tailored medical homes: the veterans health administration's homeless patient aligned care teams. *Med Care*. 2019;57:270-278.
25. O'Toole TP, Johnson EE, Aiello R, Kane V, Pape L. Tailoring care to vulnerable populations by incorporating social determinants of health: the veterans health administration's "homeless patient aligned care team" program. *Prev Chronic Dis*. 2016;13:1-12.
26. Riggs KR, Hoge AE, DeRussy AJ, et al. Prevalence of and risk factors associated with nonfatal overdose among veterans who have experienced homelessness. *JAMA Netw Open*. 2020;3:e201190.
27. Rosland A-M, Nelson K, Sun H, et al. The patient-centered medical home in the Veterans Health Administration. *Am J Manag Care*. 2013;19:e263-e272.
28. Nelson KM, Helfrich C, Sun H, et al. Implementation of the patient-centered medical home in the Veterans Health Administration: associations with patient satisfaction, quality of care, staff burnout, and hospital and emergency department use. *JAMA Intern Med*. 2014;174:1350-1358.
29. Chang ET, Zulman DM, Nelson KM, et al. Use of general primary care, specialized primary care, and other veterans affairs services among high-risk veterans. *JAMA Netw Open*. 2020;3:e208120.
30. Peterson R, Gundlapalli AV, Metraux S, et al. Identifying homelessness among veterans using VA Administrative data: opportunities to expand detection criteria. *PLoS One*. 2015;10:e0132664.
31. *National Psychosis Registry, Facility Drilldown Dashboard*. Serious Mental Illness Treatment Resource and Evaluation Center. Accessed January 1, 2021. <http://vaww.smitrec.va.gov/> (internal VA website).
32. Gelberg L, Andersen RM, Leake BD. The behavioral model for vulnerable populations: application to medical care use and outcomes for homeless people. *Health Serv Res*. 2000;34:1273-1302.
33. Gentil L, Grenier G, Bamvita JM, Fleury M-J. Satisfaction with health and community services among homeless and formerly homeless individuals in Quebec, Canada. *Health Soc Care Community*. 2020;28:22-33.
34. Petrovich JC, Hunt JJ, North CS, Pollio DE, Murphy ER. Comparing unsheltered and sheltered homeless: demographics,

- health services use and predictors of health services use. *Community Ment Health J*. 2019;56:271-279.
35. Rogers ES, Friedes R, Jakes A, Grossman E, Link A, Sherman SE. Long-term abstinence and predictors of tobacco treatment uptake among hospitalized smokers with serious mental illness enrolled in a smoking cessation trial. *J Behav Med*. 2017;40:750-759.
 36. Kelly E, Duan L, Cohen H, Kiger H, Pancake L, Brekke J. Integrating behavioral healthcare for individuals with serious mental illness: a randomized controlled trial of a peer health navigator intervention. *Schizophr Res Treatment*. 2017;182:135-141.
 37. Development of Housing and Urban Development. *Homeless Emergency Assistance and Rapid Transition to Housing: Defining "Chronically Homeless"*; 2015.
 38. Fenton JJ, Jerant AF, Bertakis KD, Franks P. The cost of satisfaction: a national study of patient satisfaction, health care utilization, expenditures, and mortality. *Arch Intern Med*. 2012;172:405-411.
 39. Brown RL, Leonard T, Saunders LA, Papasouliotis O. A two-item screening test for alcohol and other drug problems. *J Fam Pract*. 1997;44:151-160.
 40. Kertesz SG, Pollio DE, Jones RN, et al. Development of the primary care quality-homeless (PCQ-H) instrument. *Medi Care*. 2014;52:734-742.
 41. Krupski A, West II, Scharf DM, et al. Integrating primary care into community mental health centers: impact on utilization and costs of health care. *Psychiatr Serv*. 2016;67:1233-1239.
 42. Breslau J, Leckman-Westin E, Yu H, et al. Impact of a mental health based primary care program on quality of physical health care. *Adm Policy Ment Health*. 2017;45:276-285.
 43. Rosenheck R, Fontana A. A model of homelessness among male veterans of the Vietnam War generation. *Am J Psychiatry*. 1994;151:421-427.
 44. Sullivan G, Burnam A, Koegel P. Pathways to homelessness among the mentally ill. *Soc Psychiatry Psychiatr Epidemiol*. 2000;35:444-450.
 45. Tessler R, Rosenheck R, Gamache G. Comparison of homeless veterans with other homeless men in a large clinical outreach program. *Psychiat Q*. 2002;73:109-119.
 46. Hamilton AB, Poza I, Washington DL. "Homelessness and trauma go hand-in-hand": pathways to homelessness among women veterans. *Womens Health Issues*. 2011;21(suppl 4):S203-S209. doi:10.1016/j.whi.2011.04.005
 47. Miller-Matero LR, Dykuis KE, Albujoq K, et al. Benefits of integrated behavioral health services: the physician perspective. *Fam Syst Health*. 2016;34:51-55.
 48. Mandiberg JM. Commentary: The failure of social inclusion: an alternative approach through community development. *Psychiatric services*. 2012;63(5):458-460. doi:10.1176/appi.ps.201100367
 49. Teper MH, Vedel I, Yang XQ, Margo-Dermer E, Hudon C. Understanding barriers to and facilitators of case management in primary care: a systematic review and thematic synthesis. *Ann Fam Med*. 2020;18:355-363.
 50. Hudon C, Chouinard M-C, Aubrey-Bassler K, et al. Case management in primary care for frequent users of health care services: a realist synthesis. *Ann Fam Med*. 2020;18:218-226.
 51. O'Toole TP, Johnson EE, Borgia M, et al. Population-tailored care for homeless veterans and acute care use, cost, and satisfaction: a prospective quasi-experimental trial. *Prev Chronic Dis*. 2018;15:E23.
 52. Bhalla IP, Deegan D, Stefanovics EA, Rosenheck RA. Psychiatric multimorbidity in a specialized program for severely mentally ill veterans. *Psychiatr Q*. 2020;307:2493-2411.
 53. Stoeckle J, Cunningham A, Arenson C. Scaling integrated behavioral health rapidly. *Ann Fam Med*. 2018;16:464-464.
 54. Cornell PY, Halladay CW, Ader J, et al. Embedding social workers in veterans health administration primary care teams reduces emergency department visits. *Health Aff*. 2020;39:603-612.
 55. Jones AL, Hausmann LRM, Kertesz S, et al. Differences in experiences with care between homeless and nonhomeless patients in veterans affairs facilities with tailored and nontailored primary care teams. *Medi Care*. 2018;56:610-618.
 56. Brown MA, Velligan DI. Issues and developments related to assessing function in serious mental illness. *Dialogues Clin Neurosci*. 2016;18:135-144.
 57. Mausbach BT, Moore R, Bowie C, Cardenas V, Patterson TL. A review of instruments for measuring functional recovery in those diagnosed with psychosis. *Schizophr Bull*. 2009;35:307-318.
 58. Granholm E, Holden JL, Mikhael T, et al. What do people with schizophrenia do all day? ecological momentary assessment of real-world functioning in schizophrenia. *Schizophr Bull*. 2019;44:1195-1110.